

**CORRIGENDUM TO “NONLINEAR DIFFERENTIAL  
POLYNOMIALS SHARING A SMALL FUNCTION” [ARCH.  
MATH. 44 (2008), 41–56]**

ABHIJIT BANERJEE AND SONALI MUKHERJEE

This shorte note concerns the paper mentioned in the title.

The statement of **Theorem 1.1(ii)** and so that of **Corollary 1.2** given in that paper is not correct. For the uniqueness of  $f$  and  $g$  some extra conditions are required. Since the proof of that particular portion of **Theorem 1.1(ii)** depends upon **Lemma 2.11(ii)**, so the statement as well as the proof of **Lemma 2.11(ii)** should also be rectified. In the statement of **Theorem 1.1(ii)** and so in **Corollary 1.2** the extra condition “and the two expressions  $\frac{b}{n+2} g \sum_{j=0}^{n+1} \left(\frac{f}{g}\right)^j + \frac{c}{n+1} \sum_{j=0}^n \left(\frac{f}{g}\right)^j$  and  $\sum_{j=0}^{n+2} \left(\frac{f}{g}\right)^j$  have no common simple zeros” should be added. Naturally in the proof of the **Lemma 2.11(ii)** some more analysis regarding the zeros of  $\eta - u_k$  which are not the poles of  $g$  is required since the salient part of the proof is depending on the proof that  $\eta - u_k$  has multiple zeros. The corrected statements and proofs of **Theorem 1.1(ii)** and the corresponding **Lemma 2.11(ii)** are given below.

**Theorem 1.1.** *Let  $f$  and  $g$  be two transcendental meromorphic functions such that  $f^n(af^2 + bf + c)f'$  and  $g^n(ag^2 + bg + c)g'$  where  $a \neq 0$  and  $|b| + |c| \neq 0$  share “ $(\alpha, 2)$ ”. Then the following holds.*

- (ii) *If  $b \neq 0$ ,  $c \neq 0$ ,  $n > [12 - 2\Theta(\infty; f) - 2\Theta(\infty; g) - \min\{\Theta(\infty; f), \Theta(\infty; g)\}]$ , the roots of the equation  $az^2 + bz + c = 0$  are distinct, one of  $f$  and  $g$  is non entire meromorphic functions having only multiple poles and the two expressions  $\frac{b}{n+2} g \sum_{j=0}^{n+1} \left(\frac{f}{g}\right)^j + \frac{c}{n+1} \sum_{j=0}^n \left(\frac{f}{g}\right)^j$  and  $\sum_{j=0}^{n+2} \left(\frac{f}{g}\right)^j$  have no common simple zeros then  $f \equiv g$ .*

**Corollary 1.2.** *Let  $f$  and  $g$  be two transcendental meromorphic functions, one of  $f$  and  $g$  is non entire meromorphic functions having only multiple poles and the two expressions  $\frac{b}{n+2} g \sum_{j=0}^{n+1} \left(\frac{f}{g}\right)^j + \frac{c}{n+1} \sum_{j=0}^n \left(\frac{f}{g}\right)^j$  and  $\sum_{j=0}^{n+2} \left(\frac{f}{g}\right)^j$  have no common simple*

---

2000 *Mathematics Subject Classification*: primary 30D35.

*Key words and phrases*: meromorphic function, uniqueness, nonlinear differential polynomials, small function, weakly weighted sharing.

zeros, such that  $n > [12 - 2\Theta(\infty; f) - 2\Theta(\infty; g) - \min\{\Theta(\infty; f), \Theta(\infty; g)\}]$  be an integer. If  $af^n(f - \beta_1)(f - \beta_2)f'$  and  $ag^n(g - \beta_1)(g - \beta_2)g'$  share “ $(\alpha, 2)$ ”, where  $\beta_1$  and  $\beta_2$  are the distinct roots of the equation  $az^2 + bz + c = 0$  with  $|\beta_1| \neq |\beta_2|$ , then  $f \equiv g$ .

**Lemma 2.11.** *Let  $F$  and  $G$  be given as in Lemma 2.9 and  $n(\geq 6)$  be an integer. Suppose  $F \equiv G$ . Then the following holds.*

- (ii) *If  $b \neq 0, c \neq 0$ , and the roots of the equation  $az^2 + bz + c = 0$  are distinct and one of  $f$  and  $g$  is non entire meromorphic function having only multiple poles and the two expressions  $\frac{b}{n+2} g \sum_{j=0}^{n+1} \left(\frac{f}{g}\right)^j + \frac{c}{n+1} \sum_{j=0}^n \left(\frac{f}{g}\right)^j$  and  $\sum_{j=0}^{n+2} \left(\frac{f}{g}\right)^j$  have no common simple zero then  $f \equiv g$ .*

**Proof. Case 2.** Suppose  $b \neq 0$  and  $c \neq 0$ . Then  $F \equiv G$  implies

$$(1) \quad Af^{n+3} + Bf^{n+2} + Cf^{n+1} \equiv Ag^{n+3} + Bg^{n+2} + Cg^{n+1},$$

where  $A = \frac{a}{n+3}, B = \frac{b}{n+2}$  and  $C = \frac{c}{n+1}$ .

Let us assume  $f \not\equiv g$ .

**Subcase 2.1.** Suppose the roots of the equation  $az^2 + bz + c = 0$  are distinct. Since (1) implies  $f, g$  share  $(\infty, \infty)$  without loss of generality we may assume that  $g$  has some multiple poles. Putting  $\eta = \frac{f}{g}$  in (1) we get

$$Ag^2(\eta^{n+3} - 1) + Bg(\eta^{n+2} - 1) + C(\eta^{n+1} - 1) \equiv 0.$$

i.e.,

$$(2) \quad Ag^2 \equiv -Bg \frac{\eta^{n+2} - 1}{\eta^{n+3} - 1} - C \frac{\eta^{n+1} - 1}{\eta^{n+3} - 1}.$$

First we observe that since a meromorphic function can not have more than two Picard exceptional values,  $\eta$  takes at least  $n$  values among  $u_k = \exp\left(\frac{2k\pi i}{n+3}\right)$  where  $k = 1, 2, \dots, n + 2$ .

Let  $z_0$  be a pole of  $g$  with multiplicity  $p(\geq 2)$ , which is not a root of  $\eta - u_k = 0$ . Then from (2) we have

$$2p = p \quad \text{i.e.,} \quad p = 0,$$

which is impossible.

Hence from (2) we see that the poles of  $g$  are precisely the roots of  $\eta - u_k = 0$ .

Suppose  $z_1$  is a zero of  $\eta - u_k$  of multiplicity  $r$  which is a pole of  $g$  with multiplicity  $s$  (say) then from (2) we see that

$$2s = r + s$$

i.e.,

$$r = s.$$

Since  $g$  has no simple pole, it follows that such points are multiple zeros of  $\eta - u_k$ .

From (2) we know

$$(3) \quad Ag^2 \equiv - \frac{Bg \sum_{j=0}^{n+1} \eta^j + C \sum_{j=0}^n \eta^j}{\sum_{j=0}^{n+2} \eta^j}.$$

Suppose  $z_2$  be a simple zero of  $\eta - u_k$  where  $k = 1, 2, \dots, n + 2$ , which is a zero of multiplicity  $q(\geq 2)$  of the numerator of (3). Then from (3),  $z_2$  would be a zero of order  $q - 1$  of  $g^2$ . So it follows that  $z_2$  would be a zero of  $\sum_{j=0}^n \eta^j$ . Since  $\sum_{j=0}^n \eta^j$  and

$\sum_{j=0}^{n+2} \eta^j$  may have at most one common factor, we see that  $\eta - u_k$  has multiple zeros for at least  $n - 1$  values of  $k \in \{1, 2, \dots, n + 2\}$ . Hence

$$\Theta(u_k; \eta) \geq \frac{1}{2},$$

for at least  $n - 1$  values of  $k$ , which implies a contradiction as  $n \geq 6$ .  $\square$

DEPARTMENT OF MATHEMATICS, KALYANI GOVERNMENT ENGINEERING COLLEGE  
WEST BENGAL 741235, INDIA

*E-mail:* abanerjee\_kal@yahoo.co.in, abanerjee\_kal@rediffmail.com