

ON LIFTS OF SOME PROJECTABLE VECTOR FIELDS  
ASSOCIATED TO A PRODUCT PRESERVING GAUGE  
BUNDLE FUNCTOR ON VECTOR BUNDLES

A. NTYAM, G. F. WANKAP NONO, AND BITJONG NDOMBOL

ABSTRACT. For a product preserving gauge bundle functor on vector bundles, we present some lifts of smooth functions that are constant or linear on fibers, and some lifts of projectable vector fields that are vector bundle morphisms.

1. INTRODUCTION

Weil functors (product preserving bundle functors on manifolds) were used in [2] to define some lifts of geometric objects namely, smooth functions, tensor fields and linear connections on manifolds.

Product preserving gauge bundle functor on vector bundles have been classified in [7]: The set of equivalence classes of such functors are in bijection with the set of equivalence classes of pairs  $(A, V)$ , where  $A$  is a Weil algebra and  $V$  a  $A$ -module such that  $\dim_{\mathbb{R}}(V) < \infty$ .

In this paper, we adopt the approach of [2] to present some lifts of smooth functions that are constant or linear on fibers, and some lifts projectable vector fields that are vector bundle morphisms.

2. ALGEBRAIC DESCRIPTION OF WEIL FUNCTORS

Weil functors generalize through their covariant description ([3] and [6]) the classical bundle functors of velocities  $T_k^r$ . Their particular importance in differential geometry comes from the fact that there is a bijective correspondence between them and the set of product preserving bundle functors on the category of smooth manifolds.

2.1. **Weil algebra.**

A *Weil algebra* is a finite-dimensional quotient of the algebra of germs  $\mathcal{E}_p = C_0^\infty(\mathbb{R}^p, \mathbb{R})$  ( $p \in \mathbb{N}$ ).

---

2010 *Mathematics Subject Classification*: primary 58A32.

*Key words and phrases*: projectable vector field, Weil bundle, product preserving gauge bundle functor, lift.

Received March 24, 2014. Editor I. Kolář.

DOI: 10.5817/AM2014-3-161

We denote by  $\mathcal{M}_p$  the ideal of germs vanishing at 0;  $\mathcal{M}_p$  is the maximal ideal of the local algebra  $\mathcal{E}_p$ .

Equivalently, a *Weil algebra* is a real commutative unital algebra such that  $A = \mathbb{R} \cdot 1_A \oplus N$ , where  $N$  is a finite dimensional ideal of nilpotent elements.

**Example 2.1.** (1)  $\mathbb{R} = \mathbb{R} \cdot 1 \oplus \{0\} = \mathcal{E}_p / \mathcal{M}_p$  is a Weil algebra.  
 (2)  $J_0^r(\mathbb{R}^p, \mathbb{R}) = \mathbb{R} \cdot 1 \oplus J_0^r(\mathbb{R}^p, \mathbb{R})_0 = \mathcal{E}_p / \mathcal{M}_p^{r+1}$  is a Weil algebra.

**2.2. Weil functors.**

Let us recall the following lemma ([6, Lemma 35.8]): Let  $M$  be a smooth manifold and let  $\varphi: C^\infty(M, \mathbb{R}) \rightarrow A$  be an algebra homomorphism into a Weil algebra  $A$ . Then there is a point  $x \in M$  and some  $k \geq 0$  such that  $\ker \varphi$  contains the ideal of all functions which vanish at  $x$  up the order  $k$  i.e.  $\ker \varphi \supset (I_x)^{k+1}$  with  $I_x = \{f \in C^\infty(M, \mathbb{R}) / f(x) = 0\}$ . More precisely,  $\{x\} = \bigcap_{f \in \ker \varphi} f^{-1}(\{0\})$ .

One defines a functor  $F_A: \mathcal{M}f \rightarrow \mathcal{E}ns$  by:

$$F_A M := \text{Hom}(C^\infty(M, \mathbb{R}), A) \text{ and } F_A f(\varphi) := \varphi \circ f^*$$

for a manifold  $M$  and  $f \in C^\infty(M, N)$ , where  $f^*: C^\infty(N, \mathbb{R}) \rightarrow C^\infty(M, \mathbb{R})$  is the pull-back algebra homomorphism.


Equivalently, if for a manifold  $M$  and a point  $x \in M$ ,  $\text{Hom}(C_x^\infty(M, \mathbb{R}), A)$  is the set of algebra homomorphisms from  $C_x^\infty(M, \mathbb{R})$  into  $A$ , one may define a functor  $G_A: \mathcal{M}f \rightarrow \mathcal{E}ns$  by:

$$G_A M := \bigcup_{x \in M} \text{Hom}(C_x^\infty(M, \mathbb{R}), A) \text{ and } (G_A f)_x(\varphi_x) := \varphi_x \circ f_x^*$$

for a manifold  $M$  and  $f \in C^\infty(M, N)$ , where  $f_x^*: C_{f(x)}^\infty(N, \mathbb{R}) \rightarrow C_x^\infty(M, \mathbb{R})$  is the pull-back algebra homomorphism induced by  $f^*$ .

$F_A$  and  $G_A$  are equivalent functors: Indeed let  $\varphi \in \text{Hom}(C^\infty(M, \mathbb{R}), A)$  and  $\{x\} = \bigcap_{f \in \ker \varphi} f^{-1}(\{0\})$ .

By the previous lemma, there is a unique  $\varphi_x \in \text{Hom}(C_x^\infty(M, \mathbb{R}), A)$  such that the diagram



commutes; the maps  $\chi_M: F_A M \rightarrow G_A M, \varphi \mapsto \varphi_x$  are bijective and define a natural equivalence  $\chi: F_A \rightarrow G_A$ .

Now, let  $T^A = G_A$  and  $\pi_{A,M}: T^A M \rightarrow M, \varphi \ni (T^A M)_x \mapsto x$ .  $(T^A M, M, \pi_{A,M})$  is a well-defined fibered manifold. Indeed let  $c = (U, u^i), 1 \leq i \leq m$  be a chart of  $M$ ; then the map

$$\begin{aligned}
 \phi_c: (\pi_{A,M})^{-1}(U) &\longrightarrow U \times N^m \\
 \varphi_x &\longmapsto (x, \varphi_x(\text{germ}_x(u^i - u^i(x)))) ;
 \end{aligned}$$

is a local trivialization of  $T^A M$ . Given another manifold  $M'$  and a smooth map  $f: M \rightarrow M'$ , let

$$T^A f: T^A M \longrightarrow T^A M'$$

$$\varphi_x \longmapsto \varphi_x \circ f_x^* .$$

Then  $T^A f$  is a fibered map and  $T^A: \mathcal{M}f \rightarrow \mathcal{F}\mathcal{M}$  is a product preserving bundle functor called the *Weil functor* associated to  $A$ .

Let  $c = (U, u^i, \cdot)$ ,  $1 \leq i \leq m$  be a chart of  $M$ ; a fibered chart  $(\pi_{A,M}^{-1}(U), u^{i,\alpha})$ ,  $1 \leq i \leq m$ ,  $1 \leq \alpha \leq K (= \dim A)$  of  $T^A M$  is defined by  $u^{i,\alpha} = \text{pr}_\alpha \circ B \circ T^A(u^i)$ , where  $B: A \rightarrow \mathbb{R}^K$  is a linear isomorphism and  $\text{pr}_\alpha: \mathbb{R}^K \rightarrow \mathbb{R}$  the  $\alpha$ -th projection.

### 3. PRODUCT PRESERVING GAUGE BUNDLE FUNCTOR ON VECTOR BUNDLES

#### 3.1. Product preserving gauge bundle functor on $\mathcal{V}\mathcal{B}$ .

Let  $F: \mathcal{V}\mathcal{B} \rightarrow \mathcal{F}\mathcal{M}$  be a covariant functor from the category  $\mathcal{V}\mathcal{B}$  of all vector bundles and their vector bundle homomorphisms into the category  $\mathcal{F}\mathcal{M}$  of fibered manifolds and their fibered maps. Let  $B_{\mathcal{V}\mathcal{B}}: \mathcal{V}\mathcal{B} \rightarrow \mathcal{M}f$  and  $B_{\mathcal{F}\mathcal{M}}: \mathcal{F}\mathcal{M} \rightarrow \mathcal{M}f$  be the respective base functors.

**Definition 3.1.**  $F$  is a *gauge bundle functor* on  $\mathcal{V}\mathcal{B}$  when the following conditions are satisfied:

- **(Prolongation)**  $B_{\mathcal{F}\mathcal{M}} \circ F = B_{\mathcal{V}\mathcal{B}}$  i.e.  $F$  transforms a vector bundle  $E \xrightarrow{q} M$  in a fibered manifold  $FE \xrightarrow{p_E} M$  and a vector bundle morphism  $E \xrightarrow{f} G$  over  $M \xrightarrow{\bar{f}} N$  in a fibered map  $FE \xrightarrow{Ff} FG$  over  $\bar{f}$ .

- **(Localization)** For any vector bundle  $E \xrightarrow{q} M$  and any inclusion of an open vector subbundle  $i: \pi^{-1}(U) \hookrightarrow E$ , the fibered map  $F\pi^{-1}(U) \rightarrow p_E^{-1}(U)$  over  $\text{id}_U$  induced by  $Fi$  is an isomorphism; then the map  $Fi$  can be identified with the inclusion  $p_E^{-1}(U) \hookrightarrow FE$ .

Given two gauge bundle functors  $F_1, F_2$  on  $\mathcal{V}\mathcal{B}$ , by a *natural transformation*  $\tau: F_1 \rightarrow F_2$  we shall mean a system of base preserving fibered maps  $\tau_E: F_1 E \rightarrow F_2 E$  for every vector bundle  $E$  satisfying  $F_2 f \circ \tau_E = \tau_G \circ F_1 f$  for every vector bundle morphism  $f: E \rightarrow G$ .

A gauge bundle functor  $F$  on  $\mathcal{V}\mathcal{B}$  is *product preserving* if for any product projections  $E_1 \xleftarrow{\text{pr}_1} E_1 \times E_2 \xrightarrow{\text{pr}_2} E_2$  in the category  $\mathcal{V}\mathcal{B}$ ,  $FE_1 \xleftarrow{F\text{pr}_1} F(E_1 \times E_2) \xrightarrow{F\text{pr}_2} FE_2$  are product projections in the category  $\mathcal{F}\mathcal{M}$ . In other words, the map  $(F\text{pr}_1, F\text{pr}_2): F(E_1 \times E_2) \rightarrow F(E_1) \times F(E_2)$  is a fibered isomorphism over  $\text{id}_{M_1 \times M_2}$ .

**Example 3.1.** (a) Each Weil functor  $T^A$  induces a product preserving gauge bundle functor  $T^A: \mathcal{V}\mathcal{B} \rightarrow \mathcal{F}\mathcal{M}$  in a natural way.

(b) Let  $A = \mathbb{R} \cdot 1_A \oplus N$  be a Weil algebra and  $V$  a  $A$ -module such that  $\dim_{\mathbb{R}}(V) < \infty$ . For a vector bundle  $(E, M, q)$  and  $x \in M$ , let

$$T_x^{A,V} E = \{(\varphi_x, \psi_x) / \varphi_x \in \text{Hom}(C_x^\infty(M, \mathbb{R}), A) \text{ and } \psi_x \in \text{Hom}_{\varphi_x}(C_x^{\infty, \text{fl}}(E), V)\}$$

where  $\text{Hom}(C_x^\infty(M, \mathbb{R}), A)$  is the set of algebra homomorphisms  $\varphi_x$  from the algebra  $C_x^\infty(M, \mathbb{R}) = \{\text{germ}_x(g) / g \in C^\infty(M, \mathbb{R})\}$  into  $A$  and  $\text{Hom}_{\varphi_x}(C_x^{\infty, \text{fl}}(E), V)$  is the set of module homomorphisms  $\psi_x$  over  $\varphi_x$  from the  $C_x^\infty(M, \mathbb{R})$ -module  $C_x^{\infty, \text{fl}}(E, \mathbb{R}) = \{\text{germ}_x(h) / h: E \rightarrow \mathbb{R} \text{ is fibre linear}\}$  into  $\mathbb{R}$ .

Let  $T^{A,V} E = \bigcup_{x \in M} T_x^{A,V} E$  and  $p_E^{A,V}: T^{A,V} E \rightarrow M, (\varphi, \psi) \ni T_x^{A,V} E \mapsto x$ .

$(T^{A,V} E, M, p_E^{A,V})$  is a well-defined fibered manifold. Indeed, let  $c = (q^{-1}(U), x^i = u^i \circ q, y^j), 1 \leq i \leq m, 1 \leq j \leq n$  be a fibered chart of  $E$ ; then the map

$$\begin{aligned} \phi_c: (p_E^{A,V})^{-1}(U) &\longrightarrow U \times N^m \times V^n \\ (\varphi_x, \psi_x) &\longmapsto (x, \varphi_x(\text{germ}_x(u^i - u^i(x))), \psi_x(\text{germ}_x(y^j))) ; \end{aligned}$$

is a local trivialization of  $T^{A,V} E$ . Given another vector bundle  $(G, N, q')$  and a vector bundle homomorphism  $f: E \rightarrow G$  over  $\bar{f}: M \rightarrow N$ , let

$$\begin{aligned} T^{A,V} f: T^{A,V} E &\longrightarrow T^{A,V} G \\ (\varphi_x, \psi_x) &\longmapsto (\varphi_x \circ \bar{f}_x^*, \psi_x \circ f_x^*), \end{aligned}$$

where  $\bar{f}_x^*: C_{\bar{f}(x)}^\infty(N) \rightarrow C_x^\infty(M)$  and  $f_x^*: C_{\bar{f}(x)}^{\infty, \text{fl}}(G) \rightarrow C_x^{\infty, \text{fl}}(E)$  are given by the pull-back with respect to  $\bar{f}$  and  $f$ . Then  $T^{A,V} f$  is a fibered map over  $\bar{f}$ .  $T^{A,V}: \mathcal{VB} \rightarrow \mathcal{FM}$  is a product preserving gauge bundle functor (see [7]).

**Remark 3.1.** Let  $F: \mathcal{VB} \rightarrow \mathcal{FM}$  be a product preserving gauge bundle functor.

(a)  $F$  associates the pair  $(A^F, V^F)$  where  $A^F = F(\text{id}_{\mathbb{R}}: \mathbb{R} \rightarrow \mathbb{R})$  is a Weil algebra and  $V^F = F(\mathbb{R} \rightarrow pt)$  is a  $A^F$ -module such that  $\dim_{\mathbb{R}}(V^F) < \infty$ . Moreover there is a natural isomorphism  $\Theta: F \rightarrow T^{A^F, V^F}$  and equivalence classes of functors  $F$  are in bijection with equivalence classes of pairs  $(A^F, V^F)$ . In particular, the product preserving gauge bundle functor associated to the Weil functor  $T^A$  is equivalent to  $T^{A,A}$ .

(b) Let  $K = \dim A^F$  and  $L = \dim V^F$ ; if  $c = (\pi^{-1}(U), x^i, y^j), 1 \leq i \leq m, 1 \leq j \leq n$  is a fibered chart of  $E$ , a fibered chart  $(p_E^{-1}(U), x^{i,\alpha}, y^{j,\beta}), 1 \leq \alpha \leq K, 1 \leq \beta \leq L$  of  $FE$  is defined by

$$(3.1) \quad \begin{cases} x^{i,\alpha} = \lambda_B^\alpha \circ F(x^i) \\ y^{j,\beta} = \mu_C^\beta \circ F(y^j) \end{cases}$$

where  $\lambda_B^\alpha = \text{pr}_\alpha \circ B, \mu_C^\beta = \text{pr}_\beta \circ C$  and  $B: A^F \rightarrow \mathbb{R}^K, C: V^F \rightarrow \mathbb{R}^L$  are linear isomorphisms. In the particular case  $F = T, A^F = V^F = \mathbb{D} = \mathbb{R}[X] / (X^2)$  is the algebra of dual numbers and the previous coordinate system is denoted

$(x^i, \dot{x}^i, y^j, \dot{y}^j)$  where

$$x^i = \lambda_B^1 \circ F(x^i), \dot{x}^i = \lambda_B^2 \circ F(x^i), y^j = \mu_B^1 \circ F(y^j), \dot{y}^j = \mu_B^2 \circ F(y^j)$$

and  $B: \mathbb{D} \rightarrow \mathbb{R}^2$  the canonical isomorphism.

**3.2. The natural isomorphism  $\kappa: F \circ T \rightarrow T \circ F$ .**

For a product preserving gauge bundle functor  $F: \mathcal{VB} \rightarrow \mathcal{FM}$ , let  $\kappa: F \circ T \rightarrow T \circ F$  be the canonical natural isomorphism associated to the exchange isomorphism  $(\mathbb{D} \otimes_{\mathbb{R}} A^F, \mathbb{D} \otimes_{\mathbb{R}} V^F) \cong (A^F \otimes_{\mathbb{R}} \mathbb{D}, V^F \otimes_{\mathbb{R}} \mathbb{D})$  ([7, Corollary 3]). Using the definition of composed functors  $F \circ T$  and  $T \circ F$ , one can check that locally

$$\kappa_E: (x^{i,\alpha}, \dot{x}^{i,\alpha}, y^{j,\beta}, \dot{y}^{j,\beta}) \mapsto (x^{i,\alpha}, \dot{x}^{i,\alpha}, y^{j,\beta}, \dot{y}^{j,\beta})$$

with  $\dot{x}^{i,\alpha} = \dot{x}^{i,\alpha}$  and  $\dot{y}^{j,\beta} = \dot{y}^{j,\beta}$ . The following assertion is clear.

**Proposition 3.1.** (a) *The diagram*

$$\begin{array}{ccc} F(TE) & \xrightarrow{\kappa_E} & T(FE) \\ & \searrow F(\pi_E) & \downarrow \pi_{FE} \\ & & FE \end{array}$$

*commutes for any vector bundle  $E \xrightarrow{q} M$ .*

(b) *If  $(F, \pi')$  is an excellent pair (i.e.  $\pi': F \rightarrow \text{id}_{\mathcal{VB}}$  is a natural epimorphism), the diagram*

$$\begin{array}{ccc} F(TE) & \xrightarrow{\kappa_E} & T(FE) \\ & \searrow \pi'_{FE} & \downarrow T(\pi'_E) \\ & & TE \end{array}$$

*commutes for any vector bundle  $E \xrightarrow{q} M$ .*

Let  $X \in \mathfrak{X}_{\text{proj}}(E)$  be a *projectable vector field* on  $E$  i.e. a fibered map over a vector field  $\bar{X}$  on  $M$ . We assume that  $X$  is a vector bundle morphism. The commutative diagram

$$\begin{array}{ccc} FE & \xrightarrow{(\mathcal{F}_E)X} & TFE \\ FX \downarrow & \nearrow \kappa_E & \\ FTE & & \end{array}$$

defines a vector field on  $FE$  called the *complete lift* of  $X$  and denoted  $X^c$ . The natural operator  $\mathcal{F}: T \rightsquigarrow TF$  is called the *flow operator* of  $F$ .

When we restrict ourself to vector bundles of the form  $(M, M, \text{id}_M)$ ,  $\kappa$  and  $\mathcal{F}$  are exactly the canonical flow natural equivalence and the flow operator associated to the Weil functor  $T^{A^F}$ , [6].

4. PROLONGATION OF FUNCTIONS

We generalize for a product preserving gauge bundle  $F: \mathcal{VB} \rightarrow \mathcal{FM}$  some prolongations of [2].

Let  $f: E \rightarrow \mathbb{R}$  be a smooth function defined on a vector bundle  $q: E \rightarrow M$ .

**Definition 4.1.**

- (a) The  $\lambda$ -lift of  $f$  constant on fibers is  $f^{(\lambda)} := \lambda \circ Ff$ , for  $\lambda \in C^\infty(A^F, \mathbb{R})$ .
- (b) The  $\mu$ -lift of  $f$  linear on fibers (i.e.  $f \in C^\infty_\ell(E, \mathbb{R})$ ) is  $f^{(\mu)} := \mu \circ Ff$ , for  $\mu \in C^\infty(V^F, \mathbb{R})$ .

In the particular case  $(E, M, q) = (M, M, \text{id}_M)$ , lifts of functions constant on fibers correspond to lifts of functions associated to the Weil functor  $T^{A^F}$  [2]. Lifts of functions linear on fibers correspond to lifts of smooth sections of the dual bundle  $q^*: E^* \rightarrow M$ .

It is easy to check that  $(f \circ h)^{(\lambda)} = f^{(\lambda)} \circ Fh$ , for  $h: G \rightarrow E$  a vector bundle morphism and  $(f_1 + f_2)^{(\lambda)} = f_1^{(\lambda)} + f_2^{(\lambda)}$  when  $\lambda$  is a linear map. Replacing  $\lambda$  with  $\mu$ , the previous identities hold.

According to (3.1),

$$x^{i,\alpha} = (x^i)^{(\lambda_B^\alpha)} \quad \text{and} \quad y^{j,\beta} = (y^j)^{(\mu_C^\beta)},$$

hence the following result holds:

**Proposition 4.1.** *If two vector fields  $\widehat{X}, \widehat{Y}$  on  $FE$  satisfy*

$$\widehat{X}(f^{(\lambda)}) = \widehat{Y}(f^{(\lambda)}) \quad \text{and} \quad \widehat{X}(g^{(\mu)}) = \widehat{Y}(g^{(\mu)}),$$

*for any  $f$  constant on fibers,  $g$  linear on fibers,  $\lambda \in C^\infty(A^F, \mathbb{R})$  and  $\mu \in C^\infty(V^F, \mathbb{R})$ , then  $\widehat{X} = \widehat{Y}$ .*

5. PROLONGATION OF PROJECTABLE VECTOR FIELDS

5.1. **Natural transformations**  $Q(a): T \circ F \rightarrow T \circ F$ .

For a vector bundle  $q: E \rightarrow M$ , let us denote  $\mu_E: \mathbb{R} \times TE \rightarrow TE$ ,  $(\alpha, k_u) \ni \mathbb{R} \times T_u E \mapsto \alpha \cdot k_u \in T_u E$ , the fibered multiplication. This is a vector bundle morphism over the projection  $\mathbb{R} \times E \rightarrow E$ , hence for any  $a \in A^F$ , we have a natural transformation  $FT \rightarrow FT$  given by the partial maps  $F\mu_E(a, \cdot): FTE \rightarrow FTE$ . If  $\kappa: T \rightarrow TF$  is the canonical natural isomorphism (subsection 3.2), one can deduce a natural transformation  $Q(a): TF \rightarrow TF$  by

$$(5.1) \quad Q(a) = \kappa \circ F\mu(a, \cdot) \circ \kappa^{-1}.$$

The restriction of  $Q(a)$  to vector bundles of the form  $(M, M, \text{id}_M)$  is just the natural affinor associated to  $a$  [5].

Let us compute the algebra homomorphism  $\mu_a: A^{FT} \rightarrow A^{FT}$  and the module homomorphisms  $\nu_a: V^{FT} \rightarrow V^{FT}$  over  $\mu_a$  associated to the natural transformation  $F\mu(a, \cdot)$ : Let  $p_1: \mathbb{D} \rightarrow \mathbb{R}1$ ,  $p_2: \mathbb{D} \rightarrow \mathbb{R}\delta$  the canonical projections associated to the algebra of dual numbers and  $i_u: \mathbb{R} \rightarrow \mathbb{D}$ ,  $t \mapsto tu$ ,  $u \in \mathbb{D}$ ; since  $\mu_{\mathbb{R} \rightarrow \mathbb{R}}(s, u) =$

$p_1(u) + p_2(su) = T(ad_{\mathbb{R}})(p_1(u), p_2 \circ m \circ (\text{id}_{\mathbb{R}} \times \text{id}_{\mathbb{D}})(s, u))$ ,  $m(s, u) = su$  then for  $b \in A^F$

$$\begin{aligned} \mu_a(Fi_u(b)) &= F(Tad_{\mathbb{R}})(Fi_{p_1(u)}(b), F(p_2 \circ m)(a, Fi_u(b))) \\ &= F(Tad_{\mathbb{R}})(Fi_{p_1(u)}(b), F(p_2 \circ m)(F\text{id}_{\mathbb{R}}(a), Fi_u(b))) \\ &= F(Tad_{\mathbb{R}})(Fi_{p_1(u)}(b), F(p_2 \circ m \circ \text{id}_{\mathbb{R}} \times i_u)(a, b)) \\ &= F(Tad_{\mathbb{R}})(Fi_{p_1(u)}(b), F(i_{p_2(u)} \circ m_0)(a, b)), m_0(s, t) = st \\ &= Fi_{p_1(u)}(b) + Fi_{p_2(u)}(ab) \\ &= \mu_{FT}(p_1(u) \otimes b + p_2(u) \otimes ab), \end{aligned}$$

where  $\mu_{FT}: \mathbb{D} \otimes_{\mathbb{R}} A^F \rightarrow A^{FT}$  is the canonical algebra isomorphism. Hence

$$\begin{aligned} \mu_{FT}^{-1} \circ \mu_a \circ \mu_{FT}: \mathbb{D} \otimes_{\mathbb{R}} A^F &\rightarrow \mathbb{D} \otimes_{\mathbb{R}} A^F \\ u \otimes b &\mapsto p_1(u) \otimes b + p_2(u) \otimes ab. \end{aligned}$$

Similarly,

$$\begin{aligned} v_a(Fi_u(v)) &= F(Tad_{\mathbb{R}})(Fi_{p_1(u)}(v), F(p_2 \circ m)(a, Fi_u(v))) \\ &= F(Tad_{\mathbb{R}})(Fi_{p_1(u)}(v), F(p_2 \circ m)(F\text{id}_{\mathbb{R}}(a), Fi_u(v))) \\ &= F(Tad_{\mathbb{R}})(Fi_{p_1(u)}(v), F(p_2 \circ m \circ \text{id}_{\mathbb{R}} \times i_u)(a, v)) \\ &= F(Tad_{\mathbb{R}})(Fi_{p_1(u)}(v), F(i_{p_2(u)} \circ m_0)(a, v)), m_0(s, t) = st \\ &= Fi_{p_1(u)}(v) + Fi_{p_2(u)}(a \cdot v) \\ &= v_{FT}(p_1(u) \otimes v + p_2(u) \otimes a \cdot v), \end{aligned}$$

where  $v_{FT}: \mathbb{D} \otimes_{\mathbb{R}} V^F \rightarrow V^{FT}$  is the natural module isomorphism over  $\mu_{FT}$ . Hence

$$\begin{aligned} v_{FT}^{-1} \circ v_a \circ v_{FT}: \mathbb{D} \otimes_{\mathbb{R}} V^F &\rightarrow \mathbb{D} \otimes_{\mathbb{R}} V^F \\ u \otimes v &\mapsto p_1(u) \otimes v + p_2(u) \otimes a \cdot v. \end{aligned}$$

### 5.2. Prolongation of projectable vector fields.

In this subsection, all projectable vector fields are vector bundle morphisms.

**Definition 5.1.** For a projectable vector field  $X \in \mathfrak{X}_{proj}(E)$ , its  $a$ -lift is given by  $X^{(a)} = Q(a)_E \circ (\mathcal{F}_E)X$ , where  $\mathcal{F}: T \rightsquigarrow TF$  is the flow operator of  $F$ .

Let  $\lambda: A^F \rightarrow \mathbb{R}$ ,  $\mu: V^F \rightarrow \mathbb{R}$  linear maps and  $\lambda_a: A^F \rightarrow \mathbb{R}$ ,  $\mu_a: V^F \rightarrow \mathbb{R}$  given by  $\lambda_a(x) = \lambda(ax)$ ,  $\mu_a(v) = \mu(a \cdot v)$ , for  $a \in A^F$ .

**Theorem 5.1.** We have

$$X^{(a)}(f^{(\lambda)}) = (X(f))^{(\lambda_a)} \quad \text{and} \quad X^{(a)}(g^{(\mu)}) = (X(g))^{(\mu_a)},$$

for any  $f$  constant on fibers and  $g$  linear on fibers.

**Proof.** Let  $\tau \in C_{\ell}^{\infty}(T\mathbb{R}, \mathbb{R})$  such that  $\tau_x: T_x\mathbb{R} \rightarrow \mathbb{R}$ ,  $x \in \mathbb{R}$  the canonical isomorphism. Identifying the module of 1-forms on a manifold  $M$  with  $C_{\ell}^{\infty}(TM, \mathbb{R})$ , the differential of  $f \in C^{\infty}(M, \mathbb{R})$  is given by  $df = \tau \circ Tf$  and for a vector field  $\overline{X}$  on  $M$ ,  $\overline{X}(f) = \langle \overline{X}, df \rangle = \tau \circ Tf \circ \overline{X}$ .

We have

$$\begin{aligned}
 X^{(a)}(f^{(\lambda)}) &= \tau \circ Tf^{(\lambda)} \circ \kappa_E \circ F\mu_E(a, \cdot) \circ FX \\
 &= \tau \circ T\lambda \circ T(Ff) \circ \kappa_E \circ F\mu_E(a, \cdot) \circ FX \\
 &= \tau \circ T\lambda \circ \kappa_{\mathbb{R} \rightarrow \mathbb{R}} \circ F(\mu_{\mathbb{R} \rightarrow \mathbb{R}}(a, \cdot)) \circ F(Tf \circ X) \\
 &= \lambda_a \circ F(\tau) \circ F(Tf \circ X) \\
 &= (X(f))^{(\lambda_a)}.
 \end{aligned}$$

The second equality is proved in the same way.  $\square$

**Corollary 5.1.** *For any vector fields  $X, Y \in \mathfrak{X}_{\text{proj}}(E)$ , we have*

$$[X^{(a)}, Y^{(b)}] = [X, Y]^{(ab)}.$$

**Proof.** Direct consequence of the previous result and Proposition 4.1.  $\square$

This generalizes (for product preserving gauge bundle functors on vector bundles) some results of [2].

## REFERENCES

- [1] Cordero, L.A., Dobson, C.T.J., De León, M., *Differential geometry of frame bundles*, Kluwer Academic Publishers, 1989.
- [2] Gancarzewicz, J., Mikulski, W.M., Pogoda, Z., *Lifts of some tensor fields and connections to product preserving functors*, Nagoya Math. J. **135** (1994), 1–14.
- [3] Kolář, I., *Covariant approach to natural transformations of Weil bundles*, CMUC **27** (1986), 723–729.
- [4] Kolář, I., *On the natural operators on vector fields*, Ann. Global Anal. Geom. **6** (1988), 119–117.
- [5] Kolář, I., Modugno, M., *Torsions of connections on some natural bundles*, Differential Geom. Appl. **2** (1992), 1–16.
- [6] Kolář, I., Slovák, J., Michor, P.W., *Natural operations in differential geometry*, Springer-Verlag Berlin–Heidelberg, 1993.
- [7] Mikulski, W.M., *Product preserving gauge bundle functors on vector bundles*, Colloq. Math. **90** (2001), no. 2, 277–285.
- [8] Ntyam, A., Mba, A., *On natural vector bundle morphisms  $T^A \circ \otimes_s^q \rightarrow \otimes_s^q \circ T^A$  over  $\text{id}_{T^A}$* , Ann. Polon. Math. **96** (2009), no. 3, 295–301.
- [9] Ntyam, A., Wouafo, K.J., *New versions of curvature and torsion formulas for the complete lifting of a linear connection to Weil bundles*, Ann. Polon. Math. **82** (2003), no. 3, 133–140.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE,  
FACULTY OF SCIENCE, UNIVERSITY OF NGAOUNDÉRÉ,  
P.O BOX 454 NGAOUNDÉRÉ, CAMEROON  
*E-mail:* antyam@uy1.uninet.cm

DEPARTMENT OF MATHEMATICS,  
FACULTY OF SCIENCE, UNIVERSITY OF YAOUNDÉ 1,  
P.O BOX 812 YAOUNDÉ, CAMEROON  
*E-mail:* georgywan@yahoo.fr

DEPARTMENT OF MATHEMATICS,  
FACULTY OF SCIENCE, UNIVERSITY OF YAOUNDÉ 1,  
P.O BOX 812 YAOUNDÉ, CAMEROON  
*E-mail:* bitjong@uy1.uninet.cm