

## ON A SIMULTANEOUS APPROXIMATION PROBLEM CONCERNING BINARY RECURRENCES

PÉTER KISS

*Dedicated to Professor Árpád Varcza on his 60th birthday*

ABSTRACT. Let  $R_n$  ( $n = 0, 1, 2, \dots$ ) be a second order linear recursive sequence of rational integers defined by  $R_n = AR_{n-1} + BR_{n-2}$  for  $n > 1$ , where  $A$  and  $B$  are integers and the initial terms are  $R_0 = 0$ ,  $R_1 = 1$ . It is known, that if  $\alpha, \beta$  are the roots of the equation  $x^2 - Ax - B = 0$  and  $|\alpha| > |\beta|$ , then  $R_{n+1}/R_n \rightarrow \alpha$  as  $n \rightarrow \infty$ . Approximating  $\alpha$  with the rational number  $R_{n+1}/R_n$ , it was shown that  $\left| \alpha - \frac{R_{n+1}}{R_n} \right| < \frac{1}{c \cdot |R_n|^2}$  holds with a constant  $c > 0$  for infinitely many  $n$  if and only if  $|B| = 1$ . In this paper we investigate the quality of the approximation of  $\alpha$  and  $\alpha^s$  by the rational numbers  $R_{n+1}/R_n$  and  $R_{n+s}/R_n$  simultaneously.

### INTRODUCTION

Let  $A$  and  $B$  be fixed non-zero integers and let  $\{R_n\}_{n=0}^\infty$  be a second order linear recursive sequence of rational integers defined by the recursion

$$R_n = AR_{n-1} + BR_{n-2} \quad (n > 1)$$

with initial terms  $R_0 = 0$  and  $R_1 = 1$ . Denote by  $\alpha$  and  $\beta$  the roots of the characteristic equation

$$x^2 - Ax - B = 0$$

of the sequence and suppose that  $|\alpha| \geq |\beta|$ . We suppose that the sequence  $\{R_n\}$  is non degenerate, i.e.  $\alpha/\beta$  is not a root of unity. In this case the terms of the sequence can be expressed as

$$(1) \quad R_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}$$

for any  $n \geq 0$ .

If  $D = A^2 + 4B > 0$ , then  $\alpha$  and  $\beta$  are real numbers and  $|\alpha| > |\beta|$ . It implies that

$$\frac{R_{n+1}}{R_n} = \frac{\alpha^{n+1} - \beta^{n+1}}{\alpha^n - \beta^n} = \alpha \cdot \frac{1 - (\beta/\alpha)^{n+1}}{1 - (\beta/\alpha)^n} \rightarrow \alpha \text{ as } n \rightarrow \infty$$

and so  $\alpha$  can be approximated by rational numbers  $R_{n+1}/R_n$ . For the quality of this approximation in [2] we proved that the inequality

$$\left| \alpha - \frac{R_{n+1}}{R_n} \right| < \frac{1}{c \cdot |R_n|^2}$$

---

2000 *Mathematics Subject Classification.* 11B37, 11J13, 11J86.

*Key words and phrases.* Linear recursive sequence, Diophantine approximation.

Research (partially) was supported by the Hungarian OTKA foundation, grant No. T 29330 and 032898.

with some  $c > 0$  holds for infinitely many  $n$  if and only if  $|B| = 1$ , furthermore in the case  $|B| = 1$  the best approximation constant is  $c = \sqrt{D}$ . It was also proved that if  $|B| = 1$  and  $\left| \alpha - \frac{p}{q} \right| < \frac{1}{\sqrt{D}q^2}$  with a rational number  $p/q$ , then  $p/q = R_{n+1}/R_n$  for some  $n$ .

In this paper we deal with a simultaneous approximation problem. Let  $\gamma_1$  and  $\gamma_2$  be irrational numbers. It is known that there are infinitely many triples  $p_1, p_2, q$  of rational integers and a constant  $c > 0$  such that

$$\left| \gamma_1 - \frac{p_1}{q} \right| < \frac{1}{c \cdot q^{3/2}} \quad \text{and} \quad \left| \gamma_2 - \frac{p_2}{q} \right| < \frac{1}{c \cdot q^{3/2}}$$

hold simultaneously and the order  $\frac{3}{2}$  of the approximation is the best possibility in general. In the next section we show that if  $\gamma_1 = \alpha$  and  $\gamma_2 = \alpha^s$  where  $s$  is a positive integer then the order of their simultaneous approximation can be 2.

#### THE REAL CASE

In this section we investigate the case when  $D = A^2 + 4B > 0$ , so  $\alpha, \beta$  are real numbers and  $|\alpha| \neq |\beta|$ .

**Theorem 1.** *Let  $\{R_n\}$  be a second order linear recursive sequence defined in the Introduction and let  $s \geq 2$  be a positive integer. Suppose that  $D > 0$ ,  $|\alpha| > |\beta|$  and  $|B| = 1$ . Then there is a constant  $c_0 > 0$  such that the inequalities*

$$\left| \alpha - \frac{R_{n+1}}{R_n} \right| < \frac{1}{c_0 R_n^2} \quad \text{and} \quad \left| \alpha^s - \frac{R_{n+s}}{R_n} \right| < \frac{1}{c_0 \cdot R_n^2}$$

hold simultaneously for infinitely many positive integer  $n$ .

*Proof.* For an integer  $k \geq 1$  by (1) we have

$$\left| \alpha^k - \frac{R_{n+k}}{R_n} \right| = \left| \alpha^k - \frac{\alpha^{n+k} - \beta^{n+k}}{\alpha^n - \beta^n} \right| = \left| \frac{\beta^k \beta^n - \alpha^k \beta^n}{\alpha^n - \beta^n} \right| =$$

$$|\beta|^n \cdot \left| \frac{\alpha^k - \beta^k}{\alpha^n - \beta^n} \right| = |\beta|^n \frac{|R_k|}{|R_n|} = |\beta|^n \left| \frac{\alpha^n - \beta^n}{\alpha - \beta} \right| \cdot \frac{|R_k|}{R_n^2} =$$

$$(2) \quad |\alpha\beta|^n \cdot \frac{1}{|\alpha - \beta|} \cdot \left| 1 - \left( \frac{\beta}{\alpha} \right)^n \right| \cdot \frac{|R_k|}{R_n^2}.$$

But  $\left| \frac{\beta}{\alpha} \right| < 0$ ,  $|\alpha - \beta| = \sqrt{D}$  and  $|\alpha\beta| = 1$  since  $|B| = 1$ , so  $(\beta/\alpha)^n \rightarrow 0$  as  $n \rightarrow \infty$  and by (2)

$$\left| \alpha^k - \frac{R_{n+k}}{R_n} \right| < \frac{|R_k|}{\sqrt{D}} \cdot \frac{1}{R_n^2}$$

for any  $k \geq 1$  and for infinitely many positive integer  $n$  (for any  $n$  if  $\beta/\alpha > 0$  and for any even  $n$  if  $\beta/\alpha < 0$ ). From this inequality the theorem follows with

$$c_0 = \min \left( \frac{\sqrt{D}}{|R_1|}, \frac{\sqrt{D}}{|R_s|} \right) = \frac{\sqrt{D}}{|R_s|}.$$

We note that some other approximation results was obtained by F. Mátyás [6] and B. Zay [7] concerning general recurrences.  $\square$

## COMPLEX CASE

Now let  $D < 0$ . In this case  $\alpha$  and  $\beta$  are not real complex conjugate numbers with  $\left|\frac{\beta}{\alpha}\right| = 1$  and we can conclude in (2) only that  $0 < |1 - (\beta/\alpha)^n| < 2$ , so  $\lim_{n \rightarrow \infty} \frac{R_{n+k}}{R_n}$  does not exist for any  $k \geq 1$ . We can approximate only the powers of  $|\alpha|$  instead of the powers of  $\alpha$  and the quality of these approximations is much weaker than in Theorem 1. In [3] and [4] with R. F. Tichy we proved that there are positive constants  $c_1$  and  $c_2$ , depending only on the sequence  $\{R_n\}$ , such that

$$(3) \quad \left| |\alpha| - \left| \frac{R_{n+1}}{R_n} \right| \right| < \frac{1}{n^{c_1}}$$

for infinitely many  $n$  but

$$\left| |\alpha| - \left| \frac{R_{n+1}}{R_n} \right| \right| > \frac{1}{n^{c_2}}$$

for all sufficiently large  $n$ . It shows that, apart from the constant  $c_1$ , (3) is the best possibility to approximate  $|\alpha|$  by rational numbers of the form  $|R_{n+1}/R_n|$ . Now we prove:

**Theorem 2.** *For any non-degenerate second order linear recurrence  $\{R_n\}$ , defined in the Introduction, for which  $D < 0$ , there are constants  $c_4 > c_3 > 0$  such that the inequalities*

$$(4) \quad \left| |\alpha| - \left| \frac{R_{n+1}}{R_n} \right| \right| < \frac{1}{n^{c_3}}$$

and

$$(5) \quad \left| |\alpha|^s - \left| \frac{R_{n+s}}{R_n} \right| \right| < \frac{1}{n^{c_3}}$$

hold simultaneously for infinitely many pairs  $s > 1$ ,  $n$  of positive integers, but

$$(6) \quad \left| |\alpha|^s - \left| \frac{R_{n+s}}{R_n} \right| \right| > \frac{1}{n^{c_4}}$$

for any given  $s$  and sufficiently large  $n$ .

For the proof of Theorem 2 we need some auxiliary results.

First we recall some results concerning distribution properties of sequences of real numbers modulo 1. Let  $(x_n)$  ( $n = 1, 2, \dots$ ) be a sequence of real numbers. Denote by  $\{x_n\}$  the fractional part of a term  $x_n$  and denote  $I$  intervals for which  $I \in [0, 1]$ . For a positive integer  $N$  let  $A_N(x_n, I)$  be the number of indices  $n$  with  $1 \leq n \leq N$  such that the fractional part of  $x_n$  is contained in the interval  $I$ , i.e.

$$A_N(x_n, I) = \text{card}\{n \leq N : \{x_n\} \in I\}.$$

then the discrepancy of the sequence  $x_n$  is defined by

$$D_N(x_n) = \sup_I \left| \frac{A_N(x_n, I)}{N} - |I| \right|,$$

where the supremum is taken over all subintervals  $I$  of  $[0, 1]$ .

From the definition of  $D_N(x_n)$  it follows that if  $|I| \geq 2D_N(x_n)$ , then there exists an integer  $n$  with  $1 \leq n \leq N$  such that  $\{x_n\} \in I$ . For a special sequence the following estimation hold.

**Lemma 1.** *Let  $\gamma = e^{2\pi\theta i}$  be a complex number, where  $|\gamma| = 1$  and  $0 < \theta < 1$  is an irrational number. Then the discrepancy of the sequence  $(x_n) = (n\theta)$  satisfies the estimation*

$$D_N(x_n) \leq N^{-\delta}$$

for any sufficiently large  $N$ , where  $\delta (> 0)$  depends only on  $\gamma$ .

*Proof.* The lemma follows from a more general theorem of [4], but it can be proved directly using Theorem 2.5 of [5], p. 112. We need another result, too.  $\square$

**Lemma 2.** *Let*

$$\Lambda = b_1 \cdot \log \omega_1 + \cdots + b_t \cdot \log \omega_t,$$

where  $b_i$ 's are rational integers and  $\omega_i$ 's are algebraic numbers different from 0 and 1. Suppose that not all of the  $b_i$ 's are 0 and that the logarithms mean their principal values. Assume that  $\max(|b_i|) \leq B$  ( $B \geq 4$ ),  $\omega_i$  has height at most  $M_i$  ( $\geq 4$ ) and that the field generated by the  $\omega_i$ 's over the rational numbers has degree at most  $d$ . If  $\Lambda \neq 0$ , then

$$|\Lambda| > B^{-C\Omega \cdot \log \Omega'},$$

where  $\Omega = \log M_1 \cdot \log M_2 \cdots \log M_t$ ,  $\Omega' = \Omega / \log M_t$  and  $C$  is an effectively computable positive constant depending only on  $t$  and  $d$ .

*Proof.* It is a result of A. Baker, see in [1].  $\square$

*Proof of Theorem 2.*  $\alpha$  and  $\beta$  are conjugate complex numbers so we can write

$$\beta = r \cdot e^{\pi\theta i}, \quad \alpha = r \cdot e^{-\pi\theta i} \quad \text{and} \quad \frac{\beta}{\alpha} = e^{2\pi\theta i},$$

where  $0 < \theta < 1$  and  $\theta$  is an irrational number since  $\beta/\alpha$  is not a root of unity. By (1), for any  $k \geq 1$  we have

$$(7) \quad \left| \frac{R_{n+k}}{R_n} \right| = \frac{|\alpha^{n+k}(1 - (\beta/\alpha)^{n+k})|}{|\alpha^n(1 - (\beta/\alpha)^n)|} = |\alpha|^k \left| \frac{1 - e^{2\pi(n+k)\theta i}}{1 - e^{2\pi n\theta i}} \right|.$$

Let  $N$  be a positive integer large enough and denote by  $D_N$  the discrepancy of the sequence  $(x_n) = (n\theta)$ . Then, as we have seen above, there are integers  $m_{k1}$  and  $m_{k2}$  with  $1 \leq m_{k1} < m_{k2} \leq N$  such that

$$|m_{k1}\theta - p - \left(1 - \frac{\theta}{2}\right)| < 2D_N$$

and

$$|m_{k2}\theta - q - \left(1 + \frac{\theta}{2}\right)| < 2D_N,$$

where  $p$  and  $q$  are suitable integers. From these inequalities, using the notation  $z = e^{2\pi(1-\frac{\theta}{2})i}$ ,

$$e^{2\pi m_{k1}\theta i} = e^{2\pi(1-\frac{\theta}{2}+\varepsilon_0)i} = z \cdot e^{2\pi\varepsilon_0 i},$$

$$e^{2\pi(m_{k1}+1)\theta i} = e^{2\pi(1+\frac{\theta}{2}+\varepsilon_0)i} = \bar{z} \cdot e^{2\pi\varepsilon_0 i}$$

and

$$e^{2\pi m_{k2}\theta i} = e^{2\pi(1+\frac{\theta}{2}+\varepsilon_1)i} = \bar{z} \cdot e^{2\pi\varepsilon_1 i}$$

follows, where  $|\varepsilon_0|, |\varepsilon_1| = O(D_N)$ . So by (7), with  $m_{k1} = n$  and  $m_{k2} = n + s$ , we obtain the estimations

$$(8) \quad \left| \alpha^n - \frac{R_{n+1}}{R_n} \right| = |\alpha|^n \cdot \left| 1 - \frac{1 - \bar{z} \cdot e^{2\pi\varepsilon_0 i}}{1 - z \cdot e^{2\pi\varepsilon_0 i}} \right| = |\alpha|^n \cdot O(D_N)$$

and

$$(9) \quad \left| \alpha^{n+s} - \frac{R_{n+s}}{R_n} \right| = |\alpha|^{n+s} \cdot \left| 1 - \frac{1 - \bar{z} \cdot e^{2\pi\varepsilon_1 i}}{1 - z \cdot e^{2\pi\varepsilon_0 i}} \right| = |\alpha|^{n+s} \cdot O(D_N).$$

From (8) and (9) the inequalities (4) and (5) follow, since  $O(D_N) < \frac{1}{n^{c_3}}$  by Lemma 1. for any  $c'_3 < \delta$ . If we choose another  $N' (> N)$  which is sufficiently large, then we obtain another pair of integers  $m'_{k_1}$ ,  $m'_{k_2}$  and so the existence of infinitely many integers  $n, s$  can be concluded.

Now we prove inequality (6). Similary as above we obtain that

$$(10) \quad \left| |\alpha|^s - \left| \frac{R_{n+s}}{R_n} \right| \right| = |\alpha|^s \cdot \left| 1 - \left| \frac{1 - (\beta/\alpha)^{n+s}}{1 - (\beta/\theta)^n} \right| \right|.$$

Let  $z$  be a complex number defined by

$$z = e^{(\pi - \pi s \theta)i}.$$

Then for any integers  $s \geq 1$  and  $n$  we have

$$\left( \frac{\beta}{\alpha} \right)^n = e^{2\pi n \theta i} = z \cdot e^{\lambda i},$$

where  $0 < \lambda < 2\pi$  and

$$\begin{aligned} \lambda &= 2\pi n \theta - \pi + \pi s \theta - 2\pi k = \\ &= (2n + s)\pi \theta - (2k + 1)\pi = \end{aligned}$$

$$(2n + s) \cdot \arg(\beta) - (2k + 1) \cdot \arg(-1) =$$

$$(2n + s) \cdot \log \beta - (2n + s) \cdot \log |\beta| - (2k + 1) \cdot \log(-1)$$

with some integer  $k < n + s$ .  $\beta$ ,  $|\beta|$  and  $-1$  are algebraic numbers of degree at most 4, furthermore  $\lambda \neq 0$  since  $\theta$  is an irrational number, so by Lemma 2 we obtain the inequality

$$|\lambda| > n^{-c_4}$$

where  $c_4 > 0$  depends on  $s$  and the sequences  $\{R_n\}$ . It can be similary proved that

$$|\pi - \lambda| > n^{-c_5}.$$

These inequalities imply that

$$(11) \quad |Im(e^{\lambda i})| > n^{-c_6}.$$

By (10) we get

$$(12) \quad \left| |\alpha|^s - \left| \frac{R_{n+s}}{R_n} \right| \right| = |\alpha|^s \cdot \left| 1 - \left| \frac{1 - \bar{z} \cdot e^{\lambda i}}{1 - z \cdot e^{\lambda i}} \right| \right|.$$

The following estimation can be easily seen by elementary arguments (or see in [3]): If  $z$  and  $w$  are non-real complex numbers with  $zw \neq 1$ , then there is a real number  $c_7 > 0$  depending on  $z$  and  $|w|$  such that

$$(13) \quad \left| 1 - \left| \frac{1 - \bar{z}w}{1 - zw} \right| \right| > \min\{1, c_7, |Im(w)|\}.$$

So by (11), (12) and (13)

$$\left| |\alpha|^s - \left| \frac{R_{n+s}}{R_n} \right| \right| > |\alpha|^s \cdot n^{-c_8} > n^{-c_9}$$

follow which proves inequality (6).  $\square$

*Note.* We note that we obtain similar results if the initial terms of the sequence  $\{R_n\}$  are arbitrary, but in this case the constants are weaker and the expressions are more difficult.

## REFERENCES

- [1] BAKER, A., Transcendence theory and its applications, *J. Austral. Math. Soc. (ser. A)* **25** (1978), 438–444.
- [2] KISS, P., A Diophantine approximative property of the second order linear recurrences, *Period. Math. Hungar.* **11** (1980), 281–278.
- [3] KISS, P. AND TICHY, R. F., A discrepancy problem with applications to linear recurrences I, *Proc. Japan Acad. (ser. A)* **65** No. 5 (1989), 135–138.
- [4] KISS, P. AND TICHY, R. F., A discrepancy problem with applications to linear recurrences II, *Proc. Japan Acad. (ser. A)* **65** No. 6 (1989), 191–194.
- [5] KUIPERS, L. AND NIEDERREITER, H., Uniform distribution of sequences, *John Wiley and Sons*, New York, 1974.
- [6] MÁTYÁS, F., On the quotients of the elements of linear recursive sequences of second order (Hungarian) *Mat. Lapok* **27** (1976–1979), 379–389.
- [7] ZAY, B., A generalization of an approximation problem concerning linear recurrences, *Acta Acad. Paed. Agriensis*, sect. Mat. **24** (1997), 41–45.

*Received September 18, 2000.*

*E-mail address:* `kissp@ektf.hu`

DEPARTMENT OF MATHEMATICS,  
KÁROLY ESZTERHÁZY COLLEGE,  
H-3301 EGER, P.O.Box 43,  
HUNGARY