

## SYMMETRIC UNITS AND GROUP IDENTITIES IN GROUP ALGEBRAS. I

VICTOR BOVDI

*Dedicated to Professor L.G. Kovács on his 70th birthday*

ABSTRACT. We describe those group algebras over fields of characteristic different from 2 whose units symmetric with respect to the classical involution, satisfy some group identity.

### 1. INTRODUCTION

Let  $U(A)$  be the group of units of an algebra  $A$  with involution  $*$  over the field  $F$  and let  $S_*(A) = \{u \in U(A) \mid u = u^*\}$  be the set of symmetric units of  $A$ .

Algebras with involution have been actively investigated. In these algebras there are many symmetric elements, for example:  $x + x^*$  and  $xx^*$  for any  $x \in A$ . This raises natural questions about the properties of the symmetric elements and symmetric units. In [10] Ch. Lanski began to study the properties of the symmetric units in prime algebras with involution, in particular when the symmetric units commute. Using the results and methods of [4], in [5] we classified the cases when the symmetric units commute in modular group algebras of  $p$ -groups. The solution of this question for integral group rings and for some modular group rings of arbitrary groups was obtained in [6, 3].

Several results on the group of units  $U(R)$  show that if  $U(R)$  satisfies a certain group theoretical condition (for example, it is nilpotent or solvable), then  $R$ 's properties are restricted and a polynomial identity on  $R$  holds. This suggests that there may be some general underlying relationship between group identities and polynomial identities. In this topic Brian Hartley made the following:

**Conjecture 1.** *Let  $FG$  be a group algebra of a torsion group  $G$  over the field  $F$ . If  $U(FG)$  satisfies a group identity, then  $FG$  satisfies a polynomial identity.*

---

1991 *Mathematics Subject Classification.* Primary 16U60, 16W10.

*Key words and phrases.* PI-algebras, symmetric units.

The research was supported by OTKA No.T 037202 and No.T 038059.

The theory of *PI*-algebras has been established for a long time. On the contrary, the study of algebras with units satisfying a group identity has emerged only recently [11, 12]. Our goal here is to show that with a few extra assumptions, these algebras are actually *PI*-algebras. In fact, these classes of algebras are quite special, because if the group of units is too small in an algebra, a group identity condition can not limit the structure of the whole algebra. In view of Hartley's conjecture, as a natural generalization the works [5, 6, 3, 10] it is a natural question when does the symmetric units satisfy a group identity in group algebra. Note that the structure theorem of the algebras with involution whose symmetric elements satisfy a polynomial identity was obtained earlier by S.A. Amitsur in [1]. A. Giambruno, S.K. Sehgal and A. Valenti in [8] obtained the following result for group algebras of torsion groups:

**Theorem 1.** *Let  $FG$  be a group algebra of a torsion group  $G$  over an infinite field  $F$  of characteristic  $p > 2$  and assume that the involution  $*$  on  $FG$  is canonical. The symmetric units  $S_*(FG)$  satisfy a group identity if and only if  $G$  has a normal subgroup  $A$  of finite index, the commutator subgroup  $A'$  is a finite  $p$ -group and one of the following conditions holds:*

(i)  $G$  has no quaternion subgroup of order 8 and  $G'$  has of bounded exponent  $p^k$  for some  $k$ .

(ii)  $G$  has of bounded exponent  $4p^s$  for some  $s \geq 0$ , the  $p$ -Sylow subgroup of  $G$  is normal and  $G/P$  is a Hamiltonian 2-subgroup.

In the present paper we extend the result of A. Giambruno, S.K. Sehgal and A. Valenti. For non-torsion groups  $G$  we describe the group algebras  $FG$  over the field  $F$  of characteristic different from 2 whose symmetric units

$$S_*(FG) = \{u = \sum_{g \in G} \alpha_g g \in U(FG) \mid u = u^* = \sum_{g \in G} \alpha_g g^{-1}\}$$

satisfy a group identity. The present result was announced at the International Workshop Polynomial Identities in Algebras, 2002, Memorial University of Newfoundland.

## 2. MAIN RESULTS

In the sequel of this paper  $\mathfrak{d}(\omega)$  denotes a positive integer, which depends on the group identity  $\omega$  and it is defined in the next section. Our results are the following:

**Theorem 2.** *Let  $G$  be a non-torsion nonabelian group and  $\text{char}(F) = p \neq 2$  and assume that the symmetric units of  $FG$  satisfy some group identity  $\omega = 1$ . Assume that  $|F| > \mathfrak{d}(\omega)$ , where  $\mathfrak{d}(\omega)$  is an integer which depends only on the word  $\omega$ . Let  $P$  be a  $p$ -Sylow subgroup of  $G$  and let  $t(G)$  be the torsion part of  $G$ .*

(I) *If  $p > 2$  then  $P$  and  $t(G)$  are normal subgroups of  $G$  such that:*

- (a)  $B = t(G)/P$  is an abelian  $p'$ -subgroup and its subgroups are normal in  $G$ ;
  - (b) if  $B$  is noncentral in  $G/P$  then the algebraic closure  $L$  of the prime subfield  $F_p$  in  $F$  is finite and for all  $g \in G/P$  and for any  $a \in B$  there exists an  $r \in \mathbb{N}$  such that  $a^g = a^{p^r}$  and  $|L : F_p|$  is a divisor of  $r$ ;
  - (c) the  $p$ -Sylow subgroup  $P$  is a finite group;
  - (d) the  $p$ -Sylow subgroup  $P$  is infinite and  $G$  has a subgroup  $A$  of finite index, such that  $A'$  is a finite  $p$ -group and the commutator subgroup  $H'$  of  $H = AP$  is a bounded  $p$ -group. Moreover, if  $P$  is unbounded, then  $G'$  is a bounded  $p$ -group;
- (II) If  $\text{char}(F) = 0$  then  $t(G)$  is a subgroup, every subgroup of  $t(G)$  is normal in  $G$  and one of the following conditions holds:
- (a)  $t(G)$  is abelian and each idempotent of  $Ft(G)$  is central in  $FG$ ;
  - (b)  $t(G)$  is a Hamiltonian 2-group, and each symmetric idempotent of  $Ft(G)$  is central in  $FG$ .

### 3. NOTATION, PRELIMINARY RESULTS AND THE PROOF

Let  $FG$  be the group algebra of  $G$  over  $F$ . We introduce the following notation:

- $(g, h) = g^{-1}g^h = g^{-1}h^{-1}gh$  for all  $g, h \in G$ ;
- $|g|$  and  $C_G(g)$  are the order and the centralizer of  $g \in G$ , respectively;
- $G'$ ,  $\text{Syl}_p(G)$  are the commutator subgroup and the Sylow  $p$ -subgroup of  $G$ ;
- $t(G)$  is the set of elements of finite order in  $G$ ;
- $\Delta(G) = \{g \in G \mid [G : C_G(g)] < \infty\}$  is the  $FC$ -radical of  $G$ ;
- $\Delta^p(G) = \langle g \in \Delta(G) \mid g \text{ has order of a power of } p \rangle$ ;
- $T_l(G/H)$  is a left transversal of the subgroup  $H$  in  $G$ ;
- $\mathfrak{N}(FG)$  is the sum of all nilpotent ideals of the group algebra  $FG$ ;
- $A(FG)$  is the augmentation ideal of the group algebra  $FG$ .

Let  $A$  be an algebra over a field  $F$ , let  $F_0$  be the ring of integers of the field  $F$ , and suppose that  $U(A)$  satisfies a group identity  $\omega = 1$ . Then, as it was proved in Lemma 3.1 of [11], there exists a polynomial  $f(x)$  over  $F_0$  of degree  $\mathfrak{d}(\omega)$  which is determined by the word  $\omega$ . In several papers (see for example [8]) the authors assumed that the field  $F$  is infinite so they could apply the ‘‘Vandermonde determinant argument’’. We shall use some lemmas from [8], which are easy to prove using the method of the paper [11] even without the assumption that the field  $F$  is infinite.

In our proof we will use the following facts:

**Lemma 1.** ([1]) *Let  $A$  be an algebra with involution over  $F$  of  $\text{char}(F) \neq 2$ , such that the set of symmetric units of  $A$  satisfy a group identity  $\omega = 1$ . If  $I$  is a stable nil ideal of  $A$  then the symmetric units of  $A/I$  satisfy a group identity.*

**Lemma 2.** (see [8]) *Let  $A$  be an algebra over the field  $F$  of characteristic  $p \neq 2$ , such that the set of symmetric units of  $A$  satisfy a group identity  $\omega = 1$  and  $|F| > \mathfrak{d}(\omega)$ , where  $\mathfrak{d}(\omega)$  is an integer which depends only on the word  $\omega$ . Then:*

- (i) *if  $A$  is semiprime, then  $asa = 0$  for every nilpotent element  $s \in S_*(A)$  and square-zero  $a \in S_*(A)$ ;*
- (ii) *if  $a \in A$  is square-zero, then  $(aa^*)^m = 0$ , for some  $m \in \mathbb{N}$ ;*
- (iii) *if  $A$  is semiprime and  $u, v \in A$  such that  $uv = 0$ , then  $usv = 0$  for any square-zero symmetric element  $s$ ;*
- (iv) *if the subring  $L$  of  $A$  is nil, then  $L$  satisfy a polynomial identity;*
- (v) *each symmetric idempotent is central;*
- (vi) *if  $A$  is artinian, then  $A$  is isomorphic to a direct sum of division algebras and  $2 \times 2$  matrices algebras over a field with symplectic involution. Each nilpotent element of  $A$  has index at most 2;*
- (vii) *if  $A = FG$  is the group algebra of the group  $G = Q_8 \times \langle c \rangle$ , where  $Q_8$  is the quaternion group of order 8, then the order of the cyclic subgroup  $\langle c \rangle$  is finite.*

**Lemma 3.** (see [8]) *Let  $A$  be a normal abelian subgroup of  $G$  of finite index such that  $G = A \cdot H$ , where  $H$  is a finite group. Let  $\text{char}(F) = p$  and assume that the set of symmetric units of  $FG$  satisfy a group identity  $\omega = 1$ . If  $|F| > \mathfrak{d}(\omega)$ , where  $\mathfrak{d}(\omega)$  is an integer which depends only on the word  $\omega$ , then  $G'$  has bounded exponent  $p^m$ , where  $m$  depends only on  $\mathfrak{d}$ .*

Now we are ready to prove the following

**Lemma 4.** *Let  $\text{char}(F) = p > 2$  and let the set of symmetric units of  $FG$  satisfy a group identity  $\omega = 1$ . Assume that  $|F| > \mathfrak{d}(\omega)$ , where  $\mathfrak{d}(\omega)$  is an integer which depends only on the word  $\omega$ . Then the  $p$ -Sylow subgroup  $P$  of  $\Delta(G)$  is normal in  $G$  and the set of symmetric units of  $F[G/P]$  satisfy a group identity.*

*Proof.* Let  $H$  be a finite subgroup of  $\Delta(G)$  and let  $J = J(F_p H)$  be the radical of the finite group algebra  $F_p H$  over the prime subfield  $F_p$ . According to Lemma 2(vi), the factor algebra  $F_p H/J$  is isomorphic to a direct sum of fields and  $2 \times 2$  matrices algebras over a finite field with symplectic involution and a nilpotent element  $\bar{u} = u + J \in F_p H/J$  has index at most 2. Moreover, from this decomposition follows that  $\bar{u}\bar{u}^*$  is central. By Lemma 2(ii) the element  $\bar{u}\bar{u}^*$  is nilpotent and central in the semiprime algebra  $F_p H/J$ . Therefore  $\bar{u}\bar{u}^* = 0$  and  $uu^* \in J(FH)$ .

Let  $h \in H$  with  $|h| = p^t$ . Then  $u = h - 1$  is nilpotent and

$$uu^* = (h - 1)(h^{-1} - 1) \in J(FH).$$

It follows that  $h u u^* = -(h - 1)^2 \in J(FH)$ . Using Passman's result (see Lemma 5 in [8], p.453) we obtain that  $h - 1 \in J(FH)$  for all  $h \in H$  and  $H \cap (1 + J)$  is a normal  $p$ -subgroup of  $H$ , which coincides with the  $p$ -Sylow

subgroup of  $H$ . Thus the  $p$ -Sylow subgroup  $P$  of  $\Delta(G)$  is normal in  $G$ , so the proof is complete.  $\square$

**Lemma 5.** *Let  $FG$  be a semiprime group algebra over the field  $F$  with  $\text{char}(F) > 2$  such that the set of symmetric units of  $FG$  satisfy a group identity  $\omega = 1$ . Suppose that  $|F| > \mathfrak{d}(\omega)$ , where  $\mathfrak{d}(\omega)$  is an integer which depends only on the word  $\omega$ . Then one of the following conditions holds:*

- (i)  $t(G)$  is abelian and each idempotent of  $Ft(G)$  is central in  $FG$ .
- (ii)  $t(G)$  is a Hamiltonian 2-group and each symmetric idempotent of  $Ft(G)$  is central in  $FG$ .

*Proof.* (i) Let  $a \in t(G)$ , such that  $(|a|, p) = 1$ . Then, by Lemma 2(v), the symmetric idempotent  $e = \frac{1}{n}(1 + a + \dots + a^{|a|-1})$  is central in  $FG$ , so  $\langle a \rangle$  is normal in  $G$ . Now let  $p > 2$  and let  $a \in t(G)$  be of order  $p$ . If  $N_G(\langle a \rangle) = G$  then  $\overline{\langle a \rangle}$  is a central nilpotent element of the semiprime algebra  $FG$ , a contradiction.

Let us prove that each torsion element belongs to  $N_G(\langle a \rangle)$ . Pick  $h \notin N_G(\langle a \rangle)$  such that  $|h| = p^t$ . The elements  $(h - 1)(h^{-1} - 1)$  and  $\langle a \rangle$  are symmetric and  $(2 - h - h^{-1})^{p^t} = (\overline{\langle a \rangle})^2 = 0$ . By Lemma 2(i) we get  $\overline{\langle a \rangle}(2 - h - h^{-1})\overline{\langle a \rangle} = 0$  and

$$(1) \quad \overline{\langle a \rangle}h\overline{\langle a \rangle} + \overline{\langle a \rangle}h^{-1}\overline{\langle a \rangle} = 0.$$

An element of  $\text{Supp}(\overline{\langle a \rangle}h\overline{\langle a \rangle})$  can be written as  $a^i h a^j$ , where  $0 \leq i, j \leq p-1$ . If all the elements in  $\text{Supp}(\overline{\langle a \rangle}h\overline{\langle a \rangle})$  and in  $\text{Supp}(\overline{\langle a \rangle}h^{-1}\overline{\langle a \rangle})$  are distinct, then on the left-hand side of (1) each element appears at most two times, but this leads to a contradiction if  $\text{char}(F) \neq 2$ . Therefore, in the subset  $\text{Supp}(\overline{\langle a \rangle}h\overline{\langle a \rangle})$  not all elements are different, whence there exist  $i, j, k, l$  such that  $a^i h a^j = a^k h a^l$  and either  $i \neq k$  or  $j \neq l$ . If, for example,  $i > k$ , then  $h^{-1} a^{i-k} h = a^{l-j}$  and  $h \in N_G(\langle a \rangle)$ .

Now, let  $h \notin N_G(\langle a \rangle)$  be a  $p'$ -element. As we have seen before,  $\langle h \rangle$  is normal in  $G$ , so  $\langle a, h \rangle$  is a finite subgroup. By Lemma 4 the  $p$ -Sylow subgroup  $P$  of  $\langle a, h \rangle$  is normal in  $\langle a, h \rangle$  and  $(a, h) \in P \cap \langle h \rangle = \langle 1 \rangle$ , a contradiction.

Therefore, each element of finite order belongs to  $N_G(\langle a \rangle)$ . Moreover, the elements of order  $p$  in  $G$  form an elementary abelian normal  $p$ -subgroup  $E$  of  $G$ .

Finally, if  $h \notin N_G(\langle a \rangle)$ , then  $h$  has infinite order and  $h$  acts on  $E$ . The subgroups  $\langle a^h \rangle$  and  $\langle a \rangle$  are different and we can choose a subgroup  $\langle b \rangle \subset E$ , which differs from  $\langle a \rangle$ . Clearly,  $\overline{\langle a \rangle}(h + h^{-1})\overline{\langle a \rangle}$  and  $\overline{\langle b \rangle}$  are square-zero symmetric elements and according to Lemma 2(i),

$$(2) \quad \overline{\langle b \rangle}\overline{\langle a \rangle}(h + h^{-1})\overline{\langle a \rangle}\overline{\langle b \rangle} = 0.$$

Since  $hE$  and  $h^{-1}E$  are different cosets, from (2) follows that

$$(3) \quad \overline{\langle b \rangle}\overline{\langle a \rangle}h\overline{\langle a \rangle}\overline{\langle b \rangle} = 0.$$

The subgroup  $H = \langle a, b \rangle \subset E$  has order  $p^2$  and by (3) we have  $\overline{H}h_1\overline{H}h_2 = 0$  for all  $h_1, h_2$ . Since elements of finite order belong to  $N_G(H)$ , we get  $(\overline{H}FG)^2 = 0$ , which is impossible by the semiprimeness of  $FG$ . Thus  $G$  have no  $p$ -elements and all finite cyclic subgroups of  $G$  are normal in  $G$ . Applying Lemmas 6 and 7 from [8] and the fact that  $G$  have no  $p$ -elements ( $p \neq 2$ ), we obtain that  $t(G)$  is either an abelian group or a Hamiltonian 2-group.

Let  $t(G)$  be an abelian group and let  $e \in Ft(G)$  be a noncentral idempotent in  $FG$ . Set  $H = \langle Supp(e) \rangle$ . Since every subgroup of  $t(G)$  is normal in  $G$ , the subgroup  $H$  is also normal in  $G$  and  $FH$  has a primitive idempotent  $f$ , which does not commute with some  $g \in G$  of infinite order. Then  $g^{-1}fg \neq f$  is also a primitive idempotent of  $FH$  and  $(g^{-1}fg)f = 0$ , i.e.  $(fg)^2 = (gf)^2 = 0$ .

Let  $g^{-1}fg = \bar{f} \neq f^*$ . By Lemma 2(v) we have  $f \neq f^*$ , so  $g^{-1}f + f^*g$  is a square-zero symmetric element and by Lemma 2(iii), we get that

$$fg(g^{-1}f + f^*g)fg = 0.$$

It follows that  $f + g(\bar{f}f^*)gf = f = 0$ , a contradiction. Therefore,  $g^{-1}fg = f^*$ , so  $(f^*)^* = (g^{-1}fg)^* = g^{-1}f^*g = f$ . Furthermore,  $g^{-2}fg^2 = g^{-1}f^*g = f$  and  $f^*g^2 = g^2f^*$ . Since  $f^*g^2 = g^2f^*$ ,  $(gf^*)^2 = 0$  and  $gf + f^*g^{-1}$  is square-zero symmetric element, by Lemma 2(iii) we obtain that

$$gf^*(gf + f^*g^{-1})gf^* = gf^*g^2(g^{-1}fg)f^* + gf^* = gf^*g^2f^* + gf^* = 0.$$

Thus  $(g^2 + 1)f^* = 0$ , which is impossible, since  $g^2H$  and  $H$  are different cosets.  $\square$

**Lemma 6.** *Let  $F$  be a field of characteristic  $p$ , and suppose that  $G$  contains a normal locally finite  $p$ -subgroup  $P$  such that the centralizer of each element of  $P$  in every finitely generated subgroup of  $G$  is of finite index. Then  $\mathfrak{J}(P)$  is a locally nilpotent ideal.*

*Proof.* Clearly,  $\{ u(h-1) \mid u \in T_l(G/P), 1 \neq h \in P \}$  is an  $F$ -basis for the ideal  $\mathfrak{J}(P)$ . Let us show that the subalgebra  $W = \langle u_1(h_1-1), \dots, u_s(h_s-1) \rangle_F$  is nilpotent. According to our assumption, the centralizers of  $h_1, \dots, h_s$  in the subgroup  $H = \langle u_1, \dots, u_s, h_1, \dots, h_s \rangle$  have finite index. Since  $P$  is normal, its subgroup  $L = \langle h_1^u, h_2^u, \dots, h_s^u \mid u \in H \rangle$  is a finitely generated FC-group and by a Theorem of B.H. Neumann ([1], Theorem 4, p.19)  $L$  is a finite  $p$ -group. Thus the augmentation ideal  $A(FL)$  is nilpotent with index, say,  $t$ . Furthermore,  $A(FL) = u^{-1}A(FL)u$  for any  $u \in H$  and this implies that  $(A(FL) \cdot FH)^n = A^n(FL) \cdot FH$  for any  $n > 0$ , so  $W^t \subseteq A^t(FL) \cdot FH = 0$ , because  $W \subseteq A(FL) \cdot FH$ . Therefore  $W$  is a nilpotent subalgebra and  $\mathfrak{J}(P)$  is a locally nilpotent ideal.  $\square$

**Lemma 7.** *Let  $G$  be a group with a nontrivial  $p$ -Sylow subgroup  $P$  and let  $\text{char}(F) = p > 2$ . If the set of symmetric units of  $FG$  satisfy a group identity  $\omega = 1$  and  $|F| > \mathfrak{d}(\omega)$ , where  $\mathfrak{d}(\omega)$  is an integer which depends only on the word  $\omega$ , then  $P$  is normal in  $G$  and the ideal  $\mathfrak{J}(P)$  is nil.*

*Proof.* Let  $P$  be a maximal normal  $p$ -subgroup of  $G$  such that the ideal  $\mathfrak{J}(P)$  is nil. By Lemma 1 the set of symmetric units of  $F[G/P]$  satisfy a group identity. If  $F[G/P]$  is semiprime, then by (i) of the Theorem the group  $G/P$  has no  $p$ -elements and  $P$  coincides with the  $p$ -Sylow subgroup of  $G$ . Now, suppose that  $F[G/P]$  is not semiprime. According to Theorem 4.2.13 ([13], p.131) the group  $\Delta(G/P)$  has a nontrivial  $p$ -Sylow subgroup  $P_1/P$ , which is normal in  $G/P$  by Lemma 4. Clearly,  $P_1/P$  is an  $FC$ -subgroup of  $G/P$ , so by Lemma 6 the ideal  $\mathfrak{J}(P_1/P)$  is nil.

Since  $\mathfrak{J}(P_1/P) \cong \mathfrak{J}(P_1)/\mathfrak{J}(P)$  and  $P_1$  is normal in  $G$ , the ideal  $\mathfrak{J}(P_1)$  is nil and  $P \subset P_1$ , a contradiction. Thus  $P = Syl_p(G)$  and the proof is done.  $\square$

**Lemma 8.** *Let  $R$  be an algebra with involution  $*$  over a field  $F$  of characteristic  $p > 2$  and assume that  $S_*(R)$  satisfies a group identity and  $|F| > \mathfrak{d}(\omega)$ . If some nil subring  $L$  of  $R$  is  $*$ -stable, then  $L$  satisfies a non-matrix polynomial identity.*

*Proof.* Let  $A = F\langle X \rangle[[t]]$  be the ring of power series over the polynomial ring  $F\langle X \rangle$  with noncommuting indeterminates  $X = \{x_1, x_2\}$ . By a result of Magnus, the elements  $1 + x_1t, 1 + x_2t$  are units in  $A$  and  $\langle 1 + x_1t, 1 + x_2t \rangle$  is a free group.

Assume that  $S_*(R)$  satisfies the group identity  $w$ , where  $w$  is a reduced word in 2 variables. Then  $w(1 + x_1t, 1 + x_2t) \neq 1$  according to result of Magnus and it is well-known that  $(1 + x_it)^{-1} = 1 - x_it + x_i^2t^2 - \dots$ . If we substitute  $(1 + x_it)^{-1}$  in the expression  $w(1 + x_1t, 1 + x_2t) - 1$ , then it can be expanded as

$$(4) \quad \sum_{i \geq s} g_i(x_1, x_2)t^i,$$

where  $g_i(x_1, x_2) \in F\langle X \rangle$  is a homogeneous polynomial of degree  $i$ . Obviously there exists a smallest integer  $s \geq 1$  such that  $g_s(x_1, x_2) \neq 0$ .

Let  $L$  be a  $*$ -stable nil subring and let  $S(L)$  be the set of the symmetric elements of  $L$ . Take now  $r_1, r_2 \in S(L)$  and let  $\lambda \in F$ . Obviously,  $r_1, r_2$  are nilpotent elements, so each  $1 + \lambda r_i$  is a symmetric unit in  $R$  and

$$(1 + r_i\lambda)^{-1} = 1 - r_i\lambda + r_i^2\lambda^2 + \dots + (-1)^{t-1}r_i^{t-1}\lambda^{t-1}$$

for a suitable  $t$ . By evaluating  $w$  on these elements, (4) gives us a finite sum  $\sum_{i \geq s}^l g_i(r_1, r_2)\lambda^i = 0$  for some  $l$ . Since  $|F| > \mathfrak{d}(\omega)$ , we can apply the Vandermonde determinant argument to obtain  $g_i(r_1, r_2) = 0$  for all  $i$ . Therefore  $g_s(x_1, x_2)$  is a  $*$ -polynomial identity on  $S(L)$ . Finally, by [1] it follows that  $S(L)$  satisfies an ordinary polynomial identity.

Suppose that the homogeneous polynomial  $g(x_1, x_2)$  vanishes on the matrix algebra  $M_2(K)$  over a commutative ring  $K$ . Then

$$g(x_1, x_2) = h(x_1, x_2) + g_{11}(x_1, x_2) + g_{12}(x_1, x_2) + g_{21}(x_1, x_2) + g_{22}(x_1, x_2),$$

where  $h(x_1, x_2)$  consists of all monomials which contain  $x_1^2$  or  $x_2^2$  while the  $g_{ij}(x_1, x_2)$  contain all the remaining monomials beginning with  $x_i$  and ending with  $x_j$  for  $i, j \in \{1, 2\}$ . If  $a$  and  $b$  are two square-zero matrices, then  $h(a, b) = 0$ , because each term of  $h$  has  $a^2$  or  $b^2$  as a factor, so we conclude

that  $ag_{21}(a, b)b = 0$ . Clearly  $x_1g_{21}(x_1, x_2)x_2$  is some polynomial  $f(x_1x_2)$ . Then  $f(ab\lambda) = 0$  for each  $\lambda \in F$  and, by the Vandermonde determinant argument, we get  $(ab)^d = 0$  for some  $d$ . Take, for instance, the matrix units  $a = e_{12}$  and  $b = e_{21}$ , then we obtain a contradiction.  $\square$

**Lemma 9.** *Let  $R$  be an algebra over a field  $F$  of positive characteristic  $p$  satisfying a non-matrix polynomial identity. Then  $R$  satisfies also a polynomial identity of the form  $([x, y]z)^{p^l}$  and  $[x, y]^{p^l}$*

*Proof.* Let  $g(x_1, x_2, \dots, x_n)$  be a non-matrix polynomial identity in  $R$ . The variety  $W$  determined by the polynomial identity  $g(x_1, x_2, \dots, x_n)$  contains a relatively free algebra  $K$  of rank 3. Of course,  $K$  is a finitely generated *PI*-algebra, and the result of Braun and Razmyslov (Theorem 6.3.39, [14]) states that the radical  $J(K)$  of  $K$  is nilpotent. Writing  $K/J(K)$  as a subdirect sum of primitive rings  $\{L_i\}$ , we get that every primitive ring  $L_i$  satisfies the non-matrix polynomial identity  $g(x_1, x_2, \dots, x_n)$ , as a homomorphic image of  $K$ . By Theorem 2.1.4 of [9],  $L_i$  is either isomorphic to the matrix ring  $M_m(D)$  over a division ring  $D$ , or for any  $m$  the matrix ring  $M_m(D)$  is an epimorphic image of some subring of  $L_i$ .

Thus  $M_m(D)$  satisfies a non-matrix polynomial identity  $g$ , which is possible only if  $L_i$  is a commutative ring. Consequently,  $K/J(K)$  is a commutative algebra, so  $K$  satisfies a polynomial identity of the form  $([x, y]z)^{p^l}$  such that  $J(K)^{p^l} = 0$ . Since  $R$  belongs to the variety  $W$ , the algebra  $R$  also satisfies a polynomial identity  $([x, y]z)^{p^l}$ .  $\square$

**Lemma 10.** *Let  $FG$  be a non semiprime group algebra over the field  $F$  with  $\text{char}(F) > 2$ , such that the set of symmetric units of  $FG$  satisfy a group identity  $\omega = 1$  and  $|F| > \mathfrak{d}(\omega)$ , where  $\mathfrak{d}(\omega)$  is an integer which depends only on the word  $\omega$ . If  $\mathfrak{N}(FG)$  is not nilpotent then  $FG$  is a *PI*-algebra, where  $\mathfrak{N}(FG)$  is the sum of all nilpotent ideals of  $FG$ .*

*Proof.* Clearly the non nilpotent ideal  $\mathfrak{N} = \mathfrak{N}(FG)$  is invariant under the involution  $*$  and by Lemma 2(iv) the ring  $\mathfrak{N}$  satisfies a polynomial identity  $f(x_1, \dots, x_n)$ . Moreover, by Lemma 2.8 of [12] the algebra  $FG$  satisfies a non-degenerate multilinear generalized polynomial identity and hence, by Theorem 5.3.15 ([13], p.202),  $|G : \Delta(G)| < \infty$  and  $\Delta(G)'$  is finite.

Set  $P = \text{Syl}_p(G)$  and  $P_1 = \text{Syl}_p(\Delta(G)')$ . By Lemma 4,  $P \cap \Delta(G)' = P_1 \triangleleft G$  and  $P_1$  is a finite  $p$ -group. Thus  $\mathfrak{I}(P_1)$  is a nilpotent ideal and by (i) of the Theorem, the set of symmetric units of  $F[\Delta(G)'/P_1]$  satisfy a group identity, so  $\Delta(G)'/P_1$  is either an abelian  $p'$ -group or a Hamiltonian 2-group.

If  $P_1 = \Delta(G)'$ , then by Theorem 5.3.9 ([13], p.197) the algebra  $FG$  is a *PI*-algebra. If  $P_1 \subsetneq \Delta(G)'$  then we can suppose that  $G$  is a group such that  $\text{Syl}_p(\Delta(G)') = 1$  and  $\Delta(G)'$  is either an abelian  $p'$ -group or a Hamiltonian 2-group.

Set  $P_2 = \text{Syl}_p(\Delta(G))$ . Clearly,  $P_2 = P \cap \Delta(G)$  is normal in  $\Delta(G)$ . Since  $[P : P_2] < \infty$  and  $P$  is an infinite group, the group  $P_2$  is infinite, too. If



$a \in P_2, b \in \Delta(G)$ , then  $(a, b) \in P_2 \cap \Delta(G)' = 1$ , so  $(a, b) = 1$  and  $P_2$  is a central subgroup in  $\Delta(G)$ .

Let us prove that  $F\Delta(G)$  is a  $PI$ -algebra. If  $\Delta(G)$  is a torsion group, then by [8] the statement is trivial.

Since  $\mathfrak{N}(F\Delta(G)) \subseteq \mathfrak{N}(FG)$ , the ideal  $\mathfrak{N}(F\Delta(G))$  also satisfies the same polynomial identity  $f(x_1, \dots, x_n)$ . By the standard multilinearization process, we may assume that  $f(x_1, \dots, x_n)$  is multilinear.

Assume that  $P_2$  has bounded exponent. Then the maximal elementary abelian  $p$ -subgroup  $E$  of  $P_2$  is infinite. Let  $f(a_1, \dots, a_n) = \sum_i \alpha_i y_i$ , where  $a_1, \dots, a_n \in F\Delta(G), y_1, \dots, y_n \in T_1(\Delta(G)/E)$  and  $\alpha_i \in FE$ . Then there exists a finite subgroup  $B$  such that  $\alpha_i \in FB$  and  $E = B \times \prod_j \langle c_j \rangle$ . Since  $(c_k - 1)a_k \in \mathfrak{N}(F\Delta(G))$  and  $P_2$  is central, we conclude that

$$f((c_1 - 1)a_1, \dots, (c_n - 1)a_n) = (c_1 - 1) \cdots (c_n - 1)f(a_1, \dots, a_n) = 0.$$

It follows that  $f(a_1, \dots, a_n) = 0$ , because  $B \cap \prod_j \langle c_j \rangle = \langle 1 \rangle$ .

Now let  $P_2$  be of unbounded exponent and  $c \in P_2$ . Then  $(c - 1)a_k \in \mathfrak{N}(F\Delta(G))$  and also

$$f((c - 1)a_1, \dots, (c - 1)a_n) = (c - 1)^n f(a_1, \dots, a_n) = 0$$

for all  $c \in P_2$ . Then  $f(a_1, \dots, a_n)$  belongs to the annihilator of the augmentation ideal  $A(FP_2^{p^t})$ , where  $n \leq p^t$ . Since  $P_2^{p^t}$  is infinite, we have

$$Ann_l(A(FP_2^{p^t})) = 0.$$

It follows that  $f(a_1, \dots, a_n) = 0$ , so  $f(x_1, \dots, x_n)$  is a polynomial identity for  $F\Delta(G)$ . Since  $F\Delta(G)$  is a  $PI$ -algebra and  $[G : \Delta(G)] < \infty$ , the algebra  $FG$  is  $PI$ , too.  $\square$

*Proof of the theorem.* Let  $FG$  be a group algebra of a non-torsion group  $G$  over a field of positive characteristic  $p$ . By Lemma 7 the  $p$ -Sylow subgroup  $P$  is normal in  $G$  and  $F[G/P] \cong FG/\mathfrak{I}(P)$ , so the symmetric units of semiprime algebra  $F[G/P]$  satisfy a group identity. By Lemma 5  $B = t(G/P)$  is a subgroup of  $G/P$  and  $B$  is either an abelian  $p'$ -group or a Hamiltonian 2-group. If  $B$  is a Hamiltonian 2-group, then  $Q_8$  is a subgroup of  $B$ . Choose an element  $c \in G/P$  of infinite order. Since every subgroup of  $t(G)/P$  is normal in  $G/P$  and  $|Aut(Q_8)| < \infty$ , there exists a  $t \in \mathbb{N}$  such that  $c^t \in C_{G/P}(Q_8)$  and  $Q_8 \times \langle c^t \rangle \subseteq G/P$ . Then Lemma 2(vii) asserts that  $c$  has finite order, a contradiction. So  $B$  is an abelian  $p'$ -group and by Lemma 5 every idempotent of  $FB$  is central in  $F[G/P]$ . Moreover, if  $B$  is noncentral, then according to [7] the group  $B$  satisfy (i.b) of our Theorem.

Now, let  $P$  be infinite. By Corollary 8.1.14 ([13], p.312) the ideal  $\mathfrak{N}(FG)$  is non-nilpotent, so by Lemma 10, the algebra  $FG$  is a  $PI$ -algebra, i.e.  $G$  has a subgroup  $A$  with finite index such that  $A'$  is a finite  $p$ -group. According to Lemma 1, it can be assumed that  $G$  has an abelian subgroup  $A$  of finite index.

We claim that the commutator subgroup of  $H = P \cdot A$  is a bounded  $p$ -group. Clearly  $S_*(FP)$  satisfies a group identity and according to Lemma 3  $P'$  is a bounded  $p$ -group. The normal abelian  $p$ -subgroup  $P' \cap A$  has finite exponent and according to Lemma 6 the ideal  $\mathfrak{J}(P' \cap A)$  is locally nilpotent of bounded degree. The subgroup  $P' \cap A$  of  $P'$  has finite index in  $P$  and

$$\mathfrak{J}(P')/\mathfrak{J}(P' \cap A) \cong \mathfrak{J}(P'/(P' \cap A)).$$

Therefore  $\mathfrak{J}(P')$  is a locally nilpotent ideal of bounded degree  $p^t$  for some  $t$ . Clearly  $FG/\mathfrak{J}(P') \cong F[G/P']$  and put  $P' = \langle 1 \rangle$ . Since  $A$  has a finite index in  $H = P \cdot A$ , Lemma 3 ensures that  $H'$  is a  $p$ -group of bounded exponent and according to Lemma 1, we can put  $H' = \langle 1 \rangle$  again.

The  $p$ -Sylow subgroup  $P$  of  $G$  is abelian and by Lemma 8 the ideal  $\mathfrak{J}(P)$  satisfies a non-matrix polynomial identity, Moreover, by Lemma 9 the ideal  $\mathfrak{J}(P)$  satisfies polynomial identities of the following forms:  $[x, y]^{p^l}$  and  $([x, y]z)^{p^l}$ .

Let  $h \in G$  and  $a \in P$ . Clearly  $(a - 1)h, h^{-1}(a^{-1} - 1) \in \mathfrak{J}(P)$  and

$$[(a - 1)h, h^{-1}(a^{-1} - 1)]^{p^l} = (a^h)^{p^l} + (a^h)^{-p^l} - a^{p^l} - a^{-p^l} = 0$$

which implies that either  $(h, a)^{p^l} = 1$  or  $h^{-1}a^{p^l}h = a^{-p^l}$ .

Put  $z = a^{p^l}$ . From  $h^{-1}a^{p^l}h = a^{-p^l}$  it follows that  $h^{-1}zh = z^{-1}$  and  $([z - 1, (z^{-1} - 1)h])^{p^l} = 0$ . Clearly  $[z - 1, (z^{-1} - 1)h] = -z^{-2}(z + 1)(z - 1)^2h$  so

$$\begin{aligned} 0 &= ([z - 1, (z^{-1} - 1)h])^{p^l} \\ &= -((z + 1)(z - 1)^2(z^{-1} + 1)(z^{-1} - 1)^2h^2)^{\frac{p^l - 1}{2}} (z^{-2}(z + 1)(z - 1)^2h) \\ &= -z^{\frac{-3p^l - 1}{2}} \cdot (z + 1)^{p^l} \cdot (z - 1)^{2p^l} \cdot h^{p^l}. \end{aligned}$$

Since  $char(K) > 2$ , the element  $z + 1$  is a unit and  $(z - 1)^{2p^l} = (a - 1)^{2p^{2l}} = 0$  and the order of  $a$  at most  $2p^{2l}$ . Therefore  $(h, a)^{p^{2l+1}} = 1$  for all  $h \in G, a \in P$  and  $2l + 1$  depends on only the group identity. Since  $(G, P)$  is a  $p$ -group of bounded exponent, we can again make a reduction, so we can assumed that  $(G, P) = 1$  and  $P$  is central.

Let  $P$  be a central subgroup of unbounded exponent and  $h_1, h_2 \in G$ . Obviously

$$\begin{aligned} ((h_1, h_2)^{p^l} - 1)(a - 1)^{p^{3l}} &= ((h_1, h_2) - 1)^{p^l}(a - 1)^{p^{3l}} \\ &= ([h_1^{-1}(a - 1), h_2^{-1}(a - 1)]h_1h_2(a - 1))^{p^l} = 0 \end{aligned}$$

for  $a \in P$ . Since there are infinitely many element of the form  $a^{p^{3l}}$  we conclude that  $(h_1, h_2)^{p^l} = 1$  and the proof is complete.  $\square$

REFERENCES

[1] S. A. Amitsur. Rings with involution. *Israel J. Math.*, 6:99–106, 1968.  
 [2] A. A. Bovdi. *Gruppovye koltsa*. Užgorod. Gosudarstv. Univ., Uzhgorod, 1974.

- [3] V. Bovdi. On symmetric units in group algebras. *Comm. Algebra*, 29(12):5411–5422, 2001.
- [4] V. Bovdi and L. G. Kovács. Unitary units in modular group algebras. *Manuscripta Math.*, 84(1):57–72, 1994.
- [5] V. Bovdi, L. G. Kovács, and S. K. Sehgal. Symmetric units in modular group algebras. *Comm. Algebra*, 24(3):803–808, 1996.
- [6] V. Bovdi and M. M. Parmenter. Symmetric units in integral group rings. *Publ. Math. Debrecen*, 50(3-4):369–372, 1997.
- [7] S. P. Coelho. A note on central idempotents in group rings. *Proc. Edinburgh Math. Soc.* (2), 30(1):69–72, 1987.
- [8] A. Giambruno, S. K. Sehgal, and A. Valenti. Symmetric units and group identities. *Manuscripta Math.*, 96(4):443–461, 1998.
- [9] I. N. Herstein. *Noncommutative rings*. The Carus Mathematical Monographs, No. 15. Published by The Mathematical Association of America, 1968.
- [10] C. Lanski. Rings with involution whose symmetric units commute. *Canad. J. Math.*, 28(5):915–928, 1976.
- [11] C.-H. Liu. Group algebras with units satisfying a group identity. *Proc. Amer. Math. Soc.*, 127(2):327–336, 1999.
- [12] C.-H. Liu and D. S. Passman. Group algebras with units satisfying a group identity. II. *Proc. Amer. Math. Soc.*, 127(2):337–341, 1999.
- [13] D. Passman. *The algebraic structure of group rings*. Pure and Applied Mathematics. Wiley-Interscience [John Wiley & Sons], New York, 1977.
- [14] L. H. Rowen. *Ring theory. Vol. I*, volume 127 of *Pure and Applied Mathematics*. Academic Press Inc., Boston, MA, 1988.

*Received May 15, 2005.*

INSTITUTE OF MATHEMATICS,  
UNIVERSITY OF DEBRECEN,  
4010 DEBRECEN, PF. 12,  
HUNGARY

INSTITUTE OF MATHEMATICS AND COMPUTER SCIENCE,  
COLLEGE OF NYÍREGYHÁZA  
SÓSTÓI ÚT 31/B,  
H-4410 NYÍREGYHÁZA, HUNGARY  
*E-mail address:* vbovdi@math.klte.hu