

SOME UNIFYING RESULTS ON STABILITY AND STRONG CONVERGENCE FOR SOME NEW ITERATION PROCESSES

M. O. OLATINWO

ABSTRACT. In this paper, we shall establish some stability results as well as strong convergence results for a pair of nonselfmappings using some newly introduced iteration processes and two general contractive conditions. Our results are improvements, generalizations and extensions of the results in some of the references listed in the reference section of this paper as well as some other analogous ones in the literature.

1. INTRODUCTION

Let (E, d) be a complete metric space and $T : E \rightarrow E$ a selfmap of E . Suppose that $F_T = \{ p \in E \mid Tp = p \}$ is the set of fixed points of T .

There are several iteration processes in the literature for which the fixed points of operators have been approximated over the years by various authors. In a complete metric space, the Picard iteration process $\{x_n\}_{n=0}^\infty$ defined by

$$(1.1) \quad x_{n+1} = Tx_n, \quad n = 0, 1, \dots,$$

has been employed to approximate the fixed points of mappings satisfying the inequality relation

$$(1.2) \quad d(Tx, Ty) \leq \alpha d(x, y), \quad \forall x, y \in E \text{ and } \alpha \in [0, 1).$$

Condition (1.2) is called the *Banach's contraction condition*. Any operator satisfying (1.2) is called *strict contraction*. Also, condition (1.2) is significant in the celebrated Banach's fixed point theorem [3].

In the Banach space setting, we shall state some of the iteration processes generalizing (1.1) as follows:

For $x_0 \in E$, the sequence $\{x_n\}_{n=0}^\infty$ defined by

$$(1.3) \quad x_{n+1} = (1 - \alpha_n)x_n + \alpha_nTx_n, \quad n = 0, 1, \dots,$$

where $\{\alpha_n\}_{n=0}^\infty \subset [0, 1]$, is called the Mann iteration process (see Mann [20]).

2000 *Mathematics Subject Classification*. 47H06, 54H25.

Key words and phrases. Arbitrary Banach space; Jungck-Ishikawa iteration process; nonselfmappings.

For $x_0 \in E$, the sequence $\{x_n\}_{n=0}^\infty$ defined by

$$(1.4) \quad \left. \begin{aligned} x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n Tz_n \\ z_n &= (1 - \beta_n)x_n + \beta_n Tx_n \end{aligned} \right\} n = 0, 1, \dots,$$

where $\{\alpha_n\}_{n=0}^\infty$ and $\{\beta_n\}_{n=0}^\infty$ are sequences in $[0, 1]$, is called the Ishikawa iteration process (see Ishikawa [14]).

The following is the iteration process introduced by Singh et al [39] to establish some stability results: Let S and T be operators on an arbitrary set Y with values in E such that $T(Y) \subseteq S(Y)$. $S(Y)$ is a complete subspace of E . Then, for $x_0 \in Y$, the sequence $\{Sx_n\}_{n=0}^\infty$ defined by

$$(1.5) \quad Sx_{n+1} = (1 - \alpha_n)Sx_n + \alpha_n Tx_n, \quad n = 0, 1, \dots,$$

where $\{\alpha_n\}_{n=0}^\infty$ is a sequence in $[0, 1]$ is called the *Jungck-Mann* iteration process.

If $\alpha_n = 1$ and $Y = E$ in (1.5), then we obtain

$$(1.6) \quad Sx_{n+1} = Tx_n, \quad n = 0, 1, 2, \dots,$$

which is the *Jungck iteration*. See Jungck [16] for detail.

While the iteration process (1.5) extends (1.1), (1.3) and (1.6), the iteration processes (1.4) and (1.5) are independent.

Kannan [17] established an extension of the Banach's fixed point theorem by using the following contractive definition: For a selfmap T , there exists $\beta \in (0, \frac{1}{2})$ such that

$$(1.7) \quad d(Tx, Ty) \leq \beta [d(x, Tx) + d(y, Ty)], \quad \forall x, y \in E.$$

Chatterjea [8] used the following contractive condition: For a selfmap T , there exists $\gamma \in (0, \frac{1}{2})$ such that

$$(1.8) \quad d(Tx, Ty) \leq \gamma [d(x, Ty) + d(y, Tx)], \quad \forall x, y \in E.$$

Zamfirescu [40] established a nice generalization of the Banach's fixed point theorem by combining (1.2), (1.7) and (1.8). That is, for a mapping $T : E \rightarrow E$, there exist real numbers α, β, γ satisfying $0 \leq \alpha < 1$, $0 \leq \beta < \frac{1}{2}$, $0 \leq \gamma < \frac{1}{2}$ respectively such that for each $x, y \in E$, at least one of the following is true:

$$(1.9) \quad \left. \begin{aligned} (z_1) \quad & d(Tx, Ty) \leq \alpha d(x, y) \\ (z_2) \quad & d(Tx, Ty) \leq \beta [d(x, Tx) + d(y, Ty)] \\ (z_3) \quad & d(Tx, Ty) \leq \gamma [d(x, Ty) + d(y, Tx)]. \end{aligned} \right\}$$

The mapping $T : E \rightarrow E$ satisfying (1.9) is called the *Zamfirescu contraction*. Any mapping satisfying condition (z_2) of (1.9) is called a *Kannan mapping*, while the mapping satisfying condition (z_3) is called *Chatterjea operator*. The contractive condition (1.9) implies

$$(1.10) \quad \|Tx - Ty\| \leq 2\delta \|x - Tx\| + \delta \|x - y\|, \quad \forall x, y \in E,$$

where $\delta = \max \left\{ \alpha, \frac{\beta}{1-\beta}, \frac{\gamma}{1-\gamma} \right\}$, $0 \leq \delta < 1$.

Condition (1.9) was used by Rhoades [31, 32] to obtain some convergence results for Mann and Ishikawa iteration processes in a uniformly convex Banach space. The results of [31, 32] were recently extended by Berinde [6] to an arbitrary Banach space for the same fixed point iteration processes. Rafiq [30] proved a convergence result for the Noor iteration process in normed space using the Zamfirescu contraction. See Noor [2] for the Noor iteration process.

Singh et al [39] defined the following general iteration process:

Let $S, T : Y \rightarrow E$ and $T(Y) \subseteq S(Y)$. For any $x_0 \in Y$, let

$$(1.11) \quad Sx_{n+1} = f(T, x_n), \quad n = 0, 1, \dots$$

For $S = I$ (i.e. identity map on E), $f(T, x_n) = Tx_{n+1}$ and $Y = E$, then (1.11) reduces to the well-known Picard iteration process in (1.1).

If $Y = E$, and $f(T, x_n) = Tx_n$, $n = 0, 1, \dots$, then (1.11) reduces to the Jungck iteration process of (1.6). Jungck [16] established that the maps S and T satisfying

$$(1.12) \quad d(Tx, Ty) \leq kd(Sx, Sy), \quad \forall x, y \in E, \quad k \in [0, 1),$$

have a unique common fixed point in complete metric space E , provided that S and T commute, $T(Y) \subseteq S(Y)$ and S is continuous. For results which are similar to Jungck [16] in uniform space, we refer to Aamril and El Moutawakil [1] as well as Olatinwo [21, 22].

The following definition of the stability of iteration process due to Singh et al [39] shall be required in the sequel.

Definition 1.1. Let $S, T : Y \rightarrow E$, $T(Y) \subseteq S(Y)$ and z a coincidence point of S and T , that is, $Sz = Tz = p$ (say). For any $x_0 \in Y$, let the sequence $\{Sx_n\}_{n=0}^{\infty}$, generated by the iteration procedure (1.11) converge to p . Let $\{Sy_n\}_{n=0}^{\infty} \subset E$ be an arbitrary sequence, and set $\epsilon_n = d(Sy_{n+1}, f(T, y_n))$, $n = 0, 1, \dots$. Then, the iteration procedure (1.11) will be called (S, T) -stable if and only if $\lim_{n \rightarrow \infty} \epsilon_n = 0$ implies that $\lim_{n \rightarrow \infty} Sy_n = p$.

This definition reduces to that of the stability of iteration procedure due to Harder and Hicks [11] when $Y = E$ and $S = I$ (identity operator).

Several stability results established in metric space and normed linear space are available in the literature. Some of the various authors whose contributions are of colossal value in the study of stability of the fixed point iteration procedures are Ostrowski [29], Harder and Hicks [11], Rhoades [34, 36], Osilike [27], Osilike and Udomene [28], Jachymski [15], Berinde [5, 4] and Singh et al [39]. Harder and Hicks [11], Rhoades [34, 36], Osilike [27] and Singh et al [39] used the method of the summability theory of infinite matrices to prove various stability results for certain contractive definitions. The method has also been adopted to establish various stability results for certain contractive definitions in Olatinwo et al [24, 26]. Osilike and Udomene [28] introduced a shorter method of proof of stability results and this has also been employed by Berinde [5], Imoru and Olatinwo [12], Imoru et al [13], Olatinwo et al [25]

and some others. In Harder and Hicks [11], the contractive definition stated in (1.2) was used to prove a stability result for the Kirk's iteration process. The first stability result on T -stable mappings was proved by Ostrowski [29] for the Picard iteration using (1.2).

In addition to (1.2), the contractive condition in (1.9) was also employed by Harder and Hicks [11] to establish some stability results for both Picard and Mann iteration processes. Rhoades [34, 36] extended the stability results of [11] to more general classes of contractive mappings. Rhoades [34] extended the results of [11] to the following independent contractive condition: there exists $c \in [0, 1)$ such that

$$(1.13) \quad d(Tx, Ty) \leq c \max \{d(x, y), d(x, Ty), d(y, Tx)\}, \quad \forall x, y \in E.$$

Rhoades [36] used the following contractive definition: there exists $c \in [0, 1)$ such that

$$(1.14) \quad d(Tx, Ty) \leq c \max \left\{ d(x, y), \frac{d(x, Tx) + d(y, Ty)}{2}, d(x, Ty), d(y, Tx) \right\},$$

for all $x, y \in E$. Moreover, Osilike [27] generalized and extended some of the results of Rhoades [36] by using a more general contractive definition than those of Rhoades [34, 36]. Indeed, he employed the following contractive definition: there exist $a \in [0, 1]$, $L \geq 0$ such that

$$(1.15) \quad d(Tx, Ty) \leq Ld(x, Tx) + ad(x, y), \quad \forall x, y \in E.$$

Osilike and Udomene [28] introduced a shorter method to prove stability results for the various iteration processes using the condition (1.15). Berinde [5] established the same stability results for the same iteration processes using the same set of contractive definitions as in Harder and Hicks [11] but the same method of shorter proof as in Osilike and Udomene [28].

More recently, Imoru and Olatinwo [12] established some stability results which are generalizations of some of the results of Osilike [5, 11, 27, 28, 34, 36]. In Imoru and Olatinwo [12], the following contractive definition was employed: there exist

$a \in [0, 1)$ and a monotone increasing function $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ with $\varphi(0) = 0$, such that

$$(1.16) \quad d(Tx, Ty) \leq \varphi(d(x, Tx)) + ad(x, y), \quad \forall x, y \in E.$$

Condition (1.16) was also employed in Olatinwo et al [24] to establish some stability results in normed linear space setting with additional condition of continuity imposed on φ .

However, Singh et al [39] established some stability results for Jungck and Jungck-Mann iteration processes by employing two contractive definitions both of which generalize those of Osilike [27] but independent of that of Imoru and Olatinwo [12]. Singh et al [39] obtained stability results for Jungck and Jungck-Mann iterative procedures in metric space using both the contractive definition

(1.12) and the following: For $S, T : Y \rightarrow E$ and some $k \in [0, 1)$, we have

$$(1.17) \quad d(Tx, Ty) \leq kd(Sx, Sy) + Ld(Sx, Tx), \quad \forall x, y \in Y.$$

In the next section, we shall introduce the Jungck-Ishikawa iteration process to prove some stability and convergence results for nonselfmappings in normed linear space and arbitrary Banach space respectively. In establishing our results, more general contractive conditions than (1.9) will be considered.

2. PRELIMINARIES

We shall consider the following iteration processes in establishing our results:

Let $(E, \|\cdot\|)$ be a Banach space and Y an arbitrary set. Let $S, T : Y \rightarrow E$ be two nonselfmappings such that $T(Y) \subseteq S(Y)$, $S(Y)$ is a complete subspace of E and S is injective. Then, for $x_0 \in Y$, define the sequence $\{Sx_n\}_{n=0}^\infty$ iteratively by

$$(2.1) \quad \left. \begin{aligned} Sx_{n+1} &= (1 - \alpha_n)Sx_n + \alpha_n Tz_n \\ Sz_n &= (1 - \beta_n)Sx_n + \beta_n Tx_n \end{aligned} \right\}, \quad n = 0, 1, \dots,$$

where $\{\alpha_n\}_{n=0}^\infty$ and $\{\beta_n\}_{n=0}^\infty$ are sequences in $[0, 1]$. The iteration process (2.1) is called the *Jungck-Ishikawa* iteration process. See Olatinwo [23] for more detail.

The iteration processes (1.1) and (1.3)–(1.6) are special cases of (2.1). For instance,

if in (2.1), S is identity operator, $Y = E$, $\beta_n = 0$ then we obtain the Mann iteration process of (1.3).

Since S is injective, if $\beta_n = 0$, then for $x_0 \in Y$, (2.1) reduces to the Jungck-Mann iteration process of (1.5).

Also, with S and T as in (2.1), we define the following three-step iteration process which is an extension of (2.1):

For $x_0 \in Y$ and with S and T as above, define the sequence $\{Sx_n\}_{n=0}^\infty$ iteratively by

$$(2.2) \quad \left. \begin{aligned} Sx_{n+1} &= (1 - \alpha_n)Sx_n + \alpha_n Tz_n \\ Sz_n &= (1 - \beta_n)Sx_n + \beta_n Tr_n \\ Sr_n &= (1 - \gamma_n)Sx_n + \gamma_n Tx_n \end{aligned} \right\}, \quad n = 0, 1, \dots,$$

where $\{\alpha_n\}_{n=0}^\infty$, $\{\beta_n\}_{n=0}^\infty$ and $\{\gamma_n\}_{n=0}^\infty$ are sequences in $[0, 1]$. The iteration process (2.2) will be called the *Jungck-Noor* iteration process.

The iteration processes (1.1) and (1.3)–(1.6) are also special cases of (2.2). In fact, the iteration process defined in (2.2) is an extension of that of Noor [2].

Definition 2.1 (Berinde [7]). A function $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is called a *comparison function* if it satisfies the following conditions:

- (i) ψ is monotone increasing;
- (ii) $\lim_{n \rightarrow \infty} \psi^n(t) = 0, \quad \forall t \geq 0.$

Remark 2.2. Every comparison function satisfies $\psi(0) = 0$. See Rus [37] and Rus et al [38] for the Definition 2.1.

In addition to the iteration process (2.1), we shall employ the following contractive definitions:

Definition 2.3. For two nonselfmappings $S, T : Y \rightarrow E$ with $T(Y) \subseteq S(Y)$, where $S(Y)$ is a complete subspace of E , there exist:

- (a) a real number $L \geq 0$, a sublinear comparison function $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ and a monotone increasing function $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $\varphi(0) = 0$ and $\forall x, y \in Y$, we have

$$(2.3) \quad \|Tx - Ty\| \leq \frac{\varphi(\|Sx - Tx\|) + \psi(\|Sx - Sy\|)}{1 + L\|Sx - Tx\|};$$

and,

- (b) real numbers $k \geq 0$, $L \geq 0$, $a \in [0, 1)$ and a monotone increasing function $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $\varphi(0) = 0$ and $\forall x, y \in Y$, we have

$$(2.4) \quad \|Tx - Ty\| \leq \left(\frac{\varphi(\|Sx - Tx\|) + a\|Sx - Sy\|}{1 + L\|Sx - Tx\|} \right) e^{k\|Sx - Tx\|}.$$

In this paper, we shall consider the iteration processes defined in (2.1) and (2.2) to establish some stability results for nonselfmappings in normed linear space as well as obtain some strong convergence results for these nonselfmappings in an arbitrary Banach space by employing the contractive conditions (2.3) and (2.4). Our stability results are generalizations and extensions of those of Singh et al [39], some results of [5, 12, 13, 24, 26, 34, 36], while the convergence results extend, generalize and improve those of [6, 18, 19, 36, 31]. For more on the study of fixed point iteration processes and various contractive conditions, our interested readers can consult Berinde [4], Ciric [9, 10], Rhoades [33] and others in the reference section of this paper.

Definition 2.4. Let X and Y be two nonempty sets and $S, T : X \rightarrow Y$ two mappings. Then, an element $x^* \in X$ is a *coincidence point* of S and T if and only if $Sx^* = Tx^*$.

Denote the set of the coincidence points of S and F by $C(S, T)$.

There are several papers and monographs on the coincidence point theory. However, we refer our readers to Rus [37] and Rus et al [38] for the Definition 2.4 and some coincidence point results.

We shall require the following lemmas in the sequel.

Lemma 2.5 (Berinde [5, 4, 7]). *If δ is a real number such that $0 \leq \delta < 1$, and $\{\epsilon_n\}_{n=0}^{\infty}$ is a sequence of positive numbers such that $\lim_{n \rightarrow \infty} \epsilon_n = 0$, then for any sequence of positive numbers $\{u_n\}_{n=0}^{\infty}$ satisfying*

$$u_{n+1} \leq \delta u_n + \epsilon_n, \quad n = 0, 1, \dots,$$

we have $\lim_{n \rightarrow \infty} u_n = 0$.

Lemma 2.6 (Imoru et al [13]). *Let $\psi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a comparison function and $\{v_n\}_{n=0}^\infty$ a sequence of positive numbers such that $\lim_{n \rightarrow \infty} v_n = 0$. Then, we have*

$$\lim_{n \rightarrow \infty} \sum_{k=0}^n \psi^{n-k}(v_k) = 0, \text{ for each } k.$$

Proof. Since ψ is monotone increasing, there exists a convergent series $\sum a_{nk}$ of positive numbers a_{nk} , $k = 0, 1, 2, \dots, n$ such that

$$\psi^{n-k}(v_k) = a_{nk}v_k.$$

Therefore,

$$\sum_{k=0}^n \psi^{n-k}(v_k) = \sum_{k=0}^n a_{nk}v_k.$$

Let A be the lower triangular matrix with entries a_{nk} , $k = 0, 1, \dots, n$. Clearly, $\lim_{n \rightarrow \infty} a_{nk} = 0$, for each k . Since $\sum a_{nk}$ is convergent, let $\lim_{n \rightarrow \infty} \sum a_{nk} = s < \infty$. Therefore, A is multiplicative (See Rhoades [36]). Then,

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{k=0}^n \psi^{n-k}(v_k) &= \lim_{n \rightarrow \infty} \sum_{k=0}^n a_{nk}v_k = \lim_{n \rightarrow \infty} \sum_{k=0}^n a_{nk} \cdot \lim_{n \rightarrow \infty} v_n \\ &= \lim_{n \rightarrow \infty} \sum_{k=0}^n a_{nk} \lim_{n \rightarrow \infty} v_n = 0. \end{aligned}$$

□

Lemma 2.7 (Imoru et al [13]). *Let $\{\psi^k(t)\}_{k=0}^n$ be a sequence of comparison functions. Then, any linear combination $\sum_{j=0}^n c_j \psi^j(t)$ of the comparison functions is also a comparison function, where $\sum_{j=0}^n c_j = 1$ and c_0, c_1, \dots, c_n are positive constants.*

Proof. Let $\bar{\psi}(t) = c_0\psi^0(t) + c_1\psi^1(t) + c_2\psi^2(t) + \dots + c_n\psi^n(t)$. Since each $\psi^k(t)$, $k = 0, 1, \dots, n$ is a comparison function, then each $\psi^k(t)$, $k = 0, 1, \dots, n$ is monotone increasing. Also, since each $c_k > 0$, $k = 0, 1, \dots, n$, then each $c_k\psi^k(t)$ is monotone increasing, from which it follows that $\bar{\psi}(t)$ is monotone increasing. Moreover, since $\psi^k(t) \rightarrow 0$, $\forall t \geq 0$, $k = 0, 1, \dots, n$ then $c_k\psi^k(t) \rightarrow 0$, $\forall k = 0, 1, \dots, n$. Therefore, $\bar{\psi}(t) \rightarrow 0$, $\forall t \geq 0$. Hence, $\bar{\psi}(t)$ is a comparison function. □

Lemma 2.8 (Imoru et al [13]). *If $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ is a subadditive comparison function and $\{\epsilon_n\}_{n=0}^\infty$ is a sequence of positive numbers such that $\lim_{n \rightarrow \infty} \epsilon_n = 0$, then for any sequence of positive numbers $\{u_n\}_{n=0}^\infty$ satisfying*

$$(2.5) \quad u_{n+1} \leq \sum_{k=0}^m \delta_k \psi^k(u_n) + \epsilon_n, \quad n = 0, 1, \dots,$$

where $\delta_0, \delta_1, \dots, \delta_m \in [0, 1]$ with $0 \leq \sum_{k=0}^m \delta_k \leq 1$, and we have $\lim_{n \rightarrow \infty} u_n = 0$.

Proof. Let $\bar{\psi}(u_n) = \sum_{k=0}^m \delta_k \psi^k(u_n)$. Then, inequality (2.5) becomes

$$(2.6) \quad u_{n+1} \leq \bar{\psi}(u_n) + \epsilon_n, \quad n = 0, 1, 2, \dots,$$

Using Lemma 2.7, then we have that $\bar{\psi}(u_n)$ is a comparison function. Also, using (2.6) yields

$$\begin{aligned} u_1 &\leq \bar{\psi}(u_0) + \epsilon_0 \\ u_2 &\leq \bar{\psi}(u_1) + \epsilon_1 \leq \bar{\psi}(\bar{\psi}(u_0) + \epsilon_0) + \epsilon_1 = \bar{\psi}^2(u_0) + \bar{\psi}(\epsilon_0) + \epsilon_1 \\ u_3 &\leq \bar{\psi}(u_2) + \epsilon_2 \leq \bar{\psi}^3(u_0) + \bar{\psi}^2(\epsilon_0) + \bar{\psi}(\epsilon_1) + \epsilon_2, \end{aligned}$$

and in general, we have

$$(2.7) \quad u_n \leq \bar{\psi}^n(u_0) + \bar{\psi}^{n-1}(\epsilon_0) + \bar{\psi}^{n-2}(\epsilon_1) + \dots + \bar{\psi}(\epsilon_{n-2}) + \epsilon_{n-1}$$

Replacing n by $(n + 1)$ in (2.7) yields

$$(2.8) \quad \begin{aligned} u_{n+1} &\leq \bar{\psi}^{n+1}(u_0) + \bar{\psi}^n(\epsilon_0) + \bar{\psi}^{n-1}(\epsilon_1) + \dots + \bar{\psi}(\epsilon_{n-1}) + \epsilon_n \\ &= \bar{\psi}^{n+1}(u_0) + \sum_{k=0}^n \bar{\psi}^{n-k}(\epsilon_k). \end{aligned}$$

Since $\bar{\psi}$ is a comparison function, then $\bar{\psi}^{n+1}(u_0) \rightarrow 0$ as $n \rightarrow \infty$. Using Lemma 2.6, then we have that

$$\lim_{n \rightarrow \infty} \sum_{k=0}^n \bar{\psi}^{n-k}(\epsilon_k) = 0 \text{ for each } k.$$

Thus, inequality (2.8) yields $\lim_{n \rightarrow \infty} u_n = 0$. □

We establish our main results in the next two sections. Our stability results are established by using the method of Berinde [5] and Osilike and Udomene [28]. Section 3 deals with some stability results in normed linear space, while a strong convergence result is proved in section 4.

3. SOME STABILITY RESULTS IN NORMED LINEAR SPACE

Theorem 3.1. *Let $(E, \|\cdot\|)$ be a normed space and Y an arbitrary set. Suppose that $S, T : Y \rightarrow E$ are nonselfoperators such that $T(Y) \subseteq S(Y)$, $S(Y)$ a complete subspace of E , and S is an injective operator. Let z be a coincidence point of S and T (that is, $Sz = Tz = p$). Suppose that S and T satisfy condition (2.3). Let $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a continuous sublinear comparison function and $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ a monotone increasing function such that $\varphi(0) = 0$. For $x_0 \in Y$, let $\{Sx_n\}_{n=0}^\infty$ be the Jungck-Ishikawa iteration process defined by (2.1) converging to p , where $\{\alpha_n\}_{n=0}^\infty$ and $\{\beta_n\}_{n=0}^\infty$ are sequences in $[0, 1]$. Then, the Jungck-Ishikawa iteration process is (S, T) -stable.*

Proof. Suppose that $\{Sy_n\}_{n=0}^\infty \subset E$, $\epsilon_n = \|Sy_{n+1} - (1 - \alpha_n)Sy_n - \alpha_n Tb_n\|$, $n = 0, 1, \dots$, where $Sb_n = (1 - \beta_n)Sy_n + \beta_n Ty_n$ and let $\lim_{n \rightarrow \infty} \epsilon_n = 0$. Then,

we shall establish that $\lim_{n \rightarrow \infty} Sy_n = p$, using the contractive condition and the triangle inequality:

$$\begin{aligned}
(3.1) \quad \|Sy_{n+1} - p\| &\leq \epsilon_n + \|(1 - \alpha_n)(Sy_n - p) + \alpha_n(Tb_n - p)\| \\
&\leq \epsilon_n + (1 - \alpha_n)\|Sy_n - p\| + \alpha_n\|Tz - Tb_n\| \\
&\leq \epsilon_n + (1 - \alpha_n)\|Sy_n - p\| \\
&\quad + \alpha_n \left[\frac{\varphi(\|Sz - Tz\|) + \psi(\|Sz - Sb_n\|)}{1 + M\|Sz - Tz\|} \right] \\
&= (1 - \alpha_n)\|Sy_n - p\| + \alpha_n\psi(\|p - Sb_n\|) + \epsilon_n \\
&\leq (1 - \alpha_n)\|Sy_n - p\| + \alpha_n[(1 - \beta_n)\psi(\|p - Sy_n\|) \\
&\quad + \beta_n\psi(\|Tz - Ty_n\|)] + \epsilon_n \\
&\leq (1 - \alpha_n)\|Sy_n - p\| + \alpha_n(1 - \beta_n)\psi(\|Sy_n - p\|) \\
&\quad + \alpha_n\beta_n\psi^2(\|p - Sy_n\|) + \epsilon_n.
\end{aligned}$$

Using Lemma 2.8 in (3.1) yields $\lim_{n \rightarrow \infty} \|Sy_n - p\| = 0$, that is, $\lim_{n \rightarrow \infty} Sy_n = p$.

Conversely, let $\lim_{n \rightarrow \infty} Sy_n = p$. Then, by using the triangle inequality and the contractive definition, we have the following:

$$\begin{aligned}
\epsilon_n &= \|Sy_{n+1} - (1 - \alpha_n)Sy_n - \alpha_nTb_n\| \\
&\leq \|Sy_{n+1} - p\| + (1 - \alpha_n)\|Sy_n - p\| + \alpha_n\|Tz - Tb_n\| \\
&\leq \|Sy_{n+1} - p\| + (1 - \alpha_n)\|Sy_n - p\| + \alpha_n(1 - \beta_n)\psi(\|p - Sy_n\|) \\
&\quad + \alpha_n\beta_n\psi(\|Tz - Ty_n\|) \\
&\leq \|Sy_{n+1} - p\| + (1 - \alpha_n)\|Sy_n - p\| + \alpha_n(1 - \beta_n)\psi(\|Sy_n - p\|) \\
&\quad + \alpha_n\beta_n\psi^2(\|Sy_n - p\|) \rightarrow 0 \quad \text{as } n \rightarrow \infty.
\end{aligned}$$

□

Theorem 3.2. *Let $(E, \|\cdot\|)$ be a normed space and Y an arbitrary set. Suppose that $S, T : Y \rightarrow E$ are nonselfoperators such that $T(Y) \subseteq S(Y)$, $S(Y)$ a complete subspace of E , and S is an injective operator. Let z be a coincidence point of S and T (that is, $Sz = Tz = p$). Suppose that S and T satisfy condition (2.4). Let $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a monotone increasing function such that $\varphi(0) = 0$. For $x_0 \in Y$, let $\{Sx_n\}_{n=0}^\infty$ be the Jungck-Noor iteration process defined by (2.2) converging to p , where $\{\alpha_n\}_{n=0}^\infty$, $\{\beta_n\}_{n=0}^\infty$ and $\{\gamma_n\}_{n=0}^\infty$ are sequences in $[0, 1]$ such that $0 < \alpha \leq \alpha_n$, $0 < \beta \leq \beta_n$ and $0 < \gamma \leq \gamma_n$, ($n = 0, 1, \dots$). Then, the Jungck-Noor iteration process is (S, T) -stable.*

Proof. Suppose that $\{Sy_n\}_{n=0}^\infty \subset E$, $\epsilon_n = \|Sy_{n+1} - (1 - \alpha_n)Sy_n - \alpha_nTb_n\|$, $n = 0, 1, \dots$, where $Sb_n = (1 - \beta_n)Sy_n + \beta_nTc_n$, $S_c_n = (1 - \gamma_n)Sy_n + \gamma_nTy_n$ and let $\lim_{n \rightarrow \infty} \epsilon_n = 0$. Then, we shall establish that $\lim_{n \rightarrow \infty} Sy_n = p$, using the

contractive condition (2.4) and the triangle inequality:

$$\begin{aligned}
\|Sy_{n+1} - p\| &\leq (1 - \alpha_n)\|Sy_n - p\| + \alpha_n\|Tz - Tb_n\| + \epsilon_n \\
&\leq (1 - \alpha_n)\|Sy_n - p\| + a\alpha_n\|p - Sb_n\| + \epsilon_n \\
&\leq (1 - \alpha_n)\|Sy_n - p\| + a\alpha_n[(1 - \beta_n)\|Sy_n - p\| \\
&\quad + \beta_n\|Tz - Tc_n\|] + \epsilon_n \\
&= [1 - (1 - a)\alpha_n - a\alpha_n\beta_n]\|Sy_n - p\| + a\alpha_n\beta_n\|Tz - Tc_n\| + \epsilon_n \\
&\leq [1 - (1 - a)\alpha_n - a\alpha_n\beta_n]\|Sy_n - p\| + a^2\alpha_n\beta_n\|Sz - Sc_n\| + \epsilon_n \\
&= [1 - (1 - a)\alpha_n - (1 - a)a\alpha_n\beta_n \\
&\quad - (1 - a)a^2\alpha_n\beta_n\gamma_n]\|Sy_n - p\| + \epsilon_n \\
(3.2) \quad &\leq [1 - (1 - a)\alpha - (1 - a)a\alpha\beta - (1 - a)a^2\alpha\beta\gamma]\|Sy_n - p\| + \epsilon_n.
\end{aligned}$$

Since

$$0 \leq 1 - (1 - a)\alpha - (1 - a)a\alpha\beta - (1 - a)a^2\alpha\beta\gamma < 1 \text{ and } \lim_{n \rightarrow \infty} \epsilon_n = 0,$$

using Lemma 2.5 in (3.2) yields $\lim_{n \rightarrow \infty} \|Sx_n - p\| = 0$, that is, $\lim_{n \rightarrow \infty} Sx_n = p$.

Conversely, let $\lim_{n \rightarrow \infty} Sx_{n+1} = p$. Then, by using the triangle inequality and the contractive definition, we have the following:

$$\begin{aligned}
\epsilon_n &\leq \|Sy_{n+1} - p\| + (1 - \alpha_n)\|Sy_n - p\| + \alpha_n\|Tz - Tb_n\| \\
&\leq \|Sy_{n+1} - p\| + (1 - \alpha_n)\|Sy_n - p\| + a\alpha_n\|Sz - Sb_n\| \\
&\leq \|Sy_{n+1} - p\| + [1 - (1 - a)\alpha_n - a\alpha_n\beta_n]\|Sy_n - p\| + a\alpha_n\beta_n\|Tz - Tc_n\| \\
&\leq \|Sy_{n+1} - p\| + [1 - (1 - a)\alpha_n - a\alpha_n\beta_n]\|Sy_n - p\| + a^2\alpha_n\beta_n\|Sz - Sc_n\| \\
&\leq \|Sy_{n+1} - p\| + [1 - (1 - a)\alpha_n - a\alpha_n\beta_n]\|Sy_n - p\| \\
&\quad + a^2\alpha_n\beta_n[(1 - \gamma_n)\|p - Sy_n\| + \gamma_n\|p - Ty_n\|] \\
&\leq \|Sy_{n+1} - p\| + [1 - (1 - a)\alpha - (1 - a)a\alpha\beta \\
&\quad - (1 - a)a^2\alpha\beta\gamma]\|Sy_n - p\| \rightarrow 0 \text{ as } n \rightarrow \infty.
\end{aligned}$$

□

Remark 3.3. Both Theorem 3.1 and Theorem 3.2 are generalizations and extensions of Theorem 3.5 of Singh et al [39], Theorem 3 of Berinde [5], Theorem 2 of Osilike [27], Theorem 2 and Theorem 5 of Osilike and Udomene [28], Theorem 2 of Rhoades [34], Theorem 30 of Rhoades [35], Theorem 2 of Rhoades [36], Theorem 3 of Harder and Hicks [11] as well as some of the results of the author [12, 13, 24, 26, 25]. Our stability results also extend some similar ones in Berinde [7] and Olatinwo [23].

4. SOME CONVERGENCE RESULTS IN ARBITRARY BANACH SPACE

Theorem 4.1. *Let $(E, \|\cdot\|)$ be an arbitrary Banach space and Y is an arbitrary set. Suppose that $S, T : Y \rightarrow E$ are nonselfoperators such that $T(Y) \subseteq S(Y)$, $S(Y)$ a complete subspace of E , and S is an injective operator. Let z*

be a coincidence point of S and T (that is, $Sz = Tz = p$). Suppose that S and T satisfy condition (2.4). Let $\varphi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be a monotone increasing function such that $\varphi(0) = 0$. For $x_0 \in Y$, let $\{Sx_n\}_{n=0}^\infty$ be the Jungck-Noor iteration process defined by (2.2), where $\{\alpha_n\}_{n=0}^\infty$, $\{\beta_n\}_{n=0}^\infty$ and $\{\gamma_n\}_{n=0}^\infty$ are sequences in $[0, 1]$ such that $\sum_{n=0}^\infty \alpha_n = \infty$. Then, $\{Sx_n\}_{n=0}^\infty$ converges strongly to p .

Proof. Let $C(S, T)$ be the set of the coincidence points of S and T . We shall now use condition (2.4) to establish that S and T have a unique coincidence point z (i.e. $Sz = Tz = p$ (say)): Injectivity of S is sufficient.

Suppose that there exist $z_1, z_2 \in C(S, T)$ such that $Sz_1 = Tz_1 = p_1$ and $Sz_2 = Tz_2 = p_2$.

If $p_1 = p_2$, then $Sz_1 = Sz_2$ and since S is injective, it follows that $z_1 = z_2$.

If $p_1 \neq p_2$, then we have by the contractiveness condition (2.4) for S and T that

$$\begin{aligned} 0 &< \|p_1 - p_2\| = \|Tz_1 - Tz_2\| \\ &\leq \left(\frac{\varphi(\|Sz_1 - Tz_1\|) + a\|Sz_1 - Sz_2\|}{1 + L\|Sz_1 - Tz_1\|} \right) e^{k\|Sz_1 - Tz_1\|} \\ &\leq a\|p_1 - p_2\|, \end{aligned}$$

which leads to $(1 - a)\|p_1 - p_2\| \leq 0$, from which it follows that $1 - a > 0$ since $a \in [0, 1)$, but $\|p_1 - p_2\| \leq 0$, which is a contradiction since norm is nonnegative.

Therefore, we have that $\|p_1 - p_2\| = 0$, that is, $p_1 = p_2 = p$. Since $p_1 = p_2$, then we have that $p_1 = Sz_1 = Tz_1 = Sz_2 = Tz_2 = p_2$, leading to $Sz_1 = Sz_2 \Rightarrow z_1 = z_2 = z$ (since S is injective).

Hence, $z \in C(S, T)$, that is, z is a unique coincidence point of S and T .

Now, we prove that $\{Sx_n\}_{n=0}^\infty$ converges strongly to p (where $Sz = Tz = p$) using again, condition (2.4). Therefore, we have

$$\begin{aligned} \|Sx_{n+1} - p\| &\leq (1 - \alpha_n)\|Sx_n - p\| + \alpha_n\|Tz - Tz_n\| \\ (4.1) \quad &\leq (1 - \alpha_n)\|Sx_n - p\| + a\alpha_n\|p - Sz_n\|. \end{aligned}$$

Now, we have that

$$\begin{aligned} \|p - Sz_n\| &= \|(1 - \beta_n)(p - Sx_n) + \beta_n(p - Ty_n)\| \\ (4.2) \quad &\leq (1 - \beta_n)\|Sx_n - p\| + a\beta_n\|p - Sy_n\|. \end{aligned}$$

Using (4.2) in (4.1) yields

$$(4.3) \quad \|Sx_{n+1} - p\| \leq [1 - (1 - a)\alpha_n - a\alpha_n\beta_n]\|Sx_n - p\| + a^2\alpha_n\beta_n\|p - Sy_n\|.$$

Furthermore, we have

$$\begin{aligned} \|p - Sy_n\| &\leq (1 - \gamma_n)\|Sx_n - p\| + \gamma_n\|p - Tx_n\| \\ (4.4) \quad &\leq (1 - \gamma_n + a\gamma_n)\|p - Sx_n\|. \end{aligned}$$

Using (4.4) in (4.3) yields

$$\begin{aligned}
 \|Sx_{n+1} - p\| &\leq [1 - (1 - a)\alpha_n - (1 - a)a\alpha_n\beta_n - (1 - a)a^2\alpha_n\beta_n\gamma_n]\|Sx_n - p\| \\
 &\leq [1 - (1 - a)\alpha_n]\|Sx_n - p\| \\
 &\leq \prod_{j=0}^n [1 - (1 - a)\alpha_j]\|Sx_0 - p\| \\
 (4.5) \quad &\leq e^{-(1-a)\sum_{j=0}^n \alpha_j}\|Sx_0 - p\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.
 \end{aligned}$$

Therefore, we obtain from (4.5) that $\lim_{n \rightarrow \infty} \|Sx_{n+1} - p\| = 0$, i.e. $\{Sx_n\}_{n=0}^{\infty}$ converges strongly to p . \square

Remark 4.2. Theorem 4.1 is a generalization and extension of a multitude of results. In particular, Theorem 4.1 is a generalization and extension of both Theorem 1 and Theorem 2 of Berinde [6], Theorem 3 of Rafiq [30], Theorem 2 and Theorem 3 of Kannan [18], Theorem 3 of Kannan [19], Theorem 4 of Rhoades [31] as well as Theorem 8 of Rhoades [32]. Also, both Theorem 4 of Rhoades [31] and Theorem 8 of Rhoades [32] are Theorem 4.10 and Theorem 5.6 of Berinde [4] respectively. Our result also extends some similar ones in Berinde [7] and Olatinwo [23].

Remark 4.3. We have considered two new iteration processes to prove some unifying theorems for stability and convergence. These new iteration processes as well as the results obtained extend the frontiers of knowledge in the fixed point theory.

REFERENCES

- [1] M. Aamri and D. El Moutawakil. Common fixed point theorems for E -contractive or E -expansive maps in uniform spaces. *Acta Math. Acad. Paedagog. Nyházi. (N.S.)*, 20(1):83–91 (electronic), 2004.
- [2] M. Aslam Noor. Three-step iterative algorithms for multivalued quasi variational inclusions. *J. Math. Anal. Appl.*, 255(2):589–604, 2001.
- [3] S. Banach. Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *Fundamenta math.*, 3:133–181, 1922.
- [4] V. Berinde. *Iterative approximation of fixed points*. Editura Efemeride, Baia Mare, 2002.
- [5] V. Berinde. On the stability of some fixed point procedures. *Bul. Ştiinţ. Univ. Baia Mare Ser. B Fasc. Mat.-Inform.*, 18(1):7–14, 2002. Dedicated to Costică Mustăţa on his 60th anniversary.
- [6] V. Berinde. On the convergence of the Ishikawa iteration in the class of quasi contractive operators. *Acta Math. Univ. Comenian. (N.S.)*, 73(1):119–126, 2004.
- [7] V. Berinde. *Iterative approximation of fixed points*, volume 1912 of *Lecture Notes in Mathematics*. Springer, Berlin, second edition, 2007.
- [8] S. K. Chatterjea. Fixed-point theorems. *C. R. Acad. Bulgare Sci.*, 25:727–730, 1972.
- [9] L. B. Ćirić. Generalized contractions and fixed-point theorems. *Publ. Inst. Math. (Beograd) (N.S.)*, 12(26):19–26, 1971.
- [10] L. B. Ćirić. A generalization of Banach's contraction principle. *Proc. Amer. Math. Soc.*, 45:267–273, 1974.
- [11] A. M. Harder and T. L. Hicks. Stability results for fixed point iteration procedures. *Math. Japon.*, 33(5):693–706, 1988.

- [12] C. O. Imoru and M. O. Olatinwo. On the stability of Picard and Mann iteration processes. *Carpathian J. Math.*, 19(2):155–160, 2003.
- [13] C. O. Imoru, M. O. Olatinwo, and O. O. Owojori. On the stability results for Picard and Mann iteration procedures. *J. Appl. Funct. Differ. Equ. JAFDE*, 1(1):71–80, 2006.
- [14] S. Ishikawa. Fixed points by a new iteration method. *Proc. Amer. Math. Soc.*, 44:147–150, 1974.
- [15] J. R. Jachymski. An extension of A. Ostrowski's theorem on the round-off stability of iterations. *Aequationes Math.*, 53(3):242–253, 1997.
- [16] G. Jungck. Commuting mappings and fixed points. *Amer. Math. Monthly*, 83(4):261–263, 1976.
- [17] R. Kannan. Some results on fixed points. *Bull. Calcutta Math. Soc.*, 60:71–76, 1968.
- [18] R. Kannan. Some results on fixed points. III. *Fund. Math.*, 70(2):169–177, 1971.
- [19] R. Kannan. Construction of fixed points of a class of nonlinear mappings. *J. Math. Anal. Appl.*, 41:430–438, 1973.
- [20] W. R. Mann. Mean value methods in iteration. *Proc. Amer. Math. Soc.*, 4:506–510, 1953.
- [21] M. O. Olatinwo. Some common fixed point theorems for selfmappings in uniform space. *Acta Math. Acad. Paedagog. Nyházi. (N.S.)*, 23(1):47–54 (electronic), 2007.
- [22] M. O. Olatinwo. Some existence and uniqueness common fixed point theorems for self-mappings in uniform space. *Fasc. Math.*, (38):87–95, 2007.
- [23] M. O. Olatinwo. Some stability and strong convergence results for the Jungck-Ishikawa iteration process. *Creat. Math. Inform.*, 17:33–42, 2008.
- [24] M. O. Olatinwo, O. O. Owojori, and C. O. Imoru. On some stability results for fixed point iteration procedure. *J. Math. Stat.*, 2(1):339–342, 2006.
- [25] M. O. Olatinwo, O. O. Owojori, and C. O. Imoru. Some stability results for fixed point iteration processes. *Aust. J. Math. Anal. Appl.*, 3(2):Art. 8, 7 pp. (electronic), 2006.
- [26] M. O. Olatinwo, O. O. Owojori, and C. O. Imoru. Some stability results on Krasnolslseskij and Ishikawa fixed point iteration procedures. *J. Math. Stat.*, 2(1):360–362, 2006.
- [27] M. O. Osilike. Stability results for fixed point iteration procedures. *J. Nigerian Math. Soc.*, 14/15:17–29, 1995/96.
- [28] M. O. Osilike and A. Udomene. Short proofs of stability results for fixed point iteration procedures for a class of contractive-type mappings. *Indian J. Pure Appl. Math.*, 30(12):1229–1234, 1999.
- [29] A. M. Ostrowski. The round-off stability of iterations. *Z. Angew. Math. Mech.*, 47:77–81, 1967.
- [30] A. Rafiq. On the convergence of the three-step iteration process in the class of quasi-contractive operators. *Acta Math. Acad. Paedagog. Nyházi. (N.S.)*, 22(3):305–309 (electronic), 2006.
- [31] B. E. Rhoades. Fixed point iterations using infinite matrices. *Trans. Amer. Math. Soc.*, 196:161–176, 1974.
- [32] B. E. Rhoades. Comments on two fixed point iteration methods. *J. Math. Anal. Appl.*, 56(3):741–750, 1976.
- [33] B. E. Rhoades. A comparison of various definitions of contractive mappings. *Trans. Amer. Math. Soc.*, 226:257–290, 1977.
- [34] B. E. Rhoades. Fixed point theorems and stability results for fixed point iteration procedures. *Indian J. Pure Appl. Math.*, 21(1):1–9, 1990.
- [35] B. E. Rhoades. Some fixed point iteration procedures. *Internat. J. Math. Math. Sci.*, 14(1):1–16, 1991.
- [36] B. E. Rhoades. Fixed point theorems and stability results for fixed point iteration procedures. II. *Indian J. Pure Appl. Math.*, 24(11):691–703, 1993.

- [37] I. A. Rus. *Generalized contractions and applications*. Cluj University Press, Cluj-Napoca, 2001.
- [38] I. A. Rus, A. Petruşel, and G. Petruşel. *Fixed point theory: 1950–2000. Romanian contributions*. House of the Book of Science, Cluj-Napoca, 2002.
- [39] S. L. Singh, C. Bhatnagar, and S. N. Mishra. Stability of Jungck-type iterative procedures. *Int. J. Math. Math. Sci.*, (19):3035–3043, 2005.
- [40] T. Zamfirescu. Fix point theorems in metric spaces. *Arch. Math. (Basel)*, 23:292–298, 1972.
- [41] E. Zeidler. *Nonlinear functional analysis and its applications. I*. Springer-Verlag, New York, 1986. Fixed-point theorems, Translated from the German by Peter R. Wadsack.

Received March 16, 2008.

DEPARTMENT OF MATHEMATICS,
OBAFEMI AWOLOWO UNIVERSITY,
ILE-IFE,
NIGERIA.
E-mail address: polatinwo@oauife.edu.ng
E-mail address: molaposi@yahoo.com