

A NOTE ON MULTIPLICATION OPERATORS ON KÖTHE-BOCHNER SPACES

S. S. KHURANA

ABSTRACT. Let $(\Omega, \mathcal{A}, \mu)$ is a finite measure space, E an order continuous Banach function space over μ , X a Banach space and $E(X)$ the Köthe-Bochner space. A new simple proof is given of the result that a continuous linear operator $T: E(X) \rightarrow E(X)$ is a multiplication operator (by a function in L^∞) iff $T(g\langle f, x^* \rangle x) = g\langle T(f), x^* \rangle x$ for every $g \in L^\infty$, $f \in E(X)$, $x \in X$, $x^* \in X^*$.

1. INTRODUCTION AND NOTATION

In this paper all vector spaces are taken over the real field R . $(\Omega, \mathcal{A}, \mu)$ is a finite measure space and $L^\infty(\mu) = L^\infty$, $L^1(\mu) = L^1$ have their usual meanings. E is an ideal in the vector lattice L^1 , $E \supset L^\infty$, and has the norm $\|\cdot\|_E$ so that $(E, \|\cdot\|_E)$ is a Banach lattice and is called Köthe function space relative to the measure μ ([3]). The order in E is the natural order of functions in L^1 . Also the inclusions $L^\infty \subset E \subset L^1$ are continuous. $(X, \|\cdot\|_X)$ is another Banach space such that the Banach space $(E(X), \|\cdot\|_{E(X)})$ is the associated Köthe-Bochner function space relative to E . Thus $E(X)$ consists of all strongly measurable functions $f: \Omega \rightarrow X$ for which the real functions $\omega \rightarrow \|f(\omega)\|$ belongs to E and $\|f\|_{E(X)} = \|\|f(\cdot)\|_X\|_E$ ([3]). For measure theory we refer to [1]. If Y is a Banach space, Y^* will denote its dual and for a $y \in Y$, $y^* \in Y^*$, $\langle y, y^* \rangle$ will also be used for $y^*(y)$.

Received Septembr 29, 2011.

2010 *Mathematics Subject Classification*. Primary 47B38, 46B42; Secondary 28A25.

Key words and phrases. Multiplication operator; Köthe function spaces; Köthe-Bochner function spaces.



Go back

Full Screen

Close

Quit



In ([2]) a result is proved about the multiplication operators in Köthe-Bochner spaces. The proof is quite sophisticated and, besides several lemmas, makes use of Markushevich bases. In this note we give a simple elementary proof.

2. MAIN THEOREM

Now we come to the main theorem

Theorem 1. *Suppose E an order continuous Köthe function space over μ , X a Banach space and $E(X)$ the associated Köthe-Bochner space. Let $T: E(X) \rightarrow E(X)$ be a continuous linear operator. The following statements are equivalent:*

- (i) *There is a $g_0 \in L^\infty$ such that $T(f) = g_0 f$ for all $f \in E(X)$.*
- (ii) *$T(g\langle f, x^* \rangle x) = g\langle T(f), x^* \rangle x$ for every $g \in L^\infty$, $f \in E(X)$, $x \in X$, and $x^* \in X^*$.*

Proof. (i) \implies (ii): Obvious.

(ii) \implies (i): For an $h \in E$, $x \in X$, $g \in L^\infty$, we have $T((gh\langle x, x^* \rangle)x) = gh\langle T(x), x^* \rangle x$; take any $x^* \in X^*$ with $\langle x, x^* \rangle = 1$. We get $T(ghx) = ghpx$ for some $p \in E$ (note since $|\langle T(x)(\cdot), x^* \rangle| \leq \|T(x)(\cdot)\|$, we have $\langle T(x)(\cdot), x^* \rangle \in E$) and so $ghpx \in E(X)$. p may depend on x . Suppose $T(x_1) = p_1 x_1$ and $T(x_2) = p_2 x_2$. We claim $p_1 = p_2$. If x_1, x_2 are linearly dependent, there is nothing to prove; otherwise $x_1, x_1 - x_2$ are linearly independent. Take an $x^* \in X^*$ such that $\langle x_1, x^* \rangle = 1, \langle x_1 - x_2, x^* \rangle = 0$. This means $0 = T(\langle x_1 - x_2, x^* \rangle z) = \langle T(x_1 - x_2), x^* \rangle z = \langle p_1 x_1 - p_2 x_2, x^* \rangle z = (p_1 - p_2)z$, for all $z \in E$. From this it follows that $p_1 = p_2$.

Now we want to prove that p is bounded. Suppose this is not true. Select a strictly increasing sequence $\{c_n\}$ of positive real numbers such that (i) $c_n n^3$, (ii) $\mu(Q_n) > 0$ where $Q_n = |p|^{-1}(c_n, c_{n+1})$. For each n , choose positive α_n so that, for the functions $f_n = \alpha_n \chi_{Q_n}$, $\|f_n\|_E = 1$. Fix a $y \in X$ with $\|y\|_X = 1$. The function $f = \sum_{n=1}^{\infty} \frac{1}{n^2} f_n$ is in E and $f \geq \frac{1}{n^2} f_n$. This gives $f|p| \geq \frac{1}{n^2} f_n |p| \geq \frac{1}{n^2} f_n n^3$




Go back

Full Screen

Close

Quit



and so $\|f|p\|_E \geq n$ for all n . Now $\|T(fy)\|_{E(X)} = \|fpy\|_{E(X)} = \|f|p\|_E \geq n$ for all n , which is a contradiction. So $p \in L^\infty$. We put $g_0 = p$. Thus $T(gx) = gg_0x$, for all $x \in X$, $g \in L^\infty$ and so $T(h) = g_0h$ for all simple functions $h \in E(X)$. Since E is order continuous, simple functions are dense, and so the result follows. \square

1. Diestel J. and Uhl J. J., *Vector Measures*, Amer. Math. Soc. Surveys vol. 15 Amer. Math. Soc., 1977.
2. Calabuig J. M., Rodriguez J. and Sanchez-Perez, E. A., *Multiplication operators in Köthe-Bochner spaces*. J. Math. Anal. Appl. **373** (2011), 316–321.
3. Lin, P. K., *Köthe-Bochner function spaces*. Birkhauser Boston Inc., MA, 2004.

S. S. Khurana, Departem of Mathematics, Univerisity of Iowa, Iowa City, Iowa 52242, U. S. A.,
e-mail: khurana@math.uiowa.edu



Go back

Full Screen

Close

Quit