

WEAK CONVERGENCE TO OCONE MARTINGALES: A REMARK

GIOVANNI PECCATI

Laboratoire de Statistique Théorique et Appliquée, Université Pierre et Marie Curie - Paris VI, France
email: giovanni.peccati@libero.it

Submitted December 1, 2004, accepted in final form December 3, 2004

AMS 2000 Subject classification: 60G44, 60F17

Keywords: Weak Convergence, Ocone Martingales

Abstract

We show, by a simple counterexample, that the main result in a recent paper by H. Van Zanten [Electronic Communications in Probability **7** (2002), 215-222] is false. We eventually point out the origin of the error.

Throughout the following we use concepts and notation from standard semimartingale theory. The reader is referred e.g. to [3] for any unexplained notion. Every càdlàg stochastic process is defined on a given probability space $(\Omega, \mathcal{F}, \mathbb{P})$, and it is interpreted as a random element with values in $D([0, \infty))$, the Skorohod space of càdlàg functions on $[0, \infty)$. The symbol “ \Rightarrow ” indicates weak convergence (see [2]). Given a filtration \mathcal{F}_t and a real-valued càdlàg \mathcal{F}_t -local martingale started from zero, say $M = \{M_t : t \geq 0\}$, we will denote by $[M] = \{[M]_t : t \geq 0\}$ the optional quadratic variation of M . We recall that, when M is continuous, $[M] = \langle M \rangle$, where $\langle M \rangle$ is the conditional quadratic variation of M as defined in [3, Chapter III]. Moreover, by the Dambis-Dubins-Schwarz (DDS) Theorem (see [4, Ch. V]), every continuous \mathcal{F}_t -local martingale M , such that $M_0 = 0$ and $\langle M \rangle_\infty = \lim_{t \rightarrow +\infty} \langle M \rangle_t = +\infty$ a.s.- \mathbb{P} , can be represented as

$$M_t = W_{\langle M \rangle_t}^{(M)}, \quad t \geq 0, \quad (1)$$

where $W_t^{(M)}$ is a standard Brownian motion with respect to the filtration

$$\mathcal{G}_t = \mathcal{F}_{\sigma(t)}, \quad t \geq 0, \quad \text{where } \sigma(t) = \inf \{s : \langle M \rangle_s > t\}.$$

According e.g. to [7], we say that a continuous \mathcal{F}_t -martingale M_t , such that $M_0 = 0$ and $\langle M \rangle_\infty = +\infty$, is a (continuous) *Ocone martingale* if the Brownian motion $W^{(M)}$ appearing in its DDS representation (1) is independent of $\langle M \rangle$.

The following statement, concerning rescaled càdlàg martingales, appears as Theorem 4.1 in [6].

Claim 1 *Let M be a martingale with bounded jumps, and let a_n, b_n be sequences of positive numbers both increasing to infinity. For each n , define*

$$M_t^n = \frac{M_{b_n t}}{\sqrt{a_n}}. \quad (2)$$

Then, the following statements hold

(i) If $M^n \Rightarrow N$ in $D([0, \infty))$, then necessarily N is a continuous Ocone martingale.

(ii) Let N be a continuous Ocone martingale. Then, $M^n \Rightarrow N$ in $D([0, \infty))$ if, and only if, $[M^n] \Rightarrow [N]$ in $D([0, \infty))$.

Both parts (i) and (ii) of Claim 1 are false, as shown by the following counterexample. Take a standard Brownian motion started from zero $W = \{W_t : t \geq 0\}$, and define

$$\begin{aligned} M_t &= W_t^2 - t \\ M_t^n &= \frac{1}{n} M_{nt} = \left(n^{-\frac{1}{2}} W_{nt}\right)^2 - t. \end{aligned}$$

Then, M is a continuous square-integrable martingale that is *not* Ocone (since it is non-Gaussian and *pure*, see [5, Proposition 2.5] and [7, p. 423]). Moreover, $M_t^n = (a_n)^{-1/2} M_{b_n t}$, for $a_n = n^2$ and $b_n = n$, and $M^n \stackrel{\text{law}}{=} M$ for each n , due to the scaling properties of Brownian motion. It follows that $M^n \Rightarrow M$, thus contradicting point (i) of Claim 1.

As for point (ii), consider the continuous Ocone martingale (see [7, p. 427])

$$N_t = 2 \int_0^t W_s d\widetilde{W}_s$$

where \widetilde{W} is a standard Brownian motion independent of W . It is evident that

$$\begin{aligned} [N]_t &= 4 \int_0^t W_s^2 ds \\ [M^n]_t &= \frac{4}{n^2} \int_0^{nt} W_s^2 ds = 4 \int_0^t \left(n^{-1/2} W_{nu}\right)^2 du \end{aligned}$$

and therefore that $[M^n] \stackrel{\text{law}}{=} [N]$ for each n , although M^n converges weakly to the martingale M , which is not Ocone. This contradicts point (ii) of Claim 1.

The error comes from a misuse of the Skorohod almost sure representation theorem (see e.g. [1, p. 281]) in [6, Section 4]. Starting from p. 219, line 10 of [6], the author considers a sequence

$$\left\{ \left(W, \tau^{n'} \right) : n' \geq 1 \right\},$$

where W is a standard Brownian motion and $\tau^{n'}$ is an appropriate time-change, such that

$$\left(W, \tau^{n'} \right) \Rightarrow (B, [N]),$$

where B is a standard Brownian motion, and $[N]$ is a positive, continuous and increasing process. Then, the Skorohod theorem allows one to conclude that, on an auxiliary space, there exist random elements $(\overline{W}^{n'}, \overline{\tau}^{n'})$ and $(\overline{B}, \overline{[N]})$ such that

$$\left(W, \tau^{n'} \right) \stackrel{\text{law}}{=} \left(\overline{W}^{n'}, \overline{\tau}^{n'} \right) \quad \text{and} \quad (B, [N]) \stackrel{\text{law}}{=} (\overline{B}, \overline{[N]}),$$

where the Brownian motion $\overline{W}^{n'}$ depends (in general) on n' , and $(\overline{W}^{n'}, \overline{\tau}^{n'}) \xrightarrow{\text{a.s.}} (B, [N])$. On the other hand, the (fallacious) conclusion of Theorem 4.1 in [6] is obtained by supposing that, on the auxiliary space, there exists a Brownian motion \overline{W} such that $\overline{W}^{n'} = \overline{W}$ for each n' , which is clearly not the case, due to the counterexamples constructed above.

References

- [1] Grimmet G. R. and Stirzaker D. R. (1992). *Probability and Random Processes*. Oxford Science Publications. Oxford.
- [2] Jacod J. and Shiryaev A. N. (1987). *Limit theorems for stochastic processes*. Berlin: Springer-Verlag
- [3] Protter P. (1992). *Stochastic Integration and Differential Equations. A New Approach*. Berlin: Springer-Verlag
- [4] Revuz D. and Yor M. (2001). *Continuous martingales and Brownian motion*. Berlin: Springer-Verlag
- [5] Stroock D. W. and Yor M. (1981). Some remarkable martingales. In *Séminaire de probabilités XV*, 590-603. Berlin: Springer-Verlag
- [6] Van Zanten, J. H. (2002). Continuous Ocone martingales as weak limits of rescaled martingales. *Electronic Communications in Probability*, **7**, 215-222
- [7] Vostrikova, L. and Yor, M. (2000). Some invariance properties of the laws of Ocone martingales. In *Séminaire de probabilités XXXIV*, 417-431. Berlin: Springer-Verlag