

L_1 -distance for additive processes with time-homogeneous Lévy measures

Pierre Étoré*

Ester Mariucci†

Abstract

We give an explicit bound for the L_1 -distance between two additive processes of local characteristics $(f_j(\cdot), \sigma^2(\cdot), \nu_j)$, $j = 1, 2$. The cases $\sigma = 0$ and $\sigma(\cdot) > 0$ are both treated. We allow ν_1 and ν_2 to be time-homogeneous Lévy measures, possibly with infinite variation. Some examples of possible applications are discussed.

Keywords: L_1 -distance ; Total Variation ; Additive Processes.

AMS MSC 2010: 60B10 ; 60E15 ; 60G51.

Submitted to ECP on April 29, 2014, final version accepted on August 4, 2014.

Supersedes arXiv:1404.7779v2.

Supersedes HAL : hal - 00985588.

1 Introduction and main result

In this note we give an upper bound for the L_1 -distance between the laws induced on the Skorokhod space by two additive processes observed until time $T > 0$. By the L_1 -distance between two σ -finite measures μ_1 and μ_2 on (E, \mathcal{E}) such that μ_1 is absolutely continuous with respect to μ_2 we mean

$$L_1(\mu_1, \mu_2) = 2 \sup_{A \in \mathcal{E}} |\mu_1(A) - \mu_2(A)| = \int_E \left| \frac{d\mu_1}{d\mu_2} - 1 \right| d\mu_2.$$

Note that, with our definitions, the L_1 -distance is twice the so called total variation distance.

Giving bounds for the L_1 -distance is a classical problem, which, in the last decades, has been reinterpreted in more modern terms via Stein's method (see, e.g., [14, 17, 16]). This kind of problems arises in several fields such as Bayesian statistics, convergence rates of Markov chains or Monte Carlo algorithms (see [7], Section 4 and the references therein). However, to the best of our knowledge, results bounding the L_1 -distance between laws on the Skorokhod space are much less abundant. In this setting other kinds of distances have been privileged such as the Wasserstein-Kantorovich-Rubinstein metric (see [5]). More relevant to our purposes is a result due to Memin and Shirayev [11] computing the Hellinger distance between the laws of any two processes with independent increments. In particular this gives a bound for the L_1 -distance between additive processes. In order to state their result let us fix some notation.

*Laboratoire Jean Kuntzmann, Grenoble.

†Laboratoire Jean Kuntzmann, Grenoble. E-mail: Ester.Mariucci@imag.fr

Let $\{x_t\}$ be the canonical process on the Skorokhod space (D, \mathcal{D}) and denote by $P^{(f, \sigma^2, \nu)}$ the law induced on (D, \mathcal{D}) by an additive process having local characteristics $(f(\cdot), \sigma^2(\cdot), \nu)$. We will denote such a process by $(\{x_t\}, P^{(f, \sigma^2, \nu)})$ and we will write $P_T^{(f, \sigma^2, \nu)}$ for the restriction of $P^{(f, \sigma^2, \nu)}$ to the σ -algebra generated by $\{x_s : 0 \leq s \leq T\}$ (see Section 2 for the precise definitions). Our purpose is to bound $L_1(P_T^{(f_1, \sigma_1^2, \nu_1)}, P_T^{(f_2, \sigma_2^2, \nu_2)})$. From now on we will assume that $\sigma_1^2(\cdot) = \sigma_2^2(\cdot) = \sigma^2(\cdot)$, otherwise this distance is 2 (see, e.g., [13, 8]). We also need to define the following quantities:

$$\gamma^{\nu_j} = \int_{|y| \leq 1} y \nu_j(dy), \quad j = 1, 2; \quad \xi^2 = \int_0^T \frac{(f_2(r) - f_1(r) - (\gamma^{\nu_2} - \gamma^{\nu_1}))^2}{\sigma^2(r)} dr.$$

Theorem 1.1 (Memin and Shirayev). *Let $(\{x_t\}, P^{(f_1, \sigma^2, \nu_1)})$ and $(\{x_t\}, P^{(f_2, \sigma^2, \nu_2)})$ be two additive processes with ν_1 and ν_2 Lévy measures such that ν_1 is absolutely continuous with respect to ν_2 and satisfying:*

$$H^2(\nu_1, \nu_2) := \int_{\mathbb{R}} \left(\sqrt{\frac{d\nu_1}{d\nu_2}}(y) - 1 \right)^2 \nu_2(dy) < \infty. \tag{1.1}$$

The following upper bounds hold, for any $0 < T < \infty$: If $\sigma^2 > 0$ then

$$L_1(P_T^{(f_1, \sigma^2, \nu_1)}, P_T^{(f_2, \sigma^2, \nu_2)}) \leq \sqrt{8 \left(1 - \exp \left(-\frac{\xi^2}{8} - \frac{T}{2} H^2(\nu_1, \nu_2) \right) \right)}.$$

If $\sigma^2 = 0$ and $f_1 - f_2 \equiv \gamma^{\nu_1} - \gamma^{\nu_2}$, then

$$L_1(P_T^{(f_1, 0, \nu_1)}, P_T^{(f_2, 0, \nu_2)}) \leq \sqrt{8 \left(1 - \exp \left(-\frac{T}{2} H^2(\nu_1, \nu_2) \right) \right)}.$$

Observe that (1.1) implies $\gamma^{\nu_j} < \infty$, $j = 1, 2$. When $\sigma^2 = 0$ it follows from Theorem 1.2 that, for example, $L_1(P_T^{(\gamma^{\nu_1}, 0, \nu_1)}, P_T^{(\gamma^{\nu_2}, 0, \nu_2)}) \leq 2\sqrt{TL_1(\nu_1, \nu_2)}$.

The proof of Theorem 1.1, however, makes heavy use of general theory of semimartingales. This note originated from the research for a proof based only on classical results for Lévy processes, Esscher-type transformations and the Cameron-Martin formula. It turned out that this method, when applicable, gives sharper bound on the L_1 -distance. More precisely, our main result is as follows.

Theorem 1.2. *Let $(\{x_t\}, P^{(f_1, \sigma^2, \nu_1)})$ and $(\{x_t\}, P^{(f_2, \sigma^2, \nu_2)})$ be two additive processes with ν_1 and ν_2 Lévy measures such that ν_1 is absolutely continuous with respect to ν_2 and satisfying:*

$$L_1(\nu_1, \nu_2) < \infty.$$

Then, the following upper bounds hold, for any $0 < T < \infty$.

If $\sigma^2 > 0$ then

$$L_1(P_T^{(f_1, \sigma^2, \nu_1)}, P_T^{(f_2, \sigma^2, \nu_2)}) \leq 2 \sinh(TL_1(\nu_1, \nu_2)) + 2 \left[1 - 2\phi \left(-\frac{\xi}{2} \right) \right].$$

If $\sigma^2 = 0$ and $f_1 - f_2 \equiv \gamma^{\nu_1} - \gamma^{\nu_2}$, then

$$L_1(P_T^{(f_1, 0, \nu_1)}, P_T^{(f_2, 0, \nu_2)}) \leq 2 \sinh(TL_1(\nu_1, \nu_2)).$$

Remark that, in the case $\nu_1 = \nu_2 = 0$, i.e. where there are no jumps, the upper bound in Theorem 1.2 is achieved. Indeed, an explicit formula for the L_1 -distance between

Gaussian processes is well known. Denoting by ϕ the cumulative distribution function of a normal random variable $\mathcal{N}(0, 1)$, we have, for any $0 < T < \infty$:

$$L_1(P_T^{(f_1, \sigma^2, 0)}, P_T^{(f_2, \sigma^2, 0)}) = 2 \left(1 - 2\phi \left(-\frac{1}{2} \sqrt{\int_0^T \frac{(f_1(t) - f_2(t))^2}{\sigma_1^2(t)} dt} \right) \right)$$

whenever the right-hand side term makes sense (see, e.g., [1]).

The reason for our interest in the L_1 -distance lies in the fact that it is a fundamental tool in the Le Cam theory of comparison of statistical models ([9, 10]). More precisely, the presented result will be needed in a forthcoming paper by the second author, establishing an equivalence result, in the Le Cam sense, for additive processes. Similar estimations appear in many other results concerning the Le Cam Δ -distance. See for example [1, 15, 2], where the L_1 -distance between Gaussian processes is computed or [12, 4, 6] concerning diffusion processes without jumps. In recent years, however, there is a growing interest in models with jumps due to their numerous applications in econometrics, insurance theory or financial modelling. Because of that, it is useful to dispose of simple formulas for estimating distances between such processes.

Theorem 1.2 is proved in Section 3. In Section 2 we collect some preliminary results about additive processes that will play a role in the proof. Before that, we give some examples of situations where our result can be applied. The choice of these examples are inspired by the models exhibited in [3].

Example 1.3. (L_1 -distance between compound Poisson processes)

Let $\{X_t^1\}$ and $\{X_t^2\}$ be two compound Poisson processes on $[0, T]$ with intensities $\lambda_j > 0$, $j = 1, 2$ and jump size distributions G_j ; i.e. $\{X_t^j\}$ is a Lévy process of characteristic triplet $(\lambda_j \int_{|y| \leq 1} y G_j(dy), 0, \lambda_j G_j)$. Furthermore, let A be a subset of \mathbb{R} and suppose that G_j is equivalent to the Lebesgue measure restricted to A . Denote by g_j the density $\frac{dG_j}{d\text{Leb}_A}$; then, an application of Theorem 1.2 yields:

$$L_1(X^1, X^2) \leq 2 \sinh \left(T \int_A |\lambda_1 g_1(y) - \lambda_2 g_2(y)| dy \right).$$

Example 1.4. (L_1 -distance between additive processes of jump-diffusion type)

An additive process of jump-diffusion type on $[0, T]$ has the following form:

$$X_t = \int_0^t f(r) dr + \int_0^t \sigma(r) dW_r + \sum_{i=1}^{N_t} Y_i, \quad t \in [0, T],$$

where $\{W_t\}$ is a standard Brownian motion, $\{N_t\}$ is the Poisson process counting the jumps of $\{X_t\}$, and Y_i are jumps sizes (i.i.d. random variables). Consider now the additive processes of jump-diffusion type $\{X_t^j\}$ having local characteristics $(f_j(\cdot) + \lambda_j \int_{|y| \leq 1} y G_j(dy), \sigma^2(\cdot), \lambda_j G_j)$, $j = 1, 2$ and suppose again that G_j is equivalent to the Lebesgue measure restricted to some $A \subseteq \mathbb{R}$. Letting g_j denote the density of G_j as above, we have:

$$L_1(X^1, X^2) \leq 2 \sinh \left(T \int_A |\lambda_1 g_1(y) - \lambda_2 g_2(y)| dy \right) + 2 \left(1 - 2\phi \left(-\sqrt{\int_0^T \frac{(f_1(t) - f_2(t))^2}{4\sigma^2(t)} dt} \right) \right).$$

Example 1.5. (L_1 -distance between tempered stable processes)

Let $\{X_t^1\}$ and $\{X_t^2\}$ be two tempered stable processes, i.e. Lévy processes on \mathbb{R} with no gaussian component and such that their Lévy measures ν_j have densities of the form

$$\frac{d\nu_j}{d\text{Leb}}(y) = \frac{C_-}{|y|^{1+\alpha}} e^{-\lambda_-^j |y|} \mathbb{I}_{y < 0} + \frac{C_+}{y^{1+\alpha}} e^{-\lambda_+^j y} \mathbb{I}_{y > 0}, \quad j = 1, 2,$$

for some parameters $C_{\pm} > 0$, $\lambda_{\pm}^j > 0$ and $\alpha < 2$. Then the hypothesis (1.1) is satisfied and Theorem 1.2 bounds the L_1 -distance by:

$$2 \sinh \left(T \left[C_+ \int_0^{\infty} \left| \frac{e^{-\lambda_+^1 y} - e^{-\lambda_+^2 y}}{y^{1+\alpha}} \right| dy + C_- \int_{-\infty}^0 \left| \frac{e^{-\lambda_-^1 |y|} - e^{-\lambda_-^2 |y|}}{|y|^{1+\alpha}} \right| dy \right] \right).$$

2 Preliminary results

2.1 Additive processes

Definition 2.1. A stochastic process $\{X_t\} = \{X_t : t \geq 0\}$ on \mathbb{R} defined on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$ is an additive process if the following conditions are satisfied.

1. $X_0 = 0$ \mathbb{P} -a.s.
2. For any choice of $n \geq 1$ and $0 \leq t_0 < t_1 < \dots < t_n$, random variables $X_{t_0}, X_{t_1} - X_{t_0}, \dots, X_{t_n} - X_{t_{n-1}}$ are independent.
3. There is $\Omega_0 \in \mathcal{A}$ with $\mathbb{P}(\Omega_0) = 1$ such that, for every $\omega \in \Omega_0$, $X_t(\omega)$ is right-continuous in $t \geq 0$ and has left limits in $t > 0$.
4. It is stochastically continuous.

Thanks to the Lévy-Khintchine formula, the characteristic function of any additive process $\{X_t\}$ can be expressed, for all u in \mathbb{R} , as:

$$\mathbb{E}[e^{iuX_t}] = \exp \left(iu \int_0^t f(r) dr - \frac{u^2}{2} \int_0^t \sigma^2(r) dr - t \int_{\mathbb{R}} (1 - e^{iuy} + iuy \mathbb{I}_{|y| \leq 1}) \nu(dy) \right), \quad (2.1)$$

where $f(\cdot), \sigma^2(\cdot)$ are functions on $L_1[0, T]$ and ν is a measure on \mathbb{R} satisfying

$$\nu(\{0\}) = 0 \text{ and } \int_{\mathbb{R}} (|y|^2 \wedge 1) \nu(dy) < \infty.$$

In the sequel we shall refer to $(f(\cdot), \sigma^2(\cdot), \nu)$ as the local characteristics of the process $\{X_t\}$ and ν will be called Lévy measure. This data characterises uniquely the law of the process $\{X_t\}$. In the case in which $f(\cdot)$ and $\sigma(\cdot)$ are constant functions, a process $\{X_t\}$ satisfying (2.1) is said a Lévy process of characteristic triplet (f, σ^2, ν) .

Let $D = D([0, \infty), \mathbb{R})$ be the space of mappings ω from $[0, \infty)$ into \mathbb{R} that are right-continuous with left limits. Define the canonical process $x : D \rightarrow D$ by

$$\forall \omega \in D, \quad x_t(\omega) = \omega_t, \quad \forall t \geq 0.$$

Let \mathcal{D}_t and \mathcal{D} be the σ -algebras generated by $\{x_s : 0 \leq s \leq t\}$ and $\{x_s : 0 \leq s < \infty\}$, respectively (here, we use the same notations as in [18]).

Let $\{X_t\}$ be an additive process defined on $(\Omega, \mathcal{A}, \mathbb{P})$ having local characteristics $(f(\cdot), \sigma^2(\cdot), \nu)$. It is well known that it induces a probability measure $P^{(f, \sigma^2, \nu)}$ on (D, \mathcal{D}) such that $(\{x_t\}, P^{(f, \sigma^2, \nu)})$ is an additive process identical in law with $(\{X_t\}, \mathbb{P})$ (that is the local characteristics of $\{x_t\}$ under $P^{(f, \sigma^2, \nu)}$ is $(f(\cdot), \sigma^2(\cdot), \nu)$). For all $t > 0$ we will denote $P_t^{(f, \sigma^2, \nu)}$ for the restriction of $P^{(f, \sigma^2, \nu)}$ to \mathcal{D}_t . In the case where $\int_{|y| \leq 1} |y| \nu(dy) < \infty$, we set $\gamma^\nu := \int_{|y| \leq 1} y \nu(dy)$. Note that, if ν is a finite Lévy measure, then the process $(\{x_t\}, P^{(\gamma^\nu, 0, \nu)})$ is a compound Poisson process.

Here and in the sequel we will denote by Δx_r the jump of process $\{x_t\}$ at the time r :

$$\Delta x_r = x_r - \lim_{s \uparrow r} x_s.$$

Definition 2.2. Consider $(\{x_t\}, P^{(f, \sigma^2, \nu)})$ and define the jump part of $\{x_t\}$ as

$$x_t^{d, \nu} = \lim_{\varepsilon \rightarrow 0} \left(\sum_{r \leq t} \Delta x_r \mathbb{I}_{|\Delta x_r| > \varepsilon} - t \int_{\varepsilon < |y| \leq 1} y \nu(dy) \right) \quad \text{a.s.} \quad (2.2)$$

and its continuous part as

$$x_t^{c, \nu} = x_t - x_t^{d, \nu} \quad \text{a.s.} \quad (2.3)$$

We now recall the *Lévy-Itô decomposition*, i.e. the decomposition in continuous and discontinuous parts of an additive process.

Theorem 2.3 (See [18], Theorem 19.3). Consider $(\{x_t\}, P^{(f, \sigma^2, \nu)})$ and define $\{x_t^{d, \nu}\}$ and $\{x_t^{c, \nu}\}$ as in 2.2 and 2.3, respectively. Then the following hold.

- (i) There is $D_1 \in \mathcal{D}$ with $P^{(f, \sigma^2, \nu)}(D_1) = 1$ such that, for any $\omega \in D_1$, $x_t^{d, \nu}(\omega)$ is defined for all $t \in [0, T]$ and the convergence is uniform in t on any bounded interval, $P^{(f, \sigma^2, \nu)}$ -a.s. The process $\{x_t^{d, \nu}\}$ is a Lévy process on \mathbb{R} with characteristic triplet $(0, 0, \nu)$.
- (ii) There is $D_2 \in \mathcal{D}$ with $P^{(f, \sigma^2, \nu)}(D_2) = 1$ such that, for any $\omega \in D_2$, $x_t^{c, \nu}(\omega)$ is continuous in t . The process $\{x_t^{c, \nu}\}$ is an additive process on \mathbb{R} with local characteristics $(f(\cdot), \sigma^2(\cdot), 0)$.
- (iii) The two processes $\{x_t^{d, \nu}\}$ and $\{x_t^{c, \nu}\}$ are independent.

2.2 Change of measure for additive processes

For the proof of Theorem 1.2 we also need some results on the equivalence of measures for additive processes. By the notation \ll we will mean “is absolutely continuous with respect to”.

2.2.1 Case $\sigma^2 = 0$

Theorem 2.4 (See [18], Theorems 33.1–33.2 and [19] Corollary 3.18, Remark 3.19). Let $(\{x_t\}, P^{(0, 0, \tilde{\nu})})$ and $(\{x_t\}, P^{(\eta, 0, \nu)})$ be two Lévy processes on \mathbb{R} , where

$$\eta = \int_{|y| \leq 1} y(\nu - \tilde{\nu})(dy) \quad (2.4)$$

is supposed to be finite. Then $P_t^{(\eta, 0, \nu)} \ll P_t^{(0, 0, \tilde{\nu})}$ for all $t \geq 0$ if and only if $\nu \ll \tilde{\nu}$ and the density $\frac{d\nu}{d\tilde{\nu}}$ satisfies

$$\int \left(\sqrt{\frac{d\nu}{d\tilde{\nu}}}(y) - 1 \right)^2 \tilde{\nu}(dy) < \infty. \quad (2.5)$$

Remark that the finiteness in (2.5) implies that in (2.4). When $P_t^{(\eta, 0, \nu)} \ll P_t^{(0, 0, \tilde{\nu})}$, the density is

$$\frac{dP_t^{(\eta, 0, \nu)}}{dP_t^{(0, 0, \tilde{\nu})}}(x) = \exp(U_t(x)),$$

with

$$U_t(x) = \lim_{\varepsilon \rightarrow 0} \left(\sum_{r \leq t} \ln \frac{d\nu}{d\tilde{\nu}}(\Delta x_r) \mathbb{I}_{|\Delta x_r| > \varepsilon} - \int_{|y| > \varepsilon} t \left(\frac{d\nu}{d\tilde{\nu}}(y) - 1 \right) \tilde{\nu}(dy) \right), P^{(0, 0, \tilde{\nu})}\text{-a.s.} \quad (2.6)$$

The convergence in (2.6) is uniform in t on any bounded interval, $P^{(0, 0, \tilde{\nu})}$ -a.s. Besides, $\{U_t(x)\}$ defined by (2.6) is a Lévy process satisfying $\mathbb{E}_{P^{(0, 0, \tilde{\nu})}}[e^{U_t(x)}] = 1, \forall t \in [0, T]$.

2.2.2 Case $\sigma^2 > 0$

Lemma 2.5. *Let $\nu_1 \ll \nu_2$ be Lévy measures such that*

$$\int_{\mathbb{R}} \left(\sqrt{\frac{d\nu_1}{d\nu_2}}(y) - 1 \right)^2 \nu_2(dy) < \infty. \tag{2.7}$$

Define

$$\eta = \int_{|y| \leq 1} y(\nu_1 - \nu_2)(dy), \tag{2.8}$$

which is finite thanks to (2.7), and consider real functions f_1, f_2 and $\sigma > 0$ such that

$$\int_0^T \left(\frac{f_1(r) - f_2(r) - \eta}{\sigma(r)} \right)^2 dr < \infty, \quad T \geq 0. \tag{2.9}$$

Then, under $P^{(f_2, \sigma^2, \nu_2)}$,

$$M_t(x) = \exp(C_t(x) + D_t(x)) \tag{2.10}$$

is a (\mathcal{D}_t) -martingale for all t in $[0, T]$, where

$$C_t(x) := \int_0^t \frac{f_1(r) - f_2(r) - \eta}{\sigma^2(r)} (dx_r^{c, \nu_2} - f_2(r)dr) - \frac{1}{2} \int_0^t \left(\frac{f_1(r) - f_2(r) - \eta}{\sigma(r)} \right)^2 dr,$$

$$D_t(x) := \lim_{\varepsilon \rightarrow 0} \left(\sum_{r \leq t} \ln \frac{d\nu_1}{d\nu_2}(\Delta x_r) \mathbb{I}_{|\Delta x_r| > \varepsilon} - t \int_{|y| > \varepsilon} (\nu_1 - \nu_2)(dy) \right). \tag{2.11}$$

The convergence in (2.11) is uniform in t on any bounded interval, $P^{(f_2, \sigma^2, \nu_2)}$ -a.s.

Proof. The existence of the limit in (2.11) is guaranteed by (2.7) (see Theorem 2.4). Since $\int_0^t \frac{1}{\sigma(r)} (dx_r^{c, \nu_2} - f_2(r)dr)$ is a standard Brownian motion under $P^{(f_2, \sigma^2, 0)}$, we have that $\int_s^t \frac{f_1(r) - f_2(r) - \eta}{\sigma^2(r)} (dx_r^{c, \nu_2} - f_2(r)dr)$ has normal law $\mathcal{N}\left(0, \int_s^t \left(\frac{f_1(r) - f_2(r) - \eta}{\sigma(r)} \right)^2 dr\right)$, hence $\mathbb{E}_{P^{(f_2, \sigma^2, 0)}}[\exp((C_t - C_s)(x))] = 1$. Theorem 2.3 entails that $\{x_t^{c, \nu_2}\}$ and $\{x_t^{d, \nu_2}\}$ are independent under $P^{(f_2, \sigma^2, \nu_2)}$. Moreover, the law of $\{C_t(x)\}$ (resp. $\{D_t(x)\}$) is the same under $P^{(f_2, \sigma^2, \nu_2)}$ or $P^{(f_2, \sigma^2, 0)}$ (resp. $P^{(f_2, 0, \nu_2)}$ or $P^{(0, 0, \nu_2)}$). Further, using Theorem 2.4, we know that $\{D_t(x)\}$ is a Lévy process such that $\mathbb{E}_{P^{(0, 0, \nu_2)}}[\exp(D_{t-s}(x))] = 1$ for all $s < t$. These facts together with the independence of the increments of $(\{x_t\}, P^{(f_2, \sigma^2, \nu_2)})$ and the stationarity of $\{D_t(x)\}$ imply:

$$\begin{aligned} \mathbb{E}_{P^{(f_2, \sigma^2, \nu_2)}}[M_t(x) | \mathcal{D}_s] &= \mathbb{E}_{P^{(f_2, \sigma^2, \nu_2)}} \left[M_s(x) \exp\left((C_t - C_s)(x) + (D_t - D_s)(x) \right) | \mathcal{D}_s \right] \\ &= M_s(x) \mathbb{E}_{P^{(f_2, \sigma^2, \nu_2)}}[\exp((C_t - C_s)(x) + (D_t - D_s)(x))] \\ &= M_s(x) \mathbb{E}_{P^{(f_2, \sigma^2, 0)}}[\exp((C_t - C_s)(x))] \mathbb{E}_{P^{(0, 0, \nu_2)}}[\exp((D_t - D_s)(x))] \\ &= M_s(x) \mathbb{E}_{P^{(0, 0, \nu_2)}}[\exp(D_{t-s}(x))] \\ &= M_s(x). \end{aligned}$$

□

Lemma 2.6. *Suppose that the hypothesis (2.7) and (2.9) of Lemma 2.5 are satisfied. Then, using the same notations as above, $P_t^{(f_1, \sigma^2, \nu_1)} \ll P_t^{(f_2, \sigma^2, \nu_2)}$ for all t and the density is given by:*

$$\frac{dP_t^{(f_1, \sigma^2, \nu_1)}}{dP_t^{(f_2, \sigma^2, \nu_2)}}(x) = M_t(x). \tag{2.12}$$

Proof. For $s < t$, we prove that

$$\mathbb{E}_{P(f_2, \sigma^2, \nu_2)} \left[\exp(iu(x_t - x_s)) \frac{M_t}{M_s}(x) \middle| \mathcal{D}_s \right] = \mathbb{E}_{P(f_1, \sigma^2, \nu_1)} [\exp(iu(x_t - x_s))].$$

To that aim remark that, thanks again to Theorem 2.3:

$$\begin{aligned} & \mathbb{E}_{P(f_2, \sigma^2, \nu_2)} \left[e^{iu(x_t - x_s)} \frac{M_t(x)}{M_s(x)} \middle| \mathcal{D}_s \right] \\ &= \mathbb{E}_{P(f_2, \sigma^2, \nu_2)} \left[e^{iu(x_t^{c, \nu_2} - x_s^{c, \nu_2} + x_t^{d, \nu_2} - x_s^{d, \nu_2})} \frac{M_t(x)}{M_s(x)} \middle| \mathcal{D}_s \right] \\ &= \mathbb{E}_{P(f_2, \sigma^2, 0)} \left[e^{iu(x_t - x_s)} e^{(C_t - C_s)(x)} \right] \mathbb{E}_{P(0, 0, \nu_2)} \left[e^{iu(x_t - x_s)} e^{(D_t - D_s)(x)} \right]. \end{aligned} \quad (2.13)$$

Let us now compute the first factor of (2.13):

$$\begin{aligned} \mathbb{E}_{P(f_2, \sigma^2, 0)} \left[e^{iu(x_t - x_s)} e^{(C_t - C_s)(x)} \right] &= \mathbb{E}_{P(f_1 - \eta, \sigma^2, 0)} \left[e^{iu(x_t - x_s)} \right] \\ &= \exp \left(iu \int_s^t (f_1(r) - \eta) dr - \frac{u^2}{2} \int_s^t \sigma^2(r) dr \right). \end{aligned}$$

In the first equality we used the Girsanov theorem, thanks to the fact that $\int_0^t \frac{1}{\sigma(r)} (dx_r - f_2(r) dr)$ is a Brownian motion under $P(f_2, \sigma^2, 0)$, while the second one follows from (2.1). We compute the second factor of (2.13) by means of Theorem 2.4 and another application of (2.1):

$$\begin{aligned} \mathbb{E}_{P(0, 0, \nu_2)} \left[e^{iu(x_t - x_s)} e^{(D_t - D_s)(x)} \right] &= \mathbb{E}_{P(0, 0, \nu_2)} \left[e^{iu x_{t-s}} e^{D_{t-s}(x)} \right] \\ &= \mathbb{E}_{P(\eta, 0, \nu_1)} \left[e^{iu x_{t-s}} \right] \\ &= \exp \left((t - s) \left[iu\eta - \int_{\mathbb{R}} (1 - e^{iuy} + iuy \mathbb{I}_{|y| \leq 1}) \nu_1(dy) \right] \right). \end{aligned}$$

Consequently:

$$\mathbb{E}_{P(f_2, \sigma^2, \nu_2)} \left[e^{iu(x_t - x_s)} \frac{M_t(x)}{M_s(x)} \middle| \mathcal{D}_s \right] = \mathbb{E}_{P(f_1, \sigma^2, \nu_1)} [e^{iu(x_t - x_s)}] \quad \forall 0 \leq s \leq t. \quad (2.14)$$

Fix t and define a probability measure P_t on \mathcal{D}_t by $P_t(B) = \mathbb{E}_{P(f_2, \sigma^2, \nu_2)} [M_t \mathbb{I}_B]$ for $B \in \mathcal{D}_t$. As a consequence of Lemma 2.5 and the Bayes rule, the two processes given by $(\{x_s : 0 \leq s \leq t\}, P_t^{(f_1, \sigma^2, \nu_1)})$ and $(\{x_s : 0 \leq s \leq t\}, P_t)$ are identical. Indeed, by (2.14), both have independent increments and the prescribed characteristic function. Consequently, (2.12) holds. \square

3 Proof of Theorem 1.2

For the proof we will need the following three calculus lemmas.

Lemma 3.1. *Let X be a random variable with normal law $\mathcal{N}(m, \sigma^2)$. Then*

$$\mathbb{E} \left| 1 - e^X \right| = 2 \left[\phi \left(-\frac{m}{\sigma} \right) - \phi \left(-\frac{m}{\sigma} - \sigma \right) \right],$$

where $\phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{y^2}{2}} dy$.

Proof. By definition we have

$$\begin{aligned} \mathbb{E} \left| 1 - e^X \right| &= \frac{1}{\sqrt{2\pi\sigma}} \int_{-\infty}^{\infty} |1 - e^x| e^{-\frac{(x-m)^2}{2\sigma^2}} dx \\ &= \frac{1}{\sqrt{2\pi\sigma}} \left(\int_{-\infty}^0 (1 - e^x) e^{-\frac{(x-m)^2}{2\sigma^2}} dx + \int_0^{\infty} (e^x - 1) e^{-\frac{(x-m)^2}{2\sigma^2}} dx \right). \end{aligned}$$

To conclude, just split the sums inside the integrals and use the change of variables ($y = \frac{x-m}{\sigma} - \sigma$), resp. ($y = \frac{x-m}{\sigma}$). \square

Lemma 3.2. For all x, y in \mathbb{R} we have:

$$|1 - e^{x+y}| \leq \frac{1 + e^x}{2} |1 - e^y| + \frac{1 + e^y}{2} |1 - e^x|. \tag{3.1}$$

Proof. By symmetry we restrict to $x \geq 0$.

- $x, y \geq 0$: In this case we have that $|1 - e^{x+y}|$ is exactly equal to $\frac{1+e^x}{2}|1 - e^y| + \frac{1+e^y}{2}|1 - e^x|$.
- $x \geq 0, y \leq 0, x + y \geq 0$: Then the member on the right hand side of (3.1) is equal to $e^x - e^y \geq e^x - 1 \geq e^{x+y} - 1$.
- $x \geq 0, y \leq 0, x + y \leq 0$: In this case the member on the right of (3.1) is equal to $e^x - e^y \geq 1 - e^y \geq 1 - e^{x+y}$.

\square

Lemma 3.3. With the same notations as in Theorem 1.2 and Lemma 2.5, we have:

$$\mathbb{E}_{P_T^{(0,0,\nu_2)}} \left[|1 - \exp(D_T(x))| \right] = \mathbb{E}_{P_T^{(\gamma\nu_2, 0, \nu_2)}} \left[|1 - \exp(D_T(x))| \right] \leq 2 \sinh \left(T \int_{\mathbb{R}} L_1(\nu_1, \nu_2) \right). \tag{3.2}$$

Proof. Thanks to Theorem 2.3 it is clear that $\mathbb{E}_{P_T^{(0,0,\nu_2)}} [|1 - \exp(D_T(x))|] = \mathbb{E}_{P_T^{(\gamma\nu_2, 0, \nu_2)}} [|1 - \exp(D_T(x))|]$. In order to simplify the notations, let us introduce the quantities $h^+ = \left(\frac{d\nu_1}{d\nu_2}\right)^{\mathbb{I}_{\frac{d\nu_1}{d\nu_2} \geq 1}}$ and $h^- = \left(\frac{d\nu_1}{d\nu_2}\right)^{\mathbb{I}_{\frac{d\nu_1}{d\nu_2} < 1}}$ (i.e. h^+ , resp. h^- , is identically 1 where $\frac{d\nu_1}{d\nu_2} < 1$, resp. where $\frac{d\nu_1}{d\nu_2} \geq 1$). Let us define

$$\begin{aligned} A^+(x) &:= \lim_{\varepsilon \rightarrow 0} \left(\sum_{r \leq T} \ln h^+(\Delta x_r) \mathbb{I}_{|\Delta(x_r)| > \varepsilon} - T \int_{|y| > \varepsilon} (h^-(y) - 1) \nu_2(dy) \right), \\ A^-(x) &:= \lim_{\varepsilon \rightarrow 0} \left(\sum_{r \leq T} \ln h^-(\Delta x_r) \mathbb{I}_{|\Delta(x_r)| > \varepsilon} - T \int_{|y| > \varepsilon} (h^+(y) - 1) \nu_2(dy) \right). \end{aligned}$$

Remark that both $A^+(x)$ and $A^-(x)$ have the same law under $P_T^{(\gamma\nu_2, 0, \nu_2)}$ and $P_T^{(0,0,\nu_2)}$ (see Theorem 2.3), and that they are constructed in such a way that

$$D_T(x) = A^+(x) + A^-(x).$$

Using Lemma 3.2 and the fact that $A^+(x) \geq 0$ and $A^-(x) \leq 0$ we get:

$$\begin{aligned} \mathbb{E}_{P_T^{(\gamma\nu_2, 0, \nu_2)}} [|1 - D_T(x)|] &= \mathbb{E}_{P_T^{(\gamma\nu_2, 0, \nu_2)}} |1 - \exp(A^+(x) + A^-(x))| \\ &\leq \mathbb{E}_{P_T^{(\gamma\nu_2, 0, \nu_2)}} \left[\frac{1 + e^{A^+(x)}}{2} |1 - e^{A^-(x)}| + \frac{1 + e^{A^-(x)}}{2} |1 - e^{A^+(x)}| \right] \\ &= \mathbb{E}_{P_T^{(\gamma\nu_2, 0, \nu_2)}} [e^{A^+(x)} - e^{A^-(x)}]. \end{aligned}$$

In order to compute the last quantity let us consider two more Lévy measures:

$$\nu^+ := \nu_2 \mathbb{I}_{\frac{d\nu_1}{d\nu_2} < 1} + \nu_1 \mathbb{I}_{\frac{d\nu_1}{d\nu_2} \geq 1}, \quad \nu^- := \nu_2 \mathbb{I}_{\frac{d\nu_1}{d\nu_2} \geq 1} + \nu_1 \mathbb{I}_{\frac{d\nu_1}{d\nu_2} < 1}.$$

Notice that ν^+, ν^- are absolutely continuous with respect to ν_2 , with densities given by h^+, h^- , respectively. Applying Theorem 2.4 to the pairs of measures (ν^+, ν_2) and (ν^-, ν_2) we get

$$\begin{aligned} \mathbb{E}_{P_T^{(0,0,\nu_2)}} \left[e^{A^+(x)} \right] &= \exp \left(T \int_{\mathbb{R}} (h^+(y) - h^-(y)) \nu_2(dy) \right), \\ \mathbb{E}_{P_T^{(0,0,\nu_2)}} \left[e^{A^-(x)} \right] &= \exp \left(T \int_{\mathbb{R}} (h^-(y) - h^+(y)) \nu_2(dy) \right). \end{aligned}$$

Then, recalling that both $A^+(x)$ and $A^-(x)$ have the same law under $P_T^{(\gamma^{\nu_2}, 0, \nu_2)}$ and $P_T^{(0,0,\nu_2)}$ we obtain:

$$\mathbb{E}_{P_T^{(\gamma^{\nu_2}, 0, \nu_2)}} \left[e^{A^+(x)} - e^{A^-(x)} \right] = 2 \sinh \left(T \int_{\mathbb{R}} \left| 1 - \frac{d\nu_1}{d\nu_2}(y) \right| \nu_2(dy) \right).$$

□

Proof of Theorem 1.2. Case $\sigma^2 > 0$: With the same notations as in Lemma 2.5 and by means of Lemma 2.6 one can write

$$L_1(P_T^{(f_1, \sigma^2, \nu_1)}, P_T^{(f_2, \sigma^2, \nu_2)}) = \mathbb{E}_{P_T^{(f_2, \sigma^2, \nu_2)}} \left| 1 - \exp(C_T(x) + D_T(x)) \right|.$$

Now, using Lemma 3.2 and the independence between $C_T(x)$ and $D_T(x)$ (Theorem 2.3), we obtain

$$\begin{aligned} L_1(P_T^{(f_2, \sigma^2, \nu_2)}, P_T^{(f_1, \sigma^2, \nu_1)}) &\leq \mathbb{E}_{P_T^{(f_2, \sigma^2, \nu_2)}} \left(\frac{1 + e^{C_T(x)}}{2} \right) \mathbb{E}_{P_T^{(f_2, \sigma^2, \nu_2)}} \left| 1 - e^{D_T(x)} \right| \\ &\quad + \mathbb{E}_{P_T^{(f_2, \sigma^2, \nu_2)}} \left(\frac{1 + e^{D_T(x)}}{2} \right) \mathbb{E}_{P_T^{(f_2, \sigma^2, \nu_2)}} \left| 1 - e^{C_T(x)} \right|. \end{aligned}$$

We conclude the proof using Lemmas 3.3 and 3.1 together with the fact that

$$\mathbb{E}_{P_T^{(f_2, \sigma^2, \nu_2)}} e^{C_T(x)} = 1 = \mathbb{E}_{P_T^{(f_2, \sigma^2, \nu_2)}} e^{D_T(x)}.$$

Case $\sigma^2 = 0$: If $f_1 - f_2 \equiv \gamma^{\nu_1} - \gamma^{\nu_2}$, notice that, as the drift component of $(\{x_t\}, P_T^{(f_1, 0, \nu_1)})$ and $(\{x_t\}, P_T^{(f_2, 0, \nu_2)})$ is deterministic, we have

$$\frac{dP_T^{(f_1, 0, \nu_1)}}{dP_T^{(f_2, 0, \nu_2)}}(x) = \frac{dP_T^{(f_1 - f_2, 0, \nu_1)}}{dP_T^{(0, 0, \nu_2)}}(x) = D_T(x)$$

with $D_T(x)$ as in (2.11). Theorem 2.4 allows us to write the L_1 -distance between $P_T^{(f_1, 0, \nu_1)}$ and $P_T^{(f_2, 0, \nu_2)}$ as $\mathbb{E}_{P_T^{(f_2, 0, \nu_2)}} \left| 1 - D_T(x) \right|$. We then obtain the bound $2 \sinh(TL_1(\nu_1, \nu_2))$ by means of Lemma 3.3. □

References

- [1] Lawrence D. Brown and Mark G. Low. Asymptotic equivalence of nonparametric regression and white noise. *Ann. Statist.*, 24(6):2384–2398, 1996. MR-1425958
- [2] Andrew V. Carter. Deficiency distance between multinomial and multivariate normal experiments. *Ann. Statist.*, 30(3):708–730, 2002. Dedicated to the memory of Lucien Le Cam. MR-1922539

- [3] Rama Cont and Peter Tankov. *Financial modelling with jump processes*. Chapman & Hall/CRC Financial Mathematics Series. Chapman & Hall/CRC, Boca Raton, FL, 2004. MR-2042661
- [4] Arnak Dalalyan and Markus Reiß. Asymptotic statistical equivalence for scalar ergodic diffusions. *Probab. Theory Related Fields*, 134(2):248–282, 2006. MR-2222384
- [5] Jan Gairing, Michael Högele, Tetiana Kosenkova, and Alexei Kulik. Coupling distances between lévy measures and applications to noise sensitivity of sde. 2013.
- [6] Valentine Genon-Catalot and Catherine Laredo. Asymptotic equivalence of nonparametric diffusion and Euler scheme experiments. *Ann. Statist.*, to appear. MR-3224284
- [7] Alison L. Gibbs and Francis Edward Su. On choosing and bounding probability metrics. *Int. Stat. Rev.*, 70(3):419–435, 2002.
- [8] Jean Jacod and Albert N. Shiryaev. *Limit theorems for stochastic processes*, volume 288 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 2003. MR-1943877
- [9] Lucien Le Cam. *Asymptotic methods in statistical decision theory*. Springer Series in Statistics. Springer-Verlag, New York, 1986. MR-0856411
- [10] Lucien Le Cam and Grace Lo Yang. *Asymptotics in statistics*. Springer Series in Statistics. Springer-Verlag, New York, second edition, 2000. Some basic concepts. MR-1784901
- [11] J. Mémin and A. N. Shiryaev. Distance de Hellinger-Kakutani des lois correspondant à deux processus à accroissements indépendants. *Z. Wahrsch. Verw. Gebiete*, 70(1):67–89, 1985. MR-0795789
- [12] Grigori Milstein and Michael Nussbaum. Diffusion approximation for nonparametric autoregression. *Probab. Theory Related Fields*, 112(4):535–543, 1998. MR-1664703
- [13] Charles M. Newman. The inner product of path space measures corresponding to random processes with independent increments. *Bull. Amer. Math. Soc.*, 78:268–271, 1972. MR-0290453
- [14] Ivan Nourdin and Giovanni Peccati. Stein’s method on Wiener chaos. *Probab. Theory Related Fields*, 145(1-2):75–118, 2009. MR-2520122
- [15] Michael Nussbaum. Asymptotic equivalence of density estimation and Gaussian white noise. *Ann. Statist.*, 24(6):2399–2430, 1996. MR-1425959
- [16] Giovanni Peccati. The Chen-Stein method for Poisson functionals. arXiv:1112.5051
- [17] Nathan Ross. Fundamentals of Stein’s method. *Probab. Surv.*, 8:210–293, 2011. MR-2861132
- [18] Ken-iti Sato. *Lévy processes and infinitely divisible distributions*, volume 68 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999. Translated from the 1990 Japanese original, Revised by the author. MR-1739520
- [19] Ken-iti Sato. Density transformation in Lévy processes. 2000.