

Research Article

Iterative Algorithms Approach to Variational Inequalities and Fixed Point Problems

Yeong-Cheng Liou,¹ Yonghong Yao,² Chun-Wei Tseng,¹
Hui-To Lin,¹ and Pei-Xia Yang²

¹ Department of Information Management, Cheng Shiu University, Kaohsiung 833, Taiwan

² Department of Mathematics, Tianjin Polytechnic University, Tianjin 300387, China

Correspondence should be addressed to Yonghong Yao, yaoyonghong@yahoo.cn

Received 21 September 2011; Revised 16 November 2011; Accepted 17 November 2011

Academic Editor: Khalida Inayat Noor

Copyright © 2012 Yeong-Cheng Liou et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider a general variational inequality and fixed point problem, which is to find a point x^* with the property that (GVF): $x^* \in \text{GVI}(C, A)$ and $g(x^*) \in \text{Fix}(S)$ where $\text{GVI}(C, A)$ is the solution set of some variational inequality $\text{Fix}(S)$ is the fixed points set of nonexpansive mapping S , and g is a nonlinear operator. Assume the solution set Ω of (GVF) is nonempty. For solving (GVF), we suggest the following method $g(x_{n+1}) = \beta g(x_n) + (1 - \beta)SP_C[\alpha_n F(x_n) + (1 - \alpha_n)(g(x_n) - \lambda Ax_n)]$, $n \geq 0$. It is shown that the sequence $\{x_n\}$ converges strongly to $x^* \in \Omega$ which is the unique solution of the variational inequality $\langle F(x^*) - g(x^*), g(x) - g(x^*) \rangle \leq 0$, for all $x \in \Omega$.

1. Introduction

Let $A : C \rightarrow H$ and $g : C \rightarrow C$ be two nonlinear mappings. We concern the following generalized variational inequality of finding $u \in C$, $g(u) \in C$ such that

$$\langle g(v) - g(u), Au \rangle \geq 0, \quad \forall g(v) \in C. \quad (1.1)$$

The solution set of (1.1) is denoted by $\text{GVI}(C, A, g)$. It has been shown that a large class of unrelated odd-order and nonsymmetric obstacle, unilateral, contact, free, moving, and equilibrium problems arising in regional, physical, mathematical, engineering, and applied sciences can be studied in the unified and general framework of the general variational inequalities (1.1), see [1–16] and the references therein. Noor [17] has introduced a new type of variational inequality involving two nonlinear operators, which is called the general variational inequality. It is worth mentioning that this general variational inequality is

remarkably different from the so-called general variational inequality which was introduced by Noor [18] in 1988. Noor [17] proved that the general variational inequalities are equivalent to nonlinear projection equations and the Wiener-Hopf equations by using the projection technique. Using this equivalent formulation, Noor [17] suggested and analyzed some iterative algorithms for solving the special general variational inequalities and further proved that these algorithms have strong convergence.

For $g = I$, where I is the identity operator, problem (1.1) is equivalent to finding $u \in C$ such that

$$\langle v - u, Au \rangle \geq 0, \quad \forall v \in C, \quad (1.2)$$

which is known as the classical variational inequality introduced and studied by Stampacchia [19] in 1964. This field has been extensively studied due to a wide range of applications in industry, finance, economics, social, pure and applied sciences. For related works, please see [20–35]. Our main purposes in the present paper is devoted to study this topic.

Motivated and inspired by the works in this field, in this paper, we consider a general variational inequality and fixed point problem, which is to find a point x^* with the property that

$$x^* \in \text{GVI}(C, A), \quad g(x^*) \in \text{Fix}(S), \quad (\text{GVF})$$

where $\text{Fix}(S)$ is the fixed points set of nonexpansive mapping S . Assume the solution set Ω of (GVF) is nonempty. For solving (GVF), we suggest the following method

$$g(x_{n+1}) = \beta g(x_n) + (1 - \beta) SP_C [\alpha_n F(x_n) + (1 - \alpha_n)(g(x_n) - \lambda Ax_n)], \quad n \geq 0. \quad (1.3)$$

It is shown that the sequence $\{x_n\}$ converges strongly to $x^* \in \Omega$ which is the unique solution of the following variational inequality

$$\langle F(x^*) - g(x^*), g(x) - g(x^*) \rangle \leq 0, \quad \forall x \in \Omega. \quad (1.4)$$

Our results contain some interesting results as special cases.

2. Preliminaries

Let H be a real Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$, respectively. Let C be a nonempty closed convex subset of H . Recall that a mapping $S : C \rightarrow C$ is said to be nonexpansive if

$$\|Sx - Sy\| \leq \|x - y\|, \quad (2.1)$$

for all $x, y \in C$. We denote by $\text{Fix}(S)$ the set of fixed points of S . A mapping $F : C \rightarrow H$ is said to be L -Lipschitz continuous, if there exists a constant $L > 0$ such that $\|F(x) - F(y)\| \leq L\|x - y\|$

for all $x, y \in C$. A mapping $A : C \rightarrow H$ is said to be α -inverse strongly g -monotone if and only if

$$\langle Ax - Ay, g(x) - g(y) \rangle \geq \alpha \|Ax - Ay\|^2, \quad (2.2)$$

for some $\alpha > 0$ and for all $x, y \in C$. A mapping $g : C \rightarrow C$ is said to be strongly monotone if there exists a constant $\gamma > 0$ such that

$$\langle g(x) - g(y), x - y \rangle \geq \gamma \|x - y\|^2, \quad (2.3)$$

for all $x, y \in C$.

Let B be a mapping of H into 2^H . The effective domain of B is denoted by $\text{dom}(B)$, that is, $\text{dom}(B) = \{x \in H : Bx \neq \emptyset\}$. A multivalued mapping B is said to be a monotone operator on H if and only if

$$\langle x - y, u - v \rangle \geq 0, \quad (2.4)$$

for all $x, y \in \text{dom}(B)$, $u \in Bx$, and $v \in By$. A monotone operator B on H is said to be maximal if and only if its graph is not strictly contained in the graph of any other monotone operator on H . Let B be a maximal monotone operator on H and let $B^{-1}0 = \{x \in H : 0 \in Bx\}$.

It is well known that, for any $u \in H$, there exists a unique $u_0 \in C$ such that

$$\|u - u_0\| = \inf\{\|u - x\| : x \in C\}. \quad (2.5)$$

We denote u_0 by $P_C u$, where P_C is called the *metric projection* of H onto C . The metric projection P_C of H onto C has the following basic properties:

- (i) $\|P_C x - P_C y\| \leq \|x - y\|$ for all $x, y \in H$;
- (ii) $\langle x - y, P_C x - P_C y \rangle \geq \|P_C x - P_C y\|^2$ for every $x, y \in H$;
- (iii) $\langle x - P_C x, y - P_C x \rangle \leq 0$ for all $x \in H, y \in C$.

It is easy to see that the following is true:

$$u \in \text{GVI}(C, A, g) \iff g(u) = P_C(g(u) - \lambda A(u)), \quad \forall \lambda > 0. \quad (2.6)$$

We use the following notation:

- (i) $x_n \rightharpoonup x$ stands for the weak convergence of (x_n) to x ;
- (ii) $x_n \rightarrow x$ stands for the strong convergence of (x_n) to x .

We need the following lemmas for the next section.

Lemma 2.1. *Let C be a nonempty closed convex subset of a real Hilbert space H . Let $G : C \rightarrow C$ be a nonlinear mapping and let the mapping $A : C \rightarrow H$ be α -inverse strongly g -monotone. Then, for any $\lambda > 0$, one has*

$$\|P_C[g(x) - \lambda Ax] - P_C[g(y) - \lambda Ay]\|^2 \leq \|g(x) - g(y)\|^2 + \lambda(\lambda - 2\alpha)\|Ax - Ay\|^2, \quad x, y \in C. \quad (2.7)$$

Proof. Consider the following:

$$\begin{aligned}
& \|P_C[g(x) - \lambda Ax] - P_C[g(y) - \lambda Ay]\|^2 \\
& \leq \|g(x) - g(y) - \lambda(Ax - Ay)\|^2 \\
& = \|g(x) - g(y)\|^2 - 2\lambda\langle Ax - Ay, g(x) - g(y)\rangle + \lambda^2\|Ax - Ay\|^2 \quad (2.8) \\
& \leq \|g(x) - g(y)\|^2 - 2\lambda\alpha\|Ax - Ay\|^2 + \lambda^2\|Ax - Ay\|^2 \\
& \leq \|g(x) - g(y)\|^2 + \lambda(\lambda - 2\alpha)\|Ax - Ay\|^2.
\end{aligned}$$

If $\lambda \in [0, 2\alpha]$, we have

$$\|P_C[g(x) - \lambda Ax] - P_C[g(y) - \lambda Ay]\| \leq \|g(x) - g(y) - \lambda(Ax - Ay)\| \leq \|g(x) - g(y)\|. \quad (2.9)$$

□

Lemma 2.2 (see [36]). *Let C be a closed convex subset of a Hilbert space H . Let $S : C \rightarrow C$ be a nonexpansive mapping. Then $\text{Fix}(S)$ is a closed convex subset of C and the mapping $I - S$ is demiclosed at 0, that is, whenever $\{x_n\} \subset C$ is such that $x_n \rightarrow x$ and $(I - S)x_n \rightarrow 0$, then $(I - S)x = 0$.*

Lemma 2.3 (see [37]). *Let $\{x_n\}$ and $\{y_n\}$ be bounded sequences in a Banach space X and let $\{\beta_n\}$ be a sequence in $[0, 1]$ with $0 < \liminf_{n \rightarrow \infty} \beta_n \leq \limsup_{n \rightarrow \infty} \beta_n < 1$. Suppose $x_{n+1} = (1 - \beta_n)y_n + \beta_n x_n$ for all $n \geq 0$ and $\limsup_{n \rightarrow \infty} (\|y_{n+1} - y_n\| - \|x_{n+1} - x_n\|) \leq 0$. Then, $\lim_{n \rightarrow \infty} \|y_n - x_n\| = 0$.*

Lemma 2.4 (see [38]). *Assume $\{a_n\}$ is a sequence of nonnegative real numbers such that*

$$a_{n+1} \leq (1 - \gamma_n)a_n + \delta_n \gamma_n, \quad (2.10)$$

where $\{\gamma_n\}$ is a sequence in $(0, 1)$ and $\{\delta_n\}$ is a sequence such that

- (1) $\sum_{n=1}^{\infty} \gamma_n = \infty$;
- (2) $\limsup_{n \rightarrow \infty} \delta_n \leq 0$ or $\sum_{n=1}^{\infty} |\delta_n \gamma_n| < \infty$.

Then $\lim_{n \rightarrow \infty} a_n = 0$.

3. Main Results

In this section, we will prove our main results.

Theorem 3.1. *Let C be a nonempty closed and convex subset of a real Hilbert space H . Let $F : C \rightarrow H$ be an L -Lipschitz continuous mapping, $g : C \rightarrow C$ be a weakly continuous and γ -strongly monotone mapping such that $R(g) = C$. Let $A : C \rightarrow H$ be an α -inverse strongly g -monotone mapping and let $S : C \rightarrow C$ be a nonexpansive mapping. Suppose that $\Omega \neq \emptyset$. Let $\beta \in (0, 1)$ and $\gamma \in (L, 2\alpha)$. For given $x_0 \in C$, let $\{x_n\} \subset C$ be a sequence generated by*

$$g(x_{n+1}) = \beta g(x_n) + (1 - \beta)SP_C[\alpha_n F(x_n) + (1 - \alpha_n)(g(x_n) - \lambda Ax_n)], \quad n \geq 0, \quad (3.1)$$

where $\{\alpha_n\} \subset (0, 1)$ satisfies (C1): $\lim_{n \rightarrow \infty} \alpha_n = 0$ and (C2): $\sum_n \alpha_n = \infty$. Then the sequence $\{x_n\}$ generated by (3.1) converges strongly to $x^* \in \Omega$ which is the unique solution of the following variational inequality:

$$\langle F(x^*) - g(x^*), g(x) - g(x^*) \rangle \leq 0, \quad \forall x \in \Omega. \quad (3.2)$$

Proof. First, we show the solution set of variational inequality (3.2) is singleton. Assume $\tilde{x} \in \Omega$ also solves (3.2). Then, we have

$$\langle F(x^*) - g(x^*), g(\tilde{x}) - g(x^*) \rangle \leq 0, \quad \langle F(\tilde{x}) - g(\tilde{x}), g(x^*) - g(\tilde{x}) \rangle \leq 0. \quad (3.3)$$

It follows that

$$\begin{aligned} & \langle F(\tilde{x}) - g(\tilde{x}) - F(x^*) + g(x^*), g(x^*) - g(\tilde{x}) \rangle \leq 0 \\ & \implies \|g(x^*) - g(\tilde{x})\|^2 \leq \langle F(x^*) - F(\tilde{x}), g(x^*) - g(\tilde{x}) \rangle \\ & \implies \|g(x^*) - g(\tilde{x})\|^2 \leq \langle F(x^*) - F(\tilde{x}), g(x^*) - g(\tilde{x}) \rangle \leq \|F(x^*) - F(\tilde{x})\| \|g(x^*) - g(\tilde{x})\| \\ & \implies \|g(x^*) - g(\tilde{x})\| \leq \|F(x^*) - F(\tilde{x})\|. \end{aligned} \quad (3.4)$$

Since g is γ -strongly monotone, we have

$$\gamma \|x - y\|^2 \leq \langle g(x) - g(y), x - y \rangle \leq \|g(x) - g(y)\| \|x - y\|, \quad \forall x, y \in C. \quad (3.5)$$

Hence,

$$\gamma \|x - y\| \leq \|g(x) - g(y)\|, \quad \forall x, y \in C. \quad (3.6)$$

In particular, $\gamma \|x^* - \tilde{x}\| \leq \|g(x^*) - g(\tilde{x})\|$. By (3.4), we deduce

$$\gamma \|x^* - \tilde{x}\| \leq \|g(x^*) - g(\tilde{x})\| \leq \|F(x^*) - F(\tilde{x})\| \leq L \|x^* - \tilde{x}\|, \quad (3.7)$$

which implies that $\tilde{x} = x^*$ because of $L < \gamma$ by the assumption. Therefore, the solution of variational inequality (3.2) is unique.

Pick up any $u \in \Omega$. It is obvious that $u \in \text{GVI}(C, A, g)$ and $g(u) \in \text{Fix}(S)$. Set $u_n = P_C[\alpha_n F(x_n) + (1 - \alpha_n)(g(x_n) - \lambda A x_n)]$, $n \geq 0$. From (2.6), we know $g(u) = P_C[g(u) - \mu A u]$ for any $\mu > 0$. Hence, we have

$$g(u) = P_C[g(u) - (1 - \alpha_n)\lambda A u] = P_C[\alpha_n g(u) + (1 - \alpha_n)(g(u) - \lambda A u)], \quad \forall n \geq 0. \quad (3.8)$$

From (3.6), (3.8), and Lemma 2.1, we get

$$\begin{aligned}
\|u_n - g(u)\| &= \|P_C[\alpha_n F(x_n) + (1 - \alpha_n)(g(x_n) - \lambda Ax_n)] \\
&\quad - P_C[\alpha_n g(u) + (1 - \alpha_n)(g(u) - \lambda Au)]\| \\
&\leq \alpha_n \|F(x_n) - g(u)\| + (1 - \alpha_n) \|(g(x_n) - \lambda Ax_n) - (g(u) - \lambda Au)\| \\
&\leq \alpha_n \|F(x_n) - F(u)\| + \alpha_n \|F(u) - g(u)\| + (1 - \alpha_n) \|g(x_n) - g(u)\| \\
&\leq \alpha_n L \|x_n - u\| + \alpha_n \|F(u) - g(u)\| + (1 - \alpha_n) \|g(x_n) - g(u)\| \\
&\leq \frac{\alpha_n L}{\gamma} \|g(x_n) - g(u)\| + \alpha_n \|F(u) - g(u)\| + (1 - \alpha_n) \|g(x_n) - g(u)\| \\
&= \left[1 - \left(1 - \frac{L}{\gamma}\right)\alpha_n\right] \|g(x_n) - g(u)\| + \alpha_n \|F(u) - g(u)\|.
\end{aligned} \tag{3.9}$$

It follows from (3.1) that

$$\begin{aligned}
\|g(x_{n+1}) - g(u)\| &\leq \beta \|g(x_n) - g(u)\| + (1 - \beta) \|Su_n - Sg(u)\| \\
&\leq \beta \|g(x_n) - g(u)\| + (1 - \beta) \|u_n - g(u)\| \\
&\leq \beta \|g(x_n) - g(u)\| + (1 - \beta) \left[1 - \left(1 - \frac{L}{\gamma}\right)\alpha_n\right] \|g(x_n) - g(u)\| \\
&\quad + (1 - \beta)\alpha_n \|F(u) - g(u)\| \\
&= \left[1 - \left(1 - \frac{L}{\gamma}\right)(1 - \beta)\alpha_n\right] \|g(x_n) - g(u)\| \\
&\quad + \left(1 - \frac{L}{\gamma}\right)(1 - \beta)\alpha_n \frac{\|F(u) - g(u)\|}{1 - L/\gamma}.
\end{aligned} \tag{3.10}$$

This indicates by induction that

$$\|g(x_{n+1}) - g(u)\| \leq \max \left\{ \|g(x_n) - g(u)\|, \frac{\|F(u) - g(u)\|}{1 - L/\gamma} \right\}. \tag{3.11}$$

Hence, $\{g(x_n)\}$ is bounded. By (3.6), we have $\|x_n - u\| \leq (1/\gamma)\|g(x_n) - g(u)\|$. This implies that $\{x_n\}$ is bounded. Consequently, $\{F(x_n)\}$, $\{Ax_n\}$, $\{u_n\}$, and $\{Su_n\}$ are all bounded.

Note that we can rewrite (3.1) as $g(x_{n+1}) = \beta g(x_n) + (1 - \beta)Su_n$ for all n . Next, we will use Lemma 2.3 to prove that $\|x_{n+1} - x_n\| \rightarrow 0$. In fact, we firstly have

$$\begin{aligned}
\|Su_n - Su_{n-1}\| &= \|SP_C[\alpha_n F(x_n) + (1 - \alpha_n)(g(x_n) - \lambda Ax_n)] \\
&\quad - SP_C[\alpha_{n-1} F(x_{n-1}) + (1 - \alpha_{n-1})(g(x_{n-1}) - \lambda Ax_{n-1})]\| \\
&\leq \|[\alpha_n F(x_n) + (1 - \alpha_n)(g(x_n) - \lambda Ax_n)] \\
&\quad - [\alpha_{n-1} F(x_{n-1}) + (1 - \alpha_{n-1})(g(x_{n-1}) - \lambda Ax_{n-1})]\| \\
&\leq \alpha_n \|F(x_n) - F(x_{n-1})\| + |\alpha_n - \alpha_{n-1}| \|F(x_{n-1})\|
\end{aligned}$$

$$\begin{aligned}
 & + (1 - \alpha_n) \|g(x_n) - \lambda Ax_n - (g(x_{n-1}) - \lambda Ax_{n-1})\| \\
 & + |\alpha_n - \alpha_{n-1}| \|g(x_{n-1}) - \lambda Ax_{n-1}\| \\
 \leq & \alpha_n L \|x_n - x_{n-1}\| + (1 - \alpha_n) \|g(x_n) - g(x_{n-1})\| \\
 & + |\alpha_n - \alpha_{n-1}| (\|F(x_{n-1})\| + \|g(x_{n-1}) - \lambda Ax_{n-1}\|) \\
 \leq & \alpha_n \left(\frac{L}{\gamma}\right) \|g(x_n) - g(x_{n-1})\| + (1 - \alpha_n) \|g(x_n) - g(x_{n-1})\| \\
 & + |\alpha_n - \alpha_{n-1}| (\|F(x_{n-1})\| + \|g(x_{n-1}) - \lambda Ax_{n-1}\|) \\
 = & \left[1 - \left(1 - \frac{L}{\gamma}\right) \alpha_n\right] \|g(x_n) - g(x_{n-1})\| \\
 & + |\alpha_n - \alpha_{n-1}| (\|F(x_{n-1})\| + \|g(x_{n-1}) - \lambda Ax_{n-1}\|).
 \end{aligned} \tag{3.12}$$

It follows that

$$\|Su_n - Su_{n-1}\| - \|g(x_n) - g(x_{n-1})\| \leq |\alpha_n - \alpha_{n-1}| (\|F(x_{n-1})\| + \|g(x_{n-1}) - \lambda Ax_{n-1}\|). \tag{3.13}$$

Since $\alpha_n \rightarrow 0$ and the sequences $\{F(x_n)\}$, $\{g(x_n)\}$, and $\{Ax_n\}$ are bounded, we have

$$\limsup_{n \rightarrow \infty} (\|Su_n - Su_{n-1}\| - \|g(x_n) - g(x_{n-1})\|) \leq 0. \tag{3.14}$$

By Lemma 2.3, we obtain

$$\lim_{n \rightarrow \infty} \|Su_n - g(x_n)\| = 0. \tag{3.15}$$

Hence,

$$\lim_{n \rightarrow \infty} \|g(x_{n+1}) - g(x_n)\| = \lim_{n \rightarrow \infty} (1 - \beta) \|Su_n - g(x_n)\| = 0. \tag{3.16}$$

This together with (3.6) imply that

$$\lim_{n \rightarrow \infty} \|x_{n+1} - x_n\| = 0. \tag{3.17}$$

By the convexity of the norm and (3.9), we have

$$\begin{aligned}
 \|g(x_{n+1}) - g(u)\|^2 & = \|\beta(g(x_n) - g(u)) + (1 - \beta)(Su_n - Sg(u))\|^2 \\
 & \leq \beta \|g(x_n) - g(u)\|^2 + (1 - \beta) \|Su_n - Sg(u)\|^2 \\
 & \leq \beta \|g(x_n) - g(u)\|^2 + (1 - \beta) \|u_n - g(u)\|^2
 \end{aligned}$$

$$\begin{aligned}
&\leq \beta \|g(x_n) - g(u)\|^2 \\
&\quad + (1 - \beta) [\alpha_n \|F(x_n) - g(u)\| \\
&\quad\quad + (1 - \alpha_n) \|(g(x_n) - \lambda Ax_n) - (g(u) - \lambda Au)\|]^2 \\
&\leq \beta \|g(x_n) - g(u)\|^2 \\
&\quad + (1 - \beta) [\alpha_n \|F(x_n) - g(u)\|^2 \\
&\quad\quad + (1 - \alpha_n) \|(g(x_n) - \lambda Ax_n) - (g(u) - \lambda Au)\|^2].
\end{aligned} \tag{3.18}$$

From Lemma 2.1, we derive

$$\|(g(x_n) - \lambda Ax_n) - (g(u) - \lambda Au)\|^2 \leq \|g(x_n) - g(u)\|^2 + \lambda(\lambda - 2\alpha) \|Ax_n - Au\|^2. \tag{3.19}$$

Thus,

$$\begin{aligned}
\|g(x_{n+1}) - g(u)\|^2 &\leq \beta \|g(x_n) - g(u)\|^2 \\
&\quad + (1 - \beta) [\alpha_n \|F(x_n) - g(u)\|^2 \\
&\quad\quad + (1 - \alpha_n) (\|g(x_n) - g(u)\|^2 + \lambda(\lambda - 2\alpha) \|Ax_n - Au\|^2)] \\
&= (1 - \beta) \alpha_n \|F(x_n) - g(u)\|^2 + [1 - (1 - \beta) \alpha_n] \|g(x_n) - g(u)\|^2 \\
&\quad + (1 - \beta) (1 - \alpha_n) \lambda(\lambda - 2\alpha) \|Ax_n - Au\|^2.
\end{aligned} \tag{3.20}$$

So,

$$\begin{aligned}
&(1 - \beta) (1 - \alpha_n) \lambda(2\alpha - \lambda) \|Ax_n - Au\|^2 \\
&\leq (1 - \beta) \alpha_n \|F(x_n) - g(u)\|^2 + \|g(x_n) - g(u)\|^2 - \|g(x_{n+1}) - g(u)\|^2 \\
&\leq (1 - \beta) \alpha_n \|F(x_n) - g(u)\|^2 \\
&\quad + (\|g(x_n) - g(u)\| + \|g(x_{n+1}) - g(u)\|) \|g(x_{n+1}) - g(x_n)\|.
\end{aligned} \tag{3.21}$$

Since $\alpha_n \rightarrow 0$, $\|g(x_{n+1}) - g(x_n)\| \rightarrow 0$ and $\liminf_{n \rightarrow \infty} (1 - \beta) (1 - \alpha_n) \lambda(2\alpha - \lambda) > 0$, we obtain

$$\lim_{n \rightarrow \infty} \|Ax_n - Au\| = 0. \tag{3.22}$$

Set $y_n = g(x_n) - \lambda Ax_n - (g(u) - \lambda Au)$ for all n . By using the property of projection, we get

$$\begin{aligned}
\|u_n - g(u)\|^2 &= \|P_C[\alpha_n F(x_n) + (1 - \alpha_n)(g(x_n) - \lambda Ax_n)] \\
&\quad - P_C[\alpha_n g(u) + (1 - \alpha_n)(g(u) - \lambda Au)]\|^2 \\
&\leq \langle \alpha_n(F(x_n) - g(u)) + (1 - \alpha_n)y_n, u_n - g(u) \rangle \\
&= \frac{1}{2} \left\{ \|\alpha_n(F(x_n) - g(u)) + (1 - \alpha_n)y_n\|^2 + \|u_n - g(u)\|^2 \right. \\
&\quad \left. - \|\alpha_n(F(x_n) - g(u)) + (1 - \alpha_n)y_n - u_n + g(u)\|^2 \right\} \\
&\leq \frac{1}{2} \left\{ \alpha_n \|F(x_n) - g(u)\|^2 + (1 - \alpha_n) \|g(x_n) - g(u)\|^2 + \|u_n - g(u)\|^2 \right. \\
&\quad \left. - \|\alpha_n(F(x_n) - g(u) - y_n) + g(x_n) - u_n - \lambda(Ax_n - Au)\|^2 \right\} \\
&= \frac{1}{2} \left\{ \alpha_n \|F(x_n) - g(u)\|^2 + (1 - \alpha_n) \|g(x_n) - g(u)\|^2 + \|u_n - g(u)\|^2 \right. \\
&\quad - \|g(x_n) - u_n\|^2 - \lambda^2 \|Ax_n - Au\|^2 - \alpha_n^2 \|F(x_n) - g(u) - y_n\|^2 \\
&\quad + 2\lambda \alpha_n \langle Ax_n - Au, F(x_n) - g(u) - y_n \rangle + 2\lambda \langle g(x_n) - u_n, Ax_n - Au \rangle \\
&\quad \left. - 2\alpha_n \langle g(x_n) - u_n, F(x_n) - g(u) - y_n \rangle \right\}. \tag{3.23}
\end{aligned}$$

It follows that

$$\begin{aligned}
\|u_n - g(u)\|^2 &\leq \alpha_n \|F(x_n) - g(u)\|^2 + (1 - \alpha_n) \|g(x_n) - g(u)\|^2 - \|g(x_n) - u_n\|^2 \\
&\quad + 2\lambda \alpha_n \|Ax_n - Au\| \|F(x_n) - g(u) - y_n\| + 2\lambda \|g(x_n) - u_n\| \|Ax_n - Au\| \tag{3.24} \\
&\quad + 2\alpha_n \|g(x_n) - u_n\| \|F(x_n) - g(u) - y_n\|.
\end{aligned}$$

From (3.18) and (3.24), we have

$$\begin{aligned}
\|g(x_{n+1}) - g(u)\|^2 &\leq \beta \|g(x_n) - g(u)\|^2 + (1 - \beta) \|u_n - g(u)\|^2 \\
&\leq \beta \|g(x_n) - g(u)\|^2 + (1 - \beta) \alpha_n \|F(x_n) - g(u)\|^2 \\
&\quad + (1 - \alpha_n)(1 - \beta) \|g(x_n) - g(u)\|^2 - (1 - \beta) \|g(x_n) - u_n\|^2 \\
&\quad + 2\lambda(1 - \beta) \alpha_n \|Ax_n - Au\| \|F(x_n) - g(u) - y_n\| \\
&\quad + 2\lambda(1 - \beta) \|g(x_n) - u_n\| \|Ax_n - Au\|
\end{aligned}$$

$$\begin{aligned}
& + 2(1 - \beta)\alpha_n \|g(x_n) - u_n\| \|F(x_n) - g(u) - y_n\| \\
\leq & \|g(x_n) - g(u)\|^2 + \alpha_n \|F(x_n) - g(u)\|^2 - (1 - \beta) \|g(x_n) - u_n\|^2 \\
& + 2\lambda\alpha_n \|Ax_n - Au\| \|F(x_n) - g(u) - y_n\| + 2\lambda \|g(x_n) - u_n\| \|Ax_n - Au\| \\
& + 2\alpha_n \|g(x_n) - u_n\| \|F(x_n) - g(u) - y_n\|. \tag{3.25}
\end{aligned}$$

Then, we obtain

$$\begin{aligned}
(1 - \beta) \|g(x_n) - u_n\|^2 \leq & (\|g(x_n) - g(u)\| + \|g(x_{n+1}) - g(u)\|) \|g(x_{n+1}) - g(x_n)\| \\
& + \alpha_n \|F(x_n) - g(u)\|^2 + 2\lambda\alpha_n \|Ax_n - Au\| \|F(x_n) - g(u) - y_n\| \\
& + 2\lambda \|g(x_n) - u_n\| \|Ax_n - Au\| + 2\alpha_n \|g(x_n) - u_n\| \|F(x_n) - g(u) - y_n\|. \tag{3.26}
\end{aligned}$$

Since $\lim_{n \rightarrow \infty} \alpha_n = 0$, $\lim_{n \rightarrow \infty} \|g(x_{n+1}) - g(x_n)\| = 0$ and $\lim_{n \rightarrow \infty} \|Ax_n - Au\| = 0$, we have

$$\lim_{n \rightarrow \infty} \|g(x_n) - u_n\| = 0. \tag{3.27}$$

Next, we prove $\limsup_{n \rightarrow \infty} \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle \leq 0$ where x^* is the unique solution of (3.2). We take a subsequence $\{u_{n_i}\}$ of $\{u_n\}$ such that

$$\begin{aligned}
\limsup_{n \rightarrow \infty} \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle & = \lim_{i \rightarrow \infty} \langle F(x^*) - g(x^*), u_{n_i} - g(x^*) \rangle \\
& = \lim_{i \rightarrow \infty} \langle F(x^*) - g(x^*), g(x_{n_i}) - g(x^*) \rangle. \tag{3.28}
\end{aligned}$$

Since $\{x_{n_i}\}$ is bounded, there exists a subsequence $\{x_{n_{i_j}}\}$ of $\{x_{n_i}\}$ which converges weakly to some point $z \in C$. Without loss of generality, we may assume that $x_{n_{i_j}} \rightharpoonup z$. This implies that $g(x_{n_{i_j}}) \rightharpoonup g(z)$ due to the weak continuity of g . Now, we show $z \in \Omega$. First, we note that from (3.15) and (3.27) that $\|g(x_n) - Sg(x_n)\| \rightarrow 0$. Hence, $\lim_{i \rightarrow \infty} \|g(x_{n_i}) - Sg(x_{n_i})\| = 0$. By the demiclosedness principle of the nonexpansive mapping (see Lemma 2.2), we deduce $g(z) \in \text{Fix}(S)$. Next, we only need to prove $z \in \text{GVI}(C, A, g)$. Set

$$Tv = \begin{cases} Av + N_C(v), & v \in C, \\ \emptyset, & v \notin C. \end{cases} \tag{3.29}$$

By [39], we know that T is maximal g -monotone. Let $(v, w) \in G(T)$. Since $w - Av \in N_C(v)$ and $x_n \in C$, we have

$$\langle g(v) - g(x_n), w - Av \rangle \geq 0. \tag{3.30}$$

From $u_n = P_C[\alpha_n F(x_n) + (1 - \alpha_n)(g(x_n) - \lambda Ax_n)]$, we get

$$\langle g(v) - u_n, u_n - [\alpha_n F(x_n) + (1 - \alpha_n)(g(x_n) - \lambda Ax_n)] \rangle \geq 0. \quad (3.31)$$

It follows that

$$\left\langle g(v) - u_n, \frac{u_n - g(x_n)}{\lambda} + Ax_n - \frac{\alpha_n}{\lambda} (F(x_n) - g(x_n) + \lambda Ax_n) \right\rangle \geq 0. \quad (3.32)$$

Then,

$$\begin{aligned} \langle g(v) - g(x_{n_i}), w \rangle &\geq \langle g(v) - g(x_{n_i}), Av \rangle \\ &\geq \langle g(v) - g(x_{n_i}), Av \rangle - \left\langle g(v) - u_{n_i}, \frac{u_{n_i} - g(x_{n_i})}{\lambda} \right\rangle \\ &\quad - \langle g(v) - u_{n_i}, Ax_{n_i} \rangle + \frac{\alpha_{n_i}}{\lambda} \langle g(v) - u_{n_i}, F(x_{n_i}) - g(x_{n_i}) + \lambda Ax_{n_i} \rangle \\ &= \langle g(v) - g(x_{n_i}), Av - Ax_{n_i} \rangle + \langle g(v) - g(x_{n_i}), -Ax_{n_i} \rangle \\ &\quad - \left\langle g(v) - u_{n_i}, \frac{u_{n_i} - g(x_{n_i})}{\lambda} \right\rangle - \langle g(v) - u_{n_i}, Ax_{n_i} \rangle \\ &\quad + \frac{\alpha_{n_i}}{\lambda} \langle g(v) - u_{n_i}, F(x_{n_i}) - g(x_{n_i}) + \lambda Ax_{n_i} \rangle \\ &\geq - \left\langle g(v) - u_{n_i}, \frac{u_{n_i} - g(x_{n_i})}{\lambda} \right\rangle - \langle g(x_{n_i}) - u_{n_i}, Ax_{n_i} \rangle \\ &\quad + \frac{\alpha_{n_i}}{\lambda} \langle g(v) - u_{n_i}, F(x_{n_i}) - g(x_{n_i}) + \lambda Ax_{n_i} \rangle. \end{aligned} \quad (3.33)$$

Since $\|g(x_{n_i}) - u_{n_i}\| \rightarrow 0$ and $g(x_{n_i}) \rightarrow g(z)$, we deduce that $\langle g(v) - g(z), w \rangle \geq 0$ by taking $i \rightarrow \infty$ in (3.33). Thus, $z \in T^{-1}0$ by the maximal g -monotonicity of T . Hence, $z \in \text{GVI}(C, A, g)$. Therefore, $z \in \Omega$. From (3.28), we obtain

$$\begin{aligned} \limsup_{n \rightarrow \infty} \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle &= \lim_{i \rightarrow \infty} \langle F(x^*) - g(x^*), g(x_{n_i}) - g(x^*) \rangle \\ &= \langle F(x^*) - g(x^*), g(z) - g(x^*) \rangle \leq 0. \end{aligned} \quad (3.34)$$

We take $u = x^*$ in (3.23) to get

$$\begin{aligned}
\|u_n - g(x^*)\|^2 &\leq \alpha_n \langle F(x_n) - g(x^*), u_n - g(x^*) \rangle \\
&\quad + (1 - \alpha_n) \langle g(x_n) - \lambda Ax_n - (g(x^*) - \lambda Ax^*), u_n - g(x^*) \rangle \\
&\leq \alpha_n \langle F(x_n) - F(x^*), u_n - g(x^*) \rangle + \alpha_n \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle \\
&\quad + (1 - \alpha_n) \|g(x_n) - \lambda Ax_n - (g(x^*) - \lambda Ax^*)\| \|u_n - g(x^*)\| \\
&\leq \alpha_n L \|x_n - x^*\| \|u_n - g(x^*)\| + \alpha_n \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle \\
&\quad + (1 - \alpha_n) \|g(x_n) - g(x^*)\| \|u_n - g(x^*)\| \\
&\leq \alpha_n \left(\frac{L}{\gamma} \right) \|g(x_n) - g(x^*)\| \|u_n - g(x^*)\| + \alpha_n \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle \\
&\quad + (1 - \alpha_n) \|g(x_n) - g(x^*)\| \|u_n - g(x^*)\| \\
&= \left[1 - \left(1 - \frac{L}{\gamma} \right) \alpha_n \right] \|g(x_n) - g(x^*)\| \|u_n - g(x^*)\| \\
&\quad + \alpha_n \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle \\
&= \frac{1 - (1 - L/\gamma)\alpha_n}{2} \|g(x_n) - g(x^*)\|^2 + \frac{1}{2} \|u_n - g(x^*)\|^2 \\
&\quad + \alpha_n \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle. \tag{3.35}
\end{aligned}$$

It follows that

$$\|u_n - g(x^*)\|^2 \leq \left[1 - \left(1 - \frac{L}{\gamma} \right) \alpha_n \right] \|g(x_n) - g(x^*)\|^2 + 2\alpha_n \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle. \tag{3.36}$$

Therefore,

$$\begin{aligned}
\|g(x_{n+1}) - g(x^*)\|^2 &\leq \beta \|g(x_n) - g(x^*)\|^2 + (1 - \beta) \|u_n - g(x^*)\|^2 \\
&\leq \beta \|g(x_n) - g(x^*)\|^2 + (1 - \beta) \left[1 - \left(1 - \frac{L}{\gamma} \right) \alpha_n \right] \|g(x_n) - g(x^*)\|^2 \\
&\quad + 2(1 - \beta) \alpha_n \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle \\
&= \left[1 - \left(1 - \frac{L}{\gamma} \right) (1 - \beta) \alpha_n \right] \|g(x_n) - g(x^*)\|^2 \\
&\quad + 2(1 - \beta) \alpha_n \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle \\
&= \left[1 - \left(1 - \frac{L}{\gamma} \right) (1 - \beta) \alpha_n \right] \|g(x_n) - g(x^*)\|^2
\end{aligned}$$

$$\begin{aligned}
& + \left(1 - \frac{L}{\gamma}\right)(1 - \beta)\alpha_n \left(\frac{2}{1 - L/\gamma} \langle F(x^*) - g(x^*), u_n - g(x^*) \rangle\right) \\
& = (1 - \gamma_n) \|g(x_n) - g(x^*)\|^2 + \delta_n \gamma_n,
\end{aligned} \tag{3.37}$$

where $\gamma_n = (1 - L/\gamma)(1 - \beta)\alpha_n$ and $\delta_n = (2/(1 - L/\gamma))\langle F(x^*) - g(x^*), u_n - g(x^*) \rangle$. From condition (C2), we have $\sum_n \gamma_n = \infty$. By (3.34), we have $\limsup_{n \rightarrow \infty} \delta_n \leq 0$. We can therefore apply Lemma 2.4 to conclude that $g(x_n) \rightarrow g(x^*)$ and $x_n \rightarrow x^*$. This completes the proof. \square

Corollary 3.2. *Let C be a nonempty closed and convex subset of a real Hilbert space H . Let $F : C \rightarrow H$ be an L -contraction. Let $A : C \rightarrow H$ be an α -inverse strongly monotone mapping and let $S : C \rightarrow C$ be a nonexpansive mapping. Suppose that $\Omega \neq \emptyset$. Let $\beta \in (0, 1)$ and $\gamma \in (L, 2\alpha)$. For given $x_0 \in C$, let $\{x_n\} \subset C$ be a sequence generated by*

$$x_{n+1} = \beta x_n + (1 - \beta)SP_C[\alpha_n F(x_n) + (1 - \alpha_n)(x_n - \lambda A x_n)], \quad n \geq 0, \tag{3.38}$$

where $\{\alpha_n\} \subset (0, 1)$ satisfies (C1): $\lim_{n \rightarrow \infty} \alpha_n = 0$ and (C2): $\sum_n \alpha_n = \infty$. Then the sequence $\{x_n\}$ generated by (3.38) converges strongly to $x^* \in \Omega$ which is the unique solution of the following variational inequality:

$$\langle F(x^*) - x^*, x - x^* \rangle \leq 0, \quad \forall x \in \Omega. \tag{3.39}$$

Acknowledgments

The authors are very grateful to the referees for their comments and suggestions which improved the presentation of this paper. The first author was partially supported by the Program TH-1-3, Optimization Lean Cycle, of Subprojects TH-1 of Spindle Plan Four in Excellence Teaching and Learning Plan of Cheng Shiu University and was supported in part by NSC 100-2221-E-230-012. The second author was supported in part by Colleges and Universities Science and Technology Development Foundation (20091003) of Tianjin, NSFC 11071279 and NSFC 71161001-G0105.

References

- [1] M. Fukushima, "Equivalent differentiable optimization problems and descent methods for asymmetric variational inequality problems," *Mathematical Programming*, vol. 53, no. 1, pp. 99–110, 1992.
- [2] F. Giannessi and A. Maugeri, *Variational Inequalities and Network Equilibrium Problems*, New York, NY, USA, Plenum Press, 1995.
- [3] F. Giannessi, A. Maugeri, and P. M. Pardalos, *Equilibrium Problems: Nonsmooth Optimization and Variational Inequality Models*, Kluwer Academic, Dodrecht, The Netherlands, 2001.
- [4] R. Glowinski, *Numerical Methods for Nonlinear Variational Problems*, Springer, Berlin, Germany, 1984.
- [5] R. Glowinski, J.-L. Lions, and R. Trémolières, *Numerical Analysis of Variational Inequalities*, vol. 8, North-Holland, Amsterdam, The Netherlands, 1981.
- [6] R. Glowinski and P. Le Tallec, *Augmented Lagrangian and Operator-Splitting Methods in Nonlinear Mechanics*, vol. 9, Society for Industrial and Applied Mathematics, Philadelphia, Pa, USA, 1989.
- [7] D. Han and H. K. Lo, "Two new self-adaptive projection methods for variational inequality problems," *Computers & Mathematics with Applications*, vol. 43, no. 12, pp. 1529–1537, 2002.

- [8] P. T. Harker and J.-S. Pang, "Finite-dimensional variational inequality and nonlinear complementarity problems: a survey of theory, algorithms and applications," *Mathematical Programming*, vol. 48, no. 2, pp. 161–220, 1990.
- [9] Y. Yao, Y. J. Cho, and Y.-C. Liou, "Algorithms of common solutions for variational inclusions, mixed equilibrium problems and fixed point problems," *European Journal of Operational Research*, vol. 212, no. 2, pp. 242–250, 2011.
- [10] B. He, "Inexact implicit methods for monotone general variational inequalities," *Mathematical Programming*, vol. 86, no. 1, pp. 199–217, 1999.
- [11] J.-S. Pang and J. C. Yao, "On a generalization of a normal map and equation," *SIAM Journal on Control and Optimization*, vol. 33, no. 1, pp. 168–184, 1995.
- [12] J. C. Yao, "Variational inequalities with generalized monotone operators," *Mathematics of Operations Research*, vol. 19, no. 3, pp. 691–705, 1994.
- [13] Y. Yao and S. Maruster, "Strong convergence of an iterative algorithm for variational inequalities in Banach spaces," *Mathematical and Computer Modelling*, vol. 54, no. 1-2, pp. 325–329, 2011.
- [14] Y. Yao, Y.-C. Liou, S. M. Kang, and Y. Yu, "Algorithms with strong convergence for a system of nonlinear variational inequalities in Banach spaces," *Nonlinear Analysis*, vol. 74, no. 17, pp. 6024–6034, 2011.
- [15] F. Cianciaruso, G. Marino, L. Muglia, and Y. Yao, "A hybrid projection algorithm for finding solutions of mixed equilibrium problem and variational inequality problem," *Fixed Point Theory and Applications*, Article ID 383740, 19 pages, 2010.
- [16] M. Aslam Noor, "Some developments in general variational inequalities," *Applied Mathematics and Computation*, vol. 152, no. 1, pp. 199–277, 2004.
- [17] M. A. Noor, "Differentiable non-convex functions and general variational inequalities," *Applied Mathematics and Computation*, vol. 199, no. 2, pp. 623–630, 2008.
- [18] M. A. Noor, "General variational inequalities," *Applied Mathematics Letters*, vol. 1, no. 2, pp. 119–122, 1988.
- [19] G. Stampacchia, "Formes bilinéaires coercitives sur les ensembles convexes," *Comptes Rendus de l'Académie des Sciences*, vol. 258, pp. 4413–4416, 1964.
- [20] M. A. Noor, "New approximation schemes for general variational inequalities," *Journal of Mathematical Analysis and Applications*, vol. 251, no. 1, pp. 217–229, 2000.
- [21] F. Cianciaruso, G. Marino, L. Muglia, and Y. Yao, "On a two-step algorithm for hierarchical fixed point problems and variational inequalities," *Journal of Inequalities and Applications*, Article ID 208692, 13 pages, 2009.
- [22] Y. Yao, R. Chen, and H.-K. Xu, "Schemes for finding minimum-norm solutions of variational inequalities," *Nonlinear Analysis*, vol. 72, no. 7-8, pp. 3447–3456, 2010.
- [23] M. A. Noor, "Some algorithms for general monotone mixed variational inequalities," *Mathematical and Computer Modelling*, vol. 29, no. 7, pp. 1–9, 1999.
- [24] M. A. Noor and K. I. Noor, "Self-adaptive projection algorithms for general variational inequalities," *Applied Mathematics and Computation*, vol. 151, no. 3, pp. 659–670, 2004.
- [25] M. A. Noor, K. I. Noor, and T. M. Rassias, "Some aspects of variational inequalities," *Journal of Computational and Applied Mathematics*, vol. 47, no. 3, pp. 285–312, 1993.
- [26] Y. Yao, Y. C. Liou, and S. M. Kang, "Two-step projection methods for a system of variational inequality problems in Banach spaces," *Journal of Global Optimization*.
- [27] Y. Yao, R. Chen, and Y. C. Liou, "A unified implicit algorithm for solving the triplehierarchical constrained optimization problem," *Mathematical and Computer Modelling*.
- [28] Y. Yao, M. Aslam Noor, and Y. C. Liou, "Strong convergence of a modified extragradient method to the minimum-norm solution of variational inequalities," *Abstract and Applied Analysis*, p. 9, 2012.
- [29] Y. Yao, Y. C. Liou, C. L. Li, and H. T. Lin, "Extended extra-gradient methods for generalized variational inequalities," *Journal of Applied Mathematics*, p. 14, 2012.
- [30] Y. Yao, M. A. Noor, Y. C. Liou, and S. M. Kang, "Iterative algorithms for general multi-valued variational inequalities," *Abstract and Applied Analysis*, p. 10, 2012.
- [31] Y. Yao and N. Shahzad, "Strong convergence of a proximal point algorithm with general errors," *Optimization Letters*.
- [32] Y. Yao, Y.-C. Liou, and S. M. Kang, "Algorithms construction for variational inequalities," *Fixed Point Theory and Applications*, Article ID 794203, 12 pages, 2011.
- [33] V. Colao, G. L. Acedo, and G. Marino, "An implicit method for finding common solutions of variational inequalities and systems of equilibrium problems and fixed points of infinite family of nonexpansive mappings," *Nonlinear Analysis*, vol. 71, no. 7-8, pp. 2708–2715, 2009.

- [34] G. Marino and H.-K. Xu, "A general iterative method for nonexpansive mappings in Hilbert spaces," *Journal of Mathematical Analysis and Applications*, vol. 318, no. 1, pp. 43–52, 2006.
- [35] M. Aslam Noor and Z. Huang, "Wiener-Hopf equation technique for variational inequalities and non-expansive mappings," *Applied Mathematics and Computation*, vol. 191, no. 2, pp. 504–510, 2007.
- [36] K. Goebel and W. A. Kirk, *Topics in Metric Fixed Point Theory*, vol. 28 of *Cambridge Studies in Advanced Mathematics*, Cambridge University Press, Cambridge, UK, 1990.
- [37] T. Suzuki, "Strong convergence theorems for infinite families of nonexpansive mappings in general Banach spaces," *Fixed Point Theory and Applications*, no. 1, pp. 103–123, 2005.
- [38] H.-K. Xu, "Iterative algorithms for nonlinear operators," *Journal of the London Mathematical Society*, vol. 66, no. 1, pp. 240–256, 2002.
- [39] L. J. Zhang, J. M. Chen, and Z. B. Hou, "Viscosity approximation methods for nonexpansive mappings and generalized variational inequalities," *Acta Mathematica Sinica*, vol. 53, no. 4, pp. 691–698, 2010.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

