

## Research Article

# On Approximate Cubic Homomorphisms

**M. Eshaghi Gordji and M. Bavand Savadkouhi**

*Department of Mathematics, Semnan University, P. O. Box 35195-363, Semnan, Iran*

Correspondence should be addressed to M. Eshaghi Gordji, [madjid.eshaghi@gmail.com](mailto:madjid.eshaghi@gmail.com)

Received 22 October 2008; Revised 4 March 2009; Accepted 2 July 2009

Recommended by Rigoberto Medina

We investigate the generalized Hyers-Ulam-Rassias stability of the system of functional equations:  $f(xy) = f(x)f(y)$ ,  $f(2x + y) + f(2x - y) = 2f(x + y) + 2f(x - y) + 12f(x)$ , on Banach algebras. Indeed we establish the superstability of this system by suitable control functions.

Copyright © 2009 M. Eshaghi Gordji and M. Bavand Savadkouhi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## 1. Introduction

A definition of stability in the case of homomorphisms between metric groups was suggested by a problem by Ulam [2] in 1940. Let  $(G_1, \cdot)$  be a group and let  $(G_2, *)$  be a metric group with the metric  $d(\cdot, \cdot)$ . Given  $\epsilon > 0$ , does there exist a  $\delta > 0$  such that if a mapping  $h : G_1 \rightarrow G_2$  satisfies the inequality  $d(h(x \cdot y), h(x) * h(y)) < \delta$  for all  $x, y \in G_1$ , then there exists a homomorphism  $H : G_1 \rightarrow G_2$  with  $d(h(x), H(x)) < \epsilon$  for all  $x \in G_1$ ? In this case, the equation of homomorphism  $h(x \cdot y) = h(x) * h(y)$  is called stable. On the other hand, we are looking for situations when the homomorphisms are stable, that is, if a mapping is an approximate homomorphism, then there exists an exact homomorphism near it. The concept of stability for a functional equation arises when we replace the functional equation by an inequality which acts as a perturbation of the equation. In 1941, Hyers [3] gave a positive answer to the question of Ulam for Banach spaces. Let  $f : E_1 \rightarrow E_2$  be a mapping between Banach spaces such that

$$\|f(x + y) - f(x) - f(y)\| \leq \delta \quad (1.1)$$

for all  $x, y \in E_1$  and for some  $\delta \geq 0$ . Then there exists a unique additive mapping  $T : E_1 \rightarrow E_2$  satisfying

$$\|f(x) - T(x)\| \leq \delta \quad (1.2)$$

for all  $x \in E_1$ . Moreover, if  $f(tx)$  is continuous in  $t$  for each fixed  $x \in E_1$ , then the mapping  $T$  is linear. Rassias [4] succeeded in extending the result of Hyers' theorem by weakening the condition for the Cauchy difference controlled by  $(\|x\|^p + \|y\|^p)$ ,  $p \in [0, 1)$  to be unbounded. This condition has been assumed further till now, through the complete Hyers direct method, in order to prove linearity for generalized Hyers-Ulam stability problem forms. A number of mathematicians were attracted to the pertinent stability results of Rassias [4], and stimulated to investigate the stability problems of functional equations. The stability phenomenon that was introduced and proved by Rassias is called Hyers-Ulam-Rassias stability. Then the stability problems of several functional equations have been extensively investigated by a number of authors and there are many interesting results concerning this problem, see [5–13].

Bourgin [14] is the first mathematician dealing with stability of (ring) homomorphism  $f(xy) = f(x)f(y)$ . The topic of approximate homomorphisms was studied by a number of mathematicians, see [15–22] and references therein.

Jun and Kim [1] introduced the following functional equation:

$$f(2x + y) + f(2x - y) = 2f(x + y) + 2f(x - y) + 12f(x), \quad (1.3)$$

and they established the general solution and generalized Hyers-Ulam-Rassias stability problem for this functional equation. It is easy to see that the function  $f(x) = cx^3$  is a solution of the functional equation (1.3). Thus, it is natural that (1.3) is called a cubic functional equation and every solution of the cubic functional equation is said to be a cubic function.

Let  $R$  be a ring. Then a mapping  $f : R \rightarrow R$  is called a cubic homomorphism if  $f$  is a cubic function satisfying

$$f(ab) = f(a)f(b), \quad (1.4)$$

for all  $a, b \in R$ . For instance, let  $R$  be commutative, then the mapping  $f : R \rightarrow R$ , defined by  $f(a) = a^3$  ( $a \in R$ ), is a cubic homomorphism. It is easy to see that a cubic homomorphism is a ring homomorphism if and only if it is zero function. In this paper, we study the stability of cubic homomorphisms on Banach algebras. Indeed, we investigate the generalized Hyers-Ulam-Rassias stability of the system of functional equations:

$$\begin{aligned} f(xy) &= f(x)f(y), \\ f(2x + y) + f(2x - y) &= 2f(x + y) + 2f(x - y) + 12f(x), \end{aligned} \quad (1.5)$$

on Banach algebras. To this end, we need two control functions for our stability. One control function for (1.3) and an other control function for (1.4). So this is the main difference between our hypothesis (where two-degree freedom appears in the election for two control functions  $\phi_1$  and  $\phi_2$  in Theorem 2.1 in what follows), and the conditions (with one control function) that appear, for example, in [1, Theorem 3.1].

## 2. Main Results

In the following we suppose that  $A$  is a normed algebra,  $B$  is a Banach algebra, and  $f$  is a mapping from  $A$  into  $B$ , and  $\varphi, \varphi_1, \varphi_2$  are maps from  $A \times A$  into  $\mathbb{R}^+$ . Also, we put  $0^p = 0$  for  $p \leq 0$ .

**Theorem 2.1.** *Let*

$$\|f(xy) - f(x)f(y)\| \leq \varphi_1(x, y), \quad (2.1)$$

$$\|f(2x + y) + f(2x - y) - 2f(x + y) - 2f(x - y) - 12f(x)\| \leq \varphi_2(x, y) \quad (2.2)$$

for all  $x, y \in A$ . Assume that the series

$$\Psi(x, y) = \sum_{i=0}^{\infty} \frac{\varphi_2(2^i x, 2^i y)}{2^{3i}} \quad (2.3)$$

converges, and that

$$\lim_{n \rightarrow \infty} \frac{\varphi_1(2^n x, 2^n y)}{2^{6n}} = 0 \quad (2.4)$$

for all  $x, y \in A$ . Then there exists a unique cubic homomorphism  $T : A \rightarrow A$  such that

$$\|T(x) - f(x)\| \leq \frac{1}{16} \Psi(x, 0) \quad (2.5)$$

for all  $x \in A$ .

*Proof.* Setting  $y = 0$  in (2.2) yields

$$\|2f(2x) - 2^4 f(x)\| \leq \varphi_2(x, 0), \quad (2.6)$$

and then dividing by  $2^4$  in (2.6), we obtain

$$\left\| \frac{f(2x)}{2^3} - f(x) \right\| \leq \frac{\varphi_2(x, 0)}{2 \cdot 2^3} \quad (2.7)$$

for all  $x \in A$ . Now by induction we have

$$\left\| \frac{f(2^n x)}{2^{3n}} - f(x) \right\| \leq \frac{1}{2 \cdot 2^3} \sum_{i=0}^{n-1} \frac{\varphi_2(2^i x, 0)}{2^{3i}}. \quad (2.8)$$

In order to show that the functions  $T_n(x) = f(2^n x)/2^{3n}$  are a convergent sequence, we use the Cauchy convergence criterion. Indeed, replace  $x$  by  $2^m x$  and divide by  $2^{3m}$  in (2.8), where  $m$  is an arbitrary positive integer. We find that

$$\left\| \frac{f(2^{n+m}x)}{2^{3(n+m)}} - \frac{f(2^m x)}{2^{3m}} \right\| \leq \frac{1}{2 \cdot 2^3} \sum_{i=0}^{n-1} \frac{\varphi_2(2^{i+m}x, 0)}{2^{3(i+m)}} = \frac{1}{2 \cdot 2^3} \sum_{i=m}^{n+m-1} \frac{\varphi_2(2^i x, 0)}{2^{3i}} \quad (2.9)$$

for all positive integers  $m, n$ . Hence by the Cauchy criterion, the limit  $T(x) = \lim_{n \rightarrow \infty} T_n(x)$  exists for each  $x \in A$ . By taking the limit as  $n \rightarrow \infty$  in (2.8), we see that  $\|T(x) - f(x)\| \leq (1/2 \cdot 2^3) \sum_{i=0}^{\infty} (\varphi_2(2^i x, 0)/2^{3i}) = 1/16\Psi(x, 0)$  and (2.5) holds for all  $x \in A$ . If we replace  $x$  by  $2^n x$  and  $y$  by  $2^n y$ , respectively, in (2.2) and divide by  $2^{3n}$ , we see that

$$\begin{aligned} & \left\| \frac{f(2 \cdot (2^n x) + 2^n y)}{2^{3n}} + \frac{f(2 \cdot (2^n x) - 2^n y)}{2^{3n}} - 2 \frac{f(2^n x + 2^n y)}{2^{3n}} - 2 \frac{f(2^n x - 2^n y)}{2^{3n}} - 12 \frac{f(2^n x)}{2^{3n}} \right\| \\ & \leq \frac{\varphi_2(2^n x, 2^n y)}{2^{3n}}. \end{aligned} \quad (2.10)$$

Taking the limit as  $n \rightarrow \infty$ , we find that  $T$  satisfies (1.3) [1, Theorem 3.1]. On the other hand we have

$$\begin{aligned} \|T(xy) - T(x) \cdot T(y)\| &= \left\| \lim_{n \rightarrow \infty} \frac{f(2^n xy)}{2^{3n}} - \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^{3n}} \cdot \lim_{n \rightarrow \infty} \frac{f(2^n y)}{2^{3n}} \right\| \\ &= \lim_{n \rightarrow \infty} \left\| \frac{f(2^n x 2^n y)}{2^{6n}} - \frac{f(2^n x) f(2^n y)}{2^{6n}} \right\| \\ &\leq \lim_{n \rightarrow \infty} \frac{\varphi_1(2^n x, 2^n y)}{2^{6n}} = 0 \end{aligned} \quad (2.11)$$

for all  $x, y \in A$ . We find that  $T$  satisfies (1.4). To prove the uniqueness property of  $T$ , let  $T' : A \rightarrow A$  be a function satisfying  $T'(2x + y) + T'(2x - y) = 2T'(x + y) + 2T'(x - y) + 12T'(x)$  and  $\|T'(x) - f(x)\| \leq (1/16)\Psi(x, 0)$ . Since  $T, T'$  are cubic, then we have

$$T(2^n x) = 2^{3n}T(x), \quad T'(2^n x) = 2^{3n}T'(x) \quad (2.12)$$

for all  $x \in A$ , hence,

$$\begin{aligned}
 \|T(x) - T'(x)\| &= \frac{1}{2^{3n}} \|T(2^n x) - T'(2^n x)\| \\
 &\leq \frac{1}{2^{3n}} (\|T(2^n x) - f(2^n x)\| + \|T'(2^n x) - f'(2^n x)\|) \\
 &\leq \frac{1}{2^{3n}} \left( \frac{1}{2 \cdot 2^3} \Psi(2^n x, 0) + \frac{1}{2 \cdot 2^3} \Psi(2^n x, 0) \right) \\
 &= \frac{1}{2^{3(n+1)}} \Psi(2^n x, 0) = \frac{1}{2^{3(n+1)}} \sum_{i=0}^{\infty} \frac{1}{2^{3i}} \varphi_2(2^{i+n} x, 0) \\
 &= \frac{1}{2^3} \sum_{i=0}^{\infty} \frac{1}{2^{3(i+n)}} \varphi_2(2^{i+n} x, 0) = \frac{1}{2^3} \sum_{i=n}^{\infty} \frac{1}{2^{3i}} \varphi_2(2^i x, 0).
 \end{aligned} \tag{2.13}$$

By taking  $n \rightarrow \infty$  we get  $T(x) = T'(x)$ . □

**Corollary 2.2.** Let  $\theta_1$  and  $\theta_2$  be nonnegative real numbers, and let  $p \in (-\infty, 3)$ . Suppose that

$$\begin{aligned}
 \|f(xy) - f(x)f(y)\| &\leq \theta_1, \\
 \|f(2x + y) + f(2x - y) - 2f(x + y) - 2f(x - y) - 12f(x)\| &\leq \theta_2 (\|x\|^p + \|y\|^p)
 \end{aligned} \tag{2.14}$$

for all  $x, y \in A$ . Then there exists a unique cubic homomorphism  $T : A \rightarrow A$  such that

$$\|T(x) - f(x)\| \leq \frac{1}{16} \frac{\theta_2 \|x\|^p}{1 - 2^{p-3}} \tag{2.15}$$

for all  $x, y \in A$ .

*Proof.* In Theorem 2.1, let  $\varphi_1(x, y) = \theta_1$  and  $\varphi_2(x, y) = \theta_2 (\|x\|^p + \|y\|^p)$  for all  $x, y \in A$ . □

**Corollary 2.3.** Let  $\theta_1$  and  $\theta_2$  be nonnegative real numbers. Suppose that

$$\begin{aligned}
 \|f(xy) - f(x)f(y)\| &\leq \theta_1, \\
 \|f(2x + y) + f(2x - y) - 2f(x + y) - 2f(x - y) - 12f(x)\| &\leq \theta_2
 \end{aligned} \tag{2.16}$$

for all  $x, y \in A$ . Then there exists a unique cubic homomorphism  $T : A \rightarrow A$  such that

$$\|T(x) - f(x)\| \leq \frac{\theta_2}{14} \tag{2.17}$$

for all  $x \in A$ .

*Proof.* The proof follows from Corollary 2.2. □

**Corollary 2.4.** Let  $p \in (-\infty, 3)$  and let  $\theta$  be a positive real number. Suppose that

$$\lim_{n \rightarrow \infty} \frac{\varphi(2^n x, 2^n y)}{2^{6n}} = 0, \quad (2.18)$$

for all  $x, y \in A$ . Moreover, suppose that

$$\|f(xy) - f(x)f(y)\| \leq \varphi(x, y), \quad (2.19)$$

and that

$$\|f(2x + y) + f(2x - y) - 2f(x + y) - 2f(x - y) - 12f(x)\| \leq \theta \|y\|^p, \quad (2.20)$$

for all  $x, y \in A$ . Then  $f$  is a cubic homomorphism.

*Proof.* Letting  $x = y = 0$  in (2.20), we get that  $f(0) = 0$ . So by  $y = 0$ , in (2.20), we get  $f(2x) = 2^3 f(x)$  for all  $x \in A$ . By using induction we have

$$f(2^n x) = 2^{3n} f(x) \quad (2.21)$$

for all  $x \in A$  and  $n \in \mathbb{N}$ . On the other hand, by Theorem 2.1, the mapping  $T : A \rightarrow A$ , defined by

$$T(x) = \lim_{n \rightarrow \infty} \frac{f(2^n x)}{2^{3n}}, \quad (2.22)$$

is a cubic homomorphism. Therefore it follows from (2.21) that  $f = T$ . Hence it is a cubic homomorphism.  $\square$

**Corollary 2.5.** Let  $p, q, \theta \geq 0$ , and  $p + q < 3$ . Let

$$\lim_{n \rightarrow \infty} \frac{\varphi(2^n x, 2^n y)}{2^{6n}} = 0 \quad (2.23)$$

for all  $x, y \in A$ . Moreover, suppose that

$$\|f(xy) - f(x)f(y)\| \leq \varphi(x, y), \quad (2.24)$$

and that

$$\|f(2x + y) + f(2x - y) - 2f(x + y) - 2f(x - y) - 12f(x)\| \leq \theta \|x\|^q \|y\|^p \quad (2.25)$$

for all  $x, y \in A$ . Then  $f$  is a cubic homomorphism.

*Proof.* If  $q = 0$ , then by Corollary 2.4 we get the result. If  $q \neq 0$ , the following results from Theorem 2.1, by putting  $\varphi_1(x, y) = \varphi(x, y)$  and  $\varphi_2(x, y) = \theta(\|x\|^p\|y\|^p)$  for all  $x, y \in A$ .  $\square$

**Corollary 2.6.** *Let  $p \in (-\infty, 3)$  and  $\theta$  be a positive real number. Let*

$$\begin{aligned} \|f(xy) - f(x)f(y)\| &\leq \theta\|y\|^p, \\ \|f(2x + y) + f(2x - y) - 2f(x + y) - 2f(x - y) - 12f(x)\| &\leq \theta\|y\|^p \end{aligned} \tag{2.26}$$

for all  $x, y \in A$ . Then  $f$  is a cubic homomorphism.

*Proof.* Let  $\varphi(x, y) = \theta\|y\|^p$ . Then by Corollary 2.4, we get the result.  $\square$

**Theorem 2.7.** *Let*

$$\|f(xy) - f(x)f(y)\| \leq \varphi_1(x, y), \tag{2.27}$$

$$\|f(2x + y) + f(2x - y) - 2f(x + y) - 2f(x - y) - 12f(x)\| \leq \varphi_2(x, y) \tag{2.28}$$

for all  $x, y \in A$ . Assume that the series

$$\Psi(x, y) = \sum_{i=1}^{\infty} 2^{3i} \varphi_2\left(\frac{x}{2^i}, \frac{y}{2^i}\right) \tag{2.29}$$

converges and that

$$\lim_{n \rightarrow \infty} 2^{6n} \varphi_1\left(\frac{x}{2^n}, \frac{y}{2^n}\right) = 0 \tag{2.30}$$

for all  $x, y \in A$ . Then there exists a unique cubic homomorphism  $T : A \rightarrow A$  such that

$$\|T(x) - f(x)\| \leq \frac{1}{16} \Psi(x, 0) \tag{2.31}$$

for all  $x \in A$ .

*Proof.* Setting  $y = 0$  in (2.28) yields

$$\|2f(2x) - 2 \cdot 2^3 f(x)\| \leq \varphi_2(x, 0). \tag{2.32}$$

Replacing  $x$  by  $x/2$  in (2.32), we get

$$\|f(x) - 2^3 f\left(\frac{x}{2}\right)\| \leq \frac{1}{2} \varphi_2\left(\frac{x}{2}, 0\right) \tag{2.33}$$

for all  $x \in A$ . By (2.33) we use iterative methods and induction on  $n$  to prove the following relation

$$\left\| f(x) - 2^{3n} f\left(\frac{x}{2^n}\right) \right\| \leq \frac{1}{2 \cdot 2^3} \sum_{i=1}^n 2^{3i} \varphi_2\left(\frac{x}{2^i}, 0\right). \quad (2.34)$$

In order to show that the functions  $T_n(x) = 2^{3n} f(x/2^n)$  are a convergent sequence, replace  $x$  by  $x/2^m$  in (2.34), and then multiply by  $2^{3m}$ , where  $m$  is an arbitrary positive integer. We find that

$$\begin{aligned} \left\| 2^{3m} f\left(\frac{x}{2^m}\right) - 2^{3(n+m)} f\left(\frac{x}{2^{n+m}}\right) \right\| &\leq \frac{1}{2 \cdot 2^3} \sum_{i=1}^n 2^{3(i+m)} \varphi_2\left(\frac{x}{2^{i+m}}, 0\right) \\ &= \frac{1}{2 \cdot 2^3} \sum_{i=1+m}^{n+m} 2^{3i} \varphi_2\left(\frac{x}{2^i}, 0\right) \end{aligned} \quad (2.35)$$

for all positive integers. Hence by the Cauchy criterion the limit  $T(x) = \lim_{n \rightarrow \infty} T_n(x)$  exists for each  $x \in A$ . By taking the limit as  $n \rightarrow \infty$  in (2.34), we see that  $\|T(x) - f(x)\| \leq 1/2 \cdot 2^3 \sum_{i=1}^{\infty} 2^{3i} \varphi_2(x/2^i, 0) = (1/16)\Psi(x, 0)$ , and (2.31) holds for all  $x \in A$ . The rest of proof is similar to the proof of Theorem 2.1.  $\square$

**Corollary 2.8.** *Let  $p > 3$  and  $\theta$  be a positive real number. Let*

$$\lim_{n \rightarrow \infty} 2^{6n} \varphi\left(\frac{x}{2^n}, \frac{y}{2^n}\right) = 0, \quad (2.36)$$

for all  $x, y \in A$ . Moreover, suppose that

$$\|f(xy) - f(x)f(y)\| \leq \varphi(x, y), \quad (2.37)$$

$$\|f(2x + y) + f(2x - y) - 2f(x + y) - 2f(x - y) - 12f(x)\| \leq \theta \|y\|^p, \quad (2.38)$$

for all  $x, y \in A$ . Then  $f$  is a cubic homomorphism.

*Proof.* Letting  $x = y = 0$  in (2.38), we get that  $f(0) = 0$ . So by  $y = 0$ , in (2.38), we get  $f(2x) = 2^3 f(x)$  for all  $x \in A$ . By using induction, we have

$$f(x) = 2^{3n} f\left(\frac{x}{2^n}\right) \quad (2.39)$$

for all  $x \in A$  and  $n \in \mathbb{N}$ . On the other hand, by Theorem 2.8, the mapping  $T : A \rightarrow A$ , defined by

$$T(x) = \lim_{n \rightarrow \infty} 2^{3n} f\left(\frac{x}{2^n}\right), \quad (2.40)$$

is a cubic homomorphism. Therefore, it follows from (2.39) that  $f = T$ . Hence  $f$  is a cubic homomorphism.  $\square$



Example 2.9. Let

$$\mathcal{A} := \begin{bmatrix} 0 & \mathbb{R} & \mathbb{R} & \mathbb{R} \\ 0 & 0 & \mathbb{R} & \mathbb{R} \\ 0 & 0 & 0 & \mathbb{R} \\ 0 & 0 & 0 & 0 \end{bmatrix}, \tag{2.41}$$

then  $\mathcal{A}$  is a Banach algebra equipped with the usual matrix-like operations and the following norm:

$$\left\| \begin{bmatrix} 0 & a_1 & a_2 & a_3 \\ 0 & 0 & a_4 & a_5 \\ 0 & 0 & 0 & a_6 \\ 0 & 0 & 0 & 0 \end{bmatrix} \right\| = \sum_{i=1}^6 |a_i| \quad (a_i \in \mathbb{R}). \tag{2.42}$$

Let

$$a := \begin{bmatrix} 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \tag{2.43}$$

and we define  $f : \mathcal{A} \rightarrow \mathcal{A}$  by  $f(x) = x^3 + a$ , and

$$\begin{aligned} \varphi_1(x, y) &:= \|f(xy) - f(x)f(y)\| = \|a\| = 4, \\ \varphi_2(x, y) &:= \|f(2x + y) + f(2x - y) - 2f(x + y) - 2f(x - y) - 12f(x)\| = 14\|a\| = 56 \end{aligned} \tag{2.44}$$

for all  $x, y \in \mathcal{A}$ . Then we have

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{\varphi_2(2^k x, 2^k y)}{2^{3k}} &= \sum_{k=0}^{\infty} \frac{56}{2^{3k}} = 64, \\ \lim_{n \rightarrow \infty} \frac{\varphi_1(2^n x, 2^n y)}{2^{6n}} &= 0. \end{aligned} \tag{2.45}$$

Thus the limit  $T(x) = \lim_{n \rightarrow \infty} (f(2^n x)/2^{3n}) = x^3$  exists. Also,

$$T(xy) = (xy)^3 = x^3 y^3 = T(x)T(y). \tag{2.46}$$

Furthermore,

$$\begin{aligned} T(2x + y) + T(2x - y) &= (2x + y)^3 + (2x - y)^3 = 16x^3 + 12xy^2 \\ &= 2T(x + y) + 2T(x - y) + 12T(x). \end{aligned} \quad (2.47)$$

Hence  $T$  is cubic homomorphism.

Also from this example, it is clear that the superstability of the system of functional equations

$$\begin{aligned} f(xy) &= f(x)f(y), \\ f(2x + y) + f(2x - y) &= 2f(x + y) + 2f(x - y) + 12f(x), \end{aligned} \quad (2.48)$$

with the control functions in Corollaries 2.4, 2.5 and 2.6 does not hold.

## Acknowledgments

The authors would like to thank the referees for their valuable suggestions. Also, M. B. Savadkouhi would like to thank the Office of Gifted Students at Semnan University for its financial support.

## References

- [1] K. W. Jun and H. M. Kim, "The generalized Hyers-Ulam-Rassias stability of a cubic functional equation," *Journal of Mathematical Analysis and Applications*, vol. 274, no. 2, pp. 267–278, 2002.
- [2] S. M. Ulam, *A Collection of Mathematical Problems*, Interscience, New York, NY, USA, 1960.
- [3] D. H. Hyers, "On the stability of the linear functional equation," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 27, pp. 222–224, 1941.
- [4] Th. M. Rassias, "On the stability of the linear mapping in Banach spaces," *Proceedings of the American Mathematical Society*, vol. 72, pp. 297–300, 1978.
- [5] V. A. Faiziev, Th. M. Rassias, and P. K. Sahoo, "The space of  $(\psi, \gamma)$ -additive mappings on semigroups," *Transactions of the American Mathematical Society*, vol. 354, no. 11, pp. 4455–4472, 2002.
- [6] G. L. Forti, "An existence and stability theorem for a class of functional equations," *Stochastica*, vol. 4, pp. 23–30, 1980.
- [7] G. L. Forti, "Comments on the core of the direct method for proving Hyers-Ulam stability of functional equations," *Journal of Mathematical Analysis and Applications*, vol. 295, pp. 127–133, 2004.
- [8] D. H. Hyers, G. Isac, and Th. M. Rassias, *Stability of Functional Equations in Several Variables*, Birkhäuser, Boston, Mass, USA, 1998.
- [9] G. Isac and Th. M. Rassias, "On the Hyers-Ulam stability of a cubic functional equation," *Journal of Approximation Theory*, vol. 72, no. 2, pp. 131–137, 1993.
- [10] L. Maligranda, "A result of Tosio Aoki about a generalization of Hyers-Ulam stability of additive functions—a question of priority," *Aequationes Mathematicae*, vol. 75, pp. 289–296, 2008.
- [11] Th. M. Rassias and J. Tabor, *Stability of Mappings of Hyers-Ulam Type*, Hadronic Press, Palm Harbor, Fla, USA, 1994.
- [12] Th. M. Rassias, "On a modified Hyers-Ulam sequence," *Journal of Mathematical Analysis and Applications*, vol. 158, pp. 106–113, 1991.
- [13] Th. M. Rassias, "On the stability of functional equations originated by a problem of Ulam," *Mathematica*, vol. 44(67), no. 1, pp. 39–75, 2002.
- [14] D. G. Bourgin, "Classes of transformations and bordering transformations," *Bulletin of the American Mathematical Society*, vol. 57, pp. 223–237, 1951.

- [15] R. Badora, "On approximate ring homomorphisms," *Journal of Mathematical Analysis and Applications*, vol. 276, pp. 589–597, 2002.
- [16] J. Baker, J. Lawrence, and F. Zorzitto, "The stability of the equation  $f(x + y) = f(x)f(y)$ ," *Proceedings of the American Mathematical Society*, vol. 74, no. 2, pp. 242–246, 1979.
- [17] M. Eshaghi Gordji and M. Bavand Savadkouhi, "Approximation of generalized homomorphisms in quasi-Banach algebras," to appear in *Analele Stiintifice ale Universitatii Ovidius Constanta*.
- [18] M. Eshaghi Gordji, T. Karimi, and S. Kaboli Gharetapeh, "Approximately  $n$ -Jordan homomorphisms on Banach algebras," *Journal of Inequalities and Applications*, vol. 2009, Article ID 870843, 8 pages, 2009.
- [19] D. H. Hyers and Th. M. Rassias, "Approximate homomorphisms," *Aequationes Mathematicae*, vol. 44, pp. 125–153, 1992.
- [20] C. Park, "Hyers-Ulam-Rassias stability of homomorphisms in quasi-Banach algebras," *Bulletin des Sciences Mathématiques*, vol. 132, no. 2, pp. 87–96, 2008.
- [21] Th. M. Rassias, "The problem of S. M. Ulam for approximately multiplicative mappings," *Journal of Mathematical Analysis and Applications*, vol. 246, no. 2, pp. 352–378, 2000.
- [22] Th. M. Rassias, "On the stability of functional equations and a problem of Ulam," *Acta Applicandae Mathematicae*, vol. 62, no. 1, pp. 23–130, 2000.