

Research Article

Positive Solutions for Boundary Value Problems of Second-Order Functional Dynamic Equations on Time Scales

Ilkay Yaslan Karaca

Department of Mathematics, Ege University, Bornova, 35100 Izmir, Turkey

Correspondence should be addressed to Ilkay Yaslan Karaca, ilkay.karaca@ege.edu.tr

Received 3 February 2009; Accepted 26 February 2009

Recommended by Alberto Cabada

Criteria are established for existence of least one or three positive solutions for boundary value problems of second-order functional dynamic equations on time scales. By this purpose, we use a fixed-point index theorem in cones and Leggett-Williams fixed-point theorem.

Copyright © 2009 Ilkay Yaslan Karaca. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Introduction

In a recent paper [1], by applying a fixed-point index theorem in cones, Jiang and Weng studied the existence of positive solutions for the boundary value problems described by second-order functional differential equations of the form

$$\begin{aligned}y''(x) + r(x)f(y(\omega(x))) &= 0, & 0 < x < 1, \\ \alpha y(x) - \beta y'(x) &= \xi(x), & a \leq x \leq 0, \\ \gamma y(x) + \delta y'(x) &= \eta(x), & 1 \leq x \leq b.\end{aligned}\tag{1.1}$$

Aykut [2] applied a cone fixed-point index theorem in cones and obtained sufficient conditions for the existence of positive solutions of the boundary value problems of functional difference equations of the form

$$\begin{aligned}-\Delta^2 y(n-1) &= f(n, y(\omega(n))), & n \in [a, b], \\ \alpha y(n-1) - \beta \Delta y(n-1) &= \xi(n), & n \in [\tau_1, a], \\ \gamma y(n) + \delta \Delta y(n) &= \eta(n), & n \in [b, \tau_2].\end{aligned}\tag{1.2}$$

In this article, we are interested in proving the existence and multiplicity of positive solutions for the boundary value problems of a second-order functional dynamic equation of the form

$$\begin{aligned} -y^{\Delta\nabla}(t) + q(t)y(t) &= f(t, y(\omega(t))), \quad t \in [a, b], \\ \alpha y(\rho(t)) - \beta y^{\Delta}(\rho(t)) &= \xi(t), \quad t \in [\tau_1, a], \\ \gamma y(t) + \delta y^{\Delta}(t) &= \eta(t), \quad t \in [b, \tau_2]. \end{aligned} \quad (1.3)$$

Throughout this paper we let \mathbb{T} be any time scale (nonempty closed subset of \mathbb{R}) and $[a, b]$ be a subset of \mathbb{T} such that $[a, b] = \{t \in \mathbb{T}, a \leq t \leq b\}$, and for $t \in [\tau_1, a]$, t is not right scattered and left dense at the same time.

Some preliminary definitions and theorems on time scales can be found in books [3, 4] which are excellent references for calculus of time scales.

We will assume that the following conditions are satisfied.

- (H1) $q(t) \in C[a, b]$, $q(t) \geq 0$.
 (H2) $f : [a, b] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous with respect to y and $f(t, y) \geq 0$ for $y \in \mathbb{R}^+$, where \mathbb{R}^+ denotes the set of nonnegative real numbers.
 (H3) $\omega(t)$ defined on $[a, b]$ satisfies

$$\begin{aligned} c &= \inf \{ \omega(t) : a \leq t \leq b \} < b, \\ d &= \sup \{ \omega(t) : a \leq t \leq b \} > a. \end{aligned} \quad (1.4)$$

Let $E_1 := \{t \in E : a \leq \omega(t) \leq \rho(b)\}$ be nonempty subset of

$$E := \{t \in [a, b] : a \leq \omega(t) \leq b\}. \quad (1.5)$$

- (H4) $\alpha, \beta, \gamma, \delta \geq 0$, $\alpha + \beta > 0$, $\gamma + \delta > 0$;
 if $q(t) \equiv 0$ ($a \leq t \leq b$), then $\alpha + \gamma > 0$; $-\gamma/\delta \in \mathcal{R}^+$ for $\delta > 0$, where \mathcal{R}^+ denotes the set of all positively regressive and rd-continuous functions.
 (H5) $\xi(t)$ and $\eta(t)$ are defined on $[\tau_1, \sigma(a)]$ and $[b, \sigma(\tau_2)]$, respectively, where $\tau_1 := \min\{a, c\}$, $\tau_2 := \max\{b, d\}$; furthermore, $\xi(a) = \eta(b) = 0$;

$$\begin{aligned} \xi(\sigma(t)) &\geq 0, \quad \text{for } t \in [\tau_1, a] \text{ as } \beta = 0; \\ \int_t^a \xi(s) e_{\ominus(\alpha/\beta)}(s, 0) \nabla s &\geq 0, \quad \text{for } t \in [\tau_1, a] \text{ as } \beta > 0; \\ \eta(t) &\geq 0, \quad \text{for } t \in [b, \tau_2] \text{ as } \delta = 0; \\ e_{-\gamma/\delta}(t, 0) \int_b^t \eta(s) e_{\ominus(-\gamma/\delta)}(\sigma(s), 0) \Delta s &\geq 0, \quad \text{for } t \in [b, \tau_2] \text{ as } \delta > 0. \end{aligned} \quad (1.6)$$

There have been a number of works concerning of at least one and multiple positive solutions for boundary value problems recent years. Some authors have studied the problem for ordinary differential equations, while others have studied the problem for difference equations, while still others have considered the problem for dynamic equations on a time scale [5–10]. However there are fewer research for the existence of positive solutions of the boundary value problems of functional differential, difference, and dynamic equations [1, 2, 11–13].

Our problem is a dynamic analog of the BVPs (1.1) and (1.2). But it is more general than them. Moreover, conditions for the existence of at least one positive solution were studied for the BVPs (1.1) and (1.2). In this paper, we investigate the conditions for the existence of at least one or three positive solutions for the BVP (1.3). The key tools in our approach are the following fixed-point index theorem [14], and Leggett-Williams fixed-point theorem [15].

Theorem 1.1 (see [14]). *Let E be Banach space and $K \subset E$ be a cone in E . Let $r > 0$, and define $\Omega_r = \{x \in K : \|x\| < r\}$. Assume $A : \overline{\Omega}_r \rightarrow K$ is a completely continuous operator such that $Ax \neq x$ for $x \in \partial\Omega_r$.*

(i) *If $\|Au\| \leq \|u\|$ for $u \in \partial\Omega_r$, then $i(A, \Omega_r, K) = 1$.*

(ii) *If $\|Au\| \geq \|u\|$ for $u \in \partial\Omega_r$, then $i(A, \Omega_r, K) = 0$.*

Theorem 1.2 (see [15]). *Let \mathcal{D} be a cone in the real Banach space E . Set*

$$\begin{aligned} \mathcal{D}_r &:= \{x \in \mathcal{D} : \|x\| < r\}, \\ \mathcal{D}(\psi, p, q) &:= \{x \in \mathcal{D} : p \leq \psi(x), \|x\| \leq q\}. \end{aligned} \tag{1.7}$$

Suppose that $A : \overline{\mathcal{D}}_r \rightarrow \overline{\mathcal{D}}_r$ is a completely continuous operator and ψ is a nonnegative continuous concave functional on \mathcal{D} with $\psi(x) \leq \|x\|$ for all $x \in \overline{\mathcal{D}}_r$. If there exists $0 < p < q < s \leq r$ such that the following conditions hold:

(i) *$\{x \in \mathcal{D}(\psi, q, s) : \psi(x) > q\} \neq \{\}$ and $\psi(Ax) > q$ for all $x \in \mathcal{D}(\psi, q, s)$;*

(ii) *$\|Ax\| < p$ for $\|x\| \leq p$;*

(iii) *$\psi(Ax) > q$ for $x \in \mathcal{D}(\psi, q, r)$ with $\|Ax\| > s$.*

Then A has at least three fixed points x_1, x_2 , and x_3 in $\overline{\mathcal{D}}_r$ satisfying

$$\|x_1\| < p, \quad \psi(x_2) > q, \quad p < \|x_3\| \quad \text{with} \quad \psi(x_3) < q. \tag{1.8}$$

2. Preliminaries

First, we give the following definitions of solution and positive solution of BVP (1.3).

Definition 2.1. We say a function $y(t)$ is a solution of BVP (1.3) if it satisfies the following.

- (1) $y(t)$ is nonnegative on $[\rho(\tau_1), \sigma(\tau_2)]$.
 (2) $y(t) = y(\tau_1; t)$ as $t \in [\rho(\tau_1), a]$, where $y(\tau_1; t) : [\rho(\tau_1), a] \rightarrow [0, \infty)$ is defined as

$$y(\tau_1; t) = \begin{cases} e_{\alpha/\beta}(t, 0) \left[\frac{1}{\beta} \int_t^a \xi(s) e_{\ominus(\alpha/\beta)}(s, 0) \nabla s + e_{\ominus(\alpha/\beta)}(a, 0) y(a) \right], & \text{if } \beta > 0, \\ \frac{1}{\alpha} \xi(\sigma(t)), & \text{if } \beta = 0. \end{cases} \quad (2.1)$$

- (3) $y(t) = y(\tau_2; t)$ as $t \in [b, \sigma(\tau_2)]$, where $y(\tau_2; t) : [b, \sigma(\tau_2)] \rightarrow [0, \infty)$ is defined as

$$y(\tau_2; t) = \begin{cases} e_{-\gamma/\delta}(t, 0) \left[\frac{1}{\delta} \int_b^t \eta(s) e_{\ominus(-\gamma/\delta)}(\sigma(s), 0) \Delta s + e_{\ominus(-\gamma/\delta)}(b, 0) y(b) \right], & \text{if } \delta > 0, \\ \frac{1}{\gamma} \eta(t), & \text{if } \delta = 0. \end{cases} \quad (2.2)$$

- (4) y is Δ -differentiable, $y^\Delta : [\rho(a), b] \rightarrow \mathbb{R}$ is ∇ -differentiable on $[a, b]$ and $(y^\Delta)^\nabla : [a, b] \rightarrow \mathbb{R}$ is continuous.
 (5) $-y^{\Delta\nabla}(t) + q(t)y(t) = f(t, y(\omega(t)))$, for $t \in [a, b]$.

Furthermore, a solution $y(t)$ of (1.3) is called a positive solution if $y(t) > 0$ for $t \in [a, b]$. Denote by $\varphi(t)$ and $\psi(t)$ the solutions of the corresponding homogeneous equation

$$-y^{\Delta\nabla}(t) + q(t)y(t) = 0, \quad t \in [a, b], \quad (2.3)$$

under the initial conditions

$$\begin{aligned} \varphi(\rho(a)) &= \beta, & \varphi^\Delta(\rho(a)) &= \alpha, \\ \psi(b) &= \delta, & \psi^\Delta(b) &= -\gamma. \end{aligned} \quad (2.4)$$

Set

$$D = W_t(\psi, \varphi) = \psi(t)\varphi^\Delta(t) - \psi^\Delta(t)\varphi(t). \quad (2.5)$$

Since the Wronskian of any two solutions of (2.3) is independent of t , evaluating at $t = \rho(a)$, $t = b$ and using the initial conditions (2.4) yield

$$D = \alpha\psi(\rho(a)) - \beta\psi^\Delta(\rho(a)) = \gamma\psi(b) + \delta\psi^\Delta(b). \quad (2.6)$$

Using the initial conditions (2.4), we can deduce from (2.3) for $\varphi(t)$ and $\psi(t)$, the following equations:

$$\varphi(t) = \beta + \alpha(t - \rho(a)) + \int_{\rho(a)}^t \int_{\rho(a)}^{\tau} q(s)\varphi(s)\nabla s \Delta\tau, \tag{2.7}$$

$$\psi(t) = \delta + \gamma(b - t) + \int_t^b \int_{\tau}^b q(s)\psi(s)\nabla s \Delta\tau. \tag{2.8}$$

(See [8].)

Lemma 2.2 (see [8]). *Under the conditions (H1) and the first part of (H4) the solutions $\varphi(t)$ and $\psi(t)$ possess the following properties:*

$$\begin{aligned} \varphi(t) &\geq 0, & t &\in [\rho(a), \sigma(b)], & \psi(t) &\geq 0, & t &\in [\rho(a), b], \\ \varphi(t) &> 0, & t &\in (\rho(a), \sigma(b)), & \psi(t) &> 0, & t &\in [\rho(a), b), \\ \varphi^\Delta(t) &\geq 0, & t &\in [\rho(a), b], & \psi^\Delta(t) &\leq 0, & t &\in [\rho(a), b]. \end{aligned} \tag{2.9}$$

Let $G(t, s)$ be the Green function for the boundary value problem:

$$\begin{aligned} -y^{\Delta\nabla}(t) + q(t)y(t) &= 0, & t &\in [a, b], \\ \alpha y(\rho(a)) - \beta y^\Delta(\rho(a)) &= 0, \\ \gamma y(b) + \delta y^\Delta(b) &= 0, \end{aligned} \tag{2.10}$$

given by

$$G(t, s) = \frac{1}{D} \begin{cases} \psi(t)\varphi(s), & \text{if } \rho(a) \leq s \leq t \leq \sigma(b), \\ \varphi(s)\psi(t), & \text{if } \rho(a) \leq t \leq s \leq \sigma(b), \end{cases} \tag{2.11}$$

where $\varphi(t)$ and $\psi(t)$ are given in (2.7) and (2.8), respectively. It is obvious from (2.6), (H1) and (H4), that $D > 0$ holds.

Lemma 2.3. *Assume the conditions (H1) and (H4) are satisfied. Then*

- (i) $0 \leq G(t, s) \leq G(s, s)$ for $t, s \in [\rho(a), b]$,
- (ii) $G(t, s) \geq \Gamma G(s, s)$ for $t \in [a, \rho(b)]$ and $s \in [\rho(a), b]$,

where

$$\Gamma = \min \{I_1, I_2\}, \tag{2.12}$$

in which

$$\begin{aligned} I_1 &= \{\delta + (b - \rho(b))[\gamma + \delta q(b)(b - \rho(b))]\} \cdot \left\{ \delta + \gamma(b - \rho(a)) + \int_{\rho(a)}^b \int_{\tau}^b q(s)\varphi(s)\nabla s \Delta\tau \right\}^{-1}, \\ I_2 &= \{\beta + \alpha(a - \rho(a))\} \cdot \left\{ \beta + \alpha(b - \rho(a)) + \int_{\rho(a)}^b \int_{\rho(a)}^{\tau} q(s)\varphi(s)\nabla s \Delta\tau \right\}^{-1}. \end{aligned} \quad (2.13)$$

Proof. $\varphi(t) \geq 0$, for $t \in [\rho(a), \sigma(b)]$, and $\psi(t) \geq 0$, for $t \in [\rho(a), b]$. Besides, $\varphi(t)$ is nondecreasing and $\psi(t)$ is nonincreasing, for $t \in [\rho(a), b]$. Therefore, we have

$$0 \leq \frac{1}{D} \begin{cases} \varphi(t)\varphi(s), & \text{if } \rho(a) \leq s \leq t \leq \sigma(b) \\ \varphi(s)\varphi(t), & \text{if } \rho(a) \leq t \leq s \leq \sigma(b) \end{cases} \leq G(s, s). \quad (2.14)$$

So statement (i) of the lemma is proved. If $G(s, s) = 0$, for a given $s \in [\rho(a), b]$, then statement (ii) of the lemma is obvious for such values. Now, $s \in [\rho(a), b]$ and $G(s, s) \neq 0$. Consequently, $G(s, s) > 0$, for all such s . Let us take any $t \in [a, \rho(b)]$. Then we have for $s \in [\rho(a), t]$,

$$\begin{aligned} \frac{G(t, s)}{G(s, s)} &= \frac{\varphi(t)}{\varphi(s)} \geq \frac{\varphi(\rho(b))}{\varphi(\rho(a))} \\ &= \{\delta + (b - \rho(b))[\gamma + \delta q(b)(b - \rho(b))]\} \\ &\quad \cdot \left\{ \delta + \gamma(b - \rho(a)) + \int_{\rho(a)}^b \int_{\tau}^b q(s)\varphi(s)\nabla s \Delta\tau \right\}^{-1}, \end{aligned} \quad (2.15)$$

and we have for $s \in [t, b]$,

$$\frac{G(t, s)}{G(s, s)} = \frac{\varphi(t)}{\varphi(s)} \geq \frac{\varphi(a)}{\varphi(b)} = \{\beta + \alpha(a - \rho(a))\} \cdot \left\{ \beta + \alpha(b - \rho(a)) + \int_{\rho(a)}^b \int_{\rho(a)}^{\tau} q(s)\varphi(s)\nabla s \Delta\tau \right\}^{-1}. \quad (2.16)$$

Let $\mathcal{B} = C[\rho(\tau_1), \sigma(\tau_2)]$ be endowed with maximum norm $\|y\| := \max_{\rho(\tau_1) \leq t \leq \sigma(\tau_2)} |y(t)|$ for $y \in \mathcal{B}$, and let $K \subset \mathcal{B}$ be a cone defined by

$$K = \left\{ y \in \mathcal{B} : \min_{a \leq t \leq \rho(b)} y(t) \geq \Gamma \|y\| \right\}, \quad (2.17)$$

where Γ is as in (2.12).

Suppose that $y(t)$ is a solution of (1.3), then it can be written as

$$y(t) = \begin{cases} y(\tau_1; t), & \text{if } \rho(\tau_1) \leq t \leq a, \\ \int_{\rho(a)}^b G(t, s) f(s, y(\omega(s))) \nabla s, & \text{if } a \leq t \leq b, \\ y(\tau_2; t), & \text{if } b \leq t \leq \sigma(\tau_2), \end{cases} \quad (2.18)$$

where

$$\begin{aligned}
 y(\tau_1; t) &= \begin{cases} e_{\alpha/\beta}(t, 0) \left[\frac{1}{\beta} \int_t^a \xi(s) e_{\ominus(\alpha/\beta)}(s, 0) \nabla s + e_{\ominus(\alpha/\beta)}(a, 0) y(a) \right], & \text{if } \rho(\tau_1) \leq t \leq a, \beta > 0, \\ \frac{1}{\alpha} \xi(\sigma(t)), & \text{if } \rho(\tau_1) \leq t \leq a, \beta = 0, \end{cases} \\
 y(\tau_2; t) &= \begin{cases} e_{-\gamma/\delta}(t, 0) \left[\frac{1}{\delta} \int_b^t \eta(s) e_{\ominus(-\gamma/\delta)}(\sigma(s), 0) \Delta s \right. \\ \quad \left. + e_{\ominus(-\gamma/\delta)}(b, 0) y(b) \right], & \text{if } b \leq t \leq \sigma(\tau_2), \delta > 0, \\ \frac{1}{\gamma} \eta(t), & \text{if } b \leq t \leq \sigma(\tau_2), \delta = 0. \end{cases}
 \end{aligned} \tag{2.19}$$

Throughout this paper we assume that $u_0(t)$ is the solution of (1.3) with $f \equiv 0$. Clearly, $u_0(t)$ can be expressed as follows:

$$u_0(t) = \begin{cases} u_0(\tau_1; t), & \text{if } \rho(\tau_1) \leq t \leq a, \\ 0, & \text{if } a \leq t \leq b, \\ u_0(\tau_2; t), & \text{if } b \leq t \leq \sigma(\tau_2), \end{cases} \tag{2.20}$$

where

$$\begin{aligned}
 u_0(\tau_1; t) &= \begin{cases} \frac{1}{\beta} e_{\alpha/\beta}(t, 0) \int_t^a \xi(s) e_{\ominus(\alpha/\beta)}(s, 0) \nabla s, & \text{if } \rho(\tau_1) \leq t \leq a, \beta > 0, \\ \frac{1}{\alpha} \xi(\sigma(t)), & \text{if } \rho(\tau_1) \leq t \leq a, \beta = 0, \end{cases} \\
 u_0(\tau_2; t) &= \begin{cases} \frac{1}{\delta} e_{-\gamma/\delta}(t, 0) \int_b^t \eta(s) e_{\ominus(-\gamma/\delta)}(\sigma(s), 0) \Delta s, & \text{if } b \leq t \leq \sigma(\tau_2), \delta > 0, \\ \frac{1}{\gamma} \eta(t), & \text{if } b \leq t \leq \sigma(\tau_2), \delta = 0. \end{cases}
 \end{aligned} \tag{2.21}$$

Let $y(t)$ be a solution of (1.3) and $u(t) = y(t) - u_0(t)$. Noting that $u(t) \equiv y(t)$ for $t \in [a, b]$, we have

$$u(t) = \begin{cases} u(\tau_1; t), & \text{if } \rho(\tau_1) \leq t \leq a, \\ \int_{\rho(a)}^b G(t, s) f(s, u(\omega(s)) + u_0(\omega(s))) \nabla s, & \text{if } a \leq t \leq b, \\ u(\tau_2; t), & \text{if } b \leq t \leq \sigma(\tau_2), \end{cases} \tag{2.22}$$

where

$$\begin{aligned} u(\tau_1; t) &= \begin{cases} e_{\alpha/\beta}(t, a)y(a), & \text{if } \rho(\tau_1) \leq t \leq a, \beta > 0, \\ 0, & \text{if } \rho(\tau_1) \leq t \leq a, \beta = 0, \end{cases} \\ u(\tau_2; t) &= \begin{cases} e_{-\gamma/\delta}(t, b)y(b), & \text{if } b \leq t \leq \sigma(\tau_2), \delta > 0, \\ 0, & \text{if } b \leq t \leq \sigma(\tau_2), \delta = 0. \end{cases} \end{aligned} \quad (2.23)$$

Define an operator $A : K \rightarrow K$ as follows:

$$(Au)(t) := \begin{cases} (A_1u)(t), & \text{if } \rho(\tau_1) \leq t \leq a, \\ \int_{\rho(a)}^b G(t, s)f(s, u(\omega(s)) + u_0(\omega(s)))\nabla s, & \text{if } a \leq t \leq b, \\ (A_2u)(t), & \text{if } b \leq t \leq \sigma(\tau_2), \end{cases} \quad (2.24)$$

where

$$\begin{aligned} (A_1u)(t) &:= \begin{cases} e_{\alpha/\beta}(t, a) \int_{\rho(a)}^b G(a, s)f(s, u(\omega(s)) + u_0(\omega(s)))\nabla s, & \text{if } \rho(\tau_1) \leq t \leq a, \beta > 0, \\ 0, & \text{if } \rho(\tau_1) \leq t \leq a, \beta = 0, \end{cases} \\ (A_2u)(t) &:= \begin{cases} e_{-\gamma/\delta}(t, b) \int_{\rho(a)}^b G(b, s)f(s, u(\omega(s)) + u_0(\omega(s)))\nabla s, & \text{if } b \leq t \leq \sigma(\tau_2), \delta > 0, \\ 0, & \text{if } b \leq t \leq \sigma(\tau_2), \delta = 0. \end{cases} \end{aligned} \quad (2.25)$$

It is easy to derive that y is a positive solution of BVP (1.3) if and only if $u = y - u_0$ is a nontrivial fixed point u of $A : K \rightarrow K$, where u_0 be defined as before. \square

Lemma 2.4. $A(K) \subset K$.

Proof. For $u \in K$, we have $Au(t) \geq 0$, $t \in [\rho(\tau_1), \sigma(\tau_2)]$. Moreover, we have from definition of A that $Au(t) \leq Au(a)$ and $Au(t) \leq Au(b)$, for $t \in [\rho(\tau_1), a]$ and $t \in [b, \sigma(\tau_2)]$, respectively. Thus, $\|Au\| = \|Au\|_{[a, b]}$, where $\|Au\|_{[a, b]} = \max\{|Au(t)| : a \leq t \leq b\}$. It follows from the definition K and Lemma 2.3 that

$$\begin{aligned} \min_{a \leq t \leq \rho(b)} (Au)(t) &= \min_{a \leq t \leq \rho(b)} \int_{\rho(a)}^b G(t, s)f(s, u(\omega(s)) + u_0(\omega(s)))\nabla s \\ &\geq \Gamma \int_{\rho(a)}^b G(s, s)f(s, u(\omega(s)) + u_0(\omega(s)))\nabla s \\ &\geq \Gamma \max_{a \leq t \leq b} \int_{\rho(a)}^b G(t, s)f(s, u(\omega(s)) + u_0(\omega(s)))\nabla s \\ &\geq \Gamma \|Au\|_{[a, b]} = \Gamma \|Au\|, \quad t \in [a, b]. \end{aligned} \quad (2.26)$$

Thus, $A(K) \subset K$. \square

Lemma 2.5. $A : K \rightarrow K$ is completely continuous.

Lemma 2.6. If

$$\lim_{v \rightarrow 0^+} \frac{f(t, v)}{v} = \infty, \quad \lim_{v \rightarrow +\infty} \frac{f(t, v)}{v} = \infty, \quad (2.27)$$

for all $t \in [a, b]$, then there exist $0 < r_0 < R_0 < \infty$ such that $i(A, K_r, K) = 0$, for $0 < r \leq r_0$ and $i(A, K_R, K) = 0$, for $R \geq R_0$.

Proof. Choose $M > 0$ such that

$$\Gamma^2 M \int_{E_1} G(s, s) \nabla s > 1. \quad (2.28)$$

By using the first equality of (2.27), we can choose $r_0 > 0$ such that

$$f(t, v) \geq Mv, \quad 0 \leq v \leq r_0. \quad (2.29)$$

If $u \in \partial K_r$ ($0 < r \leq r_0$), then for $t_0 \in [a, \rho(b)]$, we have

$$\begin{aligned} (Au)(t_0) &= \int_{\rho(a)}^b G(t_0, s) f(s, u(\omega(s)) + u_0(\omega(s))) \nabla s \\ &\geq \Gamma \int_{\rho(a)}^b G(s, s) f(s, u(\omega(s)) + u_0(\omega(s))) \nabla s \\ &\geq \Gamma \int_{E_1} G(s, s) f(s, u(\omega(s))) \nabla s \\ &\geq \Gamma M \int_{E_1} G(s, s) u(\omega(s)) \nabla s \\ &\geq \Gamma^2 \|u\| M \int_{E_1} G(s, s) \nabla s \\ &> \|u\|. \end{aligned} \quad (2.30)$$

Therefore we get

$$\|Au\| > \|u\|, \quad \forall u \in \partial K_r. \quad (2.31)$$

Thus, we have from Theorem 1.1, $i(A, K_r, K) = 0$, for $0 < r \leq r_0$. On the other hand, the second equality of (2.27) implies for every $M > 0$, there is an $R_0 > r_0$, such that

$$f(t, v) \geq Mv, \quad v \geq \Gamma R_0. \quad (2.32)$$

Here we choose $M > 0$ satisfying (2.28). For $u \in \partial K_R$, $R \geq R_0$, we have definition of K_R that

$$u(t) \geq \Gamma \|u\| = \Gamma R, \quad t \in [a, \rho(b)]. \quad (2.33)$$

It follows from (2.32) that

$$\begin{aligned} (Au)(t_0) &= \int_{\rho(a)}^b G(t_0, s) f(s, u(\omega(s)) + u_0(\omega(s))) \nabla s \\ &\geq \Gamma \int_{\rho(a)}^b G(s, s) f(s, u(\omega(s)) + u_0(\omega(s))) \nabla s \\ &\geq \Gamma \int_{E_1} G(s, s) f(s, u(\omega(s))) \nabla s \\ &\geq \Gamma M \int_{E_1} G(s, s) u(\omega(s)) \nabla s \\ &\geq \Gamma^2 MR \int_{E_1} G(s, s) \nabla s \\ &> R = \|u\|. \end{aligned} \quad (2.34)$$

This shows that

$$\|Au\| > \|u\|, \quad \forall u \in K_R. \quad (2.35)$$

Thus, by Theorem 1.1, we conclude that $i(A, K_R, K) = 0$ for $R \geq R_0$. The proof is therefore complete. \square

3. Existence of One Positive Solution

In this section, we investigate the conditions for the existence of at least one positive solution of the BVP (1.3).

In the next theorem, we will also assume that the following condition on $f(t, v)$.
(H6):

$$\liminf_{v \rightarrow 0^+} \min_{t \in [a, b]} \frac{f(t, v)}{v} > k\lambda_1, \quad \limsup_{v \rightarrow +\infty} \max_{t \in [a, b]} \frac{f(t, v)}{v} < q\lambda_1, \quad (3.1)$$

where $k > 0$ is large enough such that

$$k\Gamma \int_{E_1} \varphi_1(s) \nabla s \geq \int_{\rho(a)}^b \varphi_1(s) \nabla s, \quad (3.2)$$

and $q > 0$ is small enough such that

$$\Gamma \int_{\rho(a)}^{\rho(b)} \varphi_1(s) \nabla s \geq q \int_{\rho(a)}^b \varphi_1(s) \nabla s, \tag{3.3}$$

where $\varphi_1(t)$ ($\varphi_1(t) > 0, t \in [a, b]$) is the eigenfunction related to the smallest eigenvalue λ_1 ($\lambda_1 > 0$) of the eigenvalue problem:

$$\begin{aligned} -\varphi_1^{\Delta \nabla}(t) + q(t)\varphi_1(t) &= \lambda \varphi_1(t), \\ \alpha \varphi_1(\rho(a)) - \beta \varphi_1^{\Delta}(\rho(a)) &= 0, \quad \gamma \varphi_1(b) + \delta \varphi_1^{\Delta}(b) = 0. \end{aligned} \tag{3.4}$$

Theorem 3.1. *If (H1)–(H6) are satisfied, then the BVP (1.3) has at least one positive solution.*

Proof. Fix $0 < m < 1 < m_1$ and let $f_1(u) = u^m + u^{m_1}$ for $u \geq 0$. Then, $f_1(u)$ satisfies (2.27). Define $\tilde{A} : K \rightarrow K$ by

$$(\tilde{A}u)(t) := \begin{cases} (\tilde{A}_1u)(t), & \text{if } \rho(\tau_1) \leq t \leq a, \\ \int_{\rho(a)}^b G(t, s) f_1(u(\omega(s)) + u_0(\omega(s))) \nabla s, & \text{if } a \leq t \leq b, \\ (\tilde{A}_2x)(t), & \text{if } b \leq t \leq \sigma(\tau_2), \end{cases} \tag{3.5}$$

where

$$\begin{aligned} (\tilde{A}_1u)(t) &= \begin{cases} e_{\alpha/\beta}(t, a) \int_{\rho(a)}^b G(a, s) f_1(u(\omega(s)) + u_0(\omega(s))) \nabla s, & \text{if } \rho(\tau_1) \leq t \leq a, \beta > 0, \\ 0, & \text{if } \rho(\tau_1) \leq t \leq a, \beta = 0, \end{cases} \\ (\tilde{A}_2u)(t) &= \begin{cases} e_{-\gamma/\delta}(t, b) \int_{\rho(a)}^b G(b, s) f_1(u(\omega(s)) + u_0(\omega(s))) \nabla s, & \text{if } b \leq t \leq \sigma(\tau_2), \delta > 0, \\ 0, & \text{if } b \leq t \leq \sigma(\tau_2), \delta = 0. \end{cases} \end{aligned} \tag{3.6}$$

Then \tilde{A} is a completely continuous operator. One has from Lemma 2.6 that there exist $0 < r_0 < R_0 < \infty$ such that

$$0 < r \leq r_0 \text{ implies } i(\tilde{A}, K_r, K) = 0, \tag{3.7}$$

$$R \geq R_0 \text{ implies } i(\tilde{A}, K_R, K) = 0. \tag{3.8}$$

Define $H : [0, 1] \times K \rightarrow K$ by $H(t, u) = (1 - t)Au + t\tilde{A}u$ then H is a completely continuous operator. By the first equality in (H6) and the definition of f_1 , there are $\epsilon > 0$ and $0 < r_1 \leq r_0$ such that

$$\begin{aligned} f(t, u) &\geq (k\lambda_1 + \epsilon)u, \quad \forall t \in [a, b], 0 \leq u \leq r_1, \\ f_1(u) &\geq (k\lambda_1 + \epsilon)u, \quad 0 \leq u \leq r_1. \end{aligned} \tag{3.9}$$

We now prove that $H(t, u) \neq u$ for all $t \in [0, 1]$ and $u \in \partial K_{r_1}$. In fact, if there exists $t_0 \in [0, 1]$ and $u_1 \in \partial K_{r_1}$ such that $H(t_0, u_1) = u_1$, then $u_1(t)$ satisfies the equation

$$\begin{aligned} & -u_1^{\Delta \nabla}(t) + q(t)u_1(t) \\ & = (1 - t_0)f(t, u_1(\omega(t)) + u_0(\omega(t))) + t_0f_1(u_1(\omega(t)) + u_0(\omega(t))), \quad a \leq t \leq b \end{aligned} \quad (3.10)$$

and the boundary conditions

$$\begin{aligned} \alpha u_1(\rho(t)) - \beta u_1^\Delta(\rho(t)) &= 0, \quad t \in [\tau_1, a], \\ \gamma u_1(t) + \delta u_1^\Delta(t) &= 0, \quad t \in [b, \tau_2]. \end{aligned} \quad (3.11)$$

Multiplying both sides of (3.10) by $\varphi_1(t)$, then integrating from a to b , and using integration by parts in the left-hand side two times, we obtain

$$\begin{aligned} & \lambda_1 \int_{\rho(a)}^b \varphi_1(s)u_1(s)\nabla s \\ & = \int_{\rho(a)}^b [(1 - t_0)f(s, u_1(\omega(s)) + u_0(\omega(s))) + t_0f_1(u_1(\omega(s)) + u_0(\omega(s)))]\varphi_1(s)\nabla s \\ & = (1 - t_0) \int_{\rho(a)}^b f(s, u_1(\omega(s)) + u_0(\omega(s)))\varphi_1(s)\nabla s \\ & \quad + t_0 \int_{\rho(a)}^b f_1(u_1(\omega(s)) + u_0(\omega(s)))\varphi_1(s)\nabla s. \end{aligned} \quad (3.12)$$

Combining (3.9) and (3.12), we get

$$\begin{aligned} \lambda_1 \int_{\rho(a)}^b \varphi_1(s)u_1(s)\nabla s &\geq (1 - t_0) \int_{E_1} f(s, u_1(\omega(s)))\varphi_1(s)\nabla s + t_0 \int_{E_1} f_1(u_1(\omega(s)))\varphi_1(s)\nabla s \\ &\geq (1 - t_0)(k\lambda_1 + \epsilon) \int_{E_1} u_1(\omega(s))\varphi_1(s)\nabla s \\ &\quad + t_0(k\lambda_1 + \epsilon) \int_{E_1} u_1(\omega(s))\varphi_1(s)\nabla s \\ &= \left(\lambda_1 + \frac{\epsilon}{k}\right)k \int_{E_1} \varphi_1(s)u_1(\omega(s))\nabla s \\ &\geq \left(\lambda_1 + \frac{\epsilon}{k}\right)k\Gamma \|u_1\| \int_{E_1} \varphi_1(s)\nabla s \\ &\geq \left(\lambda_1 + \frac{\epsilon}{k}\right)\|u_1\| \int_{\rho(a)}^b \varphi_1(s)\nabla s. \end{aligned} \quad (3.13)$$

We also have

$$\lambda_1 \int_{\rho(a)}^b \varphi_1(s) u_1(s) \nabla s \leq \lambda_1 \|u_1\| \int_{\rho(a)}^b \varphi_1(s) \nabla s. \quad (3.14)$$

Equations (3.13) and (3.14) lead to

$$\lambda_1 \geq \lambda_1 + \frac{\epsilon}{k}. \quad (3.15)$$

This is impossible. Thus $H(t, u) \neq u$ for $u \in \partial K_{r_1}$ and $t \in [0, 1]$. By (3.7) and the homotopy invariance of the fixed-point index (see [11]), we get that

$$i(A, K_{r_1}, K) = i(H(0, \cdot), K_{r_1}, K) = i(H(1, \cdot), K_{r_1}, K) = i(\tilde{A}, K_{r_1}, K) = 0. \quad (3.16)$$

On the other hand, according to the second inequality of (H6), there exist $\epsilon > 0$ and $R' > R_0$ such that

$$f(t, u) \leq (q\lambda_1 - \epsilon)u, \quad u > R', \quad t \in [a, b]. \quad (3.17)$$

We define

$$C := \max_{a \leq t \leq b, 0 \leq u \leq R'} |f(t, u) - (q\lambda_1 - \epsilon)u| + 1, \quad (3.18)$$

then it follows that

$$f(t, u) \leq (q\lambda_1 - \epsilon)u + C, \quad u \geq 0, \quad t \in [a, b]. \quad (3.19)$$

Define $H_1 : [0, 1] \times K \rightarrow K$ by $H_1(t, u) = tAu$, then H_1 is a completely continuous operator. We claim that there exists $R_1 \geq R'$ such that

$$H_1(t, u) \neq u, \quad \text{for } t \in [0, 1], \quad u \in K, \quad \|u\| \geq R_1. \quad (3.20)$$

In fact, if $H_1(t_0, u_1) = u_1$ for some $u_1 \in K$ and $0 \leq t_0 \leq 1$, then

$$\begin{aligned} \lambda_1 \int_{\rho(a)}^b u_1(s) \varphi_1(s) \nabla s &\leq \int_{\rho(a)}^b f(s, u_1(\omega(s)) + u_0(\omega(s))) \varphi_1(s) \nabla s \\ &\leq q \left(\lambda_1 - \frac{\epsilon}{q} \right) \|u_1 + u_0\| \int_{\rho(a)}^b \varphi_1(s) \nabla s + C \int_{\rho(a)}^b \varphi_1(s) \nabla s \\ &\leq q \left(\lambda_1 - \frac{\epsilon}{q} \right) \|u_1\| \int_{\rho(a)}^b \varphi_1(s) \nabla s + C_1 \int_{\rho(a)}^b \varphi_1(s) \nabla s, \end{aligned} \tag{3.21}$$

$$\begin{aligned} \lambda_1 \int_{\rho(a)}^b u_1(s) \varphi_1(s) \nabla s &\geq \lambda_1 \int_{\rho(a)}^{\rho(b)} u_1(s) \varphi_1(s) \nabla s \\ &\geq \lambda_1 \Gamma \|u_1\| \int_{\rho(a)}^{\rho(b)} \varphi_1(s) \nabla s \\ &\geq \lambda_1 q \|u_1\| \int_{\rho(a)}^b \varphi_1(s) \nabla s, \end{aligned} \tag{3.22}$$

where $C_1 = q(\lambda_1 - \epsilon/q)\|u_0\| + C$. Combining (3.21) with (3.22), we have

$$\|u_1\| \leq \frac{C_1}{\epsilon} = \tilde{R}_1. \tag{3.23}$$

Let $R_1 = \max\{R', \tilde{R}_1\} + 1$. Then we get

$$H_1(t, u) \neq u, \quad \text{for } t \in [0, 1], u \in K, \|u\| \geq R_1. \tag{3.24}$$

Consequently, by the homotopy invariance of the fixed-point index, we have

$$i(A, K_{R_1}, K) = i(H_1(1, \cdot), K_{R_1}, K) = i(H_1(0, \cdot), K_{R_1}, K) = i(\Theta, K_{R_1}, K) = 1, \tag{3.25}$$

where Θ is zero operator. Use (3.16) and (3.25) to conclude that

$$i(A, K_{R_1} \setminus K_{r_1}, K) = i(A, K_{R_1}, K) - i(A, K_{r_1}, K) = 1 - 0 = 1. \tag{3.26}$$

Hence, A has a fixed point in $(K_{R_1} \setminus K_{r_1})$.

Let $y(t) = u(t) + u_0(t)$. Since $y(t) = u(t)$ for $t \in [a, b]$ and $0 < r_1 \leq \|u\| = \|u\|_{[a,b]} = \|y\|_{[a,b]} \leq R_1$.
(H7)

$$\begin{aligned} \limsup_{v \rightarrow 0^+} \max_{t \in [a,b]} \frac{f(t, v)}{v} &< q\lambda_1, \\ \liminf_{v \rightarrow +\infty} \min_{t \in [a,b]} \frac{f(t, v)}{v} &> k\lambda_1, \quad \xi(t) \equiv 0, \eta(t) \equiv 0. \end{aligned} \tag{3.27}$$

□

Theorem 3.2. *If (H1)–(H5) and (H7) are satisfied, then the BVP (1.3) has at least one positive solution.*

Proof. Define $H_1 : [0, 1] \times K \rightarrow K$ by $H_1(t, u) = tAu$, then H_1 is a completely continuous operator. By the first inequality in (H7), there exist $\epsilon > 0$ and $0 < r_1 \leq r_0$ such that

$$f(t, v) \leq (q\lambda_1 - \epsilon)v, \quad \forall t \in [a, b], 0 \leq v \leq r_1. \tag{3.28}$$

We claim that $H_1(t, u) \neq u$ for $0 \leq t \leq 1$ and $u \in \partial K_{r_1}$. In fact, if there exist $0 \leq t_0 \leq 1$ and $u_1 \in \partial K_{r_1}$ such that $H_1(t_0, u_1) = u_1$, then $u_1(t)$ satisfies the boundary condition (3.11). Since $\xi(t) \equiv 0, \eta(t) \equiv 0$, we have $u_0(t) \equiv 0$. Then we have

$$-u_1^{\Delta \nabla}(t) + q(t)u_1(t) = t_0 f(t, u_1(\omega(t))), \quad a \leq t \leq b. \tag{3.29}$$

Multiplying the last equation by $\varphi_1(t)$ integrating from a to b , by (3.28), we obtain

$$\begin{aligned} \lambda_1 \int_{\rho(a)}^b \varphi_1(s)u_1(s) \nabla s &= t_0 \int_{\rho(a)}^b f(s, u_1(\omega(s)))\varphi_1(s) \nabla s \\ &\leq \int_{\rho(a)}^b f(s, u_1(\omega(s)))\varphi_1(s) \nabla s \\ &\leq (q\lambda_1 - \epsilon) \|u_1\| \int_{\rho(a)}^b \varphi_1(s) \nabla s, \end{aligned} \tag{3.30}$$

then we have

$$\begin{aligned} \lambda_1 \int_{\rho(a)}^b u_1(s)\varphi_1(s) \nabla s &\geq \lambda_1 \int_{\rho(a)}^{\rho(b)} u_1(s)\varphi_1(s) \nabla s \\ &\geq \lambda_1 \Gamma \|u_1\| \int_{\rho(a)}^{\rho(b)} \varphi_1(s) \nabla s \\ &\geq \lambda_1 q \|u_1\| \int_{\rho(a)}^b \varphi_1(s) \nabla s. \end{aligned} \tag{3.31}$$

Equations (3.30) and (3.31) lead to

$$\lambda_1 q \leq \lambda q_1 - \epsilon. \tag{3.32}$$

This is impossible. By homotopy invariance of the fixed-point index, we get that

$$i(A, K_{r_1}, K) = i(H_1(1, \cdot), K_{r_1}, K) = i(H_1(0, \cdot), K_{r_1}, K) = i(\Theta, K_{r_1}, K) = 1. \tag{3.33}$$

Define $H : [0, 1] \times K \rightarrow K$ by $H(t, u) = (1 - t)Au + t\tilde{A}u$, then H is a completely continuous operator. By the second inequality in (H7), and definition of f_1 , there exist $\epsilon > 0$ and $R' > R_0$ such that

$$\begin{aligned} f(t, u) &\geq (k\lambda_1 + \epsilon)u, \quad u > R', t \in [a, b], \\ f_1(u) &\geq (k\lambda_1 + \epsilon)u, \quad u > R'. \end{aligned} \quad (3.34)$$

We define

$$C := \max_{a \leq t \leq b, 0 \leq u \leq R'} |f(t, u) - (k\lambda_1 + \epsilon)u| + \max_{0 \leq u \leq R'} |f_1(u) - (k\lambda_1 + \epsilon)u| + 1, \quad (3.35)$$

then, it is obvious that

$$\begin{aligned} f(t, u) &\geq (k\lambda_1 + \epsilon)u - C, \quad \forall t \in [a, b], u \geq 0, \\ f_1(u) &\geq (k\lambda_1 + \epsilon)u - C, \quad u \geq 0. \end{aligned} \quad (3.36)$$

We claim that there exists $R_1 \geq R'$ such that

$$H(t, u) \neq u, \quad \text{for } t \in [0, 1], u \in K, \|u\| \geq R_1. \quad (3.37)$$

In fact, if $H(t_0, u_1) = u_1$ for some $u_1 \in K$ and $0 \leq t_0 \leq 1$, then using (3.36), it is analogous to the argument of (3.13) and (3.14) that

$$\begin{aligned} \lambda_1 \int_{\rho(a)}^b \varphi_1(s) u_1(s) \nabla s &= \int_{\rho(a)}^b [(1 - t_0)f(s, u_1(\omega(s)) + t_0 f_1(u_1(\omega(s))))] \varphi_1(s) \nabla s \\ &\geq \int_{E_1} [(1 - t_0)f(s, u_1(\omega(s)) + t_0 f_1(u_1(\omega(s))))] \varphi_1(s) \nabla s \\ &\geq \int_{E_1} \{ (1 - t_0)[(k\lambda_1 + \epsilon)u_1(\omega(s)) - C] + t_0[(k\lambda_1 + \epsilon)u_1(\omega(s)) - C] \} \\ &\quad \times \varphi_1(s) \nabla s \\ &= \int_{E_1} [(k\lambda_1 + \epsilon)u_1(\omega(s)) - C] \varphi_1(s) \nabla s \\ &\geq \left(\lambda_1 + \frac{\epsilon}{k} \right) k\Gamma \|u_1\| \int_{E_1} \varphi_1(s) \nabla s - C \int_{E_1} \varphi_1(s) \nabla s, \\ \lambda_1 \int_{\rho(a)}^b \varphi_1(s) u_1(s) \nabla s &\leq \lambda_1 \|u_1\| \int_{\rho(a)}^b \varphi_1(s) \nabla s \leq \lambda_1 \|u_1\| k\Gamma \int_{E_1} \varphi_1(s) \nabla s. \end{aligned} \quad (3.38)$$

Equation (3.38) leads to $\|u_1\| \leq C/\epsilon\Gamma = \tilde{R}_1$. Let $R_1 = \max\{R', \tilde{R}_1\} + 1$. Then we get

$$H_1(t, u) \neq u, \quad \text{for } t \in [0, 1], u \in K, \|u\| \geq R_1. \quad (3.39)$$

Consequently, by (3.8) and the homotopy invariance of the fixed-point index, we have

$$i(A, K_{R_1}, K) = i(H(0, \cdot), K_{R_1}, K) = i(H(1, \cdot), K_{R_1}, K) = i(\tilde{A}, K_{R_1}, K) = 0. \quad (3.40)$$

In view of (3.33) and (3.40), we obtain

$$i(A, K_{R_1} \setminus K_{r_1}, K) = i(A, K_{R_1}, K) - i(A, K_{r_1}, K) = 0 - 1 = -1. \quad (3.41)$$

Therefore, A has a fixed point in $(K_{R_1} \setminus K_{r_1})$. The proof is completed. \square

Corollary 3.3. *Using the following (H8) or (H9) instead of (H6) or (H7), the conclusions of Theorems 3.1 and 3.2 are true. For $t \in [a, b]$,*

(H8)

$$\lim_{v \rightarrow 0^+} \left(\frac{f(t, v)}{v} \right) = +\infty, \quad \lim_{v \rightarrow +\infty} \left(\frac{f(t, v)}{v} \right) = 0 \quad (\text{sublinear}); \quad (3.42)$$

(H9)

$$\lim_{v \rightarrow 0^+} \left(\frac{f(t, v)}{v} \right) = 0, \quad \lim_{v \rightarrow +\infty} \left(\frac{f(t, v)}{v} \right) = +\infty \quad (\text{superlinear}), \quad \xi(t) \equiv 0, \quad \eta(t) \equiv 0. \quad (3.43)$$

4. Existence of Three Positive Solutions

In this section, using Theorem 1.2 (the Leggett-Williams fixed-point theorem) we prove the existence of at least three positive solutions to the BVP (1.3).

Define the continuous concave functional $\psi : K \rightarrow [0, \infty)$ to be $\psi(u) := \min_{t \in [a, \rho(b)]} u(t)$, and the constants

$$M := \left[\min_{t \in [a, \rho(b)]} \int_{\rho(a)}^{\rho(b)} G(t, s) \nabla s \right]^{-1}, \quad (4.1)$$

$$N := \int_{\rho(a)}^b G(s, s) \nabla s. \quad (4.2)$$

Theorem 4.1. *Suppose there exists constants $0 < p < q < q/\Gamma \leq r$ such that*

(D1) $f(t, v) < p/N$ for $t \in [a, b], v \in [0, p]$;

(D2) $f(t, v) \geq qM$ for $t \in [a, \rho(b)], v \in [q, q/\Gamma]$;

(D3) *one of the following is satisfied:*

(a) $\limsup_{v \rightarrow \infty} \max_{t \in [a, b]} (f(t, v)/v) < 1/N$,

(b) *there exists a constant $r > q/\Gamma$ such that $f(t, v) \leq r/N$ for $t \in [a, b]$ and $v \in [0, r]$,*

where Γ , M , and R , are as defined in (2.12), (4.1), (4.2), respectively. Then the boundary value problem (1.3) has at least three positive solutions u_1, u_2 , and u_3 satisfying

$$\|u_1\| < p, \quad \min_{t \in [a, \rho(b)]} (u_2)(t) > q, \quad p < \|u_3\| \quad \text{with} \quad \min_{t \in [a, \rho(b)]} (u_3)(t) < q. \quad (4.3)$$

Proof. The technique here similar to that used in [5] Again the cone K , the operator A is the same as in the previous sections. For all $u \in K$ we have $\psi(u) \leq \|u\|$. If $u \in \bar{K}_r$, then $\|u\| \leq r$ and the condition (a) of (D3) imply that

$$\limsup_{v \rightarrow \infty} \max_{t \in [a, b]} \frac{f(t, v)}{v} < \frac{1}{N}. \quad (4.4)$$

Thus there exist a $\zeta > 0$ and $\epsilon < 1/R$ such that if $v > \zeta$, then $\max_{t \in [a, b]} (f(t, v)/v) < \epsilon$. For $\lambda := \max\{f(t, v) : v \in [0, \zeta], t \in [a, b]\}$, we have $f(t, v) \leq \epsilon v + \lambda$ for all $v \geq 0$, for all $t \in [a, b]$. Pick any

$$r > \max \left\{ \frac{q}{\Gamma}, \frac{\lambda}{1/N - \epsilon} \right\}. \quad (4.5)$$

Then $u \in \bar{K}_r$, implies that

$$\begin{aligned} \|Au\| &= \max_{t \in [a, b]} \int_{\rho(a)}^b G(t, s) f(s, u(\omega(s))) \nabla s \\ &\leq (\epsilon \|u\| + \lambda) N < \epsilon r N + r(1 - \epsilon N) = r. \end{aligned} \quad (4.6)$$

Thus $A : \bar{K}_r \rightarrow K_r$.

The condition (b) of (D3) implies that there exists a positive number r such that $f(t, v) \leq r/N$ for $t \in [a, b]$ and $v \in [0, r]$. If $u \in \bar{K}_r$, then

$$\|Au\| = \max_{t \in [a, b]} \int_{\rho(a)}^b G(t, s) f(s, u(\omega(s))) \nabla s \leq \left(\frac{r}{N}\right) N \leq r. \quad (4.7)$$

Thus $A : \bar{K}_r \rightarrow \bar{K}_r$. Consequently, the assumption (D3) holds, then there exist a number r such that $r > q/\Gamma$ and $A : \bar{K}_r \rightarrow K_r$.

The remaining conditions of Theorem 1.2 will now be shown to be satisfied.

By (D1) and argument above, we can get that $A : \bar{K}_p \rightarrow K_p$. Hence, condition (ii) of Theorem 1.2 is satisfied.

We now consider condition (i) of Theorem 1.2. Choose $u_K(t) \equiv (s + q)/2$ for $t \in [a, b]$, where $s = q/\Gamma$. Then $u_K(t) \in K(\psi, q, q/\Gamma)$ and $\psi(u_K) = \psi((s+q)/2) > q$, so that $\{K(\psi, p, q/\Gamma) : \psi(u) > q\} \neq \{\}$. For $u \in K(\psi, q, q/\Gamma)$, we have $q \leq u(t) \leq q/\Gamma, t \in [a, \rho(b)]$. Combining with (D2), we get

$$f(t, u) \geq qM, \quad (4.8)$$

for $t \in [a, \rho(b)]$. Thus, we have

$$\begin{aligned} \psi(Au) &= \min_{t \in [a, \rho(b)]} \int_{\rho(a)}^b G(t, s) f(s, u(\omega(s))) \nabla s \\ &> \min_{t \in [a, \rho(b)]} \int_{\rho(a)}^{\rho(b)} G(t, s) f(s, u(\omega(s))) \nabla s \geq \frac{qM}{M} = q. \end{aligned} \tag{4.9}$$

As a result, $u \in K(\psi, q, s)$ yields $\psi(Au) > q$.

Lastly, we consider Theorem 1.2(iii). Recall that $A : K \rightarrow K$. If $u \in K(\psi, q, r)$ and $\|Au\| > q/\Gamma$, then

$$\psi(Au) = \min_{t \in [a, \rho(b)]} Au(t) \geq \Gamma \|Au\| > \Gamma \frac{q}{\Gamma} = q. \tag{4.10}$$

Thus, all conditions of Theorem 1.2 are satisfied. It implies that the TPBVP (1.3) has at least three positive solutions u_1, u_2, u_3 with

$$\|u_1\| < p, \quad \psi(u_2) > q, \quad p < \|u_3\| \quad \text{with } \psi(u_3) < q. \tag{4.11}$$

□

5. Examples

Example 5.1. Let $\mathbb{T} = \{(n/2) : n \in \mathbb{N}_0\}$. Consider the BVP:

$$\begin{aligned} -y^{\Delta \nabla}(t) + y(t) &= \sqrt{y(2t-1)}, \quad t \in [0, 3] \subset \mathbb{T}, \\ y(\rho(t)) - 2y^\Delta(\rho(t)) &= t^2, \quad t \in [-1, 0], \\ 3y(t) + 4y^\Delta(t) &= t - 3, \quad t \in [3, 5]. \end{aligned} \tag{5.1}$$

Then $a = 0, b = 3, \tau_1 = -1, \tau_2 = 5, \alpha = 1, \beta = 2, \gamma = 3, \delta = 4$, and

$$q(t) = 1, \quad \xi(t) = t^2, \quad \eta(t) = t - 3, \quad f(t, v) = \sqrt{v}, \quad \omega(t) = 2t - 1. \tag{5.2}$$

Since $\lim_{v \rightarrow 0^+} (f(t, v)/v) = +\infty, \lim_{v \rightarrow +\infty} (f(t, v)/v) = 0$. It is clear that (H1)–(H5) and (H8) are satisfied. Thus, by Corollary 3.3, the BVP (5.1) has at least one positive solution.

Example 5.2. Let us introduce an example to illustrate the usage of Theorem 4.1. Let

$$\mathbb{T} = \left\{ \frac{4n^2}{n^2 + 9} : n \in \mathbb{N}_0 \right\} \cup \{4\}. \tag{5.3}$$

Consider the TPBVP:

$$\begin{aligned} -y^{\Delta \nabla}(t) &= \frac{500y^2(t^2 + 1)}{y^2(t^2 + 1) + 300}, \quad t \in \left[\frac{2}{5}, 2 \right], \\ y(0) &= 0, \\ y^{\Delta}(t) &= 0, \quad t \in [2, 5]. \end{aligned} \quad (5.4)$$

Then $a = 2/5$, $b = 2$, $\tau_1 = 2/5$, $\tau_2 = 5$, $\alpha = 1$, $\beta = 0$, $\delta = 1$, $\gamma = 0$, and

$$\omega(t) = t^2 + 1, \quad \xi(t) = \left(t - \frac{2}{5} \right)^2, \quad \eta(t) = 0, \quad f(t, v) = \frac{500v^2}{v^2 + 300}. \quad (5.5)$$

The Green function of the BVP (5.4) has the form

$$G(t, s) = \begin{cases} s, & \text{if } 0 \leq s \leq t \leq \frac{64}{25}, \\ t, & \text{if } 0 \leq t \leq s \leq \frac{64}{25}. \end{cases} \quad (5.6)$$

Clearly, f is continuous and increasing $[0, \infty)$. We can also see that $\alpha + \gamma > 0$. By (2.12), (4.1), and (4.2), we get $\Gamma = 1/5$, $M = 65/32$, and $N = 11496/4225$.

Now we check that (D1), (D2), and (b) of (D3) are satisfied. To verify (D1), as $f(1/10) = 0.01666611113$, we take $p = 1/10$, then

$$f(y) < \frac{p}{N} = 0.03675191371, \quad y \in [0, p] \quad (5.7)$$

and (D1) holds. Note that $f(3/2) = 3.722084367$, when we set $q = 3/2$,

$$f(y) \geq qM = 3.046875000, \quad y \in [q, 5q] \quad (5.8)$$

holds. It means that (D2) is satisfied. Let $r = 1500$, we have

$$f(y) \leq 500 < \frac{r}{N} = 551.2787056, \quad y \in [0, r], \quad (5.9)$$

from $\lim_{y \rightarrow \infty} f(y) = 500$, so that (b) of (D3) is met. Summing up, there exist constants $p = 1/10$, $q = 3/2$, and $r = 1500$ satisfying

$$0 < p < q < \frac{q}{\Gamma} < r. \quad (5.10)$$

Thus, by Theorem 4.1, the TPBVP (5.4) has at least three positive solutions y_1, y_2, y_3 with

$$\|y_1\| < \frac{1}{10}, \quad \varphi(y_2) > \frac{3}{2}, \quad \frac{1}{10} < \|y_3\| \quad \text{with } \varphi(y_3) < \frac{3}{2}. \quad (5.11)$$

References

- [1] D. Jiang and P. Weng, "Existence of positive solutions for boundary value problems of second-order functional-differential equations," *Electronic Journal of Qualitative Theory of Differential Equations*, no. 6, pp. 1–13, 1998.
- [2] N. Aykut, "Existence of positive solutions for boundary value problems of second-order functional difference equations," *Computers & Mathematics with Applications*, vol. 48, no. 3-4, pp. 517–527, 2004.
- [3] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales: An Introduction with Applications*, Birkhäuser, Boston, Mass, USA, 2001.
- [4] M. Bohner and A. Peterson, Eds., *Advances in Dynamic Equations on Time Scales*, Birkhäuser, Boston, Mass, USA, 2003.
- [5] D. R. Anderson, "Existence of solutions for nonlinear multi-point problems on time scales," *Dynamic Systems and Applications*, vol. 15, no. 1, pp. 21–33, 2006.
- [6] D. R. Anderson, R. Avery, J. Davis, J. Henderson, and W. Yin, "Positive solutions of boundary value problems," in *Advances in Dynamic Equations on Time Scales*, pp. 189–249, Birkhäuser, Boston, Mass, USA, 2003.
- [7] D. R. Anderson, R. Avery, and A. Peterson, "Three positive solutions to a discrete focal boundary value problem," *Journal of Computational and Applied Mathematics*, vol. 88, no. 1, pp. 103–118, 1998.
- [8] F. M. Atici and G. Sh. Guseinov, "On Green's functions and positive solutions for boundary value problems on time scales," *Journal of Computational and Applied Mathematics*, vol. 141, no. 1-2, pp. 75–99, 2002.
- [9] C. J. Chyan, J. M. Davis, J. Henderson, and W. K. C. Yin, "Eigenvalue comparisons for differential equations on a measure chain," *Electronic Journal of Differential Equations*, vol. 1998, no. 35, pp. 1–7, 1998.
- [10] J. M. Davis, P. W. Eloe, and J. Henderson, "Comparison of eigenvalues for discrete Lidstone boundary value problems," *Dynamic Systems and Applications*, vol. 8, no. 3-4, pp. 381–388, 1999.
- [11] C.-Z. Bai and J.-X. Fang, "Existence of multiple positive solutions for functional differential equations," *Computers & Mathematics with Applications*, vol. 45, no. 12, pp. 1797–1806, 2003.
- [12] J. M. Davis, K. R. Prasad, and W. K. C. Yin, "Nonlinear eigenvalue problems involving two classes of functional differential equations," *Houston Journal of Mathematics*, vol. 26, no. 3, pp. 597–608, 2000.
- [13] E. R. Kaufmann and Y. N. Raffoul, "Positive solutions for a nonlinear functional dynamic equation on a time scale," *Nonlinear Analysis: Theory, Methods & Applications*, vol. 62, no. 7, pp. 1267–1276, 2005.
- [14] D. J. Guo and V. Lakshmikantham, *Nonlinear Problems in Abstract Cones*, vol. 5 of *Notes and Reports in Mathematics in Science and Engineering*, Academic Press, Boston, Mass, USA, 1988.
- [15] R. W. Leggett and L. R. Williams, "Multiple positive fixed points of nonlinear operators on ordered Banach spaces," *Indiana University Mathematics Journal*, vol. 28, no. 4, pp. 673–688, 1979.