

Research Article

On Two Systems of Difference Equations

Bratislav Iričanin¹ and Stevo Stević²

¹ Faculty of Electrical Engineering, University of Belgrade, Bulevar Kralja Aleksandra 73,
11120 Belgrade, Serbia

² Mathematical Institute, Serbian Academy of Sciences, Knez Mihailova 36/III, 11000 Beograd, Serbia

Correspondence should be addressed to Stevo Stević, sstevic@ptt.rs

Received 11 January 2010; Accepted 9 March 2010

Academic Editor: Leonid Berezansky

Copyright © 2010 B. Iričanin and S. Stević. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We give very short and elegant proofs of the main results in the work of Yalcinkaya et al. (2008).

1. Introduction and a Proof of Some Resent Results

Motivated by our paper [1], the authors of [2] studied the following two systems of difference equations:

$$x_{n+1}^{(i)} = \frac{x_n^{(i+1 \pmod k)}}{x_n^{(i+1 \pmod k)} - 1}, \quad i = 1, \dots, k, \quad n \in \mathbb{N}_0, \quad (1.1)$$

$$x_{n+1}^{(i)} = \frac{x_n^{(i-1 \pmod k)}}{x_n^{(i-1 \pmod k)} - 1}, \quad i = 1, \dots, k, \quad n \in \mathbb{N}_0, \quad (1.2)$$

where we regard that $0 \pmod k = k \pmod k = k$.

Following line by line the proofs of the main results in [1] they proved the following result (see Theorems 2.1 and 2.4 in [2])

Theorem A. Assume $k \in \mathbb{N}$, then the following statements are true.

- If $k = 0 \pmod 2$, then every (well-defined) solution of systems (1.1) and (1.2) is periodic with period k .
- If $k = 1 \pmod 2$, then every (well-defined) solution of systems (1.1) and (1.2) is periodic with period $2k$.

Here we give a very short and elegant proof of Theorem A.

Proof of Theorem A. By using the change $y_n^{(i)} = x_n^{(i)} - 1$, $i = 1, \dots, k$, system (1.1) becomes

$$y_n^{(i)} = \left(y_{n-1}^{(i+1 \pmod{k})} \right)^{-1}, \quad i = 1, \dots, k, \quad n \in \mathbb{N}, \quad (1.3)$$

while system (1.2) becomes

$$y_n^{(i)} = \left(y_{n-1}^{(i-1 \pmod{k})} \right)^{-1}, \quad i = 1, \dots, k, \quad n \in \mathbb{N}. \quad (1.4)$$

From (1.3) and (1.4), for each $i \in \{1, \dots, k\}$, and $n \geq k$, we obtain correspondingly that

$$\begin{aligned} y_n^{(i)} &= \left(y_{n-k}^{(i+1+k-1 \pmod{k})} \right)^{(-1)^k} = \left(y_{n-k}^{(i)} \right)^{(-1)^k}, \\ y_n^{(i)} &= \left(y_{n-k}^{(i-1-(k-1) \pmod{k})} \right)^{(-1)^k} = \left(y_{n-k}^{(i)} \right)^{(-1)^k}. \end{aligned} \quad (1.5)$$

From (1.5), with $k = 0 \pmod{2}$, it follows that

$$y_n^{(i)} = y_{n-k}^{(i)}, \quad i = 1, \dots, k, \quad (1.6)$$

from which the statement in (a) easily follows.

If $k = 1 \pmod{2}$, we have that

$$y_n^{(i)} = \left(y_{n-k}^{(i)} \right)^{-1}, \quad i = 1, \dots, k, \quad (1.7)$$

from which it follows that

$$y_n^{(i)} = y_{n-2k}^{(i)}, \quad i = 1, \dots, k, \quad (1.8)$$

$n \geq 2k$, implying the statement in (b), as desired. \square

2. An Extension on Theorem A

Here we extend Theorem A in a natural way. Let $\gcd(k, l)$ denote the greatest common divisor of the integers k and l , $\text{lcm}(k, l)$ the least common multiple of k and l , and for $r \in \mathbb{N}$ let $f^{[r]}(x) = f(f^{[r-1]}(x))$, where $f^{[1]}(x) = f(x)$.

Theorem 2.1. *Assume that f is a real function such that $f^{[r]}(x) \equiv x$ on its domain of definition, for some $r \in \mathbb{N}$, then all well-defined solutions of the system of difference equations*

$$x_n^{(1)} = f(x_{n-1}^{(2)}), x_n^{(2)} = f(x_{n-1}^{(3)}), \dots, x_n^{(k)} = f(x_{n-1}^{(1)}), \quad n \in \mathbb{N}_0, \quad (2.1)$$

are periodic with period $T = \text{lcm}(k, r)$.

Proof. We use our method of “prolongation” described in [1]. Note that for each $s \in \mathbb{N}$, system (2.1) is equivalent to a system of ks difference equations of the same form, where

$$x_n^{(i)} = x_n^{(jk+i)}, \quad (2.2)$$

for every $n \in \mathbb{N}_0$, $i \in \{1, \dots, k\}$ and $j = 1, \dots, s$.

From (2.1) and since $f^{[r]}(x) \equiv x$, for $n \geq r - 1$ we have

$$x_n^{(i+1)} = f(x_{n-1}^{(i+2)}) = f^{[2]}(x_{n-2}^{(i+3)}) = \dots = f^{[r]}(x_{n-r}^{(i+r+1)}) = x_{n-r}^{(i+r+1)}. \quad (2.3)$$

for each $i \in \{1, 2, \dots, k\}$, and every $n \geq r - 1$.

It is clear that

$$T = k \cdot r_1 = k_1 \cdot r, \quad (2.4)$$

where $r_1, k_1 \in \mathbb{N}$ are such that $\gcd(k, r_1) = 1$ and $\gcd(k_1, r) = 1$.

From (2.3) we have

$$x_n^{(i+1)} = x_{n-r}^{(i+r+1)} = \dots = x_{n-k_1r}^{(i+k_1r+1)} = x_{n-kr_1}^{(i+kr_1+1)} = x_{n-T}^{(i+1)}, \quad (2.5)$$

for each $i = 0, 1, \dots, k - 1$, and $n \geq T - 1$, from which the result follows. \square

The following result is proved similarly. Hence we omit its proof.

Theorem 2.2. *Assume that f is a real function such that $f^{[r]}(x) \equiv x$ on its domain of definition, for some $r \in \mathbb{N}$, then all well-defined solutions of the system of difference equations*

$$x_n^{(2)} = f(x_{n-1}^{(1)}), \dots, x_n^{(k)} = f(x_{n-1}^{(k-1)}), \quad x_n^{(1)} = f(x_{n-1}^{(k)}), \quad n \in \mathbb{N}_0, \quad (2.6)$$

are periodic with period $T = \text{lcm}(k, r)$.

Remark 2.3. The proof of Theorem A follows from Theorems 2.1 and 2.2. Indeed, note that the function $f(x) = x/(x - 1)$ satisfies the condition $f^{[2]}(x) \equiv x$ on its domain of definition. By Theorems 2.1 and 2.2 we know that all well-defined solutions of systems (1.1) and (1.2) are periodic with period $T = \text{lcm}(k, 2)$, from which the result follows.

Remark 2.4. We also have to say that the main result in [3] is a trivial consequence of a result in [1] (see Remark 5 therein). Just note that the simple change of variables $x_n^{(i)} = ay_n^{(i)}$, $i \in \{1, \dots, k\}$, transforms their system (1.3) satisfying conditions $a_1 = a_2 = \dots = a_k = a$ and $b_1 = b_2 = \dots = b_k = b = a^2$, into system (4) in [1].

Acknowledgment

The results in this note were presented at the talk: S. Stević, on a class of max-type difference equations and some of our old results, *Progress on Difference Equations 2009*, Bedlewo, Poland, May 25–29, 2009.

References

- [1] B. D. Iričanin and S. Stević, "Some systems of nonlinear difference equations of higher order with periodic solutions," *Dynamics of Continuous, Discrete & Impulsive Systems. Series A*, vol. 13, no. 3-4, pp. 499–507, 2006.
- [2] İ. Yalçınkaya, C. Çinar, and M. Atalay, "On the solutions of systems of difference equations," *Advances in Difference Equations*, vol. 2008, Article ID 143943, 9 pages, 2008.
- [3] G. Papaschinopoulos, C. J. Schinas, and G. Stefanidou, "On a k -order system of Lyness-type difference equations," *Advances in Difference Equations*, vol. 2007, Article ID 31272, 13 pages, 2007.