

**PSEUDO-SASAKIAN MANIFOLDS ENDOWED  
WITH A CONTACT CONFORMAL CONNECTION**

**VLADISLAV V. GOLDBERG and RADU ROSCA**

Department of Mathematics  
N.J. Institute of Technology  
Newark, N.J. 07102 U.S.A.

(Received January 30, 1985)

**ABSTRACT.** Pseudo-Sasakian manifolds  $\tilde{M}(U, \xi, \tilde{\eta}, \tilde{g})$  endowed with a contact conformal connection are defined. It is proved that such manifolds are space forms  $\tilde{M}(K), K < 0$ , and some remarkable properties of the Lie algebra of infinitesimal transformations of the principal vector field  $\tilde{U}$  on  $\tilde{M}$  are discussed. Properties of the leaves of a co-isotropic foliation on  $\tilde{M}$  and properties of the tangent bundle manifold  $T\tilde{M}$  having  $\tilde{M}$  as a basis are studied.

**KEY WORDS AND PHRASES.** Witt frame, CICR submanifold, relative contact infinitesimal transformation, U-contact concircular pairing, differential form of Godbillon-Vey, form of E. Cartan, Finslerian form, mechanical system, dynamical system, spray, CR product.

**1980 SUBJECT CLASSIFICATION CODES.** 53C25, 53C40, 53B25

**1. INTRODUCTION.**

In the last years many papers have been concerned with Sasakian manifold  $\tilde{M}(\phi, \xi, \tilde{\eta}, \tilde{g})$  and related structures. Recently Rosca [1] has defined pseudo-Sasakian manifolds  $\tilde{M}(U, \xi, \tilde{\eta}, \tilde{g})$  and Goldberg and Rosca [2] have studied CICR submanifolds (i.e. co-isotropic CR submanifolds) of  $\tilde{M}(U, \xi, \tilde{\eta}, \tilde{g})$ .

In the present paper we study  $(2m+1)$ -dimensional pseudo-Sasakian manifolds of index  $m+1, m > 4$ , structured by a contact conformal (abr. c.c.) connection. It is proved that such manifolds are hyperbolic space forms  $\tilde{M}(K), K < 0$ , and with the c.c. connection (which in fact is a natural generalization of the connection defined by Rosca [3]) is associated (compare with Rosca [3]) a so denominated principal vector field  $\tilde{U}$ .

The paper is organized as follows. In Section 3 we develop some basic results induced by the c.c. connection and some remarkable properties of the Lie algebra of infinitesimal transformations defined by  $\tilde{U}$ . It is shown that

- (i)  $\tilde{U}$  (resp.  $U\tilde{U}$ ) is divergence free (resp. defines an infinitesimal homothety) on  $\tilde{M}$  and all connection forms on  $\tilde{M}$  are integral relations of invariance for  $U\tilde{U}$  (see Lichnerowicz [4]);
- (ii)  $\tilde{U}$  and  $U\tilde{U}$  define an U-contact concircular pairing (in the sense of Rosca [5]) and any contact extension of  $\tilde{U}$  is a relative contact

*infinitesimal transformation* (in the sense of Rosca [3]) of the canonical 1-form  $\tilde{\eta}$ ;

(11)  $\tilde{U}$  and  $U\tilde{U}$  define both infinitesimal automorphisms of  $(2q+1)$ -forms  $\tilde{\beta}_q = L^q \tilde{U}$  ( $q < m$ ) where  $\tilde{u}$  (resp.  $l$ ) is the dual form of  $\tilde{U}$  (resp. the  $(1,1)$ -operator taken with respect to the 2-form  $\tilde{\Omega} = d\tilde{\eta}/2$ ). Accordingly, if  $\Sigma_\beta$  is the exterior differential system defined by  $\{\tilde{\beta}_q\}$ ,  $\tilde{U}$  and  $U\tilde{U}$  may be considered as *isovectors* of  $\Sigma_\beta$ .

Section 4 is concerned with a *co-isotropic foliation*  $F_c$  on  $\tilde{M}$ . The leaves  $M_c$  of  $F_c$  are CICR submanifolds of  $\tilde{M}$  and if  $\text{codim } M_c = l$ , then the form of Godbillon-Vey on  $M_c$  (see Lichnerowicz [6]) is a  $(2l+1)$ -form  $w_c$  which is a *relative integral invariant* of  $U = \tilde{U}|_{M_c}$ .

Further the necessary and sufficient conditions for  $M_c$  to be *foliate* is that the isotropic component  $U^\perp$  of  $U$  vanishes. In this case  $M_c$  is a *CR product* (see Yano and Kon [7] and Rosca [8]).

Finally using some notions introduced by Yano and Ishihara [9] and also by Klein [10], we consider in Section 5 certain properties of the tangent bundle manifold  $T\tilde{M}$  having the manifold  $\tilde{M}(U, \xi, \tilde{\eta}, \tilde{g})$  as a basis.

It is proved that the *complete lifts*  $\tilde{\eta}^C$  and  $\tilde{u}^C$  of  $\tilde{\eta}$  and  $\tilde{u}$  respectively are *homogeneous* of degree one and that the form of E. Cartan  $\tilde{\Pi}$  on  $T\tilde{M}$  is a *Finslerian form*. Furthermore, we may associate with  $\tilde{\Pi}$  a *regular mechanical system* whose *dynamical system* is a *spray* on  $\tilde{M}$ .

2. PRELIMINARIES.

Let  $(\tilde{M}, \tilde{g})$  be a  $(2m+1)$ -dimensional connected pseudo-Riemannian manifold of signature  $(m+1, m)$  and suppose that  $m > 4$ .

At each point  $\tilde{p} \in \tilde{M}$  one has the standard decomposition (see Rosca [1]):

$$T_{\tilde{p}}(\tilde{M}) = H_{\tilde{p}} \oplus l_{\tilde{p}} \tag{2.1}$$

where  $T_{\tilde{p}}$ ,  $H_{\tilde{p}}$ , and  $l_{\tilde{p}}$  are the tangent space, a  $(2m)$ -dimensional *neutral* vector space, and a *time-like* line orthogonal to  $H_{\tilde{p}}$ , respectively.

Let  $S_{\tilde{p}}, S_{\tilde{p}}^* \subset H_{\tilde{p}}$  be two *self-orthogonal* (abbreviation s.o.)  $m$ -distributions which define an *involutive* automorphism  $U$  of square  $+1$  ( $U$  is the *para complex* operator defined by Libermann [11]). Let  $\xi \in T_{\tilde{p}}$  and  $\tilde{\eta} \in \Lambda^1(\tilde{M})$  be the pairing which defines a contact structure  $\sigma_c$  on  $\tilde{M}$ , and  $\tilde{\nabla}$  be the covariant differentiation operator defined by the metric tensor  $\tilde{g}$ . Then if for any vector fields  $\tilde{Z}, \tilde{Z}'$  on  $\tilde{M}$  the structure tensors  $(U, \xi, \tilde{\eta}, \tilde{g})$  satisfy

$$\begin{cases} U^2 \tilde{Z} = \tilde{Z} - \tilde{\eta}(\tilde{Z})\xi, & \tilde{g}(U\tilde{Z}, U\tilde{Z}') = -\tilde{g}(\tilde{Z}, \tilde{Z}') + \tilde{\eta}(\tilde{Z})\tilde{\eta}(\tilde{Z}'), \\ \tilde{g}(\tilde{Z}, \xi) = \tilde{\eta}(\tilde{Z}), & \tilde{\nabla}_{\tilde{Z}} \xi = U\tilde{Z}, \\ d\tilde{\eta}(\tilde{Z}, \tilde{Z}') = -2\tilde{g}(U\tilde{Z}, \tilde{Z}'), & \tilde{\eta}(\xi) = 1, \end{cases} \tag{2.2}$$

the manifold  $\tilde{M}(U, \xi, \tilde{\eta}, \tilde{g})$  has been called a *pseudo-Sasakian manifold* (see Rosca [1]).

In order to study real *co-isotropic* and *isotropic foliations* on  $\tilde{M}$  (that is *improper* immersions in  $\tilde{M}$ ), we consider an adapted field of *Witt frames*:  $\tilde{W} = \{h_A: A, B, C = 0, 1, \dots, 2m\}$ . The vectors  $h_a$  and  $h_{a^*}$  ( $a=1, \dots, m; a^*=a+m$ ) are *null* and  $h_0 = \xi$  is the *anisotropic* vector field of the  $W$ -basis  $\{h_A\}$ . We set

$$\tilde{S}_{\tilde{p}} = \{h_a\}, \quad \tilde{S}_{\tilde{p}}^* = \{h_{a^*}\} \tag{2.3}$$

and as is known, one has

$$\begin{cases} \tilde{g}(h_a, h_b^*) = \delta_{ab}, & \tilde{g}(\xi, h_a) = 0, \\ \tilde{g}(\xi, h_a^*) = 0, & \tilde{g}(\xi, \xi) = 1 \end{cases} \quad (2.4)$$

and

$$Uh_a = h_a, \quad Uh_a^* = -h_a^*, \quad U\xi = 0. \quad (2.5)$$

If  $\tilde{W}^* = \{\tilde{\omega}^\Lambda\}$  is the cobasis associated with  $\tilde{W}$ , we set  $\tilde{\omega}^0 = \tilde{\eta}$  and the line element  $d\tilde{p}$  ( $d\tilde{p}$  is a canonical vector 1-form and is independent on any connection on  $\tilde{M}$ ) is given by

$$d\tilde{p} = \tilde{\omega}^\Lambda \otimes h_\Lambda. \quad (2.6)$$

It follows from (2.4) that the metric tensor  $\tilde{g}$  is:

$$\tilde{g} = 2 \sum_a \tilde{\omega}^a \otimes \tilde{\omega}^{a^*} + \tilde{\eta} \otimes \tilde{\eta}. \quad (2.7)$$

If  $\tilde{\gamma}_B^\Lambda = \tilde{\gamma}_{BC}^\Lambda \tilde{\omega}^C$  ( $\tilde{\gamma}_{BC}^\Lambda \in C^\infty(\tilde{M})$ ) and  $\tilde{\Theta}_B^\Lambda$  are the connection forms and the curvature 2-forms on the bundle  $\tilde{W}(\tilde{M})$  respectively, then the structure equations (E. Cartan) may be written in the indexless form as follows:

$$\tilde{\nabla}h = \tilde{\theta} \otimes h, \quad (2.8)$$

$$d\tilde{\omega} = -\tilde{\theta} \wedge \tilde{\omega}, \quad (2.9)$$

$$d\tilde{\theta} = -\tilde{\theta} \wedge \tilde{\theta} + \tilde{\Theta}. \quad (2.10)$$

Referring to (2.4) and (2.8), one has

$$\begin{cases} \tilde{\theta}_b^a + \tilde{\theta}_{a^*}^{b^*} = 0, & \tilde{\theta}_b^{a^*} = 0, & \tilde{\theta}_{b^*}^a = 0, \\ \tilde{\theta}_a^0 + \tilde{\theta}_0^{a^*} = 0, & \tilde{\theta}_0^a + \tilde{\theta}_{a^*}^0 = 0 \end{cases} \quad (2.11)$$

and

$$\tilde{\theta}_a^0 = \tilde{\omega}^{a^*}, \quad \tilde{\theta}_{a^*}^0 = -\tilde{\omega}^a. \quad (2.12)$$

By virtue of (2.8), (2.9), and (2.11) one has

$$d\tilde{\eta} = 2 \sum_a \tilde{\omega}^a \wedge \tilde{\omega}^{a^*} \quad (2.13)$$

and

$$\tilde{\nabla}\xi = U d\tilde{p} \implies \langle \tilde{\nabla}_Z \xi, \tilde{Z}' \rangle + \langle \tilde{\nabla}_{Z'} \xi, Z \rangle = 0 \quad (2.14)$$

where  $Z$  and  $Z'$  are any vector fields on  $\tilde{M}$ .

In the following we agree to call the 2-form

$$\tilde{\Omega} = \sum_a \tilde{\omega}^a \wedge \tilde{\omega}^{a^*} \quad (2.15)$$

the fundamental 2-form on  $\tilde{M}$

Since by (2.11) one has

$$\tilde{\theta}_a^a + \tilde{\theta}_{a^*}^{a^*} = 0, \quad \tilde{\Theta}_a^a + \tilde{\Theta}_{a^*}^{a^*} = 0, \quad (2.16)$$

we shall call

$$\tilde{\theta}_R = \sum_a \tilde{\theta}^a_a \quad (2.17)$$

and

$$\tilde{\Theta}_R = \sum_a \tilde{\Theta}^a_a \quad (2.18)$$

the Ricci 1-form and the Ricci 2-form respectively (see Rosca [12]). As is known, the form  $\overset{\vee}{\Theta}_R$  defines the first class of Chern of  $\overset{\vee}{M}$ .

Using (2.10) and referring to (2.12) and (2.15), one quickly obtains

$$d\overset{\vee}{\Theta}_R = \overset{\vee}{\Theta}_R - \overset{\vee}{\Omega}. \tag{2.19}$$

The above equation proves that the 2-forms  $\overset{\vee}{\Theta}_R$  and  $\overset{\vee}{\Omega}$  are homologous. Hence the two cocycles  $\overset{\vee}{\Theta}_R$  and  $\overset{\vee}{\Omega}$  belong to the 2-cohomology class  $H^2(\overset{\vee}{M})$  of  $\overset{\vee}{M}$ .

Let now  $F_c$  be a coisotropic foliation on  $\overset{\vee}{M}$  and denote by  $M_c$  a maximal integral manifold (leave) of  $F_c$ . It has been shown by Goldberg and Rosca [2] that  $M_c$  is a contact CR submanifold of  $\overset{\vee}{M}$ , that is there exists a differentiable distribution  $D: p \rightarrow D_p \subset T_p(M_c)$ ,  $p \in M_c$  (one denotes the induced elements on  $M_c$  by suppressing  $\vee$ ) satisfying:

- (i)  $D$  is invariant i.e.  $UD_p \subseteq D_p$ , and
- (ii) the complementary orthogonal distribution  $D^\perp: p \rightarrow D^\perp_p \subset T_p(M_c)$  is anti-invariant i.e.  $UD^\perp_p \subseteq T^\perp_p(M_c)$ .

The distribution  $D$  (resp.  $D^\perp$ ) is called the horizontal (resp. vertical) distribution. Such type of CR submanifolds is called CICR submanifolds (see Goldberg and Rosca [2]).

3. PSEUDO-SASAKIAN MANIFOLDS ENDOWED WITH A CONTACT CONFORMAL CONNECTION.

As a natural generalization of the definition given by Rosca [3], we assume that the structure equations (2.9) are written in the form

$$\begin{cases} d\overset{\vee}{\omega}^a = (\overset{\vee}{u} + \overset{\vee}{\eta}) \wedge \overset{\vee}{\omega}^a + \overset{\vee}{t}_a \overset{\vee}{\Omega}^a, \\ d\overset{\vee}{\omega}^{a*} = (\overset{\vee}{u} - \overset{\vee}{\eta}) \wedge \overset{\vee}{\omega}^{a*} + \overset{\vee}{t}_{a*} \overset{\vee}{\Omega}^a \end{cases} \tag{3.1}$$

where  $\overset{\vee}{\Omega}^a = d\overset{\vee}{\eta}/2$ ,  $\overset{\vee}{t}_a, \overset{\vee}{t}_{a*} \in C^\infty(M)$ , and  $\overset{\vee}{u} \in \Lambda^1(\overset{\vee}{M})$  is a closed 1-form. Note that  $\overset{\vee}{t}_a$  and  $\overset{\vee}{t}_{a*}$  are the components of a vector field

$$\overset{\vee}{U} = \sum_a (\overset{\vee}{t}_a h_a + \overset{\vee}{t}_{a*} h_{a*}) \tag{3.2}$$

of constant length.

We shall say (see Rosca [3]) that in this case the pseudo-Sasakian manifold  $\overset{\vee}{M}$  is endowed with a contact conformal (abr. c.c.) connection. We also agree to call  $\overset{\vee}{U}$  the principal vector field associated with this connection.

Since  $\overset{\vee}{g}(\overset{\vee}{U}, \overset{\vee}{U}) = \text{const}$ , we may write by (3.2) that

$$\sum_a \overset{\vee}{t}_{a*} \overset{\vee}{t}_a = c, \quad c = \text{const}. \tag{3.3}$$

Taking exterior differentials of (3.1), we get

$$\begin{cases} d\overset{\vee}{t}_a = (\overset{\vee}{u} + \overset{\vee}{\eta}) \overset{\vee}{t}_a - 2\overset{\vee}{\omega}^a, \\ d\overset{\vee}{t}_{a*} = (\overset{\vee}{u} - \overset{\vee}{\eta}) \overset{\vee}{t}_{a*} - 2\overset{\vee}{\omega}^{a*}. \end{cases} \tag{3.4}$$

Denote by  $\Sigma$  the exterior differential system defined by equations (3.1) and (3.4) and by  $I$  the ideal corresponding to  $\Sigma$ . The exterior differentiation of (3.4) where  $\overset{\vee}{\omega}^a$  and  $\overset{\vee}{\omega}^{a*}$  satisfy (3.1),  $\overset{\vee}{\Omega}^a = d\overset{\vee}{\eta}/2$ ,  $d\overset{\vee}{u} = 0$ , leads to the identity. Because of this,  $dI \subset I$ , that is  $\Sigma$  is a closed system. It follows from this that the system  $\Sigma$  defining the pseudo-Sasakian manifold  $\overset{\vee}{M}$  endowed with a c.c. connection is completely integrable and its solution depends on  $2m$  constants (the number of equations in (3.4)).

From (3.4) and (3.3) we also obtain

$$\hat{c}\hat{u} = \sum_a (\hat{t}_a \hat{\omega}^a - \hat{t}_a \hat{\omega}^a) \tag{3.5}$$

and  $\hat{u}(\hat{U}) = 0$  which shows that  $\hat{u}$  is an *integral relation of invariance* for  $\hat{U}$  (see Lichnerowicz [4]). In the following we agree to call  $\hat{u}$  the *principal Pfaffian* associated with the c.c. connection.

Consider now the 1-form

$$\hat{v} = \sum_a (\hat{t}_a \hat{\omega}^a + \hat{t}_a \hat{\omega}^a). \tag{3.6}$$

Taking the exterior differential of  $\hat{v}$ , one finds with the help of (3.1) and (3.4) that  $c = 2$ . In this case we deduce

$$d\hat{v} = 2\hat{u} \wedge \hat{v}, \tag{3.7}$$

and this equation asserts that  $\hat{v}$  is *exterior recurrent* (see Datta [13] with  $2\hat{u}$  as the *recurrence 1-form*).

By (2.4) and (2.5) one easily finds

$$\hat{u}(\hat{U}\hat{U}) = \hat{v}(\hat{U}) = \hat{g}(\hat{U}, \hat{U}) = \hat{g}(\hat{U}\hat{U}, \hat{U}\hat{U}) = 2 \sum_a \hat{t}_a \hat{t}_a. \tag{3.8}$$

Hence if  $b : T(\hat{M}) \rightarrow T^*(\hat{M})$  is the *musical isomorphism* with respect to  $\hat{g}$  (see Poor [14]), we may write:  $\hat{u} = b(\hat{U}\hat{U})$ ,  $\hat{v} = b(\hat{U})$ . Since  $\hat{u}$  is closed, it follows from (3.7) that the manifold  $\hat{M}$  under consideration is foliated by 2-codimensional submanifolds orthogonal to  $\hat{U}$  and  $\hat{U}\hat{U}$ .

Next if  $\mu : \hat{Z} \rightarrow i\hat{Z}\hat{\Omega}$ ,  $T(\hat{M}) \rightarrow T^*(\hat{M})$  is the bundle isomorphism defined by  $\hat{\Omega} = d\hat{\eta}/2$ , one readily finds

$$\mu(\hat{U}) = 2\hat{\eta}. \tag{3.9}$$

In the following we agree to call the *presymplectic form*  $\hat{\Omega}$  ( $\dim \ker(\hat{\Omega}) \neq 0$ ) the *fundamental 2-form* on  $\hat{M}$ .

Let now  $\hat{U}_f = \hat{U} + \hat{\xi}\xi$  ( $\hat{\xi} \in C^\infty(\hat{M})$ ) be a *contact extension* of  $\hat{U}$  and  $\mathcal{L}_{\hat{U}_f}$  the Lie derivative with respect to  $\hat{U}_f$ . Then by (3.9) one quickly finds  $d\mathcal{L}_{\hat{U}_f} \hat{\eta} = 0$ .

Therefore according to the definition given by Rosca [3], we may say that  $\hat{U}_f$  is a *relative contact infinitesimal transformation* of  $\hat{\eta}$ .

Denote now by  $\hat{\sigma}_S$  (resp.  $\hat{\sigma}_{S^*}$ ) the simple unit form which corresponds to  $\hat{S}_p^\nu$  (resp.  $\hat{S}_p^{\nu^*}$ ). One has

$$\begin{cases} \hat{\sigma}_S = \hat{\omega}^1 \wedge \dots \wedge \hat{\omega}^m, \\ \hat{\sigma}_{S^*} = \hat{\omega}^{1^*} \wedge \dots \wedge \hat{\omega}^{m^*}, \end{cases} \tag{3.10}$$

and by (3.1) the exterior differentials of (3.10) are

$$\begin{cases} d\hat{\sigma}_S = [m(\hat{u} + \hat{\eta}) - \hat{v}] \wedge \hat{\sigma}_S, \\ d\hat{\sigma}_{S^*} = [m(\hat{u} - \hat{\eta}) + \hat{v}] \wedge \hat{\sigma}_{S^*}. \end{cases} \tag{3.11}$$

Since  $\hat{\sigma}_S$  and  $\hat{\sigma}_{S^*}$  are both exterior recurrent, it follows from a well-known property that both co-isotropic distributions  $\hat{S} + \{\xi\}$  and  $\hat{S}^* + \{\xi\}$  are *involutive* (orth.  $(\hat{S} + \{\xi\}) = \hat{S}$ ; orth.  $(\hat{S}^* + \{\xi\}) = S^*$ ). It is worth to emphasize that this property is true for any pseudo-Sasakian manifold.

Now with the help of (3.1), one finds that the connection forms are given by

$$\begin{cases} \tilde{\theta}_a^a = \tilde{\zeta}_a^{\omega^*} + \tilde{\zeta}_a^{\omega^*} + \tilde{\nu}/2 \quad (\text{no summation}), \\ \tilde{\theta}_b^a = \tilde{\zeta}_b^{\omega^*} + \tilde{\zeta}_a^{\omega^*} . \end{cases} \tag{3.12}$$

By (3.12) and (3.6) one finds

$$\tilde{\theta}_R^{\tilde{\nu}} = (m+2)\tilde{\nu}/2 \tag{3.13}$$

and (3.7) shows that  $\tilde{\theta}_R^{\tilde{\nu}}$  is exterior recurrent.

Coming back to relations (3.12), one readily finds

$$\tilde{\theta}_a^a(U\tilde{U}) = 0, \quad \tilde{\theta}_b^a(U\tilde{U}) = 0 . \tag{3.14}$$

Therefore we may say that all connection forms of the pseudo-Sasakian manifold  $\tilde{M}$  under consideration are *integral relations of invariance* for the vector field  $U\tilde{U}$ .

Denote now by  $\tilde{\tau}$  the volume element of  $\tilde{M}$ . One may take a local orientation such that

$$\tilde{\tau} = \tilde{\sigma}_S \wedge \tilde{\sigma}_S^* \wedge \tilde{\eta} \tag{3.15}$$

and denote by  $*$ :  $\Lambda^{q_T^* \tilde{M}} \rightarrow \Lambda^{2m+1-q_T^* \tilde{M}}$  the *star operator* determined by  $\tilde{\tau}$ . If, like usually,  $\mathfrak{X}^{\tilde{M}}$  means the vector space of sections over  $T\tilde{M}$ , then, as is known, for any vector field  $\tilde{Z} \in \mathfrak{X}^{\tilde{M}}$  one has

$$*\text{div } \tilde{Z} = (\text{div } \tilde{Z})\tilde{\tau} = \text{div } \tilde{Z}\tilde{\tau} = \mathfrak{L}_{\tilde{Z}}\tilde{\tau} . \tag{3.16}$$

Making use of (3.4), (3.11), (3.16), and the fact that

$$\tilde{U} = \sum_a (\tilde{\zeta}_a^h h_a + \tilde{\zeta}_a^* h_a^*), \tag{3.17}$$

one finds after some calculations:

$$\text{div } \tilde{U} = 0, \quad \text{div}(U\tilde{U}) = 2 \sum_a \tilde{\zeta}_a^h \tilde{\zeta}_a^* = 4 . \tag{3.18}$$

Hence  $\tilde{U}$  is *divergence free* and  $U\tilde{U}$  is an *infinitesimal homothety* on  $\tilde{M}$ .

Now if  $\tilde{Z} = \tilde{Z}^A h_A$ ,  $\tilde{Z}' = (\tilde{Z}')^A h_A \in \mathfrak{X}^{\tilde{M}}$  are any vector fields, then, as is known (see Poor [14]), one has

$$\tilde{\nabla}_{\tilde{Z}} \tilde{Z} = (\mathfrak{L}_{\tilde{Z}} \tilde{Z}^A) h_A + \tilde{Z}^A (\tilde{\nabla}_{\tilde{Z}} h_A) .$$

Therefore, by (2.3), (3.4), and (3.12) we get

$$\begin{cases} \tilde{\nabla}_{\tilde{Z}} \tilde{U} = (\tilde{\eta}(\tilde{Z}) + \tilde{\nu}(\tilde{Z})) U\tilde{U} - 2\tilde{U}(\tilde{Z}) \xi , \\ \tilde{\nabla}_{\tilde{Z}} U\tilde{U} = (\tilde{\eta}(\tilde{Z}) + \tilde{\nu}(\tilde{Z})) \tilde{U} + \tilde{\nu}(\tilde{Z}) \xi . \end{cases} \tag{3.19}$$

We also note that since  $b(U\tilde{U}) = \tilde{U}$  is a closed form, we may say (see Poor [14]) that  $\tilde{\nabla}_{U\tilde{U}}$  is *self-adjoint*.

According to the definition given by Rosca [5] and Rosca and Verstraelen [15], the formulae (3.19) show that the vector field  $\tilde{U}$  defines a *U-contact concircular pairing*.

Denote by  $D_U$  the 3-distribution defined by  $\{\tilde{U}, U\tilde{U}, \xi\}$ . By (2.2), (3.5), and (3.6) one readily finds from (3.19) that

$$[\tilde{U}, \xi] = 0, \quad [U\tilde{U}, \xi] = 0 . \tag{3.20}$$

Hence both vector fields  $\tilde{U}$  and  $U\tilde{U}$  commute with  $\xi$  and by (3.19) and (3.20) we see that  $D_U$  defines a *3-foliation* on  $\tilde{M}$ .

It is worth now to make the following considerations.

Let  $\tilde{Z} \in \mathfrak{X}\tilde{M}$  be any vector field on  $\tilde{M}$ . Then one has the general Bochner formula (see Poor [14]) on  $\tilde{M}$ :

$$2\langle \text{tr } \tilde{\nabla}^2 \tilde{Z}, \tilde{Z} \rangle + 2\|\tilde{\nabla} \tilde{Z}\|^2 + \tilde{\lambda} \|\tilde{Z}\|^2 = 0 \tag{3.21}$$

where  $\tilde{\Delta} = d \circ \delta + \delta \circ d$  is the Laplace-Roltrvimi operator (or Laplacian) on  $\Lambda^2 \tilde{M}$ , and the trace (abr. tr) is calculated with respect to the metric tensor  $\tilde{g}$  of  $\tilde{M}$ .

Applying formula (3.21) to the principal vector field  $\tilde{U}$  and taking into account (2.7), one has

$$\text{tr } \tilde{\nabla}^2 \tilde{U} = \sum_a \tilde{\nabla}_{h_a} \tilde{\nabla}_{h_{a^*}} \tilde{U} + \sum_a \tilde{\nabla}_{h_{a^*}} \tilde{\nabla}_{h_a} \tilde{U} + \tilde{\nabla}_\xi (\tilde{\nabla}_\xi \tilde{U}) \tag{3.22}$$

and

$$\|\tilde{\nabla} \tilde{U}\|^2 = 2 \sum_a \langle \tilde{\nabla}_{h_a} \tilde{U}, \tilde{\nabla}_{h_{a^*}} \tilde{U} \rangle + \langle \tilde{\nabla}_\xi \tilde{U}, \tilde{\nabla}_\xi \tilde{U} \rangle. \tag{3.23}$$

Now by (2.14), (3.4), (3.5), (3.16), and (3.19) one finds

$$\begin{cases} \tilde{\nabla}_{h_{a^*}} \tilde{\nabla}_{h_a} \tilde{U} = \tilde{\zeta}_a \tilde{\zeta}_{a^*} \tilde{U} + (2 - \tilde{\zeta}_a \tilde{\zeta}_{a^*} / 2) U \tilde{U} + (3 \tilde{\zeta}_a \tilde{\zeta}_{a^*} / 2 - 2) \xi + \tilde{\zeta}_{a^*} h_{a^*}, \\ \tilde{\nabla}_{h_a} \tilde{\nabla}_{h_{a^*}} \tilde{U} = \tilde{\zeta}_a \tilde{\zeta}_{a^*} \tilde{U} - (2 - \tilde{\zeta}_a \tilde{\zeta}_{a^*} / 2) U \tilde{U} + (3 \tilde{\zeta}_a \tilde{\zeta}_{a^*} / 2 - 2) \xi + \tilde{\zeta}_a h_a, \\ \tilde{\nabla}_\xi \tilde{\nabla}_\xi U = U. \end{cases} \tag{3.24}$$

Since we have found  $\sum_a \tilde{\zeta}_a \tilde{\zeta}_{a^*} = 2$ , we derive from (3.22), (3.23), (3.24), and (3.21) that  $\tilde{U}$  satisfies (3.21) and this equation is consistent with  $\|\tilde{\nabla} \tilde{U}\|^2 = 4$ .

Let  $L$  be the operator of type (1,1) defined by the fundamental 2-form  $\tilde{\Omega}$ . Denote then by  $\tilde{\beta}_q = L^q \tilde{U} = \tilde{U} \wedge (\Lambda \tilde{\Omega})^q \in \Lambda^{2q+1} \tilde{M}$ . Since  $\tilde{U}$  and  $\tilde{\Omega}$  are both closed, one finds by (3.9) and making use of the properties of the Lie derivative  $\mathfrak{L} = i \circ d + d \circ i$  that

$$\mathfrak{L} \tilde{U} \tilde{\beta}_q = 0. \tag{3.25}$$

Hence  $\tilde{U}$  is an infinitesimal automorphism of all  $(2q+1)$ -forms  $\tilde{\beta}_q$  ( $q < m$ ).

On the other hand, since  $\tilde{g}(\tilde{U}, \tilde{U}) = \text{const}$ , we may say in similar manner as in the case of a Sasakian manifold that  $\tilde{U}$  defines with  $U \tilde{U}$  an  $U$ -section.

Like usually denote by

$$R(\tilde{Z}, \tilde{Z}') = [\tilde{\nabla}_{\tilde{Z}} \tilde{\nabla}_{\tilde{Z}'} - \tilde{\nabla}_{[\tilde{Z}, \tilde{Z}']} \tilde{\nabla}_{\tilde{Z}'}, \tilde{Z}, \tilde{Z}' \in \mathfrak{X}\tilde{M} \tag{3.26}$$

the curvature operator. Then, as is known, the sectional curvature  $K(\tilde{U}, U \tilde{U})$  defined by  $\tilde{U}$  and  $U \tilde{U}$  is given by

$$K(\tilde{U}, U \tilde{U}) = \frac{R(\tilde{U}, U \tilde{U}, \tilde{U}, U \tilde{U})}{\tilde{g}(\tilde{U}, \tilde{U}) \tilde{g}(U \tilde{U}, U \tilde{U}) - (\tilde{g}(\tilde{U}, U \tilde{U}))^2} \tag{3.27}$$

where

$$R(\tilde{U}, U \tilde{U}, \tilde{U}, U \tilde{U}) = \tilde{g}(R(\tilde{U}, U \tilde{U}) U \tilde{U}, \tilde{U}). \tag{3.28}$$

Making use of (3.5), (3.6), and (3.19), one finds

$$[\tilde{U}, U \tilde{U}] = 4(\tilde{U} + 2\xi) \tag{3.29}$$

and

$$R(\tilde{U}, U \tilde{U}) U \tilde{U} = 4(5\tilde{U} + 8\xi). \tag{3.30}$$

Hence by (3.27) and (3.28) one gets  $K(\tilde{U}, U\tilde{U}) = -\frac{1}{5}$ . Now referring to (2.10) and (3.12) one finds after some calculations

$$\begin{aligned} \tilde{\Theta}_a^a &= \tilde{\nu}_S \wedge \tilde{\nu}_{S^*} + \tilde{\nu}_S \wedge \tilde{t}_a^{\nu a^*} - \tilde{\nu}_{S^*} \wedge \tilde{t}_{a^*}^{\nu a} \\ &+ \tilde{t}_a^{\nu} \tilde{t}_{a^*}^{\nu} + \omega^{a^*} \wedge \omega^a \quad (\text{no summation}) \end{aligned} \tag{3.31}$$

where we have set

$$\left\{ \begin{aligned} \tilde{\nu}_S &= \sum_a \tilde{t}_a^{\nu} \omega^a \in \Lambda^1 \tilde{S}, \\ \tilde{\nu}_{S^*} &= \sum_a \tilde{t}_a^{\nu a^*} \in \Lambda^1 \tilde{S}^*. \end{aligned} \right. \tag{3.32}$$

As is known (see Libermann [11]), the components of the Ricci tensor are given by  $\tilde{\Theta}_a^a = \tilde{R}_{bc^*}^{\nu b} \wedge \omega^{c^*} + \tilde{\Theta}_a^a + \tilde{\Theta}_{a^*}^{a^*} = 0$ . Because of this, we get from (3.31) that

$$\left\{ \begin{aligned} \tilde{R}_{bc^*}^{\nu b} &= \tilde{t}_b^{\nu} \tilde{t}_c^{\nu} \\ \tilde{R}_{aa^*}^{\nu a} &= 2\tilde{t}_a^{\nu} \tilde{t}_{a^*}^{\nu} - 1. \end{aligned} \right. \tag{3.33}$$

It follows from (3.33) that the components of the Ricci tensor are disjoint (see Rosca [16]). In addition, since the scalar curvature  $\tilde{C}_S$  is the trace of the Ricci tensor with respect to  $\tilde{g}$ , one finds by (2.7) and (3.3) that  $\tilde{C}_S = 4-m$  ( $m > 4$ ). Therefore we conclude that the pseudo-Sasakian manifold  $\tilde{M}$  under consideration is a space form  $\tilde{M}(4-m)$  of hyperbolic type.

**THEOREM 1.** Let  $\tilde{M}(U, \xi, \tilde{\eta}, \tilde{g})$  be a pseudo-Sasakian manifold endowed with a c.c. connection and let  $\tilde{U}$  (resp.  $\tilde{\eta} = d\tilde{\eta}/2$ ) be the principal vector field associated with this connection (resp. the fundamental 2-form on  $\tilde{M}$ ). One has the following properties:

- (i)  $\tilde{U}$  is divergence free, and  $U\tilde{U}$  defines an infinitesimal homothety on  $\tilde{M}$ ;
  - (ii) all the connection forms on  $\tilde{M}$  are integral relations of invariance for  $U\tilde{U}$ ;
  - (iii)  $\tilde{U}$  and  $U\tilde{U}$  define an U-contact concircular pairing, and  $\{\tilde{U}, U\tilde{U}, \xi\}$  defines a 3-foliation on  $\tilde{M}$ ;
  - (iv) any contact extension  $\tilde{U}_f = \tilde{U} + f\xi$  of  $\tilde{U}$  is a relative contact infinitesimal transformation of  $\tilde{\eta}$ ;
  - (v)  $\tilde{U}$  and  $U\tilde{U}$  define both an infinitesimal automorphism of all  $(2q+1)$ -forms  $\tilde{\beta}_q = L^q \tilde{u}$  where  $\tilde{u}$  is the dual form of  $U\tilde{U}(q < m)$ ;
  - (vi) the Ricci 1-form of  $\tilde{M}$  is exterior recurrent, and the Ricci tensor is disjoint;
  - (vii)  $\tilde{M}$  is a space-form of hyperbolic type;
  - (viii) any such submanifold  $\tilde{M}$  is defined by a completely integrable system of differential equations whose solution depends on  $2m$  arbitrary constants.
4. CO-ISOTROPIC FOLIATION ON  $\tilde{M}(U, \xi, \tilde{\eta}, \tilde{g})$ .

We shall consider on  $M$  the following three distributions:

- a) An invariant distribution  $D^r$  (i.e.  $UD^r \subseteq D^r$ ) of dimension  $2(m-l)+1$  defined by  $D^r = \{h_1, h_{1^*}, \xi; i=1, \dots, m-l; i^*=i+m\}$ .
- b) An isotropic distribution  $D^l$  (i.e.  $D^l \subseteq \text{orth } D^l$ ) of dimension  $l$  defined by  $D^l = \{h_r; r=m-l+1, \dots, m\}$ .

c) A transversal distribution  $D_t = \bigcup_{S^*} (D^T \oplus D^\perp) \cap S^*$  of dimension  $\ell$  defined by  $D_t = \{h_{r^*}; r^* = 2m - \ell + 1, \dots, 2m\}$ .

These three distributions have no common direction and they define on  $\tilde{M}$  a  $f$ -structure of rank  $2\ell$  (see Sinha [17]).

Accordingly we shall split the principal vector field  $\hat{U}$  as follows:

$$\hat{U} = \hat{U}^T \oplus \hat{U}^\perp \oplus \hat{U}_t \tag{4.1}$$

where  $\hat{U}^T \in D^T$ ,  $\hat{U}^\perp \in D^\perp$ ,  $\hat{U}_t \in D_t$ .

Denote now by

$$\tilde{\psi} = \omega^{2m-\ell+1} \wedge \dots \wedge \omega^{2m} \tag{4.2}$$

the simple unit form which corresponds to  $D_t$ . Because  $D_t$  is orientable,  $\tilde{\psi}$  is a well-defined global form. Since  $\tilde{\psi}$  annihilates  $D^T \oplus D^\perp$ , the necessary and sufficient condition for  $D^T \oplus D^\perp$  to be a *co-isotropic foliation*  $F_c$  is that  $\tilde{\psi}$  be exterior recurrent (see Lichnerowicz [18] and Yano and Kon [7]).

Hence one must write  $d\tilde{\psi} = \tilde{\gamma} \wedge \tilde{\psi}$  and if  $H^1(F_c, \mathbb{R})$  represent the 1-cohomology class of  $F_c$ , then the recurrence 1-form  $\tilde{\gamma}$  defines an element of  $H^1(F_c, \mathbb{R})$  (see Lichnerowicz [6]). In the case under discussion one finds (compare with Yano and Kon [7]) that the necessary and sufficient condition for  $\tilde{M}$  to receive a co-isotropic foliation  $F_c = D^T \oplus D^\perp$  is that the component  $\hat{U}_t$  of  $\hat{U}$  vanishes. In this case the recurrence 1-form  $\tilde{\gamma}$  of  $\tilde{\psi}$  is given by

$$\tilde{\gamma} = \ell(\tilde{u} - \tilde{\eta}). \tag{4.3}$$

Denote by  $M_c$  a  $(2m - \ell + 1)$ -dimensional leaf of  $F_c$  and suppress  $\sim$  for the induced elements on  $M_c$ .

According to the considerations of Section 1, it follows that  $M_c$  is a *CICR submanifold*. By definition we have  $du = 0$ . Because of this and (3.1), the exterior differentiation of (4.3) gives

$$d\tilde{\gamma} = -2\ell\Omega. \tag{4.4}$$

Equation (4.4) shows that the restriction  $\Omega = \tilde{\Omega}|_{M_c}$  is an *exact form*.

On the other hand, the form of Godbillon-Vey (see Lichnerowicz [6]) on  $M_c$  is the  $(2\ell + 1)$ -form  $w_G \in \Lambda^{2\ell+1}(M_c)$  given by

$$w_G = \gamma \wedge (Ad\gamma)^\ell. \tag{4.5}$$

One knows (see Lichnerowicz [18]) that the class of cohomology of  $w_G$  which is an element of  $H^{2\ell+1}(M_c; \mathbb{R})$  is an invariant of the foliation. Using the same notation as in section 3 and applying (4.4), we may write

$$w_G = c(L^\ell u - L^\ell \eta) = c(\beta_\ell - L^\ell \eta) \tag{4.6}$$

where we have set  $c = -2^\ell \ell^{\ell+1}$ .

Thus it follows from (3.22) that

$$\mathfrak{L}_U w_G = -c \mathfrak{L}_U (L^\ell \eta). \tag{4.7}$$

By means of (2.13) and (3.9) one has

$$d(L^\ell \eta) = 2(\Lambda\Omega)^{\ell+1} \tag{4.8}$$

and

$$di_U(L^\ell \eta) = 4\ell u(\Lambda\Omega)^\ell . \tag{4.9}$$

Therefore we get

$$\mathcal{L}_U(L^\ell \eta) = -u \wedge (\Lambda\Omega)^\ell = -\beta_\ell \tag{4.10}$$

and finally

$$\mathcal{L}_{U^w_G} = c\beta_\ell . \tag{4.11}$$

Since  $\beta_\ell$  is closed, the above equation gives  $d\mathcal{L}_{U^w_G} = 0$  and allows us to say that  $w_G$  is a *relative integral invariant* of  $U$ .

Further since the submanifold  $M_C$  is co-isotropic, it follows from this that the normal bundle  $T^\perp M_C$  of  $M_C$  coincides with  $D^\perp$ .

Since  $M_C$  is defined by  $\omega^{r^k} = 0$ ,  $r^i = 2m-\ell+1, \dots, 2m$ , we derive from (2.8) and (3.12) that the covariant derivatives  $\nabla h_r$  of the null normal sections  $h_r$  satisfy

$$\nabla h_r = \frac{v}{2} \otimes h_r . \tag{4.12}$$

Since  $h_r$  are null vector fields, equation (4.12) shows that  $h_r$  are *geodesic* directions. Hence according to the definition of Rosca [19], one may say that  $D^\perp$  has the *geodesic property*.

Further if  $X$  and  $Y$  are any vector fields of  $D^\perp$ , one has  $\nabla_Y X \in D^\perp$ . Thus according to a known definition, the distribution  $D^\perp$  is *autoparallel*.

Setting  $\ell_r = -\langle dp, \nabla h_r \rangle$  for the *second fundamental quadratic forms* associated with the improper immersion  $x: M_C \rightarrow \tilde{M}$  ( $\ell_r$  is a field of symmetric covariant tensors of order 2 on  $M_C$ ), we derive by a simple argument that all  $\ell_r$  vanish. Therefore according to a well-known definition, we agree to say that the improper immersion  $x: M_C \rightarrow \tilde{M}$  is *improper totally geodesic*.

It was proved by Goldberg and Rosca [2] that the distribution  $D^\perp$  is always involutive. If  $M^\perp$  are the leaves of  $D^\perp$ , then in a similar manner as for  $M_C$  one easily finds that the improper immersion  $x: M^\perp \rightarrow \tilde{M}$  is *improper totally geodesic*. Since  $x: M^\perp \rightarrow \tilde{M}$  is a proper immersion, it is *totally geodesic*.

Next as it was proved (Goldberg and Rosca [2]) the necessary and sufficient condition for the manifold  $M_C$  to be *foliate* is that the simple unit form  $\psi$  which corresponds to  $D^\perp$  be exterior recurrent.

Since obviously one has  $\phi = \omega^{m-\ell+1} \wedge \dots \wedge \omega^m$ , then by (3.1) one finds that the property of exterior recurrency for  $\phi$  is equivalent to the condition  $U^\perp = 0$ .

Since by definition in this case  $D^\perp$  is involutive, let us denote by  $M^\perp$  a  $(2(m-\ell)+1)$ -dimensional leaf of  $D^\perp$ . Because  $M_C$  is a CICR submanifold,  $M^\perp$  is as is known an invariant submanifold of  $\tilde{M}$ , and this implies (see Rosca [1]) that  $M^\perp$  is *minimal*.

Coming back to the case under discussion, using (2.8), (3.12) and the fact that on  $M$  one has  $U_t = 0$ ,  $U^\perp = 0$ , we can show by means of a simple calculation that  $M^\perp$  is also totally geodesic.

Hence  $M_C$  is foliated by two families of orthogonal totally geodesic submanifolds  $M^\perp$  and  $M^\perp$ .

On the other hand, let  $X \in \mathfrak{X}_{M_c}$  be any vector field on  $M_c$ . According to Rosca [1], one has  $UX = PX + FX$  where  $PX$  (resp.  $FX$ ) is the tangential (resp. the normal) component of  $UX$ . By virtue of the total geodesicity of  $M^T$ , one easily finds that  $\nabla PX \in M^T$ .

Therefore the tangential component  $PX$  of  $X$  is *parallel*. According to Yano and Kon [7], it follows from this that  $M_c$  is a CR product i.e.  $M_c = M^\perp \times M^T$ .

Since  $M_c$  is connected, this property can be checked by *de Rham decomposition theorem*.

It is worth to note that this situation is quite similar to that of co-isotropic CR submanifolds of a para Kaehlerian manifold structured by a *geodesic connection* (Rosca [20]).

**THEOREM 2.** Let  $M$  be a pseudo-Sasakian manifold structured by a c.c. connection and let  $\hat{U}$  be the principal vector field associated with this connection. Then the necessary and sufficient condition for  $\hat{M}$  to receive a co-isotropic foliation  $F_c$  is that the transversal component  $\hat{U}_c$  of  $\hat{U}$  vanishes. In this case the leaves  $M_c$  of  $F_c$  are CICR submanifolds of  $M_c$ , and if  $\text{codim } M_c = \ell$ , the form of Godbillon-Vey on  $M_c$  is a  $(2\ell+1)$ -form  $w_G^c$  which is a relative integral invariant of  $U = \hat{U}|_{M_c}$ .

In addition, one has the following properties:

- (i) the improper immersion  $x: M_c \rightarrow \hat{M}$  is improper totally geodesic;
- (ii)  $M_c$  is foliated by anti-invariant submanifolds  $M^\perp$  which are improper totally geodesic and have the geodesic property.

Further the necessary and sufficient condition for  $M_c$  to be foliate is that the vertical (or isotropic) component  $U^\perp$  of  $U = \hat{U}|_{M_c}$  vanishes. In this case  $M_c$  is a CR product.

5. TANGENT BUNDLE MANIFOLD  $\hat{TM}$ .

Let  $\hat{TM}$  be the *tangent bundle manifold* having the pseudo-Sasakian manifold discussed in Section 3 as a basis.

Denote by  $\hat{\nu}_L^{\nu\Lambda}$  ( $\hat{\nu}^\Lambda$ ) the *canonical vector field* (or *the vector field of Liouville*) on  $\hat{TM}$ . Accordingly we may consider the set  $B^* = \{\omega^\Lambda, d\nu^\Lambda\}$  as an adapted cobasis on  $\hat{TM}$ . Following Godbillon [21], we shall designate by  $d_\nu$  and  $i_\nu$  the *vertical differentiation* and the *vertical derivation operators*, respectively taken with respect to  $B^*$  ( $d_\nu$  is an *antiderivation* of degree 1 of  $\Lambda\hat{TM}$  and  $i_\nu$  is a *derivative* of degree 0 of  $\Lambda\hat{TM}$ ).

Let  $T_s^{r\hat{M}}$  be the set of all tensor fields of type  $(r,s)$  on  $\hat{M}$ . In general the *vertical* and *complete* lifts are linear mappings of  $T_s^r \hat{M}$  into  $T_s^{r\hat{M}}$ , and for complete lifts one has:

$$(T_1^r \otimes T_2^s)^C = T_1^{rV} \otimes T_2^{rC} + T_1^{rC} \otimes T_2^{rV}.$$

With respect to  $B^*$  the complete lift of the fundamental form  $\hat{\Omega} = d\hat{\eta}/2$  is given by

$$\hat{\Omega}^C = \sum_a (d\nu^a \wedge \omega^a + \omega^a \wedge d\nu^a). \tag{5.1}$$

The exterior differentiation of (5.1) by means of (3.1) gives

$$d\tilde{\Omega}^C = \tilde{u} \wedge \tilde{\Omega}^C + \sum_a (\tilde{t}_a^v d\tilde{v}^{a*} - \tilde{t}_a^* d\tilde{v}^a) \wedge \tilde{\Omega}^C + \tilde{\eta} \wedge (\tilde{\omega}^a \wedge d\tilde{\omega}^{a*} + \tilde{\omega}^{a*} \wedge d\tilde{v}^a) . \tag{5.2}$$

Using (5.2), we find

$$\mathcal{L}_{\tilde{V}_L} \tilde{\Omega}^C = \tilde{\Omega}^C . \tag{5.3}$$

As is known (see Godbillon [21]), equation (5.3) shows that  $\tilde{\Omega}^C$  is homogeneous of degree 1.

We will now take the complete lift  $\tilde{u}^C$  of the principal Pfaffian  $\tilde{u}$  associated with the c.c. connection with structures  $\tilde{M}$ . For this purpose we shall denote by  $\partial_B(\tilde{t}_A^v) = h_B(\tilde{t}_A^*)$  the Pfaffian derivatives of  $\tilde{t}_A^*$  ( $A=0,1,\dots,2m$ ) with respect to cobasis  $\tilde{W}^*$ . Then according to the general theory (Yano and Ishihara [7]) one has

$$\tilde{u}^C = \tilde{u}_A d\tilde{v}^A + \partial_B(\tilde{u}_A) \tilde{v}^B \tilde{\omega}^A \tag{5.4}$$

where we have set  $\tilde{u} = \tilde{u}_A \tilde{\omega}^A$ . Referring to (3.4) and (3.5) ( $c=2$ ), after some calculations one finds

$$\tilde{u}^C = \frac{1}{2} \sum (\tilde{t}_a^* \tilde{v}^a - \tilde{t}_a^v d\tilde{v}^{a*}) + \frac{1}{2} \sum (\tilde{t}_a^* \tilde{v}^a - \tilde{t}_a^v \tilde{v}^{a*}) \tilde{u} + \sum_a (\tilde{v}^a \tilde{\omega}^a + \tilde{v}^{a*} \tilde{\omega}^{a*}) - \frac{1}{2} \tilde{v}^0 \tilde{v} . \tag{5.5}$$

The exterior differentiation of (5.5) by means of (3.1) gives

$$d\tilde{u}^C = \frac{1}{2} \sum (\tilde{t}_a^* \tilde{v}^{a*} + \tilde{t}_a^v \tilde{v}^a) \tilde{u} + \sum (\tilde{v}^a \tilde{\omega}^{a*} - \tilde{v}^{a*} \tilde{\omega}^a) + \sum_a (\tilde{t}_a^v d\tilde{v}^a + \tilde{t}_a^* d\tilde{v}^{a*}) \wedge \tilde{\eta} . \tag{5.6}$$

Using (5.5) and (5.6), one finds  $\mathcal{L}_{\tilde{V}_L} \tilde{u}^C = \tilde{u}^C$ . Hence  $\tilde{u}^C$  is also a homogeneous form of degree 1.

Consider now the following scalar field on  $\tilde{TM}$ :

$$\tilde{\Upsilon} = \sum \tilde{v}^a \tilde{v}^{a*} + (\tilde{v}^0)^2 / 2 \tag{5.7}$$

and apply the vertical differentiation of  $\tilde{\Upsilon}$ . According to Godbillon [21], one has

$$\tilde{v} = d_{\tilde{V}} \tilde{\Upsilon} = \sum_a (\tilde{v}^a \tilde{\omega}^{a*} + \tilde{v}^{a*} \tilde{\omega}^a) + \tilde{v}^0 \tilde{\eta} \tag{5.8}$$

and by means of (3.1) one gets

$$\tilde{\eta} = d\tilde{v} = \tilde{v}^0 \tilde{\eta} + \tilde{u} \wedge \sum (\tilde{v}^a \tilde{\omega}^{a*} + \tilde{v}^{a*} \tilde{\omega}^a) + \tilde{\eta} \wedge (\sum (\tilde{v}^a \tilde{\omega}^a - \tilde{v}^{a*} \tilde{\omega}^{a*}) - d\tilde{v}^0) + 2\tilde{v}^0 \tilde{\eta} \tag{5.9}$$

In (5.9)  $\iota : \Lambda^1 \tilde{M} \rightarrow \infty \tilde{TM}$  is the operator of Yano and Ishihara [7], that is with respect to  $B^*$  one has by (3.6)

$$\iota \tilde{v} = \sum_a (\tilde{t}_a^* \tilde{v}^a + \tilde{t}_a^v \tilde{v}^{a*}) . \tag{5.10}$$

One quickly finds

$$\iota \tilde{v}_L \tilde{\eta} = \tilde{v} , \tag{5.11}$$

and since  $\tilde{\eta}$  is closed, it follows from (5.11) that

$$\mathcal{L}_{\tilde{V}_L} \tilde{\eta} = \tilde{\eta} \tag{5.12}$$

i.e.  $\tilde{\eta}$  is homogeneous of degree 1. Moreover, taking the vertical derivation of  $\tilde{\eta}$ , one has (see Godbillon [21]):

$$i_{\tilde{V}}\tilde{\eta} = 0 . \tag{5.13}$$

On the other hand, it is easy to see from (5.9) that  $\tilde{\eta}$  is of maximal rank (see Godbillon [21]) on  $\tilde{M}$ . Accordingly, as is known, equations (5.11) and (5.13) prove that  $\tilde{\eta}$  is a *Finslerian form* (See Klein and Voutier [22]). Since the vertical differentiation  $d_{\tilde{V}}$  is an anti-derivation of square zero, one easily derives from (5.8) that

$$d_{\tilde{V}}\tilde{\nu} = 0, \quad i_{\tilde{V}_L}\tilde{\nu} = 0 . \tag{5.14}$$

Thus according to Godbillon [21],  $\tilde{\nu}$  is a *semibasic form*.

In the following we shall call  $\tilde{T}$  (resp.  $\tilde{\nu}$ ) the *Liouville function* (resp. *the Liouville 1-form*) on  $\tilde{M}$  (see Rosca [16]). Further one may call  $\tilde{\eta}$  the *2-form of Cartan* on  $\tilde{M}$  (see Rosca [19]).

Denote now by  $B = \{h_{\Lambda}, \frac{\partial}{\partial v^{\Lambda}}\}$  the vectorial basis dual to  $B^*$  on  $\tilde{M}$ . Then as is known (see Yano and Ishihara [9] or Godbillon [21]) the vertical lift  $(\tilde{Z})^V$  of  $\tilde{Z}$  is expressed by

$$(\tilde{Z})^V = z^{\Lambda} \frac{\partial}{\partial v^{\Lambda}} . \tag{5.15}$$

Coming back to the case under consideration and using that  $U\tilde{U} = b^{-1}(\tilde{u})$  (see Section 3), we find by (5.15) that

$$(U\tilde{U})^V = \sum_a (\tilde{\xi}_a \frac{\partial}{\partial v^a} - \tilde{\xi}_{a^*} \frac{\partial}{\partial v^{a^*}}) . \tag{5.16}$$

Now, taking the dual  $\mu(U\tilde{U})^V$  of  $(U\tilde{U})^V$  with respect to  $\tilde{\eta}$  and referring to (3.5) (c=2), we quickly find

$$\mu(U\tilde{U})^V = 2\tilde{u} . \tag{5.17}$$

Since  $\tilde{u}$  and  $\tilde{\eta}$  are both closed, it follows from this that  $\mathcal{L}_{(U\tilde{U})^V}\tilde{\eta} = 0$ , i.e.  $(U\tilde{U})^V$  is an infinitesimal automorphism of  $\tilde{\eta}$ .

Consider now on  $\tilde{M}$  the *mechanical system*  $\mathcal{M} = (\tilde{M}, \tilde{T}, \tilde{\pi})$  where  $\tilde{T}$  and

$$\tilde{\pi} = 2\tilde{T}\tilde{u} \tag{5.18}$$

are the *kinetic energy* and the *field of forces* of  $\mathcal{M}$  (see Godbillon [21]).

Since  $\tilde{u}$  is closed, one has  $d\tilde{\pi} = \frac{d\tilde{T}}{v} \wedge \tilde{\eta}$  and referring to (5.7), one quickly finds

$$\begin{aligned} \tilde{V}_L \tilde{T} &= 2\tilde{T} , \\ \tilde{V}_L \tilde{\pi} &= 2\tilde{\pi} . \end{aligned} \tag{5.19}$$

Equations (5.19) show that  $\tilde{T}$  and  $\tilde{\pi}$  are *homogeneous of degree 2*. On the other hand, since  $\tilde{\pi}$  is an exact 2-form of maximal rank, it defines a *potential symplectic structure* on  $\tilde{M}$ . Hence, according to the definition given by Klein (see Godbillon [21]) the system  $\mathcal{M}$  is *regular*.

Denote now by  $\tilde{Z}_d$  the *dynamical system* associated with  $\mathcal{M}$ . As is known,  $\tilde{Z}_d$

is defined via formula

$$i_{\tilde{Z}_d} \tilde{\eta} = d(\tilde{\gamma} - \tilde{\gamma}_L \tilde{\gamma}) + \tilde{\pi} . \quad (5.20)$$

Then:

- a) Since  $\tilde{\gamma}$  and  $\tilde{\pi}$  are both homogeneous and of the same degree,  $\tilde{Z}_d$  is a spray on  $\tilde{M}$ , i.e.  $[\tilde{V}_L, \tilde{Z}_d] = \tilde{Z}_d$ .
- b) Since  $\tilde{\gamma}$  is of degree 2, the 2-form  $\tilde{\eta} - (d\tilde{\gamma} - \tilde{\pi}) \wedge dt \in \Lambda^2(T\tilde{M} \times \mathbb{R})$  is an integral relation of invariance for  $\tilde{Z}_d + \frac{\partial}{\partial t}$  (Lichnerowicz [5]).

THEOREM 3. Let  $T\tilde{M}$  be the tangent bundle manifold having as a basis the manifold  $\tilde{M}(U, \xi, \tilde{\eta}, \tilde{g})$  defined in Section 3 and let  $\tilde{U}$  (resp.  $\tilde{\eta}$ ) be the principal vector field (resp. the fundamental 2-form) on  $\tilde{M}$ . Then:

- (i) the complete lifts  $\tilde{\eta}^C$  and  $\tilde{u}^C$  of  $\tilde{\eta}$  and  $\tilde{u} = \mathbf{b}(U\tilde{U})$  are homogeneous of degree one;
- (ii) the 2-form of Cartan  $\tilde{\eta}$  on  $T\tilde{M}$  is a Finslerian form;
- (iii) one may associate with  $\tilde{\eta}$  a regular mechanical system whose dynamical system is a spray on  $\tilde{M}$ .

#### REFERENCES

1. ROSCA, R. On Pseudo-Sasakian Manifolds, Rend. Mat. 1984 (to appear).
2. GOLDBERG, V.V. and ROSCA, R. Contact Co-isotropic CR Submanifolds of a Pseudo-Sasakian Manifold, Intern. J. Math. Math. Sci. 7(1984), No. 2, 339-350.
3. ROSCA, R. Variétés Sasakienne à Connexion Conforme de Contact, C.R. Acad. Sci. Paris Sér. I Math. 294(1982), 43-46.
4. LICHNEROWICZ, A. Les Relations Integrales d'Invariance et Leurs Applications à la Dynamique, Bull. Sci. Math. 70(1946), 82-95.
5. ROSCA, R. Variétés Lorentziennes à Structure Sasakienne et Admettant un Champ Vectoriel Isotrope  $\psi$ -quasi Concirculaire, C.R. Acad. Sci. Paris Sér. A. 291(1980), 45-47.
6. LICHNEROWICZ, A. Variétés de Poisson et Feuilletages, Ann. Fac. Sci. Toulouse Math (5) 4(1982), 195-262.
7. YANO, K. and KON, M. CR Submanifolds of Kaehlerian and Sasakian Manifolds, Birkhäuser, Boston-Basel-Stuttgart, 1983.
8. ROSCA, R. CR-sous-variétés Co-isotropes d'une Variété Parakählerienne, C.R. Acad. Sci Paris Sér. I Math. 298(1984), 149-151.
9. YANO, K. and ISHIHARA, S. Differential Geometry of Tangent and Cotangent Bundles, Marcel Dekker Inc., New York, 1973.
10. KLEIN, I. Espaces Variationels et Mécanique, Ann. Inst. Fourier (Grenoble) 12(1962), 1-124.
11. LIBERMANN, P. Sur le Problème d'Équivalence de Certaines Structures Infinitésimales, Ann. Mat. Pura Appl. 36(1951), 27-120.
12. ROSCA, R. Codimension 2 CR Submanifolds with Null Covariant Decomposable Vertical Distribution of a Neutral Manifold, Rend. Mat. (4) 2(1982), 787-796.
13. DATTA, D.K. Exterior Recurrent Forms on a Manifold, Tensor (N.S.) 36(1982), No. 1, 115-120.
14. POOR, W.A. Differential Geometric Structures, McGraw-Hill Book Comp., New York, 1981.
15. ROSCA, R. and VERSTRAELEN, L. On Submanifolds Admitting a Normal Section Which is Quasi-concircular w.r.t. a Corresponding Principal Tangent Section, Bull. Math. Soc. Math. R.S. Roumanie 20(68)(1976), No. 3-4, 399-402.

16. ROSCA, R. On Parallel Conformal Connections, Kodai Math. J. 2(1979), No. 1, 1-10.
17. SINHA, B.B. A Differentiable Manifold with Para  $f$ -Structure of Rank  $r$ , Ann. Fac. Sci. Univ. Nat. Zaire (Kinshasa) Sect. Math.-Phys. 6(1980), No. 1-2, 79-94.
18. LICHNEROWICZ, A. Feuilletages, Géométrie Riemannienne et Géométrie Symplectique, C.R. Acad. Sci. Paris Ser. I Math. 296(1983), 205-210.
19. ROSCA, R. Espace-temps Ayant la Propriété Géodésique, C.R. Acad. Sci. Paris Sér. A 285(1977), 305-308.
20. ROSCA, R. Sous-variétés Anti-invariantes d'une Variété Parakählerienne Structurée par une Connexion Géodésique, C.R. Acad. Sci. Paris Sér. A 287(1978), 539-541.
21. GODBILLON, C. Géométrie Différentielle et Mécanique Analytique, Hermann, Paris, 1969.
22. KLEIN, I. and VOUTIER, A. Formes extérieures génératrices de sprays, Ann. Inst. Fourier (Grenoble) 18(1968), 241-268.