

*Research Article*

## Improving Results on Convergence of AOR Method

**Guangbin Wang, Ting Wang, and Yanli Du**

*Department of Mathematics, Qingdao University of Science and Technology, Qingdao 266061, China*

Correspondence should be addressed to Guangbin Wang, wguangbin750828@sina.com

Received 16 June 2012; Revised 27 August 2012; Accepted 4 October 2012

Academic Editor: Carlos J. S. Alves

Copyright © 2012 Guangbin Wang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We present some sufficient conditions on convergence of AOR method for solving  $Ax = b$  with  $A$  being a strictly doubly  $\alpha$  diagonally dominant matrix. Moreover, we give two numerical examples to show the advantage of the new results.

### 1. Introduction

Let us denote all complex square matrices by  $C^{n \times n}$  and all complex vectors by  $C^n$ .

For  $A = (a_{ij}) \in C^{n \times n}$ , we denote by  $\rho(A)$  the spectral radius of matrix  $A$ .

Let us consider linear system  $Ax = b$ , where  $b \in C^n$  is a given vector and  $x \in C^n$  is an unknown vector. Let  $A = D - T - S$  be given and  $D$  is the diagonal matrix,  $-T$  and  $-S$  are strictly lower and strictly upper triangular parts of  $A$ , respectively, and denote

$$L = D^{-1}T, \quad U = D^{-1}S, \quad (1.1)$$

where  $\det(D) \neq 0$ .

Then the AOR method [1] can be written as

$$x^{k+1} = M_{\sigma, \omega} x^k + d, \quad k = 0, 1, \dots, x^0 \in C^n, \quad (1.2)$$

where

$$\begin{aligned} M_{\sigma, \omega} &= (I - \sigma L)^{-1}[(1 - \omega)I + (\omega - \sigma)L + \omega U], \\ d &= \omega(I - \sigma L)^{-1}b, \quad \omega, \sigma \in R. \end{aligned} \quad (1.3)$$

## 2. Preliminaries

We denote

$$R_i(A) = \sum_{j \neq i} |a_{ij}|, \quad S_i(A) = \sum_{j \neq i} |a_{ji}|, \quad P_{i,\alpha}(A) = \alpha R_i(A) + (1 - \alpha) S_i(A), \quad \forall i \in N. \quad (2.1)$$

For any matrix  $A = (a_{ij}) \in C^{n \times n}$ , the comparison matrix  $M(A) = (m_{ij}) \in R^{n \times n}$  is defined by

$$m_{ii} = |a_{ii}|, \quad m_{ij} = -|a_{ij}|, \quad i, j \in N, \quad i \neq j, \quad N = \{1, 2, \dots, n\}. \quad (2.2)$$

*Definition 2.1* (see [2]). A matrix  $A \in C^{n \times n}$  is called a strictly diagonally dominant matrix (SD) if

$$|a_{ii}| > R_i(A), \quad \forall i \in N. \quad (2.3)$$

A matrix  $A \in C^{n \times n}$  is called a strictly doubly diagonally dominant matrix (DD) if

$$|a_{ii}| |a_{jj}| > R_i(A) R_j(A), \quad \forall i, j \in N, \quad i \neq j. \quad (2.4)$$

*Definition 2.2* (see [3]). A matrix  $A \in C^{n \times n}$  is called a strictly  $\alpha$  diagonally dominant matrix ( $D(\alpha)$ ) if there exists  $\alpha \in [0, 1]$ , such that

$$|a_{ii}| > \alpha R_i(A) + (1 - \alpha) S_i(A), \quad \forall i \in N. \quad (2.5)$$

*Definition 2.3* (see [4]). Let  $A = (a_{ij}) \in C^{n \times n}$ , if there exists  $\alpha \in [0, 1]$  such that

$$|a_{ii}| |a_{jj}| > \alpha R_i(A) R_j(A) + (1 - \alpha) S_i(A) S_j(A), \quad \forall i, j \in N, \quad i \neq j. \quad (2.6)$$

then  $A$  is called a strictly doubly  $\alpha$  diagonally dominant matrix ( $DD(\alpha)$ ).

In [3, 5, 6], some people studied the convergence of AOR method for solving linear system  $Ax = b$  and gave the areas of convergence. In [5], Cvetković and Herceg studied the convergence of AOR method for strictly diagonally dominant matrices. In [3], Huang and Wang studied the convergence of AOR method for strictly  $\alpha$  diagonally dominant matrices. In [6], Gao and Huang studied the convergence of AOR method for strictly doubly diagonally dominant matrices.

**Theorem 2.4** (see [3]). Let  $A \in D(\alpha)$ , then AOR method converges for

$$\begin{aligned}
 \text{(I)} \quad & 0 \leq \sigma < \frac{2}{(1 + \rho(M_{0,1}(M(A))))} = s, \\
 & 0 < \omega < \max \left\{ \frac{2}{(1 + \max_i P_{i,\alpha}(L+U))} = t, \frac{2\sigma}{(1 + \rho(M_{\sigma,\omega}))} \right\}, \quad \text{or} \\
 \text{(II)} \quad & \max_i \frac{(-\omega(1 - P_{i,\alpha}(L+U)) + 2 \max(0, \omega - 1))}{2P_{i,\alpha}(L)} < \sigma < 0, \quad 0 < \omega < t, \quad \text{or} \\
 \text{(III)} \quad & t \leq \sigma < \min_i \frac{(\omega(1 + P_{i,\alpha}(L) - P_{i,\alpha}(U)) + 2 \min(0, 1 - \omega))}{2P_{i,\alpha}(L)}, \quad 0 < \omega < t.
 \end{aligned} \tag{2.7}$$

**Theorem 2.5** (see [6]). Let  $A \in DD$ , then AOR method converges for

$$\begin{aligned}
 \text{(I)} \quad & 0 \leq \sigma < \frac{2}{(1 + \rho(M_{0,1}(M(A))))} = s, \\
 & 0 < \omega < \max \left\{ \frac{2\sigma}{(1 + \rho(M_{\sigma,\omega}))}, \min_{\substack{i,j \\ i \neq j}} \frac{2}{1 + \sqrt{R_i(L+U)R_j(L+U)}} = t \right\}, \quad \text{or} \\
 \text{(II)} \quad & \max_{\substack{i,j \\ i \neq j}} \frac{\omega P_1 - \sqrt{\omega^2 P_2^2 + P_3 \min(\omega^2, (\omega-2)^2)}}{P_3} < \sigma < 0, \quad 0 < \omega < t, \quad \text{or} \\
 \text{(III)} \quad & t \leq \sigma < \min_{\substack{i,j \\ i \neq j}} \frac{\omega P_4 + \sqrt{\omega^2 P_5^2 + P_3 \min(\omega^2, (\omega-2)^2)}}{P_3}, \quad 0 < \omega < t,
 \end{aligned} \tag{2.8}$$

where

$$\begin{aligned}
 P_1 &= R_i(L)R_j(L+U) + R_i(L+U)R_j(L), \\
 P_2 &= R_i(L)R_j(L+U) - R_i(L+U)R_j(L), \\
 P_3 &= 4R_i(L)R_j(L), \\
 P_4 &= R_i(L)(R_j(L) - R_j(U)) + R_j(L)(R_i(L) - R_i(U)), \\
 P_5 &= R_i(L)(R_j(L) - R_j(U)) - R_j(L)(R_i(L) - R_i(U)).
 \end{aligned} \tag{2.9}$$

### 3. Upper Bound for Spectral Radius of $M_{\sigma,\omega}$

In the following, we present an upper bound for spectral radius of AOR iterative matrix  $M_{\sigma,\omega}$  for strictly doubly  $\alpha$  diagonally dominant coefficient matrix.

**Lemma 3.1** (see [4]). If  $A \in DD(\alpha)$ , then  $A$  is a nonsingular H-matrix.

**Theorem 3.2.** Let  $A \in DD(\alpha)$ , if  $1 - \sigma^2[\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] > 0$ , for all  $j \in N$ ,  $i \neq j$ , then

$$\rho(M_{\sigma,\omega}) \leq \max_{\substack{i,j \\ i \neq j}} \frac{A_2 + \sqrt{A_2^2 - 4A_1A_3}}{2A_1}, \quad (3.1)$$

where

$$\begin{aligned} A_1 &= 1 - \sigma^2 [\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)], \\ A_2 &= 2|1 - \omega| + 2|\omega - \sigma||\sigma| [\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] \\ &\quad + |\omega||\sigma| [\alpha R_i(L)R_j(U) + (1 - \alpha)S_i(U)S_j(L)] \\ &\quad + |\omega||\sigma| [\alpha R_i(U)R_j(L) + (1 - \alpha)S_i(L)S_j(U)], \\ A_3 &= (1 - \omega)^2 - \alpha[|\omega - \sigma|R_i(L) + |\omega|R_i(U)][|\omega - \sigma|R_j(L) + |\omega|R_j(U)] \\ &\quad - (1 - \alpha)[|\omega - \sigma|S_i(L) + |\omega|S_i(U)][|\omega - \sigma|S_j(L) + |\omega|S_j(U)]. \end{aligned} \quad (3.2)$$

*Proof.* Let  $\lambda$  be an eigenvalue of  $M_{\sigma,\omega}$  such that

$$\det(\lambda I - M_{\sigma,\omega}) = 0, \quad (3.3)$$

that is,

$$\det(\lambda(I - \sigma L) - ((1 - \omega)I + (\omega - \sigma)L + \omega U)) = 0. \quad (3.4)$$

If  $\lambda(I - \sigma L) - ((1 - \omega)I + (\omega - \sigma)L + \omega U) \in DD(\alpha)$ , then by Lemma 3.1,  $\lambda(I - \sigma L) - ((1 - \omega)I + (\omega - \sigma)L + \omega U)$  is nonsingular and  $\lambda$  is not an eigenvalue of iterative matrix  $M_{\sigma,\omega}$ , that is, if

$$\begin{aligned} |\lambda - (1 - \omega)|^2 &> \alpha \left[ \sum_{t \neq i} |\lambda \sigma l_{it} - (\omega - \sigma)l_{it} - \omega u_{it}| \right] \left[ \sum_{l \neq j} |\lambda \sigma l_{jl} - (\omega - \sigma)l_{jl} - \omega u_{jl}| \right] \\ &\quad + (1 - \alpha) \left[ \sum_{k \neq i} |\lambda \sigma l_{ki} - (\omega - \sigma)l_{ki} - \omega u_{ki}| \right] \\ &\quad \times \left[ \sum_{s \neq j} |\lambda \sigma l_{sj} - (\omega - \sigma)l_{sj} - \omega u_{sj}| \right], \quad \forall i, j \in N, i \neq j, \end{aligned} \quad (3.5)$$

then  $\lambda$  is not an eigenvalue of  $M_{\sigma,\omega}$ . Especially, if

$$\begin{aligned}
 (|\lambda| - |1 - \omega|)^2 &> \alpha \left[ \sum_{t \neq i} (|\lambda||\sigma||l_{it}| + |\omega - \sigma||l_{it}| + |\omega||u_{it}|) \right] \\
 &\quad \times \left[ \sum_{l \neq j} (|\lambda||\sigma||l_{jl}| + |\omega - \sigma||l_{jl}| + |\omega||u_{jl}|) \right] \\
 &\quad + (1 - \alpha) \left[ \sum_{k \neq i} (|\lambda||\sigma||l_{ki}| + |\omega - \sigma||l_{ki}| + |\omega||u_{ki}|) \right] \\
 &\quad \times \left[ \sum_{s \neq j} (|\lambda||\sigma||l_{sj}| + |\omega - \sigma||l_{sj}| + |\omega||u_{sj}|) \right], \tag{3.6}
 \end{aligned}$$

then  $\lambda$  is not an eigenvalue of  $M_{\sigma,\omega}$ .

If  $\lambda$  is an eigenvalue of  $M_{\sigma,\omega}$ , then there exists at least a couple of  $i, j$  ( $i \neq j$ ), such that

$$\begin{aligned}
 (|\lambda| - |1 - \omega|)^2 &\leq \alpha \left[ \sum_{t \neq i} (|\lambda||\sigma||l_{it}| + |\omega - \sigma||l_{it}| + |\omega||u_{it}|) \right] \\
 &\quad \times \left[ \sum_{l \neq j} (|\lambda||\sigma||l_{jl}| + |\omega - \sigma||l_{jl}| + |\omega||u_{jl}|) \right] \\
 &\quad + (1 - \alpha) \left[ \sum_{k \neq i} (|\lambda||\sigma||l_{ki}| + |\omega - \sigma||l_{ki}| + |\omega||u_{ki}|) \right] \\
 &\quad \times \left[ \sum_{s \neq j} (|\lambda||\sigma||l_{sj}| + |\omega - \sigma||l_{sj}| + |\omega||u_{sj}|) \right], \tag{3.7}
 \end{aligned}$$

that is,

$$A_1|\lambda|^2 - A_2|\lambda| + A_3 \leq 0. \tag{3.8}$$

Since  $A_1 = 1 - \sigma^2[\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] > 0$ , and the discriminant  $\Delta$  of the quadratic in  $|\lambda|$  satisfies  $\Delta \geq 0$ , then the solution of (3.8) satisfies

$$\frac{A_2 - \sqrt{A_2^2 - 4A_1A_3}}{2A_1} \leq |\lambda| \leq \frac{A_2 + \sqrt{A_2^2 - 4A_1A_3}}{2A_1}. \tag{3.9}$$

So

$$\rho(M_{\sigma,\omega}) \leq \max_{\substack{i,j \\ i \neq j}} \frac{A_2 + \sqrt{A_2^2 - 4A_1A_3}}{2A_1}. \quad (3.10)$$

□

#### 4. Improving Results on Convergence of AOR Method

In this section, we present new results on convergence of AOR method.

**Theorem 4.1.** *Let  $A \in DD(\alpha)$ , then AOR method converges if  $\omega, \sigma$  satisfy either*

$$(I) \quad 0 \leq \sigma < \frac{2}{(1 + \rho(M_{0,1}(M(A))))} = s, \\ 0 < \omega < \max \left\{ \frac{2\sigma}{(1 + \rho(M_{\sigma,\sigma}))}, \right. \\ \left. \min_{\substack{i,j \\ i \neq j}} \frac{2}{1 + \alpha R_i(L+U)R_j(L+U) + (1-\alpha)S_i(L+U)S_j(L+U)} = t \right\}, \quad \text{or} \\ (II) \quad \max_{\substack{i,j \\ i \neq j}} \frac{\omega P'_1 - \sqrt{\omega^2 P'^2_2 + P'_3 \min(\omega^2, (\omega-2)^2)}}{P'_3} < \sigma < 0, \quad 0 < \omega < t, \quad \text{or} \\ (III) \quad t \leq \sigma < \min_{\substack{i,j \\ i \neq j}} \frac{\omega P'_4 + \sqrt{\omega^2 P'^2_5 + P'_3 \min(\omega^2, (\omega-2)^2)}}{P'_3}, \quad 0 < \omega < t, \quad (4.1)$$

where

$$\begin{aligned} P'_1 &= \alpha R_i(L)R_j(L+U) + (1-\alpha)S_i(L)S_j(L+U) + \alpha R_i(L+U)R_j(L) \\ &\quad + (1-\alpha)S_i(L+U)S_j(L), \\ P'_2 &= \alpha R_i(L)R_j(L+U) + (1-\alpha)S_i(L)S_j(L+U) - \alpha R_i(L+U)R_j(L) \\ &\quad - (1-\alpha)S_i(L+U)S_j(L), \\ P'_3 &= 4[\alpha R_i(L)R_j(L) + (1-\alpha)S_i(L)S_j(L)], \\ P'_4 &= \alpha R_i(L)[R_j(L) - R_j(U)] + (1-\alpha)S_i(L)[S_j(L) - S_j(U)] \\ &\quad + \alpha R_j(L)[R_i(L) - R_i(U)] + (1-\alpha)S_j(L)[S_i(L) - S_i(U)], \\ P'_5 &= \alpha R_i(L)[R_j(L) - R_j(U)] + (1-\alpha)S_i(L)[S_j(L) - S_j(U)] \\ &\quad - \alpha R_j(L)[R_i(L) - R_i(U)] - (1-\alpha)S_j(L)[S_i(L) - S_i(U)]. \end{aligned} \quad (4.2)$$

*Proof.* It is easy to verify that for each  $\sigma$ , which satisfies one of the conditions (I)–(III), we have

$$1 - \sigma^2 [\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] > 0, \quad \forall i \neq j; \quad i, j \in N. \quad (4.3)$$

Firstly, we consider case (I). Since  $A$  be a  $\alpha$  diagonally dominant matrix, then by Lemma 3.1, we know that  $A$  is a nonsingular  $H$ -matrix; therefore,  $M(A)$  is a nonsingular  $M$ -matrix, and it follows that from paper [7],  $\rho(M_{\sigma,\sigma}) < 1$  holds for  $0 \leq \sigma < s$  and for  $\sigma \neq 0$ ,

$$M_{\sigma,\omega} = \left(1 - \frac{\omega}{\sigma}\right)I + \left(\frac{\omega}{\sigma}\right)M_{\sigma,\sigma}. \quad (4.4)$$

If  $0 < \omega < \max\{2\sigma/(1 + \rho(M_{\sigma,\sigma})), t\} = 2\sigma/(1 + \rho(M_{\sigma,\sigma}))$ , then

$$0 < \frac{\omega}{\sigma} < \frac{2}{(1 + \rho(M_{\sigma,\sigma}))}, \quad (4.5)$$

by extrapolation theorem [6], we have  $\rho(M_{\sigma,\omega}) < 1$ .

If  $0 < \omega < \max\{2\sigma/(1 + \rho(M_{\sigma,\sigma})), t\} = t$ , then it remains to analyze the case

$$\frac{2\sigma}{(1 + \rho(M_{\sigma,\sigma}))} \leq \omega < t, \quad 0 \leq \sigma < s. \quad (4.6)$$

Since when  $\rho(M_{\sigma,\sigma}) < 1$ ,

$$\sigma < \frac{2\sigma}{(1 + \rho(M_{\sigma,\sigma}))}, \quad (4.7)$$

then  $0 \leq \sigma < \omega$ . From

$$\begin{aligned} A_2 - A_3 &< A_1 & (a) \\ A_2 &< 2A_1 \leq 2 & (b) \iff \frac{A_2 + \sqrt{A_2^2 - 4A_1A_3}}{2A_1} < 1, \\ \sqrt{A_2^2 - 4A_1A_3} &< 2A_1 \leq 2 & (c) \end{aligned} \quad (4.8)$$

we have  $\rho(M_{\sigma,\omega}) \leq \max_{i,j} ((A_2 + \sqrt{A_2^2 - 4A_1A_3})/2A_1) < 1$ .

(1) When  $\omega \leq 1$ , it is easy to verify that (4.8) holds.

(2) When  $\omega > 1$ , since

$$\begin{aligned} A_1 &= 1 - \sigma^2 [\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)], \\ A_2 &= 2(\omega - \sigma) [\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] + \omega\sigma [\alpha R_i(L)R_j(U) + (1 - \alpha)S_i(U)S_j(L)] \\ &\quad + \omega\sigma [\alpha R_i(U)R_j(L) + (1 - \alpha)S_i(L)S_j(U)] + 2|1 - \omega|, \\ A_3 &= (1 - \omega)^2 - \alpha[(\omega - \sigma)R_i(L) + \omega R_i(U)][(\omega - \sigma)R_j(L) + \omega R_j(U)] \\ &\quad - (1 - \alpha)[(\omega - \sigma)S_i(L) + \omega S_i(U)][(\omega - \sigma)S_j(L) + \omega S_j(U)], \end{aligned} \tag{4.9}$$

then by  $A_2 - A_3 < A_1$  and  $1 - \sigma^2 [\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] > 0$ , for all  $i, j \in N$ ,  $i \neq j$ , we have

$$\omega^2 [1 - \alpha R_i(L+U)R_j(L+U) - (1 - \alpha)S_i(L+U)S_j(L+U)] - 4\omega + 4 > 0. \tag{4.10}$$

It is easy to verify that the discriminant  $\Delta$  of the quadratic in  $\omega$  satisfies  $\Delta > 0$ , and so there holds

$$\omega_1 > \frac{2}{1 - \alpha R_i(L+U)R_j(L+U) - (1 - \alpha)S_i(L+U)S_j(L+U)}, \tag{4.11}$$

or

$$\omega_2 < \frac{2}{1 + \alpha R_i(L+U)R_j(L+U) + (1 - \alpha)S_i(L+U)S_j(L+U)}, \quad \forall i \neq j. \tag{4.12}$$

For  $\omega_1$ , we have  $A_2 > 2$ , it is in contradiction with ((4.8)b). Therefore,  $\omega_1$  should be deleted. Secondly, we prove (II).

(1) When  $0 < \omega \leq 1$ ,  $\sigma < 0$ ,

$$\begin{aligned} A_2 &= 2(1 - \omega) - 2(\omega - \sigma)\sigma [\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] \\ &\quad - \omega\sigma [\alpha R_i(L)R_j(U) + (1 - \alpha)S_i(U)S_j(L)] \\ &\quad - \omega\sigma [\alpha R_i(U)R_j(L) + (1 - \alpha)S_i(L)S_j(U)], \end{aligned} \tag{4.13}$$

$$\begin{aligned} A_3 &= (1 - \omega)^2 - \alpha[(\omega - \sigma)R_i(L) + \omega R_i(U)][(\omega - \sigma)R_j(L) + \omega R_j(U)] \\ &\quad - (1 - \alpha)[(\omega - \sigma)S_i(L) + \omega S_i(U)][(\omega - \sigma)S_j(L) + \omega S_j(U)]. \end{aligned}$$

By  $A_2 - A_3 < A_1$ , we have

$$\begin{aligned} & 4\sigma^2 [\alpha R_i(L)R_j(L) + (1-\alpha)S_i(L)S_j(L)] \\ & - 2\omega\sigma [\alpha R_i(L)R_j(L+U) + (1-\alpha)S_i(L)S_j(L+U) \\ & + \alpha R_i(L+U)R_j(L) + (1-\alpha)S_i(L+U)S_j(L)] \\ & + \omega^2 [\alpha R_i(L+U)R_j(L+U) + (1-\alpha)S_i(L+U)S_j(L+U) - 1] < 0. \end{aligned} \quad (4.14)$$

It is easy to verify that the discriminant  $\Delta$  of the quadratic in  $\sigma$  satisfies  $\Delta > 0$ , and so there holds

$$\frac{\omega P'_1 - \omega \sqrt{P'_2 + P'_3}}{P'_3} < \sigma < \frac{\omega P'_1 + \omega \sqrt{P'_2 + P'_3}}{P'_3}. \quad (4.15)$$

By  $\sigma < 0$  and  $1 - \sigma^2 [\alpha R_i(L)R_j(L) + (1-\alpha)S_i(L)S_j(L)] > 0$ , for all  $i, j \in N, i \neq j$ , we obtain

$$\max_{\substack{i,j \\ i \neq j}} \frac{\omega P'_1 - \omega \sqrt{P'_2 + P'_3}}{P'_3} < \sigma < 0. \quad (4.16)$$

(2) When  $1 < \omega < t, \sigma < 0$ ,

$$\begin{aligned} A_2 &= 2(\omega - 1) - 2(\omega - \sigma)\sigma [\alpha R_i(L)R_j(L) + (1-\alpha)S_i(L)S_j(L)] \\ &\quad - \omega\sigma [\alpha R_i(L)R_j(U) + (1-\alpha)S_i(U)S_j(L)] - \omega\sigma [\alpha R_i(U)R_j(L) + (1-\alpha)S_i(L)S_j(U)], \\ A_3 &= (1-\omega)^2 - \alpha[(\omega - \sigma)R_i(L) + \omega R_i(U)][(\omega - \sigma)R_j(L) + \omega R_j(U)] \\ &\quad - (1-\alpha)[(\omega - \sigma)S_i(L) + \omega S_i(U)][(\omega - \sigma)S_j(L) + \omega S_j(U)]. \end{aligned} \quad (4.17)$$

By  $A_2 - A_3 < A_1$ , we have

$$\begin{aligned} & 4\sigma^2 [\alpha R_i(L)R_j(L) + (1-\alpha)S_i(L)S_j(L)] \\ & - 2\omega\sigma [\alpha R_i(L)R_j(L+U) + (1-\alpha)S_i(L)S_j(L+U) \\ & + \alpha R_i(L+U)R_j(L) + (1-\alpha)S_i(L+U)S_j(L)] \\ & + \omega^2 [\alpha R_i(L+U)R_j(L+U) + (1-\alpha)S_i(L+U)S_j(L+U)] \\ & - \omega^2 + 4\omega - 4 < 0. \end{aligned} \quad (4.18)$$

It is easy to verify that the discriminant  $\Delta$  of the quadratic in  $\sigma$  satisfies  $\Delta > 0$ , and so there holds

$$\frac{\omega P'_1 - \sqrt{\omega^2 P'_2 + (\omega - 2)^2 P'_3}}{P'_3} < \sigma < \frac{\omega P'_1 + \sqrt{\omega^2 P'_2 + (\omega - 2)^2 P'_3}}{P'_3}. \quad (4.19)$$

By  $\sigma < 0$  and  $1 - \sigma^2[\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] > 0$ , for all  $i, j \in N, i \neq j$ , we obtain

$$\max_{i,j \atop i \neq j} \frac{\omega P'_1 - \sqrt{\omega^2 P'_2 + (\omega - 2)^2 P'_3}}{P'_3} < \sigma < 0. \quad (4.20)$$

Therefore, by (4.16) and (4.20), we get

$$\max_{i,j \atop i \neq j} \frac{\omega P'_1 - \sqrt{\omega^2 P'_2 + P'_3 \min(\omega^2, (\omega - 2)^2)}}{P'_3} < \sigma < 0. \quad (4.21)$$

Finally, we prove (III).

(1) When  $0 < \omega \leq 1, \sigma \geq t$ ,

$$\begin{aligned} A_2 &= 2(1 - \omega) + 2(\omega - \sigma)\sigma[\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] \\ &\quad + \omega\sigma[\alpha R_i(L)R_j(U) + (1 - \alpha)S_i(U)S_j(L)] + \omega\sigma[\alpha R_i(U)R_j(L) + (1 - \alpha)S_i(L)S_j(U)], \\ A_3 &= (1 - \omega)^2 - \alpha[(\sigma - \omega)R_i(L) + \omega R_i(U)][(\sigma - \omega)R_j(L) + \omega R_j(U)] \\ &\quad - (1 - \alpha)[(\sigma - \omega)S_i(L) + \omega S_i(U)][(\sigma - \omega)S_j(L) + \omega S_j(U)]. \end{aligned} \quad (4.22)$$

By  $A_2 - A_3 < A_1$ , we have

$$\begin{aligned} &4\sigma^2[\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] \\ &\quad - 2\omega\sigma[\alpha R_i(L)(R_j(L) - R_j(U)) + (1 - \alpha)S_i(L)(S_j(L) - S_j(U))] \\ &\quad - 2\omega\sigma[\alpha R_j(L)(R_i(L) - R_i(U)) + (1 - \alpha)S_j(L)(S_i(L) - S_i(U))] \\ &\quad + \omega^2[\alpha(R_i(L) - R_i(U))(R_j(L) - R_j(U)) \\ &\quad + (1 - \alpha)(S_i(L) - S_i(U))(S_j(L) - S_j(U))] - \omega^2 < 0. \end{aligned} \quad (4.23)$$

It is easy to verify that the discriminant  $\Delta$  of the quadratic in  $\sigma$  satisfies  $\Delta > 0$ , and so there holds

$$\frac{\omega P'_4 - \omega \sqrt{P'_5^2 + P'_3}}{P'_3} < \sigma < \frac{\omega P'_4 + \omega \sqrt{P'_5^2 + P'_3}}{P'_3}. \quad (4.24)$$

By  $\sigma \geq t$ , we obtain

$$t \leq \sigma < \min_{\substack{i,j \\ i \neq j}} \frac{\omega P'_4 + \omega \sqrt{P'_5^2 + P'_3}}{P'_3}. \quad (4.25)$$

(2) When  $1 < \omega < t$ ,  $\sigma \geq t$ ,

$$\begin{aligned} A_2 = & 2(\omega - 1) + 2(\sigma - \omega)\sigma [\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] \\ & + \omega\sigma [\alpha R_i(L)R_j(U) + (1 - \alpha)S_i(U)S_j(L)] \\ & + \omega\sigma [\alpha R_i(U)R_j(L) + (1 - \alpha)S_i(L)S_j(U)], \end{aligned} \quad (4.26)$$

$$\begin{aligned} A_3 = & (1 - \omega)^2 - \alpha[(\sigma - \omega)R_i(L) + \omega R_i(U)][(\sigma - \omega)R_j(L) + \omega R_j(U)] \\ & - (1 - \alpha)[(\sigma - \omega)S_i(L) + \omega S_i(U)][(\sigma - \omega)S_j(L) + \omega S_j(U)]. \end{aligned}$$

By  $A_2 - A_3 < A_1$ , we have

$$\begin{aligned} & 4\sigma^2 [\alpha R_i(L)R_j(L) + (1 - \alpha)S_i(L)S_j(L)] \\ & - 2\omega\sigma [\alpha R_i(L)(R_j(L) - R_j(U)) + (1 - \alpha)S_i(L)(S_j(L) - S_j(U))] \\ & - 2\omega\sigma [\alpha R_j(L)(R_i(L) - R_i(U)) + (1 - \alpha)S_j(L)(S_i(L) - S_i(U))] \\ & + \omega^2 [\alpha(R_i(L) - R_i(U))(R_j(L) - R_j(U)) \\ & + (1 - \alpha)(S_i(L) - S_i(U))(S_j(L) - S_j(U))] - \omega^2 + 4\omega - 4 < 0. \end{aligned} \quad (4.27)$$

It is easy to verify that the discriminant  $\Delta$  of the quadratic in  $\sigma$  satisfies  $\Delta > 0$ , and so there holds

$$\frac{\omega P'_4 - \sqrt{\omega^2 P'_5^2 + (\omega - 2)^2 P'_3}}{P'_3} < \sigma < \frac{\omega P'_4 + \sqrt{\omega^2 P'_5^2 + (\omega - 2)^2 P'_3}}{P'_3}. \quad (4.28)$$

By  $\sigma \geq t$ , we obtain

$$t \leq \sigma < \min_{\substack{i,j \\ i \neq j}} \frac{\omega P'_4 + \sqrt{\omega^2 P'_5^2 + (\omega - 2)^2 P'_3}}{P'_3}. \quad (4.29)$$

Therefore, by (4.25) and (4.29), we obtain

$$t \leq \sigma < \min_{\substack{i,j \\ i \neq j}} \frac{\omega P'_4 + \sqrt{\omega^2 P_5'^2 + P'_3 \min(\omega^2, (\omega-2)^2)}}{P'_3}. \quad (4.30)$$

□

We can obtain the following results easily.

**Theorem 4.2.** Let  $A \in DD(\alpha)$ . If  $R_i(L+U)R_j(L+U) - S_i(L+U)S_j(L+U) > 0$ , when  $0 < \omega \leq 1$ , the following conditions hold:

$$\begin{aligned} \text{(I)} \quad 0 \leq \alpha &< \frac{\sqrt{R_i(L+U)R_j(L+U)} - S_i(L+U)S_j(L+U)}{R_i(L+U)R_j(L+U) - S_i(L+U)S_j(L+U)}, \\ \text{(II)} \quad \frac{\omega P'_1 - \sqrt{\omega^2 P_2'^2 + \omega^2 P'_3}}{P'_3} &< \frac{\omega P_1 - \sqrt{\omega^2 P_2^2 + \omega^2 P_3}}{P_3}, \\ \text{(III)} \quad \frac{\omega P'_4 + \sqrt{\omega^2 P_5'^2 + \omega^2 P'_3}}{P'_3} &> \frac{\omega P_4 + \sqrt{\omega^2 P_5^2 + \omega^2 P_3}}{P_3}, \end{aligned} \quad (4.31)$$

or when  $1 < \omega < t$ , the following conditions hold:

$$\begin{aligned} \text{(I)} \quad 0 \leq \alpha &< \frac{\sqrt{R_i(L+U)R_j(L+U)} - S_i(L+U)S_j(L+U)}{R_i(L+U)R_j(L+U) - S_i(L+U)S_j(L+U)}, \\ \text{(II)} \quad \frac{\omega P'_1 - \sqrt{\omega^2 P_2'^2 + (\omega-2)^2 P'_3}}{P'_3} &< \frac{\omega P_1 - \sqrt{\omega^2 P_2^2 + (\omega-2)^2 P_3}}{P_3}, \\ \text{(III)} \quad \frac{\omega P'_4 + \sqrt{\omega^2 P_5'^2 + (\omega-2)^2 P'_3}}{P'_3} &> \frac{\omega P_4 + \sqrt{\omega^2 P_5^2 + (\omega-2)^2 P_3}}{P_3}, \end{aligned} \quad (4.32)$$

then the area of convergence of AOR method obtained by Theorem 4.1 is larger than that obtained by Theorem 2.5.

**Theorem 4.3.** Let  $A \in DD(\alpha)$ . If  $R_i(L+U)R_j(L+U) - S_i(L+U)S_j(L+U) < 0$ , when  $0 < \omega \leq 1$ , the following conditions hold:

$$\begin{aligned} \text{(I)} \quad & \frac{\sqrt{R_i(L+U)R_j(L+U)} - S_i(L+U)S_j(L+U)}{R_i(L+U)R_j(L+U) - S_i(L+U)S_j(L+U)} < \alpha \leq 1, \\ \text{(II)} \quad & \frac{\omega P'_1 - \sqrt{\omega^2 P_2'^2 + \omega^2 P_3'}}{P'_3} < \frac{\omega P_1 - \sqrt{\omega^2 P_2^2 + \omega^2 P_3}}{P_3}, \\ \text{(III)} \quad & \frac{\omega P'_4 + \sqrt{\omega^2 P_5'^2 + \omega^2 P_3'}}{P'_3} > \frac{\omega P_4 + \sqrt{\omega^2 P_5^2 + \omega^2 P_3}}{P_3}, \end{aligned} \quad (4.33)$$

or when  $1 < \omega < t$ , the following conditions hold:

$$\begin{aligned} \text{(I)} \quad & \frac{\sqrt{R_i(L+U)R_j(L+U)} - S_i(L+U)S_j(L+U)}{R_i(L+U)R_j(L+U) - S_i(L+U)S_j(L+U)} < \alpha \leq 1, \\ \text{(II)} \quad & \frac{\omega P'_1 - \sqrt{\omega^2 P_2'^2 + (\omega-2)^2 P_3'}}{P'_3} < \frac{\omega P_1 - \sqrt{\omega^2 P_2^2 + (\omega-2)^2 P_3}}{P_3}, \\ \text{(III)} \quad & \frac{\omega P'_4 + \sqrt{\omega^2 P_5'^2 + (\omega-2)^2 P_3'}}{P'_3} > \frac{\omega P_4 + \sqrt{\omega^2 P_5^2 + (\omega-2)^2 P_3}}{P_3}, \end{aligned} \quad (4.34)$$

then the area of convergence of AOR method obtained by Theorem 4.1 is larger than that obtained by Theorem 2.5.

## 5. Examples

In the following examples, we give the areas of convergence of AOR method to show that our results are better than ones obtained by Theorems 2.4 and 2.5.

*Example 5.1* (see [6]). Let

$$A = \begin{pmatrix} 5 & 3 & 2 \\ 2 & 6 & 3 \\ 2 & 1 & 9 \end{pmatrix} = D - T - S, \quad (5.1)$$

where

$$\begin{aligned}
 D &= \begin{pmatrix} 5 & 0 & 0 \\ 0 & 6 & 0 \\ 0 & 0 & 9 \end{pmatrix}, & T &= \begin{pmatrix} 0 & 0 & 0 \\ -2 & 0 & 0 \\ -2 & -1 & 0 \end{pmatrix}, & S &= \begin{pmatrix} 0 & -3 & -2 \\ 0 & 0 & -3 \\ 0 & 0 & 0 \end{pmatrix}, \\
 L = D^{-1}T &= \begin{pmatrix} 0 & 0 & 0 \\ -\frac{1}{3} & 0 & 0 \\ -\frac{2}{9} & -\frac{1}{9} & 0 \end{pmatrix}, & U = D^{-1}S &= \begin{pmatrix} 0 & -\frac{3}{5} & -\frac{2}{5} \\ 0 & 0 & -\frac{1}{2} \\ 0 & 0 & 0 \end{pmatrix}, & L + U &= \begin{pmatrix} 0 & -\frac{3}{5} & -\frac{2}{5} \\ -\frac{1}{3} & 0 & -\frac{1}{2} \\ -\frac{2}{9} & -\frac{1}{9} & 0 \end{pmatrix}.
 \end{aligned} \tag{5.2}$$

Obviously,  $A \notin SD$ , but  $A \in DD(1/2)$ .

By Theorem 4.1, we have the following area of convergence:

$$\begin{aligned}
 (1) \quad 0 \leq \sigma < 1.1896, \quad 0 < \omega < \max \left\{ 1.2390, \frac{2\sigma}{(1 + \rho(M_{\sigma,\sigma}))} \right\}, \quad \text{or} \\
 (2) \quad 0 < \omega \leq 1, \quad -1.0266\omega < \sigma < 0, \quad \text{or} \\
 1 < \omega < 1.2390, \quad \frac{(22\omega - 3\sqrt{201\omega^2 - 800\omega + 800})}{20} < \sigma < 0, \quad \text{or} \\
 (3) \quad 0 < \omega \leq 1, \quad 1.2390 < \sigma < 2.0266\omega, \quad \text{or} \\
 1 < \omega < 1.2390, \quad 1.2390 \leq \sigma < \frac{(-2\omega + 3\sqrt{201\omega^2 - 800\omega + 800})}{20}.
 \end{aligned} \tag{5.3}$$

Obviously,  $A \in DD$ .

By Theorem 2.5, we have the following area of convergence:

$$\begin{aligned}
 (1) \quad 0 \leq \sigma < 1.1896, \quad 0 < \omega < \max \left\{ 1.0455, \frac{2\sigma}{(1 + \rho(M_{\sigma,\sigma}))} \right\}, \quad \text{or} \\
 (2) \quad 0 < \omega \leq 1, \quad -0.6712\omega < \sigma < 0, \quad \text{or} \\
 1 < \omega < 1.0455, \quad \frac{(7\omega - 3\sqrt{17\omega^2 - 64\omega + 64})}{8} < \sigma < 0, \quad \text{or} \\
 (3) \quad 0 < \omega \leq 1, \quad 1.0455 \leq \sigma < 1.6712\omega, \quad \text{or} \\
 1 < \omega < 1.0455, \quad 1.0455 \leq \sigma < \frac{(\omega + 3\sqrt{17\omega^2 - 64\omega + 64})}{8}.
 \end{aligned} \tag{5.4}$$

In addition,  $A \in D(1/2)$ .




Figure 1




Figure 2

By Theorem 2.4, we have the following area of convergence:

- (1)  $0 \leq \sigma < 1.1896, \quad 0 < \omega < \max\left\{1.1250, \frac{2\sigma}{(1 + \rho(M_{\sigma,\sigma}))}\right\}, \quad \text{or}$
  - (2)  $0 < \omega \leq 1, \quad -0.6\omega < \sigma < 0, \quad \text{or} \quad 1 < \omega < 1.1250, \quad \frac{7}{5}\omega - \frac{9}{5} < \sigma < 0, \quad \text{or}$
  - (3)  $0 < \omega \leq 1, \quad 1.1250 < \sigma < 1.31\omega, \quad \text{or} \quad 1 < \omega < 1.1250, \quad 1.1250 \leq \sigma < 1.8 - 0.49\omega.$
- (5.5)

Now we give two figures. In Figure 1, we can see that the area of convergence obtained by Theorem 4.1 (real line) is larger than that obtained by Theorem 2.5 (virtual line). In Figure 2, we can see that the area of convergence obtained by Theorem 4.1 (real line)

is larger than that obtained by Theorem 2.4 (virtual line). From above we know that the area of convergence obtained by Theorem 4.1 is larger than ones obtained by Theorems 2.5 and 2.4.

*Example 5.2.* Let

$$A = \begin{pmatrix} 3 & 1 & 1 \\ 1 & 2 & 1 \\ 2 & 1 & 3 \end{pmatrix}. \quad (5.6)$$

Obviously,  $A \in DD(1/3)$ ,  $A \notin D(\alpha)$ ,  $A \notin DD$ . So we cannot use Theorems 2.4 and 2.5. By Theorem 4.1, we have the following area of convergence:

$$\begin{aligned} (1) \quad 0 \leq \sigma < 1.0724, \quad 0 < \omega < \max \left\{ 1.0693, \frac{2\sigma}{(1 + \rho(M_{\sigma,\sigma}))} \right\}, \quad \text{or} \\ (2) \quad 0 < \omega \leq 1, \quad -0.1973\omega < \sigma < 0, \quad \text{or} \\ 1 < \omega < 1.0693, \quad \frac{(37\omega - \sqrt{1945\omega^2 - 7776\omega + 7776})}{36} < \sigma < 0, \quad \text{or} \\ (3) \quad 0 < \omega \leq 1, \quad 1.0693 < \sigma < 1.1973\omega, \quad \text{or} \\ 1 < \omega < 1.0693, \quad 1.0693 \leq \sigma < \frac{(\sqrt{1945\omega^2 - 7776\omega + 7776} - \omega)}{36}. \end{aligned} \quad (5.7)$$

## Acknowledgments


The authors would like to thank the referees for their valuable comments and suggestions, which greatly improved the original version of this paper. This work was supported by the National Natural Science Foundation of China (Grant no. 11001144) and the Science and Technology Program of Shandong Universities of China (J10LA06).

## References


- [1] A. Hadjidimos, "Accelerated overrelaxation method," *Mathematics of Computation*, vol. 32, no. 141, pp. 149–157, 1978.
- [2] B. Li and M. J. Tsatsomeros, "Doubly diagonally dominant matrices," *Linear Algebra and Its Applications*, vol. 261, pp. 221–235, 1997.
- [3] T. Z. Huang and G. B. Wang, "Convergence theorems for the AOR method," *Applied Mathematics and Mechanics*, vol. 23, no. 11, pp. 1183–1187, 2002.
- [4] M. Li and Y. X. Sun, "Discussion on  $\alpha$ -diagonally dominant matrices and their applications," *Chinese Journal of Engineering Mathematics*, vol. 26, no. 5, pp. 941–945, 2009.
- [5] L. Cvetković and D. Herceg, "Convergence theory for AOR method," *Journal of Computational Mathematics*, vol. 8, no. 2, pp. 128–134, 1990.
- [6] Z.-X. Gao and T.-Z. Huang, "Convergence of AOR method," *Applied Mathematics and Computation*, vol. 176, no. 1, pp. 134–140, 2006.
- [7] A. Hadjidimos and A. Yeyios, "The principle of extrapolation in connection with the accelerated overrelaxation method," *Linear Algebra and Its Applications*, vol. 30, pp. 115–128, 1980.




Advances in  
Operations Research



Advances in  
Decision Sciences




Mathematical Problems  
in Engineering




Algebra



Journal of  
Probability and Statistics



The Scientific  
World Journal




Hindawi

Submit your manuscripts at  
<http://www.hindawi.com>




International Journal of  
Differential Equations




International Journal of  
Combinatorics




Advances in  
Mathematical Physics




Journal of  
Complex Analysis




Journal of  
Mathematics




International Journal of  
Stochastic Analysis



Abstract and  
Applied Analysis



Discrete Dynamics in  
Nature and Society




International  
Journal of  
Mathematics and  
Mathematical  
Sciences



Journal of  
Discrete Mathematics



Journal of  
Function Spaces



Journal of  
Applied Mathematics



Journal of  
Optimization