

HYERS-ULAM-RASSIAS STABILITY OF JORDAN HOMOMORPHISMS ON BANACH ALGEBRAS

TAKESHI MIURA, SIN-EI TAKAHASI, AND GO HIRASAWA

Received 18 July 2003

We prove that a Jordan homomorphism from a Banach algebra into a semisimple commutative Banach algebra is a ring homomorphism. Using a signum effectively, we can give a simple proof of the Hyers-Ulam-Rassias stability of a Jordan homomorphism between Banach algebras. As a direct corollary, we show that to each approximate Jordan homomorphism f from a Banach algebra into a semisimple commutative Banach algebra there corresponds a unique ring homomorphism near to f .

1. Introduction and statement of results

It seems that the stability problem of functional equations had been first raised by Ulam (cf. [11, Chapter VI] and [12]): For what metric groups G is it true that an ε -automorphism of G is necessarily near to a strict automorphism?

An answer to the above problem has been given as follows. Suppose E_1 and E_2 are two real Banach spaces and $f : E_1 \rightarrow E_2$ is a mapping. If there exist $\delta \geq 0$ and $p \geq 0$, $p \neq 1$ such that

$$\|f(x+y) - f(x) - f(y)\| \leq \delta(\|x\|^p + \|y\|^p) \quad (1.1)$$

for all $x, y \in E_1$, then there is a unique additive mapping $T : E_1 \rightarrow E_2$ such that $\|f(x) - T(x)\| \leq 2\delta\|x\|^p/|2 - 2^p|$ for every $x \in E_1$. This result is called the Hyers-Ulam-Rassias stability of the additive Cauchy equation $g(x+y) = g(x) + g(y)$. Indeed, Hyers [5] obtained the result for $p = 0$. Then Rassias [8] generalized the above result of Hyers to the case where $0 \leq p < 1$. Gajda [4] solved the problem for $1 < p$, which was raised by Rassias; In the same paper, Gajda also gave an example that a similar result to the above does not hold for $p = 1$ (cf. [9]). If $p < 0$, then $\|x\|^p$ is meaningless for $x = 0$; In this case, if we assume that $\|0\|^p$ means ∞ , then the proof given in [8] also works for $x \neq 0$. Moreover, with minor changes in the proof, the result is also valid for $p < 0$. Thus, the Hyers-Ulam-Rassias stability of the additive Cauchy equation holds for $p \in \mathbb{R} \setminus \{1\}$. Here and after, the letter \mathbb{R} denotes the real number field and \mathbb{C} stands for the complex number field.

Suppose A and B are two Banach algebras. We say that a mapping $\tau : A \rightarrow B$ is a Jordan homomorphism if

$$\begin{aligned}\tau(a+b) &= \tau(a) + \tau(b) \quad (a, b \in A), \\ \tau(a^2) &= \tau(a)^2 \quad (a \in A).\end{aligned}\tag{1.2}$$

If, in addition, τ is multiplicative, that is

$$\tau(ab) = \tau(a)\tau(b) \quad (a, b \in A),\tag{1.3}$$

we say that τ is a ring homomorphism. The study of ring homomorphisms between Banach algebras A and B is of interest even if $A = B = \mathbb{C}$. For example, the zero mapping, the identity and the complex conjugate are ring homomorphisms on \mathbb{C} , which are all continuous. On the other hand, the existence of a discontinuous ring homomorphism on \mathbb{C} is well-known (cf. [6]). More explicitly, if G is the set of all surjective ring homomorphisms on \mathbb{C} , then $\#G = 2^{\#\mathbb{C}}$, where $\#S$ denotes the cardinal number of a set S . In fact, Charnow [3, Theorem 3] proved that there exist $2^{\#k}$ automorphisms for every algebraically closed field k ; It is also known that if \mathcal{A} is a uniform algebra on a compact metric space, then there are exactly $2^{\#\mathbb{C}}$ complex-valued ring homomorphisms on \mathcal{A} whose kernels are non-maximal prime ideals (see [7, Corollary 2.4]).

By definition, it is obvious that ring homomorphisms are Jordan homomorphisms. Conversely, under a certain condition, Jordan homomorphisms are ring homomorphisms. For example, each Jordan homomorphism τ from a commutative Banach algebra \mathcal{B} into \mathbb{C} is a ring homomorphism: Fix $a, b \in \mathcal{B}$ arbitrarily. Since $\tau((a+b)^2) = \tau(a+b)^2$, a simple calculation shows that $\tau(ab+ba) = 2\tau(a)\tau(b)$. The commutativity of \mathcal{B} implies $\tau(ab) = \tau(a)\tau(b)$, and hence τ is a ring homomorphism. This simple example leads us to the following general result.

THEOREM 1.1. *Suppose A is a Banach algebra, which need not be commutative, and suppose B is a semisimple commutative Banach algebra. If $\tau : A \rightarrow B$ is a Jordan homomorphism, then $\tau(ab) = \tau(a)\tau(b)$ for all $a, b \in A$, that is, τ is a ring homomorphism.*

Next, we consider the stability, in the sense of Hyers-Ulam-Rassias, of Jordan homomorphisms. Bourgin [2] proved the following stability result of ring homomorphisms between two unital Banach algebras.

THEOREM 1.2. *Suppose A and B are unital Banach algebras. If $f : A \rightarrow B$ is a surjective mapping such that*

$$\begin{aligned}\|f(a+b) - f(a) - f(b)\| &\leq \varepsilon \quad (a, b \in A), \\ \|f(ab) - f(a)f(b)\| &\leq \delta \quad (a, b \in A)\end{aligned}\tag{1.4}$$

for some $\varepsilon \geq 0$ and $\delta \geq 0$, then f is a ring homomorphism.

Applying a theorem of Hyers [5], Rassias [8] and Gajda [4], Badora [1] proved the Hyers-Ulam-Rassias stability of ring homomorphisms, which generalizes the above result of Bourgin. We will prove the Hyers-Ulam-Rassias stability of Jordan homomorphisms.

We emphasize that the introduction of the signum $s = |1 - p|/(1 - p)$ made it possible to give a simple proof of our stability results.

THEOREM 1.3. *Suppose A and B are Banach algebras. If $f : A \rightarrow B$ is a mapping such that*

$$\|f(a + b) - f(a) - f(b)\| \leq \delta(\|a\|^p + \|b\|^p) \quad (a, b \in A), \tag{1.5}$$

$$\|f(a^2) - f(a)^2\| \leq \delta\|a\|^{2p} \quad (a \in A) \tag{1.6}$$

for some $\delta \geq 0$ and $p \geq 0, p \neq 1$, then there is a unique Jordan homomorphism $\tau : A \rightarrow B$ such that

$$\|f(a) - \tau(a)\| \leq \frac{2\delta}{|2 - 2^p|} \|a\|^p \quad (a \in A). \tag{1.7}$$

For $p < 0$, we can also give a similar result to Theorem 1.3, under an additional condition that $f(0) = 0$. The hypothesis $f(0) = 0$ seems to be natural. It follows from (1.5) that $f(0) = 0$ whenever $p > 0$; On the other hand, if $p < 0$ then the inequalities (1.5) and (1.6) give no information for $f(0)$.

THEOREM 1.4. *Suppose A and B are Banach algebras. If $f : A \rightarrow B$ is a mapping, with $f(0) = 0$, such that the inequalities (1.5) and (1.6) are valid for some $\delta \geq 0$ and $p < 0$, then there is a unique Jordan homomorphism $\tau : A \rightarrow B$ such that*

$$\|f(a) - \tau(a)\| \leq \frac{2\delta}{|2 - 2^p|} \|a\|^p \quad (a \in A). \tag{1.8}$$

As an easy corollary to Theorems 1.1, 1.3, and 1.4, we obtain the following stability result.

COROLLARY 1.5. *Suppose A is a Banach algebra and suppose B is a semisimple commutative Banach algebra. If $f : A \rightarrow B$ is a mapping such that*

$$\|f(a + b) - f(a) - f(b)\| \leq \delta(\|a\|^p + \|b\|^p) \quad (a, b \in A), \tag{1.9}$$

$$\|f(a^2) - f(a)^2\| \leq \delta\|a\|^{2p} \quad (a \in A)$$

for some $\delta \geq 0$ and $p \in \mathbb{R}$. If $p \geq 0$ and $p \neq 1$, or $p < 0$ and $f(0) = 0$, then there is a unique ring homomorphism $\tau : A \rightarrow B$ such that

$$\|f(a) - \tau(a)\| \leq \frac{2\delta}{|2 - 2^p|} \|a\|^p \quad (a \in A). \tag{1.10}$$

2. Proof of results

Before we turn to the proof of Theorem 1.1, we need the following lemma. It should be mentioned that the following proof is just a slight modification of [13, Proof of Theorem 1] by Żelazko.

LEMMA 2.1. *Suppose A is a Banach algebra, which need not be commutative. Then each Jordan homomorphism $\phi : A \rightarrow \mathbb{C}$ is a ring homomorphism.*

Proof. Recall that ϕ is an additive mapping such that $\phi(a^2) = \phi(a)^2$ for all $a \in A$. Replacement of a by $x + y$ results in

$$\phi(xy + yx) = 2\phi(x)\phi(y) \quad (x \in A, y \in A). \quad (2.1)$$

Then (2.1), with $x = x^2$, implies

$$\phi(x^2y + yx^2) = 2\phi(x)^2\phi(y). \quad (2.2)$$

Taking $y = xy + yx$ in (2.1), we see that

$$\phi(x(xy + yx) + (xy + yx)x) = 2\phi(x)\phi(xy + yx), \quad (2.3)$$

and hence, by (2.1)

$$\phi(x^2y + 2xyx + yx^2) = 4\phi(x)^2\phi(y) \quad (x \in A, y \in A). \quad (2.4)$$

Subtraction (2.4) from (2.2) gives

$$\phi(xyx) = \phi(x)^2\phi(y) \quad \text{if } x \in A, y \in A. \quad (2.5)$$

Fix $a \in A$ and $b \in A$ arbitrarily, and put

$$2t = \phi(ab - ba). \quad (2.6)$$

It follows from (2.1) and (2.6) that

$$\phi(ab) = \phi(a)\phi(b) + t, \quad \phi(ba) = \phi(a)\phi(b) - t. \quad (2.7)$$

By (2.5), (2.6), (2.7),

$$\begin{aligned} 4t^2 &= \phi((ab - ba)^2) \\ &= \phi(ab)^2 - \phi(ab^2a) - \phi(ba^2b) + \phi(ba)^2 \\ &= \{\phi(a)\phi(b) + t\}^2 - 2\phi(a)^2\phi(b)^2 + \{\phi(a)\phi(b) - t\}^2 \\ &= 2t^2; \end{aligned} \quad (2.8)$$

hence $t = 0$, which proves $\phi(ab) = \phi(ba)$. It follows from (2.1) that $\phi(ab) = \phi(a)\phi(b)$, and the proof is complete. \square

Proof of Theorem 1.1. We show that τ is multiplicative. Let M_B be the maximal ideal space of B . We associate to each $\varphi \in M_B$ a function $\tau_\varphi : A \rightarrow \mathbb{C}$ defined by

$$\tau_\varphi(a) = \varphi(\tau(a)) \quad (a \in A). \quad (2.9)$$

Pick $\varphi \in M_B$ arbitrarily. We see that $\tau_\varphi(a^2) = \tau_\varphi(a)^2$ for all $a \in A$, and so Lemma 2.1, applied to τ_φ , implies that τ_φ is multiplicative. By the definition of τ_φ , we get $\varphi(\tau(ab)) = \varphi(\tau(a)\tau(b))$ for all $a, b \in A$. Since $\varphi \in M_B$ was arbitrary and since B is assumed to be semisimple, we obtain $\tau(ab) = \tau(a)\tau(b)$ for all $a, b \in A$. We thus conclude that τ is a ring homomorphism, and the proof is complete. \square

Proof of Theorem 1.3. It follows from [8] and [4] (cf. [5]) that there is an additive mapping $\tau : A \rightarrow B$ such that

$$\|f(a) - \tau(a)\| \leq \frac{2\delta}{|2 - 2^p|} \|a\|^p \quad (a \in A). \tag{2.10}$$

We first show that $\tau(a^2) = \tau(a)^2$ for all $a \in A$. Pick $a \in A$ arbitrarily, and put $s = |1 - p|/(1 - p)$. Note that $s = 1$ if $0 \leq p < 1$ and that $s = -1$ if $p > 1$. Since τ is additive, it follows from (2.10) that

$$\begin{aligned} \|n^{-2s}f(n^{2s}a^2) - \tau(a^2)\| &= \|n^{-2s}f(n^{2s}a^2) - n^{-2s}\tau(n^{2s}a^2)\| \\ &\leq n^{-2s} \frac{2\delta}{|2 - 2^p|} \|n^{2s}a^2\|^p \end{aligned} \tag{2.11}$$

for all $n \in \mathbb{N}$, and hence

$$\|n^{-2s}f(n^{2s}a^2) - \tau(a^2)\| \leq n^{2s(p-1)} \frac{2\delta}{|2 - 2^p|} \|a^2\|^p \tag{2.12}$$

for all $n \in \mathbb{N}$. A similar argument to the above shows for each $n \in \mathbb{N}$ that

$$\|n^{-s}f(n^s a) - \tau(a)\| \leq n^{s(p-1)} \frac{2\delta}{|2 - 2^p|} \|a\|^p. \tag{2.13}$$

Since $s(p - 1) < 0$, it follows from (2.12) and (2.13) that

$$\tau(a^2) = \lim_{n \rightarrow \infty} n^{-2s}f(n^{2s}a^2), \quad \tau(a) = \lim_{n \rightarrow \infty} n^{-s}f(n^s a). \tag{2.14}$$

By (1.6), we get $\|f(n^{2s}a^2) - f(n^s a)^2\| \leq \delta \|n^s a\|^{2p}$ for all $n \in \mathbb{N}$. So,

$$\lim_{n \rightarrow \infty} n^{-2s} \left(f(n^{2s}a^2) - f(n^s a)^2 \right) \leq \lim_{n \rightarrow \infty} n^{2s(p-1)} \delta \|a\|^{2p} = 0, \tag{2.15}$$

since $s(p - 1) < 0$. Now it follows from (2.14) and (2.15) that

$$\begin{aligned} \tau(a^2) &= \lim_{n \rightarrow \infty} n^{-2s}f(n^{2s}a^2) \\ &= \lim_{n \rightarrow \infty} \left\{ n^{-2s}f(n^{2s}a^2) - n^{-2s} \left(f(n^{2s}a^2) - f(n^s a)^2 \right) \right\} \\ &= \left\{ \lim_{n \rightarrow \infty} n^{-s}f(n^s a) \right\}^2 = \tau(a)^2. \end{aligned} \tag{2.16}$$

Since $a \in A$ was arbitrary, we obtain $\tau(a^2) = \tau(a)^2$ for all $a \in A$, and hence τ is a Jordan homomorphism.

Finally, suppose that $\tau^* : A \rightarrow B$ is another Jordan homomorphism such that $\|f(a) - \tau^*(a)\| \leq 2\delta \|a\|^p / |2 - 2^p|$ for all $a \in A$. Then (2.13), with $\tau = \tau^*$, is also valid. We thus obtain

$$\begin{aligned} \|\tau(a) - \tau^*(a)\| &\leq \|\tau(a) - n^{-s}f(n^s a)\| + \|n^{-s}f(n^s a) - \tau^*(a)\| \\ &\leq n^{s(p-1)} \frac{4\delta}{|2 - 2^p|} \|a\|^p \end{aligned} \tag{2.17}$$

for all $a \in A$ and $n \in \mathbb{N}$. Since $s(p - 1) < 0$, it follows that $\tau = \tau^*$, and hence the uniqueness have been proved. \square

Proof of Theorem 1.4. It follows from [8] that there exists an additive mapping $\tau : A \rightarrow B$ such that

$$\|f(a) - \tau(a)\| \leq \frac{2\delta}{|2 - 2^p|} \|a\|^p \quad (a \in A), \tag{2.18}$$

where we assume $\|0\|^p = \infty$. It suffices to show that $\tau(a^2) = \tau(a)^2$ for all $a \in A$. Since τ is additive, we obtain $\tau(0) = 0$, and so the case $a = 0$ is omitted. Pick $a \in A \setminus \{0\}$ arbitrarily. There are now two possibilities. Either $a^2 = 0$ or $a^2 \neq 0$, in which case the proof of Theorem 1.3 works well, and so $\tau(a^2) = \tau(a)^2$. Thus we need consider only the case $a^2 = 0$ (In this case, we cannot apply the proof of Theorem 1.3. In fact, if $a^2 = 0$, then $\|a^2\|^p = \infty$ and hence (2.13), with $a = a^2$, is meaningless). We will show that $\tau(a)^2 = 0$ whenever $a^2 = 0$.

Pick $a \in A \setminus \{0\}$ such that $a^2 = 0$. It follows from (1.6), with the hypothesis $f(0) = 0$, that

$$\|n^{-2} f(na)^2\| \leq n^{-2} \delta \|na\|^{2p} = n^{2(p-1)} \delta \|a\|^{2p}. \tag{2.19}$$

Since $a \neq 0$ and since $p - 1 < 0$, we obtain

$$\lim_{n \rightarrow \infty} n^{-2} f(na)^2 = 0. \tag{2.20}$$

Note also that

$$\|n^{-1} f(na) - \tau(a)\| \leq n^{-1} \frac{2\delta}{|2 - 2^p|} \|na\|^p = n^{p-1} \frac{2\delta}{|2 - 2^p|} \|a\|^p \tag{2.21}$$

for all $n \in \mathbb{N}$, and hence

$$\tau(a) = \lim_{n \rightarrow \infty} n^{-1} f(na). \tag{2.22}$$

It follows from (2.20) and (2.22) that

$$\tau(a)^2 = \lim_{n \rightarrow \infty} n^{-2} f(na)^2 = 0, \tag{2.23}$$

which proves $\tau(a^2) = 0 = \tau(a)^2$ whenever $a^2 = 0$. This completes the proof. \square

In this paper, we have proved the Hyers-Ulam-Rassias stability of Jordan homomorphisms for $p \in \mathbb{R} \setminus \{1\}$. On the other hand, Šemrl [10] gave an example that the stability result fails for $p = 1$: In fact, to each $\delta > 0$ there corresponds a multiplicative continuous function $f : \mathbb{C} \rightarrow \mathbb{C}$ satisfying $f(ia) = if(a)$ for all $a \in \mathbb{C}$ such that

$$|f(a + b) - f(a) - f(b)| \leq \delta(|a| + |b|) \quad (a, b \in \mathbb{C}) \tag{2.24}$$

and that

$$\sup_{a \in \mathbb{C} \setminus \{0\}} \frac{|f(a) - \tau(a)|}{|a|} \geq 1 \quad (2.25)$$

for all ring homomorphism $\tau : \mathbb{C} \rightarrow \mathbb{C}$.

Acknowledgment

The first and second authors are partly supported by the Grant-in-Aid for Scientific Research, Japan Society for the Promotion of Science.

References

- [1] R. Badora, *On approximate ring homomorphisms*, J. Math. Anal. Appl. **276** (2002), no. 2, 589–597.
- [2] D. G. Bourgin, *Approximately isometric and multiplicative transformations on continuous function rings*, Duke Math. J. **16** (1949), 385–397.
- [3] A. Charnow, *The automorphisms of an algebraically closed field*, Canad. Math. Bull. **13** (1970), 95–97.
- [4] Z. Gajda, *On stability of additive mappings*, Int. J. Math. Math. Sci. **14** (1991), no. 3, 431–434.
- [5] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. U.S.A. **27** (1941), 222–224.
- [6] H. Kestelman, *Automorphisms of the field of complex numbers*, Proc. London Math. Soc. (2) **53** (1951), 1–12.
- [7] T. Miura, S.-E. Takahasi, and N. Niwa, *Prime ideal which is the kernel of some complex ring homomorphism on a commutative complex algebra*, preprint.
- [8] T. M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), no. 2, 297–300.
- [9] T. M. Rassias and P. Šemrl, *On the behavior of mappings which do not satisfy Hyers-Ulam stability*, Proc. Amer. Math. Soc. **114** (1992), no. 4, 989–993.
- [10] P. Šemrl, *Nonlinear perturbations of homomorphisms on $C(X)$* , Quart. J. Math. Oxford Ser. (2) **50** (1999), no. 197, 87–109.
- [11] S. M. Ulam, *A collection of Mathematical Problems*, Interscience Tracts in Pure and Applied Mathematics, no. 8, Interscience Publishers, New York, 1960.
- [12] ———, *Sets, Numbers, and Universes: Selected Works*, Mathematicians of Our Time, vol. 9, MIT Press, Massachusetts, 1974.
- [13] W. Żelazko, *A characterization of multiplicative linear functionals in complex Banach algebras*, Studia Math. **30** (1968), 83–85.

Takeshi Miura: Department of Basic Technology, Applied Mathematics and Physics, Yamagata University, Yonezawa 992-8510, Japan
E-mail address: miura@yz.yamagata-u.ac.jp

Sin-Ei Takahasi: Department of Basic Technology, Applied Mathematics and Physics, Yamagata University, Yonezawa 992-8510, Japan
E-mail address: sin-ei@emperor.yz.yamagata-u.ac.jp

Go Hirasawa: Department of Mathematics, Nippon Institute of Technology, Miyashiro, Saitama 345-8501, Japan
E-mail address: hirasawa1@muh.biglobe.ne.jp