

## Research Article

# On Strong Law of Large Numbers for Dependent Random Variables

**Zhongzhi Wang**

*Faculty of Mathematics and Physics, Anhui University of Technology, Ma'anshan 243002, China*

Correspondence should be addressed to Zhongzhi Wang, wzz30@ahut.edu.cn

Received 16 December 2010; Accepted 3 March 2011

Academic Editor: Vijay Gupta

Copyright © 2011 Zhongzhi Wang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We discuss strong law of large numbers and complete convergence for sums of uniformly bounded negatively associate (NA) random variables (RVs). We extend and generalize some recent results. As corollaries, we investigate limit behavior of some other dependent random sequence.

## 1. Introduction

Throughout this paper, let  $\mathbb{N}$  denote the set of nonnegative integer, let  $\{X, X_n, n \in \mathbb{N}\}$  be a sequence of random variables defined on probability space  $(\Omega, \mathcal{F}, P)$ , and put  $S_n = \sum_{k=1}^n X_k$ . The symbol  $C$  will denote a generic constant ( $0 < C < \infty$ ) which is not necessarily the same one in each appearance.

In [1], Jajte studied a large class of summability method as follows: a sequence  $\{X_n, n \geq 1\}$  is summable to  $X$  by the method  $(h, g)$  if

$$\frac{1}{g(n)} \sum_{k=1}^n \frac{X_k}{h(k)} \rightarrow X, \quad \text{as } n \rightarrow \infty. \quad (1.1)$$

The main result of Jajte is as follows.

**Theorem 1.1.** *Let  $g(\cdot)$  be a positive, increasing function and  $h(\cdot)$  a positive function such that  $\phi(y) \equiv g(y)h(y)$  satisfies the following conditions.*

- (1) *For some  $d \geq 0$ ,  $\phi(\cdot)$  is strictly increasing on  $[d, +\infty)$  with range  $[0, +\infty)$ .*
- (2) *There exist  $C$  and a positive integer  $k_0$  such that  $\phi(y+1)/\phi(y) \leq C$ ,  $y \geq k_0$ .*
- (3) *There exist constants  $a$  and  $b$  such that  $\phi^2(s) \int_s^\infty (1/\phi^2(x)) dx \leq as + b$ ,  $s > d$ .*

Then, for i.i.d. random variables  $\{X_n, n \in \mathbb{N}\}$ ,

$$\frac{1}{g(n)} \sum_{k=1}^n \frac{X_k - EX_k \mathbf{1}_{[|X_k| \leq \phi(k)]}}{h(k)} \rightarrow 0 \quad \text{a.s. iff } E[\phi^{-1}(|X|)] < \infty, \quad (1.2)$$

where  $\phi^{-1}$  is the inverse of function  $\phi$ ,  $\mathbf{1}_A$  is the indicator of event  $A$ .

Motivated by Jajte [1], the present paper is devoted to the study of the limiting behavior of sums when  $\{X, X_n, n \in \mathbb{N}\}$  are dependent RVs. In particular, we will consider the case when  $\{X, X_n, n \in \mathbb{N}\}$  are NA RVs and obtain some general results on the complete convergence of dependent RVs. First, we shall give some definitions.

*Definition 1.2.* A finite family of random variables  $\{X_n, 1 \leq i \leq n\}$  is said to be negatively associated (abbreviated NA) if, for every pair of disjoint subsets  $A$  and  $B$  of  $\{1, 2, \dots, n\}$ , we have

$$\text{Cov}(f_1(X_i, i \in A), f_2(X_j, j \in B)) \leq 0, \quad (1.3)$$

whenever  $f_1$  and  $f_2$  are coordinatewise increasing and the covariance exists.

*Definition 1.3.* A finite family of random variables  $\{X_n, 1 \leq i \leq n\}$  is said to be positively associated (abbreviated PA) if

$$\text{Cov}(f(X_1, \dots, X_n), g(X_1, \dots, X_n)) \geq 0, \quad (1.4)$$

whenever  $f$  and  $g$  are coordinatewise increasing and the covariance exists.

An infinite family of random variables is NA (resp., PA) if every finite subfamily is NA (resp., PA).

Let  $\mathcal{F}_m^n$  be the  $\sigma$ -algebra generated by RVs  $X_i, m \leq i \leq n$ .

*Definition 1.4.* A sequence of random variables  $\{X_n, n \in \mathbb{N}\}$  is said to be  $m$ -dependence if  $\mathcal{F}_1^r$  and  $\mathcal{F}_{r'}^\infty$  are independent for all  $r$  and  $r'$  such that  $1 \leq r < r' < \infty, r' - r > m$ .

*Definition 1.5.* A sequence of random variables  $\{X_n, n \in \mathbb{N}\}$  is said to be  $\varphi$ -mixing (or uniformly strong mixing), if

$$\varphi(\tau) = \sup_{t \in \mathbb{N}} \sup_{A \in \mathcal{F}_1^t, B \in \mathcal{F}_{t+\tau}^\infty, P(A) > 0} |P(B | A) - P(B)| \rightarrow 0 \quad \text{as } \tau \rightarrow \infty. \quad (1.5)$$

These concepts of dependence were introduced by Esary et al. [2] and Joag-Dev and Proschan [3]. Their basic properties may be found in [2, 3] and the references therein.

*Definition 1.6.* Let  $\{X_n, n \in \mathbb{N}\}$  be a sequence of random variables which is said to be: uniformly bounded by a random variable  $X$  (we write  $\{X_n, n \in \mathbb{N}\} < X$ ) if there exists a constant  $C > 0$ , for almost every  $\omega \in \Omega$ , such that

$$\sup_{n \geq 1} P\{|X_n| > t\} \leq CP\{|X| > t\} \quad \forall t > 0. \quad (1.6)$$

*Remark 1.7.* The uniformly bounded random variables in (1.6) can be insured by moment conditions. For example, if

$$\sup_n E|X_n|^{2+\delta} < \infty \quad \text{for some } \delta > 0, \quad (1.7)$$

then there exists a uniformly bounded random variable  $X$  such that  $EX^2 < \infty$ .

The structure of this paper is as follows. Some needed technical results will be presented in Section 2. The strong law of large numbers for NA RVs will be established in Section 3. The Spitzer and Hus-Robbins-type law of large numbers will be presented in Sections 4 and 5, respectively.

## 2. Preliminaries

We now present some terminologies and lemmas. The following six properties are listed for reference in obtaining the main results in the next sections. Detailed proofs can be founded in the cited references.

**Lemma 2.1** (cf. [4]) (three-series theorem for NA). *Let  $\{X_n, n \in \mathbb{N}\}$  be NA. Let  $C > 0$  and let  $X_n^C = X_n \mathbf{1}_{\{|X_n| \leq C\}}$ . In order that  $\sum_{n=1}^{\infty} X_n$  converges a.s., it is sufficient that*

- (1)  $\sum_{n=1}^{\infty} P\{|X_n| > C\} < \infty$ ,
- (2)  $\sum_{n=1}^{\infty} EX_n^C$  converges,
- (3)  $\sum_{n=1}^{\infty} \text{var}(X_n^C) < \infty$ .

**Lemma 2.2** (cf. [4]). *Let  $\{X_n, n \in \mathbb{N}\}$  be NA with  $EX_n = 0$ ,  $EX_n^2 < \infty$ , then for  $p > 2$ , for all  $\varepsilon > 0$*

$$\begin{aligned} P\left\{\max_{1 \leq k \leq n} |S_k| \geq \varepsilon\right\} &\leq 2\varepsilon^{-2} \sum_{i=1}^n EX_i^2, \\ E\left(\max_{1 \leq k \leq n} |S_k|^2\right) &\leq C \sum_{i=1}^n EX_i^2, \\ E\left(\max_{1 \leq k \leq n} |S_k|^p\right) &\leq C \left\{ \sum_{i=1}^n EX_i^p + \left(\sum_{i=1}^n EX_i^2\right)^{p/2} \right\}. \end{aligned} \quad (2.1)$$

**Lemma 2.3** (cf. [5]). Let  $\{X_n, n \in \mathbb{N}\}$  be  $m$ -dependence with  $EX_n = 0$ ,  $EX_n^2 < \infty$ , then

$$E(S_n)^2 \leq (m+1) \sum_{i=1}^n EX_i^2. \quad (2.2)$$

**Lemma 2.4** (cf. [5]). Let  $\{X_n, n \in \mathbb{N}\}$  be  $\varphi$ -mixing with  $EX_n = 0$ ,  $EX_n^2 < \infty$ , then

$$E(S_n)^2 \leq \left(1 + 4 \sum_{r=1}^{n-1} \varphi^{1/2}(r)\right) \sum_{i=1}^n EX_i^2. \quad (2.3)$$

**Lemma 2.5** (cf. [6]). Let  $\{X_n, n \in \mathbb{N}\}$  be PA with  $EX_n = 0$ ,  $EX_n^2 < \infty$ , then

$$E \max_{1 \leq k \leq n} S_k^2 \leq 2ES_n^2. \quad (2.4)$$

Furthermore, if

$$\sum_{k=1}^n \mu^{1/2}(2^k) < \infty, \quad (2.5)$$

then

$$E \max_{1 \leq k \leq n} S_k^2 \leq Cn \max_{1 \leq k \leq n} EX_k^2, \quad (2.6)$$

where  $\mu(n) = \sup_{i \geq 1} \sum_{j: j-i \geq n} \text{cov}(X_i, X_j)$ .

**Lemma 2.6.** Let  $\{X_n, n \geq 1\}$  be a sequence of random variables and  $X$  a random variable. If  $\{X_n\} < X$ , then for all  $t > 0$ ,  $p \geq 2$

$$\mathbb{E}X_n^p \mathbf{1}_{[|X_n| \leq t]} \leq C [t^p \mathbb{P}\{|X| > t\} + \mathbb{E}X^p \mathbf{1}_{[|X| \leq t]}]. \quad (2.7)$$

*Proof.* By the integral equality

$$p \int_0^t s^{p-1} \mathbb{P}(|X| > s) ds = t^p \mathbb{P}(|X| > t) + \mathbb{E}|X|^p \mathbf{1}_{[|X| \leq t]}, \quad (2.8)$$

it follows that

$$\begin{aligned} \mathbb{E}|X_n|^p \mathbf{1}_{[|X_n| \leq t]} &\leq p \int_0^t s^{p-1} \mathbb{P}(X_n > s) ds \\ &\leq Cp \int_0^t s^{p-1} \mathbb{P}(|X| > s) ds = C [t^p \mathbb{P}(|X| > t) + \mathbb{E}|X|^p \mathbf{1}_{[|X| \leq t]}]. \quad \square \end{aligned} \quad (2.9)$$

**Lemma 2.7** (cf. [7]). Let  $\{A_n, n \geq 1\}$  be a sequence of events defined on  $(\Omega, \mathcal{F}, P)$ . If  $\sum_{n=1}^{\infty} P(A_n) < \infty$ , then  $P(\limsup_n A_n) = 0$ , if  $\sum_{n=1}^{\infty} P(A_n) = \infty$  and  $P(A_k \cap A_m) \leq P(A_k)P(A_m)$  for  $k \neq m$ , then  $P(\limsup_n A_n) = 1$ .

### 3. Strong Law of Large Numbers

**Theorem 3.1.** Let  $g(\cdot), h(\cdot)$ , and  $\phi(\cdot)$  be as in Theorem 1.1, and let  $\{X, X_n, n \in \mathbb{N}\}$  be a sequence of negatively associated random variables with  $EX_n = 0$ . Assume that  $\{X_n, n \in \mathbb{N}\} \prec X$ . If  $E[\phi^{-1}(|X|)] < \infty$ , then

$$\lim_n \frac{1}{g(n)} \sum_{k=1}^n \frac{X_k}{h(k)} = 0, \quad \text{a.s.} \quad (3.1)$$

Conversely, let  $\{X, X_n, n \in \mathbb{N}\}$  be a sequence of identically distributed NA random variables, if (3.1) is true, then  $E[\phi^{-1}(|X|)] < \infty$ .

*Proof.* Assume that  $E[\phi^{-1}(X)] < \infty$ . To prove (3.1) by applying the Kronecker lemma, it suffices to show that

$$\text{the series } \sum_{k=1}^{\infty} \frac{X_k}{\phi(k)} \text{ converges a.s.} \quad (3.2)$$

Here, we shall use the three-series theorem for NA RVs.

Let  $Y_k = X_k / \phi(k) \mathbf{1}_{\{|X_k / \phi(k)| \leq 1\}}$ . Then, by  $E[\phi^{-1}(|X|)] < \infty$ , we have

$$\begin{aligned} \sum_{k=1}^{\infty} P\left\{\left|\frac{X_k}{\phi(k)}\right| > 1\right\} &\leq C \sum_{k=1}^{\infty} P\left\{\left|\frac{X}{\phi(k)}\right| > 1\right\} \\ &= C \sum_{k=1}^{\infty} P\left\{\phi^{-1}(|X|) > k\right\} \leq CE\left[\phi^{-1}(|X|)\right] < \infty, \end{aligned} \quad (3.3)$$

which shows that  $P\{|X_k / \phi(k)| > 1, i.o.\} = 0$ , and

$$\sum_{k=1}^{\infty} \left|\frac{X_k}{\phi(k)}\right| \mathbf{1}_{\{|X_k / \phi(k)| > 1\}} < \infty \quad \text{a.s.} \quad (3.4)$$

Therefore, from  $EX_k = 0$ , it follows that

$$\sum_{k=1}^{\infty} |E(Y_k)| \leq \sum_{k=1}^{\infty} E\left|\frac{X_k}{\phi(k)}\right| \mathbf{1}_{\{|X_k / \phi(k)| > 1\}} < \infty. \quad (3.5)$$

To this end we estimate the series

$$\begin{aligned}
\sum_{k=1}^{\infty} \text{Var}(Y_k) &\leq \sum_{k=1}^{\infty} E(Y_k^2) = E \sum_{k=1}^{\infty} \frac{X_k^2}{\phi^2(k)} \mathbf{1}_{[|X_k/\phi(k)| \leq 1]} \\
&\leq C \sum_{k=1}^{\infty} \left[ E \mathbf{1}_{[|X| > \phi(k)]} + E \frac{X^2}{\phi^2(k)} \mathbf{1}_{[|X/\phi(k)| \leq 1]} \right] \\
&\leq C \sum_{k=1}^{\infty} P\{|X| > \phi(k)\} + C \left[ k_0 + C \sum_{k=k_0+1}^{\infty} \frac{EX^2}{\phi^2(k+1)} \mathbf{1}_{[|X| \leq \phi(k)]} \right] \quad (3.6) \\
&\leq CE[\phi^{-1}(|X|)] + Ck_0 + CE \left[ X^2 \int_{\phi^{-1}(|X|)}^{\infty} \frac{1}{\phi^2(x)} dx \right] \\
&\leq CE[\phi^{-1}(|X|)] + Ck_0 + CaE[\phi^{-1}(|X|)] + Cb < \infty.
\end{aligned}$$

Conversely, since  $\{X, X_n, n \in \mathbb{N}\}$  are identically NA RVs. If (3.1) holds, that is,

$$\frac{1}{\phi(n)} \sum_{k=1}^n X_k = o(1), \quad \text{a.s.} \quad (3.7)$$

It follows that

$$\frac{X_n}{\phi(n)} = \frac{\sum_{k=1}^n X_k}{\phi(n)} - \frac{\phi(n-1)}{\phi(n)} \frac{\sum_{k=1}^{n-1} X_k}{\phi(n-1)} \rightarrow o(1) \quad \text{a.s.} \quad (3.8)$$

which shows that  $\phi^{-1}(n)X_n \rightarrow 0$  a.s. Hence,

$$\phi^{-1}(n)X_n^{\pm} \rightarrow 0 \quad \text{a.s.}, \quad (3.9)$$

where  $x^+ = \max(0, x)$  and  $x^- = \max(0, -x)$ .

Since  $\{X_n^{\pm}, n \in \mathbb{N}\}$  is still an NA sequence. Defining the following events,

$$A_n = \left\{ X_n^+ > \frac{\varepsilon \phi(n)}{3} \right\}, \quad B_n = \left\{ X_n^- > \frac{\varepsilon \phi(n)}{3} \right\}, \quad (3.10)$$

we have  $P(A_k \cap A_l) \leq P(A_k)P(A_l)$ , and  $P(B_k \cap B_l) \leq P(B_k)P(B_l)$ , for  $k \neq l$ . By Lemma 2.7, if  $\phi^{-1}(n)X_n^{\pm} \rightarrow 0$  a.s., then  $\sum_{n=1}^{\infty} P\{X_n^+ > (1/3)\varepsilon\phi(n)\} < \infty$  and  $\sum_{n=1}^{\infty} P\{X_n^- > (1/3)\varepsilon\phi(n)\} < \infty$ . Therefore,

$$\sum_{n=1}^{\infty} P\{|X_n| > \varepsilon\phi(n)\} \leq \sum_{n=1}^{\infty} P\left\{X_n^+ > \frac{1}{3}\varepsilon\phi(n)\right\} + \sum_{n=1}^{\infty} P\left\{X_n^- > \frac{1}{3}\varepsilon\phi(n)\right\} < \infty, \quad \forall \varepsilon > 0, \quad (3.11)$$

which is equivalent to  $E[\phi^{-1}(|X|)] < \infty$ .

These complete the proof of Theorem 3.1.  $\square$

Theorem 3.1 also includes a particular case of logarithmic means, we can establish the following.

**Corollary 3.2.** *Let  $\{X, X_n, n \in \mathbb{N}\}$  be a sequence of NA RVs with  $EX_n = 0$  and  $\{X_n, n \in \mathbb{N}\} < X$ . If  $E|X| < \infty$ , then, one has*

$$\lim_n \frac{1}{\log n} \sum_{k=1}^n \frac{X_k}{k} = 0, \quad a.s. \quad (3.12)$$

*Proof.* Let  $h(y) = y$ ,  $g(y) = \log y$ , that is,  $\phi(y) = y \log y$ . In this case,  $\phi^{-1}(y) \sim y / \log y$  as  $y \rightarrow \infty$ , therefore  $E(|X|^\alpha) \leq E[\phi^{-1}(|X|)] \leq E(|X|)$ , for  $0 < \alpha < 1$ .  $\square$

**Corollary 3.3.** *Let  $\{X, X_n, n \in \mathbb{N}\}$  be a sequence of NA RVs with  $EX_n = 0$  and  $\{X_n, n \in \mathbb{N}\} < X$ . If  $E|X| < \infty$ , then, for every  $a, b \geq 0$ ,  $a + b > 1/2$ , one has*

$$\lim_n \frac{1}{n^a} \sum_{k=1}^n \frac{X_k}{k^b} = 0, \quad a.s. \quad (3.13)$$

*Remark 3.4.* As pointed out by Jajte [1], Theorem 3.1 includes several regular summability methods such as (1) the Kolmogorov SLLN [ $h(y) = 1$ ,  $g(y) = y$ ]; (2) the classical MZ SLLN [ $h(y) = 1$ ,  $g(y) = y^{1/\alpha}$ ,  $1 \leq \alpha \leq 2$ ].

#### 4. Spitzer Type Law of Large Numbers

Since the definition of complete convergence was introduced by Hsu and Robbins, there have been many authors who devote themselves to the study of the complete convergence for sums of independent and dependent RVs and obtain a series of elegant results, see [4, 8] and reference therein.

We say that the Hsu-Robbins [9] law of large numbers (LLN) is valid if, for all  $\varepsilon > 0$ ,

$$\sum_{n=1}^{\infty} P\{|S_n| \geq n\varepsilon\} < \infty, \quad (4.1)$$

and the Spitzer [10] LLN is valid if, for all  $\varepsilon > 0$ ,

$$\sum_{n=1}^{\infty} \frac{1}{n} P\{|S_n| \geq n\varepsilon\} < \infty. \quad (4.2)$$

**Theorem 4.1.** *Let  $\phi(\cdot)$  be as in Theorem 1.1, and let  $\{X, X_n, n \in \mathbb{N}\}$  be a sequence of NA random variables with  $EX_n = 0$ . Assume that  $\{X_n, n \in \mathbb{N}\} < X$ . If  $E[\phi^{-1}(|X|)] < \infty$ , then for all  $\varepsilon > 0$ ,*

$$\sum_{n=1}^{\infty} \frac{1}{n} P\{|S_n| \geq \varepsilon\phi(n)\} < \infty. \quad (4.3)$$

Conversely, let  $\{X, X_n, n \in \mathbb{N}\}$  be a sequence of identically distributed NA random variables, if (4.3) is true, then  $E[\phi^{-1}(|X|)] < \infty$  and  $EX = 0$ .

*Proof.* For  $1 \leq k \leq n$ , let  $U_k^{(n)} = X_k \mathbf{1}_{[|X_k| \leq \phi(n)]}$ ,  $V_k^{(n)} = X_k - U_k^{(n)}$ , then for every  $n$

$$|S_n| = \left| \sum_{k=1}^n (U_k^{(n)} + V_k^{(n)}) \right| \leq \left| \sum_{k=1}^n (U_k^{(n)} - EU_k^{(n)}) \right| + \left| \sum_{k=1}^n EU_k^{(n)} \right| + \left| \sum_{k=1}^n V_k^{(n)} \right|. \quad (4.4)$$

Note that

$$\begin{aligned} \{|S_n| \geq \varepsilon \phi(n)\} &\subset \left\{ \left| \sum_{k=1}^n (U_k^{(n)} - EU_k^{(n)}) \right| > \frac{\varepsilon \phi(n)}{2} \right\} \cup \left\{ \left| \sum_{k=1}^n V_k^{(n)} \right| > \frac{\varepsilon \phi(n)}{2} \right\} \\ &\cup \left\{ \frac{1}{\phi(n)} \max_{1 \leq i \leq n} \left| \sum_{k=1}^n EU_k^{(n)} \right| \rightarrow 0 \right\}. \end{aligned} \quad (4.5)$$

For the first term on the RHS of (4.5), by Markov inequality and Lemma 2.2 and (3.6), we have

$$\begin{aligned} &\sum_{n=1}^{\infty} \frac{1}{n} P \left\{ \left| \sum_{k=1}^n (U_k^{(n)} - EU_k^{(n)}) \right| > \frac{\varepsilon \phi(n)}{2} \right\} \\ &\leq \sum_{n=1}^{\infty} \frac{C}{n \phi^2(n)} \sum_{k=1}^n E (U_k^{(n)})^2 \quad (\text{Since } U_k^{(n)}, 1 \leq k \leq n \text{ is also NA}) \\ &= C \sum_{n=1}^{\infty} \frac{1}{n \phi^2(n)} \sum_{k=1}^n EX_k^2 \mathbf{1}_{[|X_k| \leq \phi(n)]} \\ &\leq C \sum_{n=1}^{\infty} \frac{1}{n} \sum_{k=1}^n \left[ E \mathbf{1}_{[|X| \geq \phi(n)]} + \frac{EX^2 \mathbf{1}_{[|X| \leq \phi(n)]}}{\phi^2(n)} \right] \\ &= C \sum_{n=1}^{\infty} \left[ E \mathbf{1}_{[|X| \geq \phi(n)]} + \frac{EX^2 \mathbf{1}_{[|X| \leq \phi(n)]}}{\phi^2(n)} \right] < \infty. \end{aligned} \quad (4.6)$$

For the second term on the RHS of (4.5), since

$$\left\{ \left| \sum_{k=1}^n V_k^{(n)} \right| > \frac{\varepsilon \phi(n)}{2} \right\} \subset \bigcup_{k=1}^n \{|X_k| > \phi(n)\}, \quad (4.7)$$



hence,

$$\begin{aligned}
 & \sum_{n=1}^{\infty} \frac{1}{n} P \left\{ \left| \sum_{k=1}^n V_k^{(n)} \right| > \frac{\varepsilon \phi(n)}{2} \right\} \\
 & \leq \sum_{n=1}^{\infty} \frac{1}{n} \sum_{k=1}^n P \{ |X_k| > \phi(n) \} \\
 & \leq \sum_{n=1}^{\infty} \frac{C}{n} \sum_{k=1}^n P \{ |X| > \phi(n) \} \\
 & = C \sum_{n=1}^{\infty} P \{ |X| > \phi(n) \} < \infty.
 \end{aligned} \tag{4.8}$$

For the third term on the RHS of (4.5), we have, by (3.5),

$$\begin{aligned}
 \frac{1}{\phi(n)} \sum_{k=1}^n |EU_k^{(n)}| &= \frac{1}{\phi(n)} \sum_{k=1}^n |EV_k^{(n)}| \quad (\text{By } EX_k = 0) \\
 &= \frac{\sum_{k=1}^n |EX_k \mathbf{1}_{[|X_k| > \phi(n)]}|}{\phi(n)} \\
 &\leq \frac{\sum_{k=1}^n E|X_k| \mathbf{1}_{[|X_k| > \phi(n)]}}{\phi(n)} \\
 &\leq CE \left[ \phi^{-1}(|X|) \right] < \infty.
 \end{aligned} \tag{4.9}$$

Therefore, (4.3) follows.

Conversely, since  $\sum_{n=1}^{\infty} (1/n) P \{ |S_n| \geq \varepsilon \phi(n) \} < \infty$  imply that  $S_n/\phi(n) = o(1)$  a.s., hence from Theorem 3.1, we have  $E[\phi^{-1}(|X|)] < \infty$ . These complete the proof of Theorem 4.1.  $\square$

**Corollary 4.2.** *Under the assumptions of Theorem 4.1, one has*

$$\sum_{n=1}^{\infty} \frac{1}{n} P \left\{ \max_{1 \leq k \leq n} |S_k| \geq \varepsilon \phi(n) \right\} < \infty. \tag{4.10}$$

*Proof.* Denote  $\bar{S}_k^{(n)} = \sum_{i=1}^k U_i^{(n)}$ , and noticing that

$$\left\{ \max_{1 \leq k \leq n} |S_k| \geq \varepsilon \phi(n) \right\} \subset \left\{ \max_{1 \leq k \leq n} \left| \bar{S}_k^{(n)} \right| \geq \varepsilon \phi(n) \right\} \cup \left\{ \bigcup_{k=1}^n (|X_k| > \phi(n)) \right\}, \tag{4.11}$$

hence, similarly to the proof of Theorem 4.1, we obtain (4.10).  $\square$

Analogously, we can prove the following corollaries, and omit the details.

**Corollary 4.3.** Let  $\{X, X_n, n \in \mathbb{N}\}$  be a sequence of  $\varphi$ -mixing random variables with  $EX_n = 0$ . Assume that  $\{X_n, n \in \mathbb{N}\} < X$ . If  $E[\phi^{-1}(|X|)] < \infty$ , and

$$\sum_{r=1}^{\infty} \varphi^{1/2}(r) < \infty, \quad (4.12)$$

then (4.3) holds.

**Corollary 4.4.** Let  $\{X, X_n, n \in \mathbb{N}\}$  be a sequence of  $m$ -dependent random variables with  $EX_n = 0$ . Assume that  $\{X_n, n \in \mathbb{N}\} < X$ . If  $E[\phi^{-1}(|X|)] < \infty$ , then (4.3) holds.

**Corollary 4.5.** Let  $\{X, X_n, n \in \mathbb{N}\}$  be a sequence of PA random variables with  $EX_n = 0$ . Assume that  $\{X_n, n \in \mathbb{N}\} < X$ . If  $E[\phi^{-1}(|X|)] < \infty$ , and

$$\sum_{n=1}^{\infty} \mu \frac{1}{2} (2^n) < \infty, \quad (4.13)$$

then (4.3) holds.

## 5. Hsu-Robbins Type Law of Large Numbers

**Theorem 5.1.** Let  $\phi(\cdot)$ , be define as in Theorem 1.1, but the following condition (3) is replaced by

(3') There exist constants  $a$  such that  $\phi^p(s) \int_s^{\infty} (x^{p/2} / \phi^p(x)) dx \leq as^{1+p/2}$ ,  $s > d$ ,  $p > 2$ , and let  $\{X, X_n, n \in \mathbb{N}\}$  be a sequence of NA random variables with  $EX_n = 0$ . Assume that  $\{X_n, n \in \mathbb{N}\} < X$ . If  $E[\phi^{-1}(|X|)]^{1+p/2} < \infty$ , then for all  $\varepsilon > 0$ ,

$$\sum_{n=1}^{\infty} P\{|S_n| \geq \varepsilon \phi(n)\} < \infty. \quad (5.1)$$

*Proof.* From the previous section, we know that to prove Theorem 5.1, we need only to prove the convergence of the following three series.

First, note that  $E[\phi^{-1}(|X|)]^{1+p/2} < \infty \Rightarrow E[\phi^{-1}(|X|)]^2 < \infty \Rightarrow nP(|X| > \phi(n)) \rightarrow 0$ , we have

$$\begin{aligned} \frac{1}{\phi(n)} \sum_{k=1}^n |EU_k^{(n)}| &= \frac{1}{\phi(n)} \sum_{k=1}^n |EX_k \mathbf{1}_{[|X_k| \leq \phi(n)]}| \\ &= \frac{1}{\phi(n)} \sum_{k=1}^n |EX_k \mathbf{1}_{[|X_k| > \phi(n)]}| \quad (\text{Since } EX_k = 0) \\ &\leq \frac{1}{\phi(n)} \sum_{k=1}^n E|X_k| \mathbf{1}_{[|X_k| > \phi(n)]} \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{\phi(n)} \sum_{k=1}^n \int_0^\infty P\{|X_k| \mathbf{1}_{\{|X_k| > \phi(n)\}} > t\} dt \\
&\leq \frac{Cn}{\phi(n)} \cdot \phi(n) P\{|X| > \phi(n)\} + \frac{Cn}{\phi(n)} \int_n^\infty P(\phi^{-1}(|X|) > t) dt \\
&\leq Cn P\{|X| > \phi(n)\} + \frac{Cn}{\phi(n)} \int_n^\infty \frac{E[\phi^{-1}(|X|)]^2}{t^2} dt \\
&= Cn P\{|X| > \phi(n)\} + \frac{Cn E[\phi^{-1}(|X|)]^2}{\phi(n)} \cdot \frac{1}{n} \\
&= Cn P\{|X| > \phi(n)\} + \frac{CE[\phi^{-1}(|X|)]^2}{\phi(n)} \rightarrow 0.
\end{aligned} \tag{5.2}$$

Hence,  $(1/\phi(n)) \max_j \sum_{k=1}^j E|U_k^{(n)}| \rightarrow 0$  as  $n \rightarrow \infty$ . Next, by (4.4), we have

$$\begin{aligned}
\sum_{n=1}^\infty P\left\{\left|\sum_{k=1}^n V_k^{(n)}\right| > \frac{\varepsilon \phi(n)}{2}\right\} &\leq \sum_{n=1}^\infty \sum_{k=1}^n P\{|X_k| > \phi(n)\} \leq \sum_{n=1}^\infty C \sum_{k=1}^n P\{|X| > \phi(n)\} \\
&\leq C \sum_{n=1}^\infty n P\{|X| > \phi(n)\} < CE[\phi^{-1}(|X|)]^2 < \infty.
\end{aligned} \tag{5.3}$$

Last, from the definition of  $U_k^{(n)}$  and the NA's property, we know that  $\{U_k^{(n)}, 1 \leq k \leq n, n \geq 1\}$  remains a sequence of NA RVs. By applying Lemma 2.2 and  $C_r$  inequality, we have

$$\begin{aligned}
\sum_{n=1}^\infty P\left\{\left|\sum_{k=1}^n (U_k^{(n)} - EU_k^{(n)})\right| > \frac{\varepsilon \phi(n)}{2}\right\} &\leq \sum_{n=1}^\infty \frac{C}{\phi^p(n)} E\left|\sum_{k=1}^n (U_k^{(n)} - EU_k^{(n)})\right|^p \\
&= \sum_{n=1}^\infty \frac{C}{\phi^p(n)} \sum_{k=1}^n E(U_k^{(n)} - EU_k^{(n)})^p \\
&\quad + \sum_{n=1}^\infty \frac{C}{\phi^p(n)} \left[\sum_{k=1}^n E(U_k^{(n)} - EU_k^{(n)})^2\right]^{p/2} \\
&:= I_1 + I_2.
\end{aligned} \tag{5.4}$$

It is easy to see that

$$\begin{aligned}
I_1 &\leq C \sum_{n=1}^\infty \frac{1}{\phi^p(n)} \sum_{k=1}^n E(U_k^{(n)})^p \\
&\leq C \sum_{n=1}^\infty \frac{1}{\phi^p(n)} \left[ \sum_{k=1}^n \phi^p(n) P(|X| > \phi(n)) + \sum_{k=1}^n E|X|^p \mathbf{1}_{\{|X| \leq \phi(n)\}} \right]
\end{aligned}$$

$$\begin{aligned}
&= C \sum_{n=1}^{\infty} n P\{|X| > \phi(n)\} + C \sum_{n=1}^{\infty} \frac{n}{\phi^p(n)} E|X|^p \mathbf{1}_{\{|X| \leq \phi(n)\}} \\
&\leq CE \left[ \phi^{-1}(|X|) \right]^2 + \frac{1}{2} k_0 (k_0 + 1) C + C \sum_{k=k_0+1}^{\infty} \frac{(k+1)^{p/2}}{\phi^p(k+1)} E|X|^p \mathbf{1}_{\{|X| \leq \phi(k)\}} \\
&\leq CE \left[ \phi^{-1}(|X|) \right]^{1+p/2} + Ck_0^2 + CE \left[ |X|^p \int_{\phi^{-1}(|X|)}^{\infty} \frac{x^{p/2}}{\phi^p(x)} dx \right] \\
&\leq CE \left[ \phi^{-1}(|X|) \right]^{1+p/2} + Ck_0^2 < \infty,
\end{aligned} \tag{5.5}$$

$$\begin{aligned}
I_2 &\leq C \sum_{n=1}^{\infty} \frac{1}{\phi^p(n)} \left[ \sum_{k=1}^n E \left( U_k^{(n)} \right)^2 \right]^{p/2} \\
&\leq C \sum_{n=1}^{\infty} \frac{1}{\phi^p(n)} \left[ n\phi^2(n) P\{|X| > \phi(n)\} + nEX^2 \mathbf{1}_{\{|X| \leq \phi(n)\}} \right]^{p/2} \\
&\leq C \sum_{n=1}^{\infty} \frac{1}{\phi^p(n)} \left[ \phi^p(n) n^{p/2} P\{|X| > \phi(n)\} \right] + C \sum_{n=1}^{\infty} \frac{n^{p/2}}{\phi^p(n)} \left[ EX^2 \mathbf{1}_{\{|X| \leq \phi(n)\}} \right]^{p/2} \\
&\leq C \sum_{n=1}^{\infty} n^{p/2} P\{|X| > \phi(n)\} + C \sum_{n=1}^{\infty} \frac{n^{p/2}}{\phi^p(n)} E|X|^p \mathbf{1}_{\{|X| \leq \phi(n)\}} \\
&\leq CE \left[ \phi^{-1}(|X|) \right]^{1+p/2} < \infty.
\end{aligned} \tag{5.6}$$

These complete the proof of Theorem 5.1.  $\square$

## Acknowledgments

This work is supported by the National Nature Science Foundation of China (no. 11071104) and the Anhui high Education Research Grant (no. KJ2010A337). The author expresses his sincere gratitude to the referees and the editors for their hospitality.

## References

- [1] R. Jajte, "On the strong law of large numbers," *The Annals of Probability*, vol. 31, no. 1, pp. 409–412, 2003.
- [2] J. D. Esary, F. Proschan, and D. W. Walkup, "Association of random variables, with applications," *Annals of Mathematical Statistics*, vol. 38, pp. 1466–1474, 1967.
- [3] K. Joag-Dev and F. Proschan, "Negative association of random variables, with applications," *The Annals of Statistics*, vol. 11, no. 1, pp. 286–295, 1983.
- [4] C. Su and Y. B. Wang, "Strong convergence for identically distributed negatively associated sequences," *Chinese Journal of Applied Probability and Statistics*, vol. 14, no. 2, pp. 131–140, 1998 (Chinese).
- [5] J. Sunklodas, "On the law of large numbers for weakly dependent random variables," *Lietuvos Matematikos Rinkinys*, vol. 44, no. 3, pp. 359–371, 2004.

- [6] S. C. Yang, "Moment inequalities for partial sums of random variables," *Science in China Series A*, vol. 44, no. 3, pp. 218–223, 2003.
- [7] P. Matuła, "A note on the almost sure convergence of sums of negatively dependent random variables," *Statistics & Probability Letters*, vol. 15, no. 3, pp. 209–213, 1992.
- [8] J. Bingyi and H. Y. Liang, "Strong limit theorems for weighted sums of negatively associated random sequence," *Journal of Theoretical Probability*, vol. 21, no. 4, pp. 890–909, 2008.
- [9] P. L. Hsu and H. Robbins, "Complete convergence and the law of large numbers," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 33, pp. 25–31, 1947.
- [10] F. Spitzer, "A combinatorial lemma and its application to probability theory," *Transactions of the American Mathematical Society*, vol. 82, pp. 323–339, 1956.