

Journal of Inequalities in Pure and Applied Mathematics

AN ALGEBRAIC INEQUALITY

FENG QI

Department of Mathematics
Jiaozuo Institute of Technology
Jiaozuo City, Henan 454000
THE PEOPLE'S REPUBLIC OF CHINA
E-Mail: qifeng@jzit.edu.cn
URL: <http://rgmia.vu.edu.au/qi.html>

©2000 Victoria University
ISSN (electronic): 1443-5756
006-00



volume 2, issue 1, article 13,
2000.

*Received 5 April, 2000;
accepted 15 January 2001.*

Communicated by: K. B. Stolarsky

Abstract

Contents



Home Page

Go Back

Close

Quit

Abstract

In this short note, an algebraic inequality related to those of Alzer, Minc and Sathre is proved by using analytic arguments and Cauchy's mean-value theorem. An open problem is proposed.


2000 Mathematics Subject Classification: 26D15.

Key words: Algebraic Inequality, Cauchy's Mean-Value Theorem, Alzer's Inequality.

The author was supported in part by NSF of Henan Province (#004051800), SF for Pure Research of the Education Department of Henan Province (#1999110004), Doctor Fund of Jiaozuo Institute of Technology, and NNSF (#10001016) of China.

Contents

1	An Algebraic Inequality	3
	References	



An Algebraic Inequality

Feng Qi

Title Page

Contents



Go Back

Close

Quit

Page 2 of 8

1. An Algebraic Inequality

In this note, we prove the following algebraic inequality

Theorem 1.1. *Let $b > a > 0$ and $\delta > 0$ be real numbers. Then for any given positive $r \in \mathbb{R}$, we have*

$$(1.1) \quad \left(\frac{b + \delta - a}{b - a} \cdot \frac{b^{r+1} - a^{r+1}}{(b + \delta)^{r+1} - a^{r+1}} \right)^{1/r} > \frac{b}{b + \delta}.$$

The lower bound in (1.1) is best possible.

Proof. The inequality (1.1) is equivalent to

$$\frac{b^{r+1} - a^{r+1}}{b - a} \Big/ \frac{(b + \delta)^{r+1} - a^{r+1}}{b + \delta - a} > \left(\frac{b}{b + \delta} \right)^r,$$

that is,


$$(1.2) \quad \frac{b^{r+1} - a^{r+1}}{b^r(b - a)} > \frac{(b + \delta)^{r+1} - a^{r+1}}{(b + \delta)^r(b + \delta - a)}.$$

Therefore, it is sufficient to prove that the function $(s^{r+1} - a^{r+1})/s^r(s - a)$ is decreasing for $s > a$. By direct computation, we have

$$\left(\frac{s^{r+1} - a^{r+1}}{s^r(s - a)} \right)'_s = \frac{(r + 1)(s - a)s^{2r} - s^{r-1}(s^{r+1} - a^{r+1})[(r + 1)s - ra]}{[s^r(s - a)]^2}.$$

So, it suffices to prove

$$(1.3) \quad (r + 1)(s - a)s^{r+1} - [(r + 1)s - ra](s^{r+1} - a^{r+1}) \leq 0.$$



An Algebraic Inequality

Feng Qi

Title Page

Contents



Go Back

Close

Quit

Page 3 of 8

A straightforward calculation shows that the inequality (1.3) reduces to

$$(1.4) \quad \frac{s^r - a^r}{r(s - a)} > \frac{a^r}{s}.$$

From Cauchy's mean-value theorem, there exists a point $\xi \in (a, s)$ such that

$$\frac{s^r - a^r}{r(s - a)} = \xi^{r-1} = \frac{\xi^r}{\xi} > \frac{a^r}{\xi} > \frac{a^r}{s}.$$

Hence, the inequality (1.4) holds.

The L'Hospital rule yields

$$(1.5) \quad \lim_{r \rightarrow +\infty} \left(\frac{b + \delta - a}{b - a} \cdot \frac{b^{r+1} - a^{r+1}}{(b + \delta)^{r+1} - a^{r+1}} \right)^{1/r} = \frac{b}{b + \delta},$$

so the lower bound in (1.1) is best possible. The proof is complete. \square


Remark 1.1. *The inequality (1.1) can be rewritten as*

$$(1.6) \quad \frac{b}{b + \delta} < \left(\frac{1}{b - a} \int_a^b x^r dx \Big/ \frac{1}{b + \delta - a} \int_a^{b+\delta} x^r dx \right)^{1/r}.$$

It is easy to see that inequality (1.6) is indeed an integral analogue of the following inequality

$$(1.7) \quad \frac{n + k}{n + m + k} < \left(\frac{1}{n} \sum_{i=k+1}^{n+k} i^r \Big/ \frac{1}{n + m} \sum_{i=k+1}^{n+m+k} i^r \right)^{1/r},$$

where r is a given positive real number, n and m are natural numbers, and k is a nonnegative integer. The lower bound in (1.7) is best possible.



An Algebraic Inequality

Feng Qi

Title Page

Contents



Go Back

Close

Quit

Page 4 of 8

The inequality (1.7) was presented in [5] by the author using Cauchy's mean-value theorem and mathematical induction. It generalizes the inequality of Alzer in [1].

Using the same method as in [5], the author in [9] further generalized the inequality of Alzer and obtained that, if $a = (a_1, a_2, \dots)$ is a positive and increasing sequence satisfying

$$(1.8) \quad a_{k+1}^2 \geq a_k a_{k+2},$$

$$(1.9) \quad \frac{a_{k+1} - a_k}{a_{k+1}^2 - a_k a_{k+2}} \geq \max \left\{ \frac{k+1}{a_{k+1}}, \frac{k+2}{a_{k+2}} \right\}$$

for $k \in \mathbb{N}$, then we have


$$(1.10) \quad \frac{a_n}{a_{n+m}} < \left(\frac{\frac{1}{n} \sum_{i=1}^n a_i^r}{\frac{1}{n+m} \sum_{i=1}^{n+m} a_i^r} \right)^{1/r},$$

where n and m are natural numbers. The lower bound in (1.10) is best possible.

Recently, some new inequalities related to those of Alzer, Minc and Sathre were obtained by many mathematician. These inequalities involve ratios for the sum of powers of positive numbers (see [2, 12]) and for the geometric mean of natural numbers (see [4, 6, 7, 10, 11]). Many of them can be deduced from monotonicity and convexity considerations (see [8]). Moreover, inequality (1.1) has been generalised to an inequality for linear positive functionals in [3].

Here L'Hospital's rule yields

$$(1.11) \quad \lim_{r \rightarrow 0^+} \left(\frac{b + \delta - a}{b - a} \cdot \frac{b^{r+1} - a^{r+1}}{(b + \delta)^{r+1} - a^{r+1}} \right)^{1/r} = \frac{[b^b/a^a]^{1/(b-a)}}{[(b + \delta)^{b+\delta}/a^a]^{1/(b+\delta-a)}}.$$



An Algebraic Inequality

Feng Qi

Title Page

Contents



Go Back

Close

Quit

Page 5 of 8


Hence, we propose the following

Open Problem. Let $b > a > 0$ and $\delta > 0$ be real numbers. Then for any positive $r \in \mathbb{R}$, we have

$$(1.12) \quad \left(\frac{b + \delta - a}{b - a} \cdot \frac{b^{r+1} - a^{r+1}}{(b + \delta)^{r+1} - a^{r+1}} \right)^{1/r} < \frac{[b^b/a^a]^{1/(b-a)}}{[(b + \delta)^{b+\delta}/a^a]^{1/(b+\delta-a)}}.$$

The upper bound in (1.12) is best possible.

Remark 1.2. The inequalities in this paper are related to the study of monotonicity of the ratios and differences of mean values.



An Algebraic Inequality

Feng Qi

Title Page

Contents



Go Back


Close

Quit

Page 6 of 8

References

- [1] H. ALZER, On an inequality of H. Minc and L. Sathre, *J. Math. Anal. Appl.*, **179** (1993), 396–402.
- [2] S.S. DRAGOMIR AND J. VAN DER HOEK, Some new analytic inequalities and their applications in guessing theory, *J. Math. Anal. Appl.*, **225** (1998), 542–556.
- [3] B. GAVREA AND I. GAVREA, An inequality for linear positive functionals, *J. Inequal. Pure Appl. Math.*, **1**(1) (2000), Article 5. [ONLINE] http://jipam.vu.edu.au/v1n1/004_99.html.
- [4] J.-C. KUANG, Some extensions and refinements of Minc-Sathre inequality, *Math. Gaz.*, **83** (1999), 123–127.
- [5] F. QI, Generalization of H. Alzer's inequality, *J. Math. Anal. Appl.*, **240** (1999), 294–297.
- [6] F. QI, Generalizations of Alzer's and Kuang's inequality, *Tamkang Journal of Mathematics*, **31**(3) (2000), 223–227. Preprint available from the *RGMIA Research Report Collection*, **2**(6) (1999), Article 12. <http://rgmia.vu.edu.au/v2n6.html>.
- [7] F. QI, Inequalities and monotonicity of sequences involving $\sqrt[n]{(n+k)!/k!}$, *RGMIA Research Report Collection*, **2**(5) (1999), Article 8. [ONLINE] <http://rgmia.vu.edu.au/v2n5.html>.



An Algebraic Inequality

Feng Qi

Title Page

Contents




Go Back

Close

Quit

Page 7 of 8

- [8] F. QI, Monotonicity of sequences involving convex function and sequence, *RGMI*A Research Report Collection, **3**(2) (2000), Article 14. [ONLINE] <http://rgmia.vu.edu.au/v3n2.html>.
- [9] F. QI AND L. DEBNATH, On a new generalization of Alzer's inequality, *Int. J. Math. Math. Sci.*, **23**(12) (2000), 815–818.
- [10] F. QI AND B.-N. GUO, Some inequalities involving geometric mean of natural numbers and ratio of gamma functions, *RGMI*A Research Report Collection, **4**(1) (2001), Article 3. [ONLINE] <http://rgmia.vu.edu.au/v4n1.html>.
- [11] F. QI AND Q.-M. LUO, Generalization of H. Minc and J. Sathre's inequality, *Tamkang J. Math.*, **31**(2) (2000), 145–148. Preprint available from the *RGMI*A Research Report Collection, **2**(6) (1999), Article 14. <http://rgmia.vu.edu.au/v2n6.html>.
- [12] J. SÁNDOR, Comments on an inequality for the sum of powers of positive numbers, *RGMI*A Research Report Collection, **2**(2) (1999), 259–261. [ONLINE] <http://rgmia.vu.edu.au/v2n2.html>.



An Algebraic Inequality

Feng Qi

Title Page

Contents



Go Back

Close

Quit

Page 8 of 8