



## A PRIORI ESTIMATE FOR A SYSTEM OF DIFFERENTIAL OPERATORS

CHIKH BOUZAR

DÉPARTEMENT DE MATHÉMATIQUES.  
UNIVERSITÉ D'ORAN-ESSENIA.ALGÉRIE.  
bouzarchikh@hotmail.com

*Received 23 October, 2000; accepted 22 January, 2001.*

*Communicated by S. Saitoh*

---

ABSTRACT. We characterize in algebraic terms an inequality in Sobolev spaces for a system of differential operators with constant coefficients.

---

*Key words and phrases:* Differential operators, a priori estimate.

2000 *Mathematics Subject Classification.* 35B45.

### 1. INTRODUCTION

We are interested in the following inequality

$$(1.1) \quad \exists C > 0, \|R(D)u\| \leq C \sum_{j=1}^k \|P_j(D)u\|, \forall u \in C_0^\infty(\Omega),$$

where  $S = \{P_j(D); j = 1, \dots, k\}$ ,  $R(D)$  are linear differential operators of order  $\leq m$  with constant complex coefficients and  $C_0^\infty(\Omega)$  is the space of infinitely differentiable functions with compact supports in a bounded open set  $\Omega$  of the Euclidian space  $\mathbb{R}^n$ . By  $\|\cdot\|$  we denote the norm of the Hilbert space  $L^2(\Omega)$  of square integrable functions.

Each differential operator  $P_j(D)$  has a complete symbol  $P_j(\xi)$  such that

$$(1.2) \quad P_j(\xi) = p_j(\xi) + q_j(\xi) + r_j(\xi) + \dots,$$

where  $p_j(\xi)$ ,  $q_j(\xi)$  and  $r_j(\xi)$  are the homogeneous polynomial parts of  $P_j(\xi)$  in  $\xi \in \mathbb{R}^n$  of orders, respectively,  $m$ ,  $m - 1$  and  $m - 2$ .

It is well-known that the system  $S$  satisfies the inequality (1.1) for all differential operators  $R(D)$  of order  $\leq m$  if and only if it is elliptic, i.e.

$$(1.3) \quad \sum_{j=1}^k |p_j(\xi)| \neq 0, \forall \xi \in \mathbb{R}^n \setminus 0.$$

In this paper we give an necessary and sufficient algebraic condition on the system  $S$  such that it satisfies the inequality (1.1) for all differential operators  $R(D)$  of order  $\leq m - 1$ .

The estimate (1.1) has been used in our work [1], without proof, in the study of local estimates for certain classes of pseudodifferential operators.

## 2. THE RESULTS

To prove the main theorem we need some lemmas. The first one gives an algebraic characterization of the inequality (1.1) based on a well-known result of Hörmander [3].

Recall the Hörmander function

$$(2.1) \quad \tilde{P}_j(\xi) = \left( \sum_{\alpha} |P_j^{(\alpha)}(\xi)|^2 \right)^{\frac{1}{2}},$$

where  $P_j^{(\alpha)}(\xi) = \frac{\partial^{|\alpha|}}{\partial \xi_1^{\alpha_1} \dots \partial \xi_n^{\alpha_n}} P_j(\xi)$ , (see [3]).

**Lemma 2.1.** *The inequality (1.1) holds for every  $R(D)$  of order  $\leq m - 1$  if and only if*

$$(2.2) \quad \exists C > 0, \quad |\xi|^{m-1} \leq C \sum_{j=1}^k \tilde{P}_j(\xi), \quad \forall \xi \in \mathbb{R}^n.$$

*Proof.* The proof of this lemma follows essentially from the classical one in the case of  $k = 1$ , and it is based on Hörmander's inequality (see [3, p. 7]).  $\square$

The scalar product in the complex Euclidian space  $C^k$  of  $A = (a_1, \dots, a_k)$  and  $B = (b_1, \dots, b_k)$  is denoted as usually by  $A \cdot B = \sum_{i=1}^k a_i \bar{b}_i$ , and the norm of  $C^k$  by  $|\cdot|$ .

Let, by definition,

$$(2.3) \quad |A \wedge B|^2 = \sum_{i < j}^k |a_i b_j - b_i a_j|^2.$$

The next lemma is a consequence of the classical Lagrange's identity (see [2]).

**Lemma 2.2.** *Let  $A = (a_1, \dots, a_k) \in C^k$  and  $B = (b_1, \dots, b_k) \in C^k$ , then*

$$(2.4) \quad |At + B|^2 = \left( |A|t + \frac{Re(A \cdot B)}{|A|} \right)^2 + \frac{|Im(A \cdot B)|^2 + |A \wedge B|^2}{|A|^2}, \quad \forall t \in \mathbb{R}.$$

*Proof.* We have

$$\begin{aligned} |At + B|^2 &= (|A|t)^2 + 2t Re(A \cdot B) + |B|^2 \\ &= \left( |A|t + \frac{Re(A \cdot B)}{|A|} \right)^2 + |B|^2 - \left( \frac{Re(A \cdot B)}{|A|} \right)^2. \end{aligned}$$

We obtain (2.4) from the next classical Lagrange's identity

$$|A|^2 |B|^2 = |Re(A \cdot B)|^2 + |Im(A \cdot B)|^2 + |A \wedge B|^2.$$

$\square$

For  $\xi \in \mathbb{R}^n$  we define the vector functions

$$(2.5) \quad A(\xi) = (p_1(\xi), \dots, p_k(\xi)) \text{ and } B(\xi) = (q_1(\xi), \dots, q_k(\xi)).$$

Let

$$(2.6) \quad \Xi = \left\{ \omega \in S^{n-1} : |A(\omega)|^2 = \sum_{j=1}^k |p_j(\omega)|^2 \neq 0 \right\},$$

where  $S^{n-1}$  is the unit sphere of  $\mathbb{R}^n$ , and

$$(2.7) \quad F(t, \xi) = |\text{grad}A(\xi)|^2 + |A(\xi)t + B(\xi)|^2,$$

where  $|\text{grad}A(\xi)|^2 = \sum_{j=1}^k |\text{grad}p_j(\xi)|^2$ .

**Lemma 2.3.** *The inequality (2.2) holds if and only if there exist no sequences of real numbers  $t_j \rightarrow +\infty$  and  $\omega_j \in S^{n-1}$  such that*

$$(2.8) \quad F(t_j, \omega_j) \rightarrow 0.$$

*Proof.* Let  $t_j$  be a sequence of real numbers and  $\omega_j$  a sequence of  $S^{n-1}$ , using the homogeneity of the functions  $p, q$  and  $r$ , then (2.2) is equivalent to

$$\frac{|t_j \omega_j|^{2(m-1)}}{\sum_{l=1}^k \tilde{P}_l(t_j \omega_j)^2} = \frac{1}{F(t_j, \omega_j) + 2 \sum_{l=1}^k \text{Re}(p_l(\omega_j) \cdot \bar{r}_l(\omega_j)) + \chi(\omega_j) \cdot O(\frac{1}{t_j})} \leq C,$$

where  $\chi$  is a bounded function. Hence it is easy to see Lemma 2.3. □

If  $\omega \in \Xi$  we define the function  $G$  by

$$G(\omega) = |\text{grad}A(\omega)|^2 + \frac{|\text{Im}(A(\omega) \cdot B(\omega))|^2 + |A(\omega) \wedge B(\omega)|^2}{|A(\omega)|^2}.$$

**Theorem 2.4.** *The estimate (1.1) holds if and only if*

$$(2.9) \quad \exists C > 0, G(\omega) \geq C, \forall \omega \in \Xi$$

*Proof.* All positive constants are denoted by  $C$ . If (2.9) holds then from (2.4) and (2.7) we have

$$(2.10) \quad F(t, \omega) = \left( |A(\omega)|t + \frac{\text{Re}(A(\omega) \cdot B(\omega))}{|A(\omega)|} \right)^2 + G(\omega) \geq C, \forall \omega \in \Xi, \forall t \geq 0.$$

The vector function  $A$  is analytic and the set  $\Xi$  is dense in  $S^{n-1}$ , therefore by continuity we obtain

$$(2.11) \quad F(t, \omega) \geq C, \forall t \geq 0, \forall \omega \in S^{n-1}.$$

For  $\xi \in \mathbb{R}^n$ , set  $\omega = \frac{\xi}{|\xi|}$  and  $t = |\xi|$  in (2.11), as the vector functions  $A$  and  $B$  are homogeneous, we obtain

$$|A(\xi) + B(\xi)|^2 + |\text{grad}A(\xi)|^2 \geq C |\xi|^{2(m-1)}, \forall \xi \in \mathbb{R}^n,$$

and then, for  $|\xi| \geq C$ , we have

$$(2.12) \quad \sum_{j=1}^k (|P_j(\xi)|^2 + |\text{grad}P_j(\xi)|^2) + O\left((1 + |\xi|^2)^{m-2}\right) \geq C |\xi|^{2(m-1)}.$$

From the last inequality we easily get (2.2) of Lemma 2.1.

Suppose that (2.9) does not hold, then there exists a sequence  $\omega_j \in \Xi$  such that  $G(\omega_j) \rightarrow 0$ , i.e.

$$(2.13) \quad |\text{grad}A(\omega_j)|^2 \rightarrow 0,$$

and

$$(2.14) \quad \frac{|Im(A(\omega_j).B(\omega_j))|^2 + |A(\omega_j) \wedge B(\omega_j)|^2}{|A(\omega_j)|^2} \rightarrow 0.$$

As  $S^{n-1}$  is compact we can suppose that  $\omega_j \rightarrow \omega_0 \in S^{n-1}$ . Hence, from (2.14) and (2.4) with  $t = 0$ , we obtain

$$(2.15) \quad \frac{Re(A(\omega_j).B(\omega_j))}{|A(\omega_j)|} \rightarrow \pm |B(\omega_0)|.$$

From (2.13), due to Euler's identity for homogeneous functions,

$$(2.16) \quad A(\omega_0) = \vec{0}.$$

Now if  $B(\omega_0) = 0$  then  $F(t, \omega_0) \equiv 0$ , which contradicts (2.8).

Let  $B(\omega_0) \neq 0$ , and suppose that

$$(2.17) \quad \frac{Re(A(\omega_j).B(\omega_j))}{|A(\omega_j)|} \rightarrow -|B(\omega_0)|,$$

then setting  $t_j = \frac{|B(\omega_j)|}{|A(\omega_j)|}$  in (2.10), it is clear that  $t_j \rightarrow +\infty$ , so, with  $G(\omega_j) \rightarrow 0$ ,  $F(t_j, \omega_j)$  will converge to 0, which contradicts (2.8).

If

$$\frac{Re(A(\omega_j).B(\omega_j))}{|A(\omega_j)|} \rightarrow +|B(\omega_0)|,$$

then changing  $\omega_j$  to  $-\omega_j$  and using the homogeneity of the functions  $A$  and  $B$ , we obtain the same conclusion.  $\square$

## REFERENCES

- [1] C. BOUZAR, Local estimates for pseudodifferential operators, *Doklady Nats. Akad. Nauk Belarusi*, **44**(4) (2000), 18–20. (in Russian)
- [2] G. HARDY, J. LITTLEWOOD AND G. POLYA, *Inequalities*, Cambridge Univ. Press, 2nd Ed., 1967.
- [3] L. HÖRMANDER, *The Analysis of Partial Differential Operators*, T.II, Springer-Verlag. 1983.