

Journal of Inequalities in Pure and Applied Mathematics

ON A NEW GENERALISATION OF OSTROWSKI'S INEQUALITY

B.G. PACHPATTE

57 Shri Niketan Colony
Near Abhinay Talkies
Aurangabad 431 001
(Maharashtra) India.

EMail: bgpachpatte@hotmail.com

©2000 Victoria University
ISSN (electronic): 1443-5756
088-02



volume 5, issue 2, article 36,
2004.

*Received 21 July, 2002;
accepted 08 July, 2003.*

Communicated by: P. Cerone

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

The aim of this note is to establish a new integral inequality involving two functions and their derivatives. Our result for particular cases yields the well known Ostrowski inequality and its generalization given by Milovanović and Pečarić.

2000 Mathematics Subject Classification: Primary 26D15; Secondary 41A55.

Key words: Ostrowski's inequality, Functions and their derivatives, Differentiable functions, Empty sum, Taylor's formula.

Contents

1	Introduction	3
2	Main Result	5
	References	



On a New Generalisation of Ostrowski's Inequality

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 2 of 10

1. Introduction

In 1938, Ostrowski [4] (see [3, p. 468]) proved the following integral inequality.

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous on $[a, b]$ and differentiable on (a, b) , and whose derivative $f' : (a, b) \rightarrow \mathbb{R}$ is bounded on (a, b) , i.e., $\|f'\|_\infty = \sup_{t \in (a, b)} |f'(t)| < \infty$. Then

$$(1.1) \quad \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty,$$


for all $x \in [a, b]$.

In 1976, Milovanović and Pečarić [2] (see [3, p. 469]) proved the following generalization of Ostrowski's inequality.

Let $f : [a, b] \rightarrow \mathbb{R}$ be an n -times differentiable function, $n \geq 1$ and such that $\|f^{(n)}\|_\infty = \sup_{t \in (a, b)} |f^{(n)}(t)| < \infty$. Then

$$(1.2) \quad \left| \frac{1}{n} \left(f(x) + \sum_{k=1}^{n-1} \frac{n-k}{k!} \times \frac{f^{(k-1)}(a)(x-a)^k - f^{(k-1)}(b)(x-b)^k}{b-a} - \frac{1}{b-a} \int_a^b f(t) dt \right) \right| \leq \frac{\|f^{(n)}\|_\infty}{n(n+1)!} \cdot \frac{(x-a)^{n+1} - (b-x)^{n+1}}{b-a},$$

for all $x \in [a, b]$.



On a New Generalisation of Ostrowski's Inequality

B.G. Pachpatte

Title Page

Contents




Go Back

Close

Quit

Page 3 of 10

The main purpose of this note is to establish a new generalization of Ostrowski's integral inequality involving two functions and their derivatives by using fairly elementary analysis. Our result in special cases yield the inequalities given in (1.1) and (1.2). For some other extensions, generalizations and similar results, see [3] and the references cited therein.



On a New Generalisation of Ostrowski's Inequality

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 4 of 10

2. Main Result

In what follows, \mathbb{R} and n denote the set of real numbers and a positive integer respectively. The n th derivative of a function $f : [a, b] \rightarrow \mathbb{R}$ is denoted by $f^{(n)}(t)$, $t \in [a, b]$. For n -times differentiable functions $f, g : [a, b] \rightarrow \mathbb{R}$, we use the following notation to simplify the details of presentation:

$$F_k(x) = \frac{n-k}{k!} \cdot \frac{f^{(k-1)}(a)(x-a)^k - f^{(k-1)}(b)(x-b)^k}{b-a},$$

$$G_k(x) = \frac{n-k}{k!} \cdot \frac{g^{(k-1)}(a)(x-a)^k - g^{(k-1)}(b)(x-b)^k}{b-a},$$

$$I_k = \frac{1}{k!} \int_a^b f^{(k)}(y)(x-y)^k dy, \quad I_0 = \int_a^b f(y) dy,$$


$$J_k = \frac{1}{k!} \int_a^b g^{(k)}(y)(x-y)^k dy, \quad J_0 = \int_a^b g(y) dy,$$

for $1 \leq k \leq n-1$. We use the usual convention that an empty sum is taken to be zero.

Our main result is given in the following theorem.

Theorem 2.1. *Let $f, g : [a, b] \rightarrow \mathbb{R}$ be continuous functions on $[a, b]$ and n -times differentiable on (a, b) , and whose derivatives $f^{(n)}, g^{(n)} : (a, b) \rightarrow \mathbb{R}$ are bounded on (a, b) , i.e.,*

$$\|f^{(n)}\|_{\infty} = \sup_{t \in (a,b)} |f^{(n)}(t)| < \infty, \quad \|g^{(n)}\|_{\infty} = \sup_{t \in (a,b)} |g^{(n)}(t)| < \infty.$$



On a New Generalisation of
Ostrowski's Inequality

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 5 of 10

Then

$$(2.1) \quad \left| f(x)g(x) - \frac{1}{2(b-a)} [g(x)I_0 + f(x)J_0] - \frac{1}{2(b-a)} \left[g(x) \sum_{k=1}^{n-1} I_k + f(x) \sum_{k=1}^{n-1} J_k \right] \right| \leq \frac{1}{2(n+1)!} [|g(x)| \|f^{(n)}\|_\infty + |f(x)| \|g^{(n)}\|_\infty] \left[\frac{(x-a)^{n+1} + (b-x)^{n+1}}{b-a} \right],$$

for all $x \in [a, b]$.


Proof. Let $x \in [a, b]$, $y \in (a, b)$. With the stipulation on f, g and using Taylor's formula with the Lagrange form of the remainder (see [2, p. 156]) we have

$$(2.2) \quad f(x) = f(y) + \sum_{k=1}^{n-1} f^{(k)}(y) (x-y)^k + \frac{1}{n!} f^{(n)}(\xi) (x-y)^n,$$

$$(2.3) \quad g(x) = g(y) + \sum_{k=1}^{n-1} g^{(k)}(y) (x-y)^k + \frac{1}{n!} g^{(n)}(\sigma) (x-y)^n,$$

where $\xi = y + \alpha(x-y)$ ($0 < \alpha < 1$) and $\sigma = y + \beta(x-y)$ ($0 < \beta < 1$). From the definitions of I_k, J_k and integration by parts (see [2]) we have the relations

$$(2.4) \quad I_0 + \sum_{k=1}^{n-1} I_k = nI_0 - (b-a) \sum_{k=1}^{n-1} F_k(x),$$



On a New Generalisation of Ostrowski's Inequality

B.G. Pachpatte

Title Page

Contents




Go Back

Close

Quit

Page 6 of 10



$$(2.5) \quad J_0 + \sum_{k=1}^{n-1} J_k = nJ_0 - (b-a) \sum_{k=1}^{n-1} G_k(x).$$

Multiplying (2.2) and (2.3) by $g(x)$ and $f(x)$ respectively, adding the resulting identities and rewriting, we have

$$(2.6) \quad f(x)g(x) = \frac{1}{2}g(x)f(y) + \frac{1}{2}f(x)g(y) \\ + \frac{1}{2}g(x) \sum_{k=1}^{n-1} \frac{1}{k!} f^{(k)}(y) (x-y)^k + \frac{1}{2}f(x) \sum_{k=1}^{n-1} \frac{1}{k!} g^{(k)}(y) (x-y)^k \\ + \frac{1}{2} \cdot \frac{1}{n!} g(x) f^{(n)}(\xi) (x-y)^n + \frac{1}{2} \cdot \frac{1}{n!} f(x) g^{(n)}(\sigma) (x-y)^n.$$

Integrating (2.6) with respect to y on (a, b) and rewriting, we obtain

$$(2.7) \quad f(x)g(x) = \frac{1}{2(b-a)} [g(x)I_0 + f(x)J_0] \\ + \frac{1}{2(b-a)} \left[g(x) \sum_{k=1}^{n-1} I_k + f(x) \sum_{k=1}^{n-1} J_k \right] \\ + \frac{1}{2(b-a)} \cdot \frac{1}{n!} \left[g(x) \int_a^b f^{(n)}(\xi) (x-y)^n dy \right. \\ \left. + f(x) \int_a^b g^{(n)}(\sigma) (x-y)^n dy \right].$$

On a New Generalisation of
Ostrowski's Inequality

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit


Page 7 of 10

From (2.7) and using the properties of modulus, we have

$$\begin{aligned}
 & \left| f(x)g(x) - \frac{1}{2(b-a)} [g(x)I_0 + f(x)J_0] \right. \\
 & \quad \left. - \frac{1}{2(b-a)} \left[g(x) \sum_{k=1}^{n-1} I_k + f(x) \sum_{k=1}^{n-1} J_k \right] \right| \\
 & \leq \frac{1}{2(b-a)} \cdot \frac{1}{n!} \left[|g(x)| \int_a^b |f^{(n)}(\xi)| |x-y|^n dy \right. \\
 & \quad \left. + |f(x)| \int_a^b |g^{(n)}(\sigma)| |x-y|^n dy \right] \\
 & \leq \frac{1}{2(b-a)} \cdot \frac{1}{n!} [|g(x)| \|f^{(n)}\|_\infty + |f(x)| \|g^{(n)}\|_\infty] \int_a^b |x-y|^n dy \\
 & = \frac{1}{2(b-a)} \cdot \frac{1}{(n+1)!} [|g(x)| \|f^{(n)}\|_\infty + |f(x)| \|g^{(n)}\|_\infty] \\
 & \quad \times \left[\frac{(x-a)^{n+1} + (b-x)^{n+1}}{b-a} \right],
 \end{aligned}$$

which is the required inequality in (2.1). The proof is complete. \square

Corollary 2.2. Let $f, g : [a, b] \rightarrow \mathbb{R}$ be continuous functions on $[a, b]$ and differentiable on (a, b) and whose derivatives $f', g' : (a, b) \rightarrow \mathbb{R}$ are bounded on (a, b) , i.e., $\|f'\|_\infty = \sup_{t \in (a, b)} |f'(t)| < \infty$, $\|g'\|_\infty = \sup_{t \in (a, b)} |g'(t)| < \infty$.



On a New Generalisation of
Ostrowski's Inequality

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 8 of 10

Then


$$(2.8) \quad \left| f(x) g(x) - \frac{1}{2(b-a)} [g(x) I_0 + f(x) J_0] \right| \\ \leq \frac{1}{2} [\|g(x)\| \|f'\|_\infty + |f(x)| \|g'\|_\infty] \left[\frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a),$$

for all $x \in [a, b]$.

We note that in the special cases, if we take

- (i) $g(x) = 1$ and hence $g^{(n)}(x) = 0$ in (2.1) and
- (ii) $g(x) = 1$ and hence $g'(x) = 0$ in (2.8),

we get the inequalities (1.2) and (1.1) respectively. Further, we note that, here we have used Taylor's formula with the Lagrange form of remainder to prove our result. Instead of this, one can use as in [1] the Taylor formula with integral remainder to establish a variant of Theorem A in [1] in the framework of our main result given above. Here we omit the details.



**On a New Generalisation of
Ostrowski's Inequality**

B.G. Pachpatte

Title Page

Contents



Go Back


Close

Quit

Page 9 of 10

References

- [1] A.M. FINK, Bounds on the deviation of a function from its averages, *Czechoslovak Math. J.*, **42** (1992), 289–310.
- [2] G.V. MILOVANOVIĆ AND J.E. PEČARIĆ, On generalization of the inequality of A. Ostrowski and some related applications, *Univ. Beograd Publ. Elek. Fak. Ser. Mat. Fiz.*, No. 544 – No. 576 (1976), 155–158.
- [3] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Inequalities for Functions and Their Integrals and Derivatives*, Kluwer Academic Publishers, 1994.
- [4] A. OSTROWSKI, Über die Absolutabweichung einer differentiebaren Funktion van ihrem Integralmittelwert, *Comment Math. Helv.*, **10** (1938), 226–227.



On a New Generalisation of
Ostrowski's Inequality

B.G. Pachpatte

Title Page

Contents



Go Back

Close

Quit

Page 10 of 10