

FRANK HANSEN

Institute of Economics
Copenhagen University
Studiestraede 6
1455 Copenhagen K, Denmark.

*E*Mail: Frank.Hansen@econ.ku.dk



volume 6, issue 5, article 133,
2005.

*Received 23 March, 2005;
accepted 23 September, 2005.*

Communicated by: S.S. Dragomir

Abstract

Contents




Home Page

Go Back

Close

Quit



Abstract

We study conditional expectations generated by an abelian C^* -subalgebra in the centralizer of a positive functional. We formulate and prove Jensen's inequality for functions of several variables with respect to this type of conditional expectations, and we obtain as a corollary Jensen's inequality for expectation values.

2000 Mathematics Subject Classification: 47A63, 26A51.

Key words: Trace function, Jensen's inequality, Conditional expectation.

This paper is based on the talk given by the author within the "International Conference of Mathematical Inequalities and their Applications, I", December 06-08, 2004, Victoria University, Melbourne, Australia [<http://rgmia.vu.edu.au/conference>]

Contents

1	Preliminaries	3
2	Conditional Expectations	4
3	Jensen's inequality	6
References		

Jensen's Inequality for Conditional Expectations

Frank Hansen

Title Page

Contents



Go Back

Close

Quit

Page 2 of 12

1. Preliminaries

An n -tuple $\underline{x} = (x_1, \dots, x_n)$ of elements in a C^* -algebra \mathcal{A} is said to be abelian if the elements x_1, \dots, x_n are mutually commuting. We say that an abelian n -tuple \underline{x} of self-adjoint elements is in the domain of a real continuous function f of n variables defined on a cube of real intervals $\underline{I} = I_1 \times \dots \times I_n$ if the spectrum $\sigma(x_i)$ of x_i is contained in I_i for each $i = 1, \dots, n$. In this situation $f(\underline{x})$ is naturally defined as an element in \mathcal{A} in the following way. We may assume that \mathcal{A} is realized as operators on a Hilbert space and let

$$x_i = \int \lambda dE_i(\lambda) \quad i = 1, \dots, n$$


denote the spectral resolutions of the operators x_1, \dots, x_n . Since the n -tuple $\underline{x} = (x_1, \dots, x_n)$ is abelian, the spectral measures E_1, \dots, E_n are mutually commuting. We may thus set

$$E(S_1 \times \dots \times S_n) = E_1(S_1) \cdots E_n(S_n)$$

for Borel sets S_1, \dots, S_n in \mathbb{R} and extend E to a spectral measure on \mathbb{R}^n with support in \underline{I} . Setting

$$f(\underline{x}) = \int f(\lambda_1, \dots, \lambda_n) dE(\lambda_1, \dots, \lambda_n)$$

and since f is continuous, we finally realize that $f(\underline{x})$ is an element in \mathcal{A} .



Jensen's Inequality for
Conditional Expectations

Frank Hansen

Title Page

Contents



Go Back

Close

Quit

Page 3 of 12

2. Conditional Expectations

Let \mathcal{C} be a separable abelian C^* -subalgebra of a C^* -algebra \mathcal{A} , and let φ be a positive functional on \mathcal{A} such that \mathcal{C} is contained in the centralizer

$$\mathcal{A}^\varphi = \{y \in \mathcal{A} \mid \varphi(xy) = \varphi(yx) \quad \forall x \in \mathcal{A}\}.$$

The subalgebra is of the form $\mathcal{C} = C_0(S)$ for some locally compact metric space S .

Theorem 2.1. *There exists a positive linear mapping*

$$(2.1) \quad \Phi: M(\mathcal{A}) \rightarrow L^\infty(S, \mu_\varphi)$$

on the multiplier algebra $M(\mathcal{A})$ such that

$$\Phi(xy) = \Phi(yx) = \Phi(x)y, \quad x \in M(\mathcal{A}), y \in \mathcal{C}$$

almost everywhere, and a finite Radon measure μ_φ on S such that


$$\int_S z(s)\Phi(x)(s) d\mu_\varphi(s) = \varphi(zx), \quad z \in \mathcal{C}, x \in M(\mathcal{A}).$$

Proof. By the Riesz representation theorem there is a finite Radon measure μ_φ on S such that

$$\varphi(y) = \int_S y(s) d\mu_\varphi(s), \quad y \in \mathcal{C} = C_0(S).$$

For each positive element x in the multiplier algebra $M(\mathcal{A})$ we have

$$0 \leq \varphi(yx) = \varphi(y^{1/2}xy^{1/2}) \leq \|x\|\varphi(y), \quad y \in \mathcal{C}_+.$$



Jensen's Inequality for
Conditional Expectations

Frank Hansen

Title Page

Contents



Go Back

Close

Quit

Page 4 of 12

The functional $y \rightarrow \varphi(yx)$ on \mathcal{C} consequently defines a Radon measure on S which is dominated by a multiple of μ_φ , and it is therefore given by a unique element $\Phi(x)$ in $L^\infty(S, \mu_\varphi)$. By linearization this defines a positive linear mapping defined on the multiplier algebra

$$(2.2) \quad \Phi: M(\mathcal{A}) \rightarrow L^\infty(S, \mu_\varphi)$$

such that


$$\int_S z(s)\Phi(x)(s) d\mu_\varphi(s) = \varphi(zx), \quad z \in \mathcal{C}, x \in M(\mathcal{A}).$$

Furthermore, since

$$\int_S z(s)\Phi(yx)(s) d\mu_\varphi(s) = \varphi(zyx) = \int_S z(s)y(s)\Phi(x)(s) d\mu_\varphi(s)$$

for $x \in M(\mathcal{A})$ and $z, y \in \mathcal{C}$ we derive $\Phi(yx) = y\Phi(x) = \Phi(x)y$ almost everywhere. Since \mathcal{C} is contained in the centralizer \mathcal{A}^φ and thus $\varphi(zxy) = \varphi(yzx)$, we similarly obtain $\Phi(xy) = \Phi(x)y$ almost everywhere. \square

Note that $\Phi(z)(s) = z(s)$ almost everywhere in S for each $z \in \mathcal{C}$, cf. [6, 4, 5]. With a slight abuse of language we call Φ a conditional expectation even though its range is not a subalgebra of $M(\mathcal{A})$.



Jensen's Inequality for Conditional Expectations

Frank Hansen

Title Page

Contents



Go Back

Close

Quit

Page 5 of 12

3. Jensen's inequality

Following the notation in [5] we consider a separable C^* -algebra \mathcal{A} of operators on a (separable) Hilbert space \mathfrak{H} , and a field $(a_t)_{t \in T}$ of operators in the multiplier algebra

$$M(\mathcal{A}) = \{a \in \mathbb{B}(\mathfrak{H}) \mid a\mathcal{A} + \mathcal{A}a \subseteq \mathcal{A}\}$$

defined on a locally compact metric space T equipped with a Radon measure ν . We say that the field $(a_t)_{t \in T}$ is weak*-measurable if the function $t \rightarrow \varphi(a_t)$ is ν -measurable on T for each $\varphi \in \mathcal{A}^*$; and we say that the field is continuous if the function $t \rightarrow a_t$ is continuous [4].

As noted in [5] the field $(a_t)_{t \in T}$ is weak*-measurable, if and only if for each vector $\xi \in \mathfrak{H}$ the function $t \rightarrow a_t \xi$ is weakly (equivalently strongly) measurable. In particular, the composed field $(a_t^* b_t)_{t \in T}$ is weak*-measurable if both $(a_t)_{t \in T}$ and $(b_t)_{t \in T}$ are weak*-measurable fields.


If for a weak*-measurable field $(a_t)_{t \in T}$ the function $t \rightarrow |\varphi(a_t)|$ is integrable for every state $\varphi \in S(\mathcal{A})$ and the integrals

$$\int_T |\varphi(a_t)| d\nu(t) \leq K, \quad \forall \varphi \in S(\mathcal{A})$$

are uniformly bounded by some constant K , then there is a unique element (a C^* -integral in Pedersen's terminology [8, 2.5.15]) in the multiplier algebra $M(\mathcal{A})$, designated by

$$\int_T a_t d\nu(t),$$

such that



Jensen's Inequality for Conditional Expectations

Frank Hansen

Title Page

Contents



Go Back

Close

Quit

Page 6 of 12

$$\varphi \left(\int_T a_t d\nu(t) \right) = \int_T \varphi(a_t) d\nu(t), \quad \forall \varphi \in \mathcal{A}^*.$$

We say in this case that the field $(a_t)_{t \in T}$ is integrable. Finally we say that a field $(a_t)_{t \in T}$ is a unital column field [1, 4, 5], if it is weak*-measurable and

$$\int_T a_t^* a_t d\nu(t) = 1.$$

We note that a C^* -subalgebra of a separable C^* -algebra is automatically separable.

Theorem 3.1. *Let \mathcal{C} be an abelian C^* -subalgebra of a separable C^* -algebra \mathcal{A} , φ be a positive functional on \mathcal{A} such that \mathcal{C} is contained in the centralizer \mathcal{A}^φ and let*


$$\Phi: M(\mathcal{A}) \rightarrow L^\infty(S, \mu_\varphi)$$

be the conditional expectation defined in (2.1). Let furthermore $f: \underline{I} \rightarrow \mathbb{R}$ be a continuous convex function of n variables defined on a cube, and let $t \rightarrow a_t \in M(\mathcal{A})$ be a unital column field on a locally compact Hausdorff space T with a Radon measure ν . If $t \rightarrow \underline{x}_t$ is an essentially bounded, weak-measurable field on T of abelian n -tuples of self-adjoint elements in \mathcal{A} in the domain of f , then*

$$(3.1) \quad f(\Phi(y_1), \dots, \Phi(y_n)) \leq \Phi \left(\int_T a_t^* f(\underline{x}_t) a_t d\nu(t) \right)$$

almost everywhere, where the n -tuple \underline{y} in $M(\mathcal{A})$ is defined by setting

$$\underline{y} = (y_1, \dots, y_n) = \int_T a_t^* \underline{x}_t a_t d\nu(t).$$



Jensen's Inequality for Conditional Expectations

Frank Hansen

Title Page

Contents



Go Back

Close

Quit

Page 7 of 12

Proof. The subalgebra \mathcal{C} is as noted above of the form $\mathcal{C} = C_0(S)$ for some locally compact metric space S , and since the C^* -algebra $C_0(\underline{I})$ is separable we may for almost every s in S define a Radon measure μ_s on \underline{I} by setting

$$\mu_s(g) = \int_{\underline{I}} g(\underline{\lambda}) d\mu_s(\underline{\lambda}) = \Phi \left(\int_T a_t^* g(\underline{x}_t) a_t d\mu(t) \right) (s), \quad g \in C_0(\underline{I}).$$

Since

$$\mu_s(1) = \Phi \left(\int_T a_t^* a_t d\mu(t) \right) = \Phi(1) = 1$$


we observe that μ_s is a probability measure. If we put $g_i(\underline{\lambda}) = \lambda_i$ then

$$\int_{\underline{I}} g_i(\underline{\lambda}) d\mu_s(\underline{\lambda}) = \Phi \left(\int_T a_t^* x_{it} a_t d\mu(t) \right) (s) = \Phi(y_i)(s)$$

for $i = 1, \dots, n$ and since f is convex we obtain

$$\begin{aligned} f(\Phi(y_1)(s), \dots, \Phi(y_n)(s)) &= f \left(\int_{\underline{I}} g_1(\underline{\lambda}) d\mu_s(\underline{\lambda}), \dots, \int_{\underline{I}} g_n(\underline{\lambda}) d\mu_s(\underline{\lambda}) \right) \\ &\leq \int_{\underline{I}} f(g_1(\underline{\lambda}), \dots, g_n(\underline{\lambda})) d\mu_s(\underline{\lambda}) \\ &= \int_{\underline{I}} f(\underline{\lambda}) d\mu_s(\underline{\lambda}) \\ &= \Phi \left(\int_T a_t^* f(\underline{x}_t) a_t d\mu(t) \right) (s) \end{aligned}$$

for almost all s in S . □



Jensen's Inequality for Conditional Expectations

Frank Hansen

Title Page

Contents



Go Back

Close

Quit

Page 8 of 12

The following corollary is known as “Jensen’s inequality for expectation values”. It was formulated (for continuous fields) in the reference [3], where a more direct proof is given.

Corollary 3.2. *Let $f : \underline{I} \rightarrow \mathbb{R}$ be a continuous convex function of n variables defined on a cube, and let $t \rightarrow a_t \in B(H)$ be a unital column field on a locally compact Hausdorff space T with a Radon measure ν . If $t \rightarrow \underline{x}_t$ is a bounded weak*-measurable field on T of abelian n -tuples of self-adjoint operators on H in the domain of f , then*

$$(3.2) \quad f((y_1\xi | \xi), \dots, (y_n\xi | \xi)) \leq \left(\int_T a_t^* f(\underline{x}_t) a_t d\nu(t) \xi | \xi \right)$$

for any unit vector $\xi \in H$, where the n -tuple \underline{y} is defined by setting


$$\underline{y} = (y_1, \dots, y_n) = \int_T a_t^* \underline{x}_t a_t d\nu(t).$$

Proof. The statement follows from Theorem 3.1 by choosing φ as the trace and letting \mathcal{C} be the C^* -algebra generated by the orthogonal projection P on the vector ξ . Then $\mathcal{C} = C_0(S)$ where $S = \{0, 1\}$, and an element $z \in \mathcal{C}$ has the representation

$$z = z(0)P + z(1)(1 - P).$$

The measure $d\mu_\varphi$ gives unit weight in each of the two points, and the conditional expectation Φ is given by

$$\Phi(x)(s) = \begin{cases} (x\xi | \xi) & s = 0 \\ \text{Tr}(x - Px) & s = 1. \end{cases}$$



Jensen's Inequality for Conditional Expectations

Frank Hansen

Title Page

Contents



Go Back

Close

Quit

Page 9 of 12

Indeed,

$$\begin{aligned}\varphi(zx) &= \text{Tr}\left(\left(z(0)P + z(1)(1 - P)\right)x\right) \\ &= z(0)\Phi(x)(0) + z(1)\Phi(x)(0) \\ &= \int_S z(s)\Phi(x)(s) ds\end{aligned}$$

as required. The statement follows by evaluating the functions appearing on each side of the inequality (3.1) at the point $s = 0$. \square

Remark 1. If we choose ν as a probability measure on T , then the trivial field $a_t = 1$ for $t \in T$ is unital and (3.2) takes the form


$$f\left(\left(\int_T x_{1t} d\nu(t)\xi \mid \xi\right), \dots, \left(\int_T x_{nt} d\nu(t)\xi \mid \xi\right)\right) \leq \left(\int_T f(\underline{x}_t) d\nu(t)\xi \mid \xi\right)$$

for bounded weak*-measurable fields of abelian n -tuples $\underline{x}_t = (x_{1t}, \dots, x_{nt})$ of self-adjoint operators in the domain of f and unit vectors ξ . By choosing ν as an atomic measure with one atom we get a version

$$(3.3) \quad f((x_1\xi \mid \xi), \dots, (x_n\xi \mid \xi)) \leq (f(\underline{x})\xi \mid \xi)$$

of the Jensen inequality by Mond and Pečarić [7]. By further considering a direct sum

$$\xi = \bigoplus_{j=1}^m \xi_j \quad \text{and} \quad x = (x_1, \dots, x_n) = \bigoplus_{j=1}^m (x_{1j}, \dots, x_{nj})$$



Jensen's Inequality for Conditional Expectations

Frank Hansen

Title Page

Contents



Go Back

Close


Quit

Page 10 of 12

we obtain the familiar version

$$f \left(\sum_{j=1}^m (x_{1j} \xi_j \mid \xi_j), \dots, \sum_{j=1}^m (x_{nj} \xi_j \mid \xi_j) \right) \leq \sum_{j=1}^m \left(f(x_{1j}, \dots, x_{nj}) \xi_j \mid \xi_j \right)$$

valid for abelian n -tuples (x_{1j}, \dots, x_{nj}) , $j = 1, \dots, m$ of self-adjoint operators in the domain of f and vectors ξ_1, \dots, ξ_m with $\|\xi_1\|^2 + \dots + \|\xi_m\|^2 = 1$.



Jensen's Inequality for Conditional Expectations

Frank Hansen

Title Page

Contents



Go Back

Close

Quit

Page 11 of 12

References

- [1] H. ARAKI AND F. HANSEN, Jensen's operator inequality for functions of several variables, *Proc. Amer. Math. Soc.*, **128** (2000), 2075–2084.
- [2] L.G. BROWN AND H. KOSAKI, Jensen's inequality in semi-finite von Neumann algebras, *J. Operator Theory*, **23** (1990), 3–19.
- [3] F. HANSEN, Monotone trace functions of several variables, *International Journal of Mathematics*, **16** (2005), 777–785.
- [4] F. HANSEN AND G.K. PEDERSEN, Jensen's operator inequality, *Bull. London Math. Soc.*, **35** (2003), 553–564.
- [5] F. HANSEN AND G.K. PEDERSEN, Jensen's trace inequality in several variables, *International Journal of Mathematics*, **14** (2003), 667–681.
- [6] E. LIEB AND G.K. PEDERSEN, Convex multivariable trace functions, *Reviews in Mathematical Physics*, **14** (2002), 631–648.
- [7] B. MOND AND J.E. PEČARIĆ, On some operator inequalities, *Indian Journal of Mathematics*, **35** (1993), 221–232.
- [8] G.K. PEDERSEN, *Analysis Now, Graduate Texts in Mathematics*, Vol. 118, Springer Verlag, Heidelberg, 1989, reprinted 1995.
- [9] G.K. PEDERSEN, Convex trace functions of several variables on C^* -algebras, *J. Operator Theory*, **50** (2003), 157–167.



Jensen's Inequality for
Conditional Expectations

Frank Hansen

Title Page

Contents



Go Back

Close

Quit

Page 12 of 12