

GENERALIZED NONLINEAR MIXED QUASI-VARIATIONAL INEQUALITIES INVOLVING MAXIMAL η -MONOTONE MAPPINGS

MAO-MING JIN

Department of Mathematics
Fuling Teachers College
Fuling, Chongqing 408003
People's Republic of China.

EMail: mmj1898@163.com



volume 7, issue 3, article 114,
2006.

*Received 27 July, 2004;
accepted 09 May, 2006.*

Communicated by: S.S. Dragomir

[Abstract](#)

[Contents](#)



[Home Page](#)

[Go Back](#)

[Close](#)

[Quit](#)

Abstract

In this paper, we introduce and study a new class of generalized nonlinear mixed quasi-variational inequalities involving maximal η -monotone mapping. Using the resolvent operator technique for maximal η -monotone mapping, we prove the existence of solution for this kind of generalized nonlinear multi-valued mixed quasi-variational inequalities without compactness and the convergence of iterative sequences generated by the new algorithm. We also discuss the convergence and stability of the iterative sequence generated by the perturbed iterative algorithm for solving a class of generalized nonlinear mixed quasi-variational inequalities.

2000 Mathematics Subject Classification: 49J40; 47H10.


Key words: Mixed quasi-variational inequality, Maximal η -monotone mapping, Resolvent operator, Convergence, stability.

This work was supported by the National Natural Science Foundation of China (69903012) and the Educational Science Foundation of Chongqing, Chongqing of China (021301).

Contents

1	Introduction	3
2	Preliminaries	5
3	Iterative Algorithms	12
4	Main Results	15

References



Generalized Nonlinear Mixed Quasi-Variational Inequalities Involving Maximal η -Monotone Mappings

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit


Page 2 of 29

1. Introduction

In recent years, variational inequalities have been generalized and extended in many different directions using novel and innovative techniques. These have been used to study wider classes of unrelated problems arising in optimization and control, economics and finance, transportation and electrical networks, operations research and engineering sciences in a general and unified framework, see [1] – [15], [18] – [27] and the references therein. An important and useful generalization of variational inequality is called the variational inclusion. It is well known that one of the most important and interesting problems in the theory of variational inequalities is the development of an efficient and implementable algorithm for solving variational inequalities. For the past years, many numerical methods have been developed for solving various classes of variational inequalities, such as the projection method and its variant forms, linear approximation, descent, and Newton's methods.

Recently, Huang and Fang [10] introduced a new class of maximal η -monotone mappings and proved the Lipschitz continuity of the resolvent operator for maximal η -monotone mappings in Hilbert spaces. They also introduced and studied a new class of generalized variational inclusions involving maximal η -monotone mappings and constructed a new algorithm for solving this class of generalized variational inclusions by using the resolvent operator technique for maximal η -monotone mappings.

The main purpose of this paper is to introduce and study a new class of generalized nonlinear mixed quasi-variational inequalities involving maximal η -monotone mappings. Using the resolvent operator technique for maximal η -monotone mappings, we prove the existence of a solution for this kind of



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents




Go Back

Close

Quit

Page 3 of 29

generalized nonlinear multivalued mixed quasi-variational inequalities without compactness and the convergence of iterative sequences generated by the new algorithm. We also discuss the convergence and stability of the iterative sequence generated by the perturbed iterative algorithm for solving a class of generalized nonlinear mixed quasi-variational inequalities. The results shown in this paper improve and extend the previously known results in this area.



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 4 of 29

2. Preliminaries

Let H be a real Hilbert space endowed with a norm $\|\cdot\|$ and an inner product $\langle \cdot, \cdot \rangle$, respectively. Let $2^H, CB(H)$, and $\mathcal{H}(\cdot, \cdot)$ denote the family of all the nonempty subsets of H , the family of all the nonempty closed bounded subsets of H , and the Hausdorff metric on $CB(H)$, respectively. Let $\eta, N : H \times H \rightarrow H$ be two single-valued mappings with two variables and $g : H \rightarrow H$ be a single-valued mapping. Let $S, T, G : H \rightarrow CB(H)$ be three multivalued mappings and $M : H \times H \rightarrow 2^H$ be a multivalued mapping such that for each $t \in H$, $M(\cdot, t)$ is maximal η -monotone with $\text{Ran}(g) \cap \text{Dom } M(\cdot, t) \neq \emptyset$. Now we consider the following problem:

Find $u \in H, x \in Su, y \in Tu$, and $z \in Gu$ such that $g(u) \in \text{Dom}(M(\cdot, z))$ and

$$(2.1) \quad 0 \in N(x, y) + M(g(u), z).$$

Problem (2.1) is called a generalized nonlinear multivalued mixed quasi-variational inequality.

Some special cases of the problem (2.1):

- (I) If $\eta(x, y) = x - y$ for all x, y in H and G is the identity mapping, then problem (2.1) reduces to finding $u \in H, x \in Su, y \in Tu$ such that $g(u) \in \text{Dom}(M(\cdot, u))$ and

$$(2.2) \quad 0 \in N(x, y) + M(g(u), u).$$

Problem (2.2) is called the multivalued quasi-variational inclusion, which was studied by Noor [18] – [22].



Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 5 of 29

(II) If S, T are single-valued mappings and G is the identity mapping, then problem (2.1) is equivalent to finding $u \in H$ such that $g(u) \in \text{Dom}(M(\cdot, u))$ and

$$(2.3) \quad 0 \in N(Su, Tu) + M(g(u), u).$$

Problem (2.3) is called a generalized nonlinear mixed quasi-variational inequality.


(III) If $M(\cdot, t) = \partial\varphi(\cdot, t)$, where $\varphi : H \times H \rightarrow R \cup \{+\infty\}$ is a functional such that for each fixed t in H , $\varphi(\cdot, t) : H \rightarrow R \cup \{+\infty\}$ is lower semicontinuous and η -subdifferentiable on H , and $\partial\varphi(\cdot, t)$ denotes the η -subdifferential of $\varphi(\cdot, t)$, then problem (2.1) reduces to the following problem:

Find $u \in H, x \in Su$ and $y \in Tu$ such that

$$(2.4) \quad \langle N(x, y), \eta(v, g(u)) \rangle \geq \varphi(g(u), z) - \varphi(v, z)$$

for all v in H , which appears to be a new one. Furthermore, if $N(x, y) = x - y$ for all x, y in H , S, T are single-valued mappings and G is the identity mapping, then problem (2.4) reduces to the general quasi-variational-like inclusion considered by Ding and Luo [3].

(IV) If $S, T : H \rightarrow H$ are single-valued mappings, g is an identity mapping, $N(x, y) = x - y$ for all x, y in H , and $M(\cdot, t) = \partial\varphi$ for all t in H , where $\partial\varphi$ denotes the η -subdifferential of a proper convex lower semicontinuous function $\varphi : H \rightarrow R \cup \{+\infty\}$, then problem (2.1) reduces to the following problem:



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 6 of 29

Find $u \in H$ such that

$$(2.5) \quad \langle Su - Tu, \eta(v, u) \rangle \geq \varphi(u) - \varphi(v)$$


for all v in H , which is called the strongly nonlinear variational-like inclusion problem considered by Lee et al. [15].

- (V) If G is an identity mapping, $\eta(x, y) = x - y$ and $M(\cdot, t) = \partial\varphi$ for each $t \in H$, where $\varphi : H \rightarrow R \cup \{+\infty\}$ is a proper convex lower semicontinuous function on H and $g(H) \cap \text{Dom}(\partial\varphi(\cdot, t)) \neq \emptyset$ for each $t \in H$ and $\partial\varphi(\cdot, t)$ denotes the subdifferential of function $\varphi(\cdot, t)$, then problem (2.1) reduces to finding $u \in H, x \in Su$ and $y \in Tu$ such that $g(u) \in \text{Dom}(\partial\varphi(\cdot, t))$ and

$$(2.6) \quad \langle N(x, y), v - g(u) \rangle \geq \varphi(g(u)) - \varphi(v)$$

for all v in H . Furthermore, if $N(x, y) = x - y$ for all x, y in H , and g is an identity mapping, then the problem (2.6) is equivalent to the set-valued nonlinear generalized variational inclusion considered by Huang [6] and, in turn, includes the variational inclusions studied by Hassouni and Moudafi [5] and Kazmi [14] as special cases.

For a suitable choice of N, η, M, S, T, G, g , and for the space H , one can obtain a number of known and new classes of variational inclusions, variational inequalities, and corresponding optimization problems from the general set-valued variational inclusion problem (2.1). Furthermore, these types of variational inclusions enable us to study many important problems arising in the mathematical, physical, and engineering sciences in a general and unified framework.



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 7 of 29

Definition 2.1. Let T be a selfmap of H , $x_0 \in H$ and let $x_{n+1} = f(T, x_n)$ define an iteration procedure which yields a sequence of points $\{x_n\}_{n=0}^{\infty}$ in H . Suppose that $\{x \in H : Tx = x\} \neq \emptyset$ and $\{x_n\}_{n=0}^{\infty}$ converges to a fixed point x^* of T . Let $\{y_n\} \subset H$ and let $\epsilon_n = \|y_{n+1} - f(T, y_n)\|$. If $\lim_{n \rightarrow \infty} \epsilon_n = 0$ implies that $\lim_{n \rightarrow \infty} y_n = x^*$, then the iteration procedure defined by $x_{n+1} = f(T, x_n)$ is said to be T -stable or stable with respect to T .

Lemma 2.1 ([16]). Let $\{a_n\}$, $\{b_n\}$, and $\{c_n\}$ be three sequences of nonnegative numbers satisfying the following conditions: there exists n_0 such that

$$a_{n+1} \leq (1 - t_n)a_n + b_nt_n + c_n,$$

for all $n \geq n_0$, where $t_n \in [0, 1]$, $\sum_{n=0}^{\infty} t_n = \infty$, $\lim_{n \rightarrow \infty} b_n = 0$ and $\sum_{n=0}^{\infty} c_n < \infty$. Then $\lim_{n \rightarrow \infty} a_n = 0$.

Definition 2.2. A mapping $g : H \rightarrow H$ is said to be

(i) α -strongly monotone if there exists a constant $\alpha > 0$ such that


$$\langle g(u_1) - g(u_2), u_1 - u_2 \rangle \geq \alpha \|u_1 - u_2\|^2,$$

for all $u_i \in H$, $i = 1, 2$;

(ii) β -Lipschitz continuous if there exists a constant $\beta > 0$ such that

$$\|g(u_1) - g(u_2)\| \leq \beta \|u_1 - u_2\|,$$

for all $u_i \in H$, $i = 1, 2$.



Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 8 of 29

Definition 2.3. A multivalued mapping $S : H \rightarrow CB(H)$ is said to be

(i) \mathcal{H} -Lipschitz continuous if there exists a constant $\gamma > 0$ such that

$$\mathcal{H}(Su_1, Su_2) \leq \gamma \|u_1 - u_2\|,$$

for all $u_i \in H, i = 1, 2$;

(ii) strongly monotone with respect to the first argument of $N(\cdot, \cdot) : H \times H \rightarrow H$, if there exists a constant $\mu > 0$ such that

$$\langle N(x_1, \cdot) - N(x_2, \cdot), u_1 - u_2 \rangle \geq \mu \|u_1 - u_2\|^2,$$

for all $x_i \in Su_i, u_i \in H, i = 1, 2$.

Definition 2.4. A mapping $N(\cdot, \cdot) : H \times H \rightarrow H$ is said to be Lipschitz continuous with respect to the first argument if there exists a constant $\nu > 0$ such that

$$\|N(u_1, \cdot) - N(u_2, \cdot)\| \leq \nu \|u_1 - u_2\|,$$


for all $u_i \in H, i = 1, 2$.

In a similar way, we can define Lipschitz continuity of $N(\cdot, \cdot)$ with respect to the second argument.

Definition 2.5. Let $\eta : H \times H \rightarrow H$ be a single-valued mapping. A multivalued mapping $M : H \rightarrow 2^H$ is said to be

(i) η -monotone if

$$\langle x - y, \eta(u, v) \rangle \geq 0 \quad \text{for all } u, v \in H, x \in Mu, y \in Mv;$$



Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 9 of 29

(ii) strictly η -monotone if

$$\langle x - y, \eta(u, v) \rangle \geq 0 \quad \text{for all } u, v \in H, x \in Mu, y \in Mv$$

and equality holds if and only if $u = v$;

(iii) strongly η -monotone if there exists a constant $r > 0$ such that

$$\langle x - y, \eta(u, v) \rangle \geq r\|u - v\|^2 \quad \text{for all } u, v \in H, x \in Mu, y \in Mv;$$


(iv) maximal η -monotone if M is η -monotone and $(I + \lambda M)(H) = H$ for any $\lambda > 0$.

Remark 1.

1. If $\eta(u, v) = u - v$ for all u, v in H , then (i)-(iv) of Definition 2.5 reduce to the classical definitions of monotonicity, strict monotonicity, strong monotonicity, and maximal monotonicity, respectively.
2. Huang and Fang gave one example of maximal η -monotone mapping in [10].

Lemma 2.2 ([10]). Let $\eta : H \times H \rightarrow H$ be strictly monotone and $M : H \rightarrow 2^H$ be a maximal η -monotone mapping. Then the following conclusions hold:

1. $\langle x - y, \eta(u, v) \rangle \geq 0$ for all $(v, y) \in \text{Graph}(M)$ implies that $(u, x) \in \text{Graph}(M)$, where $\text{Graph}(M) = \{(u, x) \in H \times H : x \in Mu\}$;
2. the inverse mapping $(I + \lambda M)^{-1}$ is single-valued for any $\lambda > 0$.



Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 10 of 29


Based on Lemma 2.2, we can define the resolvent operator for a maximal η -monotone mapping M as follows:

$$(2.7) \quad J_{\rho}^M(z) = (I + \rho M)^{-1}(z) \quad \text{for all } z \in H,$$

where $\rho > 0$ is a constant and $\eta : H \times H \rightarrow H$ is a strictly monotone mapping.

Lemma 2.3 ([10]). *Let $\eta : H \times H \rightarrow H$ be strongly monotone and Lipschitz continuous with constants $\delta > 0$ and $\tau > 0$, respectively. Let $M : H \rightarrow 2^H$ be a maximal η -monotone mapping. Then the resolvent operator J_{ρ}^M for M is Lipschitz continuous with constant τ/δ , i.e.,*

$$\|J_{\rho}^M(u) - J_{\rho}^M(v)\| \leq \frac{\tau}{\delta} \|u - v\| \quad \text{for all } u, v \in H.$$



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 11 of 29

3. Iterative Algorithms

We first transfer problem (2.1) into a fixed point problem.

Lemma 3.1. For given $u \in H$, $x \in Su$, $y \in Tu$, and $z \in Gu$, (u, x, y, z) is a solution of the problem (2.1) if and only if

$$(3.1) \quad g(u) = J_{\rho}^{M(\cdot, z)}(g(u) - \rho N(x, y)),$$

where $J_{\rho}^{M(\cdot, z)} = (I + \rho M(\cdot, z))^{-1}$ and $\rho > 0$ is a constant.

Proof. This directly follows from the definition of $J_{\rho}^{M(\cdot, z)}$. □


Remark 2. Equality (3.1) can be written as

$$u = (1 - \lambda)u + \lambda[u - g(u) + J_{\rho}^{M(\cdot, z)}(g(u) - \rho N(x, y))],$$

where $0 < \lambda \leq 1$ is a parameter and $\rho > 0$ is a constant. This fixed point formulation enables us to suggest the following iterative algorithm for problem (2.1) as follows:

Algorithm 1. Let $\eta, N : H \times H \rightarrow H$, $g : H \rightarrow H$ be single-valued mappings and $S, T, G : H \rightarrow CB(H)$ be multivalued mappings. Let $M : H \times H \rightarrow 2^H$ be such that for each fixed $t \in H$, $M(\cdot, t) : H \rightarrow 2^H$ is a maximal η -monotone mapping satisfying $g(u) \in \text{Dom}(M(\cdot, z))$. For given $\lambda \in [0, 1]$, $u_0 \in H$, $x_0 \in Su_0$, $y_0 \in Tu_0$ and $z_0 \in Gu_0$, let

$$u_1 = (1 - \lambda)u_0 + \lambda [u_0 - g(u_0) + J_{\rho}^{M(\cdot, z_0)}(g(u_0) - \rho N(x_0, y_0))].$$



Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 12 of 29

By Nadler [17], there exist $x_1 \in Su_1$, $y_1 \in Tu_1$ and $z_1 \in Gu_1$ such that

$$\begin{aligned}\|x_0 - x_1\| &\leq (1 + 1)\mathcal{H}(Su_0, Su_1), \\ \|y_0 - y_1\| &\leq (1 + 1)\mathcal{H}(Tu_0, Tu_1), \\ \|z_0 - z_1\| &\leq (1 + 1)\mathcal{H}(Gu_0, Gu_1).\end{aligned}$$

Let

$$u_2 = (1 - \lambda)u_1 + \lambda [u_1 - g(u_1) + J_\rho^{M(\cdot, z_1)}(g(u_1) - \rho N(x_1, y_1))].$$


By induction, we can obtain sequences $\{u_n\}$, $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$ satisfying

$$(3.2) \quad \begin{cases} u_{n+1} = (1 - \lambda)u_n \\ \quad + \lambda [u_n - g(u_n) + J_\rho^{M(\cdot, z_n)}(g(u_n) - \rho N(x_n, y_n))], \\ x_n \in Su_n, \|x_n - x_{n+1}\| \leq (1 + (1 + n)^{-1})\mathcal{H}(Su_n, Su_{n+1}), \\ y_n \in Tu_n, \|y_n - y_{n+1}\| \leq (1 + (1 + n)^{-1})\mathcal{H}(Tu_n, Tu_{n+1}), \\ z_n \in Gu_n, \|z_n - z_{n+1}\| \leq (1 + (1 + n)^{-1})\mathcal{H}(Gu_n, Gu_{n+1}), \end{cases}$$

for $n = 1, 2, 3, \dots$, where $0 < \lambda \leq 1$ and $\rho > 0$ are both constants.

Now we construct a new pertured iterative algorithm for solving the generalized nonlinear mixed quasi-variational inequality (2.3) as follows:

Algorithm 2. Let $\eta, N : H \times H \rightarrow H$ and $S, T : H \rightarrow H$ be single-valued mappings. Let $M : H \times H \rightarrow 2^H$ be such that for each fixed $t \in H$, $M(\cdot, t) : H \rightarrow 2^H$ is a maximal η -monotone mapping satisfying $g(u) \in \text{Dom}(M(\cdot, u))$.



Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 13 of 29

For given $u_0 \in H$, the perturbed iterative sequence $\{u_n\}$ is defined by

$$(3.3) \quad \begin{cases} u_{n+1} = (1 - \alpha_n)u_n + \alpha_n[v_n - g(v_n) \\ \quad + J_\rho^{M(\cdot, v_n)}(g(v_n) - \rho N(Sv_n, Tv_n))] + \alpha_n e_n, \\ v_n = (1 - \beta_n)u_n + \beta_n[u_n - g(u_n) \\ \quad + J_\rho^{M(\cdot, u_n)}(g(u_n) - \rho N(Su_n, Tu_n))] + \beta_n f_n, \end{cases}$$


for $n = 0, 1, 2, \dots$, where $\{e_n\}$ and $\{f_n\}$ are two sequences of the elements of H introduced to take into account possible inexact computation and the sequences $\{\alpha_n\}$, $\{\beta_n\}$ satisfy the following conditions:

$$0 \leq \alpha_n, \beta_n \leq 1 (n \geq 0), \quad \text{and} \quad \sum_{n=0}^{\infty} \alpha_n = \infty.$$

Let $\{y_n\}$ be any sequence in H and define $\{\epsilon_n\}$ by

$$(3.4) \quad \begin{cases} \epsilon_n = \left\| y_{n+1} - \left\{ (1 - \alpha_n)y_n + \alpha_n \left[x_n - g(x_n) \right. \right. \right. \\ \quad \left. \left. \left. + J_\rho^{M(\cdot, x_n)}(g(x_n) - \rho N(Sx_n, Tx_n)) \right] + \alpha_n e_n \right\} \right\|, \\ x_n = (1 - \beta_n)y_n + \beta_n \left[y_n - g(y_n) \right. \\ \quad \left. + J_\rho^{M(\cdot, y_n)}(g(y_n) - \rho N(Sy_n, Ty_n)) \right] + \beta_n f_n, \end{cases}$$

for $n = 0, 1, 2, \dots$



Title Page

Contents



Go Back

Close

Quit

Page 14 of 29

4. Main Results


In this section, we first prove the existence of a solution of problem (2.1) and the convergence of an iterative sequence generated by Algorithm 1.

Theorem 4.1. *Let $\eta : H \times H \rightarrow H$ be strongly monotone and Lipschitz continuous with constants δ and τ , respectively. Let $S, T, G : H \rightarrow CB(H)$ be \mathcal{H} -Lipschitz continuous with constants α, β, γ , respectively, $g : H \rightarrow H$ be μ -Lipschitz continuous and ν -strongly monotone. Let $N : H \times H \rightarrow H$ be Lipschitz continuous with respect to the first and second arguments with constants ξ and ζ , respectively, and $S : H \rightarrow CB(H)$ be strongly monotone with respect to the first argument of $N(\cdot, \cdot)$ with constant r . Let $M : H \times H \rightarrow 2^H$ be a multivalued mapping such that for each fixed $t \in H$, $M(\cdot, t)$ is maximal η -monotone. Suppose that there exist constants $\rho > 0$ and $\kappa > 0$ such that for each $x, y, z \in H$,*

$$(4.1) \quad \left\| J_{\rho}^{M(\cdot, x)}(z) - J_{\rho}^{M(\cdot, y)}(z) \right\| \leq \kappa \|x - y\|,$$

and

$$(4.2) \quad \left\{ \begin{array}{l} \left| \rho - \frac{\tau r - \delta(1-h)\zeta\beta}{\tau(\xi^2\alpha^2 - \zeta^2\beta^2)} \right| < \frac{\sqrt{[\tau r - \delta(1-h)\zeta\beta]^2 - (\xi^2\alpha^2 - \zeta^2\beta^2)(\tau^2 - \delta^2(1-h)^2)}}{\tau(\xi^2\alpha^2 - \zeta^2\beta^2)}, \\ \tau r > \delta(1-h)\zeta\beta \\ \quad + \sqrt{(\xi^2\alpha^2 - \zeta^2\beta^2)(\tau^2 - \delta^2(1-h)^2)}, \quad \xi\alpha > \zeta\beta, \\ h = (1 + \delta\tau^{-1})\sqrt{1 - 2\nu + \mu^2} \\ \quad + \kappa\gamma, \quad \rho\tau\zeta\beta < \delta(1-h), \quad h < 1. \end{array} \right.$$



Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings

Mao-Ming Jin

Title Page

Contents



Go Back

Close


Quit

Page 15 of 29

Then the iterative sequences $\{u_n\}$, $\{x_n\}$, $\{y_n\}$ and $\{z_n\}$ generated by Algorithm 1 converge strongly to u^* , x^* , y^* and z^* , respectively and (u^*, x^*, y^*, z^*) is a solution of problem (2.1).

Proof. It follows from (3.2), (4.1) and Lemma 2.3 that

$$\begin{aligned}
 & \|u_{n+1} - u_n\| \\
 &= \left\| (1 - \lambda)(u_n - u_{n-1}) + \lambda \left[u_n - u_{n-1} - (g(u_n) - g(u_{n-1})) \right. \right. \\
 &\quad \left. \left. + J_\rho^{M(\cdot, z_n)}(g(u_n) - \rho N(x_n, y_n)) \right. \right. \\
 &\quad \left. \left. - J_\rho^{M(\cdot, z_{n-1})}(g(u_{n-1}) - \rho N(x_{n-1}, y_{n-1})) \right) \right\| \\
 &\leq (1 - \lambda) \|u_n - u_{n-1}\| + \lambda \|u_n - u_{n-1} - (g(u_{n+1}) - g(u_n))\| \\
 &\quad + \lambda \left\| J_\rho^{M(\cdot, z_n)}(g(u_n) - \rho N(x_n, y_n)) \right. \\
 &\quad \left. - J_\rho^{M(\cdot, z_n)}(g(u_{n-1}) - \rho N(x_{n-1}, y_{n-1})) \right\| \\
 &\quad + \lambda \left\| J_\rho^{M(\cdot, z_n)}(g(u_{n-1}) - \rho N(x_{n-1}, y_{n-1})) \right. \\
 &\quad \left. - J_\rho^{M(\cdot, z_{n-1})}(g(u_{n-1}) - \rho N(x_{n-1}, y_{n-1})) \right\| \\
 &\leq (1 - \lambda) \|u_n - u_{n-1}\| + \lambda \|u_n - u_{n-1} - (g(u_n) - g(u_{n-1}))\| \\
 &\quad + \lambda \frac{\tau}{\delta} \|g(u_n) - g(u_{n-1}) - \rho(N(x_n, y_n) - N(x_{n-1}, y_{n-1}))\| \\
 &\quad + \lambda \kappa \|z_n - z_{n-1}\| \\
 &\leq (1 - \lambda) \|u_n - u_{n-1}\| \\
 &\quad + \lambda \left(1 + \frac{\tau}{\delta} \right) \|u_n - u_{n-1} - (g(u_n) - g(u_{n-1}))\|
 \end{aligned}$$



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 16 of 29

$$(4.3) \quad \begin{aligned} & + \lambda \frac{\tau}{\delta} \|u_n - u_{n-1} - \rho(N(x_n, y_n) - N(x_{n-1}, y_n))\| \\ & + \lambda \rho \frac{\tau}{\delta} \|N(x_{n-1}, y_n) - N(x_{n-1}, y_{n-1})\| + \lambda \kappa \|z_n - z_{n-1}\|. \end{aligned}$$

Since g is strongly monotone and Lipschitz continuous, we obtain


$$(4.4) \quad \begin{aligned} & \|u_n - u_{n-1} - (g(u_n) - g(u_{n-1}))\|^2 \\ & = \|u_n - u_{n-1}\|^2 \\ & \quad - 2\langle u_n - u_{n-1}, g(u_n) - g(u_{n-1}) \rangle + \|g(u_n) - g(u_{n-1})\|^2 \\ & \leq (1 - 2\nu + \mu^2) \|u_n - u_{n-1}\|^2. \end{aligned}$$

Since S is \mathcal{H} -Lipschitz continuous and strongly monotone with respect to the first argument of $N(\cdot, \cdot)$ and N is Lipschitz continuous with respect to the first argument, we have

$$(4.5) \quad \begin{aligned} & \|u_n - u_{n-1} - \rho(N(x_n, y_n) - N(x_{n-1}, y_n))\|^2 \\ & = \|u_n - u_{n-1}\|^2 - 2\rho\langle u_n - u_{n-1}, N(x_n, y_n) - N(x_{n-1}, y_n) \rangle \\ & \quad + \rho^2 \|N(x_n, y_n) - N(x_{n-1}, y_n)\|^2 \\ & \leq (1 - 2\rho r + \rho^2 \xi^2 (1 + n^{-1})^2 \alpha^2) \|u_n - u_{n-1}\|^2. \end{aligned}$$

Further, since T, G are \mathcal{H} -Lipschitz continuous and N is Lipschitz continuous with respect to the second argument, we get

$$(4.6) \quad \begin{aligned} \|N(x_{n-1}, y_n) - N(x_{n-1}, y_{n-1})\| & \leq \zeta \|y_n - y_{n-1}\| \\ & \leq \zeta \beta (1 + n^{-1}) \|u_n - u_{n-1}\|, \end{aligned}$$



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 17 of 29

$$(4.7) \quad \|z_n - z_{n-1}\| \leq \gamma(1 + n^{-1})\|u_n - u_{n-1}\|.$$

By (4.3) – (4.7), we obtain

$$(4.8) \quad \begin{aligned} \|u_n - u_{n-1}\| &\leq (1 - \lambda + \lambda(1 + \tau\delta^{-1})\sqrt{1 - 2\nu + \mu^2} \\ &\quad + \lambda\tau\delta^{-1}\sqrt{1 - 2\rho r + \rho^2\xi^2(1 + n^{-1})^2\alpha^2} \\ &\quad + \lambda\rho\tau\delta^{-1}\zeta\beta(1 + n^{-1}) + \lambda\kappa\gamma(1 + n^{-1}))\|u_n - u_{n-1}\| \\ &= (1 - \lambda + \lambda h_n + \lambda t_n(\rho))\|u_n - u_{n-1}\| \\ &= \theta_n\|u_n - u_{n-1}\|, \end{aligned}$$


where

$$\begin{aligned} h_n &= (1 + \tau\delta^{-1})\sqrt{1 - 2\nu + \mu^2} + \kappa\gamma(1 + n^{-1}), \\ t_n(\rho) &= \tau\delta^{-1}\sqrt{1 - 2\rho r + \rho^2\xi^2(1 + n^{-1})^2\alpha^2} + \rho\tau\delta^{-1}\zeta\beta(1 + n^{-1}) \quad \text{and} \\ \theta_n &= 1 - \lambda + \lambda h_n + \lambda t_n(\rho). \end{aligned}$$

Letting $\theta = 1 - \lambda + \lambda h + \lambda t(\rho)$, where

$$\begin{aligned} h &= (1 + \tau\delta^{-1})\sqrt{1 - 2\nu + \mu^2} + \kappa\gamma \quad \text{and} \\ t(\rho) &= \tau\delta^{-1}\sqrt{1 - 2\rho r + \rho^2\xi^2\alpha^2} + \rho\tau\delta^{-1}\zeta\beta, \end{aligned}$$

we have that $h_n \rightarrow h$, $t_n(\rho) \rightarrow t(\rho)$ and $\theta_n \rightarrow \theta$ as $n \rightarrow \infty$. It follows from condition (4.2) that $\theta < 1$. Hence $\theta_n < 1$ for n sufficiently large. Therefore,



Title Page

Contents



Go Back

Close

Quit

Page 18 of 29

(4.8) implies that $\{u_n\}$ is a Cauchy sequence in H and so we can assume that $u_n \rightarrow u^* \in H$ as $n \rightarrow \infty$. By the Lipschitz continuity of S, T and G we obtain

$$\begin{aligned} \|x_n - x_{n-1}\| &\leq (1 + (1 + n)^{-1})\mathcal{H}(Su_n, Su_{n-1}) \\ &\leq \alpha(1 + (1 + n)^{-1})\|u_n - u_{n-1}\|, \\ \|y_n - y_{n-1}\| &\leq (1 + (1 + n)^{-1})\mathcal{H}(Tu_n, Tu_{n-1}) \\ &\leq \beta(1 + (1 + n)^{-1})\|u_n - u_{n-1}\|, \\ \|z_n - z_{n-1}\| &\leq (1 + (1 + n)^{-1})\mathcal{H}(Gu_n, Gu_{n-1}) \\ &\leq \gamma(1 + (1 + n)^{-1})\|u_n - u_{n-1}\|. \end{aligned}$$

It follows that $\{x_n\}, \{y_n\}$ and $\{z_n\}$ are also Cauchy sequences in H . We can assume that $x_n \rightarrow x^*, y_n \rightarrow y^*$ and $z_n \rightarrow z^*$, respectively. Note that for $x_n \in Su_n$, we have

$$\begin{aligned} d(x^*, Su^*) &\leq \|x^* - x_n\| + d(x_n, Su^*) \\ &\leq \|x^* - x_n\| + \mathcal{H}(Su_n, Su^*) \\ &\leq \|x^* - x_n\| + \alpha\|u_n - u^*\| \rightarrow 0, \end{aligned}$$


as $n \rightarrow \infty$. Hence we must have $x^* \in Su^*$. Similarly, we can show that $y^* \in Tu^*$ and $z^* \in Gu^*$. From

$$u_{n+1} = (1 - \lambda)u_n + \lambda [u_n - g(u_n) + J_\rho^{M(\cdot, z_n)}(g(u_n) - \rho N(x_n, y_n))],$$

it follows that

$$g(u^*) = J_\rho^{M(\cdot, z^*)}(g(u^*) - \rho N(x^*, y^*)).$$

By Lemma 3.1, (u^*, x^*, y^*, z^*) is a solution of problem (2.1). This completes the proof. \square



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 19 of 29

Remark 3. For an appropriate and suitable choice of the mappings η, N, S, T, G, g, M and the space H , we can obtain several known results in [1], [3], [5] – [8], [14], [18] – [22], [24] – [26] as special cases of Theorem 4.1.


Now we prove the convergence and stability of the iterative sequence generated by the Algorithm 2.

Theorem 4.2. Let $\eta : H \times H \rightarrow H$ be strongly monotone and Lipschitz continuous with constants δ and τ , respectively. Let $S, T : H \rightarrow H$ be Lipschitz continuous with constants α, β , respectively, $g : H \rightarrow H$ be μ -Lipschitz continuous and ν -strongly monotone. Let $N : H \times H \rightarrow H$ be Lipschitz continuous with respect to the first and second arguments with constants ξ and ζ , respectively, and $S : H \rightarrow H$ be strongly monotone with respect to the first argument of $N(\cdot, \cdot)$ with constant r . Let $M : H \times H \rightarrow 2^H$ be a multivalued mapping such that for each fixed $t \in H$, $M(\cdot, t)$ is maximal η -monotone. Suppose that there exist constants $\rho > 0$ and $\kappa > 0$ such that for each $x, y, z \in H$,

$$(4.9) \quad \left\| J_{\rho}^{M(\cdot, x)}(z) - J_{\rho}^{M(\cdot, y)}(z) \right\| \leq \kappa \|x - y\|,$$

and

$$(4.10) \quad \left\{ \begin{array}{l} \left| \rho - \frac{\tau r - \delta(1-h)\zeta\beta}{\tau(\xi^2\alpha^2 - \zeta^2\beta^2)} \right| < \frac{\sqrt{[\tau r - \delta(1-h)\zeta\beta]^2 - (\xi^2\alpha^2 - \zeta^2\beta^2)(\tau^2 - \delta^2(1-h)^2)}}{\tau(\xi^2\alpha^2 - \zeta^2\beta^2)}, \\ \tau r > \delta(1-h)\zeta\beta \\ \quad + \sqrt{(\xi^2\alpha^2 - \zeta^2\beta^2)(\tau^2 - \delta^2(1-h)^2)}, \quad \xi\alpha > \zeta\beta, \\ h = (1 + \delta\tau^{-1})\sqrt{1 - 2\nu + \mu^2} + \kappa, \\ \rho\tau\zeta\beta < \delta(1-h), \quad h < 1. \end{array} \right.$$



Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 20 of 29

If $\lim_{n \rightarrow \infty} \|e_n\| = 0$, $\lim_{n \rightarrow \infty} \|f_n\| = 0$, then

(I) The sequence $\{u_n\}$ defined by Algorithm 2 converges strongly to the unique solution u^* of problem (2.3).

(II) If $\sum_{n=0}^{\infty} \epsilon_n < \infty$, then $\lim_{n \rightarrow \infty} y_n = u^*$.


(III) If $\lim_{n \rightarrow \infty} y_n = u^*$, then $\lim_{n \rightarrow \infty} \epsilon_n = 0$.

Proof. (I) It follows from Theorem 4.1 that there exists $u^* \in H$ which is a solution of problem (2.3) and so

$$(4.11) \quad g(u^*) = J_{\rho}^{M(\cdot, u^*)}(g(u^*) - \rho N(Su^*, Tu^*)).$$

From (4.9), (4.11) and Algorithm 2, it follows that

$$\begin{aligned} & \|u_{n+1} - u^*\| \\ &= \left\| (1 - \alpha_n)(u_n - u^*) - \alpha_n \left[v_n - u^* - (g(v_n) - g(u^*)) \right. \right. \\ & \quad \left. \left. + J_{\rho}^{M(\cdot, v_n)}(g(v_n) - \rho N(Sv_n, Tv_n)) \right. \right. \\ & \quad \left. \left. - J_{\rho}^{M(\cdot, u^*)}(g(u^*) - \rho N(Su^*, Tu^*)) \right] + \alpha_n e_n \right\| \\ &\leq (1 - \alpha_n) \|u_n - u^*\| + \alpha_n \|v_n - u^* - (g(v_n) - g(u^*))\| + \alpha_n \|e_n\| \\ & \quad + \alpha_n \left\| J_{\rho}^{M(\cdot, v_n)}(g(v_n) - \rho N(Sv_n, Tv_n)) \right. \\ & \quad \left. - J_{\rho}^{M(\cdot, v_n)}(g(u^*) - \rho N(Su^*, Tu^*)) \right\| \\ & \quad + \alpha_n \left\| J_{\rho}^{M(\cdot, v_n)}(g(u^*) - \rho N(Su^*, Tu^*)) \right\| \end{aligned}$$



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents




Go Back

Close

Quit

Page 21 of 29



Title Page

Contents



Go Back

Close

Quit

Page 22 of 29

$$\begin{aligned}
 & -J_{\rho}^{M(\cdot, u^*)}(g(u^*) - \rho N(Su^*, Tu^*))\| \\
 \leq & (1 - \alpha_n)\|u_n - u^*\| + \alpha_n\|v_n - u^* - (g(v_n) - g(u^*))\| + \alpha_n\|e_n\| \\
 & + \alpha_n \frac{\tau}{\delta} \|g(v_n) - g(u^*) - \rho(N(Sv_n, Tv_n) - N(Su^*, Tu^*))\| \\
 & + \alpha_n \kappa \|v_n - u^*\| \\
 \leq & (1 - \alpha_n)\|u_n - u^*\| \\
 & + \alpha_n \left(1 + \frac{\tau}{\delta}\right) \|u_n - u^* - (g(v_n) - g(u^*))\| + \alpha_n\|e_n\| \\
 & + \alpha_n \frac{\tau}{\delta} \|v_n - u^* - \rho(N(Sv_n, Tv_n) - N(Su^*, Tu^*))\| \\
 (4.12) \quad & + \alpha_n \rho \frac{\tau}{\delta} \|N(Su^*, Tv_n) - N(Su^*, Tu^*)\| + \alpha_n \kappa \|v_n - u^*\|.
 \end{aligned}$$

By the Lipschitz continuity of N, S, T, g and the strong monotonicity of S and g , we obtain

$$(4.13) \quad \|v_n - u^* - (g(v_n) - g(u^*))\|^2 \leq (1 - 2\nu + \mu^2)\|v_n - u^*\|^2,$$

$$\begin{aligned}
 (4.14) \quad \|v_n - u^* - \rho(N(Sv_n, Tv_n) - N(Su^*, Tu^*))\|^2 \\
 \leq (1 - 2\rho r + \rho^2 \xi^2 \alpha^2)\|v_n - u^*\|^2,
 \end{aligned}$$

$$(4.15) \quad \|N(Su^*, Tv_n) - N(Su^*, Tu^*)\| \leq \zeta \beta \|v_n - u^*\|.$$

It follows from (4.12) – (4.15) that

$$(4.16) \quad \|u_{n+1} - u^*\| \leq (1 - \alpha_n)\|u_n - u^*\| + \theta \alpha_n \|v_n - u^*\| + \alpha_n \|e_n\|,$$

where

$$\theta = \kappa + (1 + \tau\delta^{-1})\sqrt{1 - 2\nu + \mu^2} + \tau\delta^{-1}\sqrt{1 - 2\rho r + \rho^2\xi^2\alpha^2} + \rho\tau\delta^{-1}\zeta\beta.$$

Similarly, we have

$$(4.17) \quad \|v_n - u^*\| \leq (1 - \beta_n)\|u_n - u^*\| + \theta\beta_n\|u_n - u^*\| + \beta_n\|f_n\|.$$

From (4.16) and (4.17), we have

$$\|u_{n+1} - u^*\| \leq [1 - \alpha_n(1 - \theta)(1 + \theta\beta_n)]\|u_n - u^*\| + \alpha_n\beta_n\theta\|f_n\| + \alpha_n\|e_n\|$$

Condition (4.10) implies that $0 < \theta < 1$, and so

$$(4.18) \quad \|u_{n+1} - u^*\| \leq [1 - \alpha_n(1 - \theta)]\|u_n - u^*\| + \alpha_n(1 - \theta)d_n,$$

where $d_n = (\beta_n\theta\|f_n\| + \|e_n\|)(1 - \theta)^{-1} \rightarrow 0$, as $n \rightarrow \infty$. It follows from (4.18) and Lemma 2.1 that $u_n \rightarrow u^*$ as $n \rightarrow \infty$.


Now we prove that u^* is a unique solution of problem (2.3). In fact, if u is also a solution of problem (2.3), then

$$g(u) = J_\rho^{M(\cdot, u)}(g(u) - \rho N(Su, Tu)),$$

and, as the proof of (4.16), we have

$$\|u^* - u\| \leq \theta\|u^* - u\|,$$

where $0 < \theta < 1$ and so $u^* = u$. This completes the proof of Conclusion (I).



Title Page

Contents



Go Back

Close

Quit

Page 23 of 29

Next we prove Conclusion (II). Using (3.4) we obtain

$$\begin{aligned}
 & \|y_{n+1} - u^*\| \\
 & \leq \|y_{n+1} - \{(1 - \alpha_n)y_n + \alpha_n[x_n - g(x_n) \\
 & \quad + J_\rho^{M(\cdot, x_n)}(g(x_n) - \rho N(Sx_n, Tx_n))]\} + \alpha_n e_n\| \\
 & \quad + \|(1 - \alpha_n)y_n + \alpha_n[x_n - g(x_n) \\
 & \quad + J_\rho^{M(\cdot, x_n)}(g(x_n) - \rho N(Sx_n, Tx_n))]\} + \alpha_n e_n - u^*\| \\
 (4.19) \quad & = \|(1 - \alpha_n)y_n + \alpha_n[x_n - g(x_n) \\
 & \quad + J_\rho^{M(\cdot, x_n)}(g(x_n) - \rho N(Sx_n, Tx_n))]\} + \alpha_n e_n - u^*\| + \epsilon_n.
 \end{aligned}$$


As the proof of inequality (4.18), we have

$$\begin{aligned}
 (4.20) \quad & \|(1 - \alpha_n)y_n + \alpha_n[x_n - g(x_n) \\
 & \quad + J_\rho^{M(\cdot, x_n)}(g(x_n) - \rho N(Sx_n, Tx_n))]\} + \alpha_n e_n - u^*\| \\
 & \leq (1 - \alpha_n(1 - \theta))\|y_n - u^*\| + \alpha_n(1 - \theta)d_n.
 \end{aligned}$$

It follows from (4.19) and (4.20) that

$$(4.21) \quad \|y_{n+1} - u^*\| \leq (1 - \alpha_n(1 - \theta))\|y_n - u^*\| + \alpha_n(1 - \theta)d_n + \epsilon_n.$$

Since $\sum_{n=0}^{\infty} \epsilon_n < \infty$, $d_n \rightarrow 0$, and $\sum_{n=0}^{\infty} \alpha_n < \infty$. It follows that (4.21) and Lemma 2.1 that $\lim_{n \rightarrow \infty} y_n = u^*$.



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents



Go Back

Close


Quit

Page 24 of 29

Now we prove Conclusion (III). Suppose that $\lim_{n \rightarrow \infty} y_n = u^*$. Then we have

$$\begin{aligned} \epsilon_n &= \left\| y_{n+1} - (1 - \alpha_n)y_n + \alpha_n \left[x_n \right. \right. \\ &\quad \left. \left. - g(x_n) + J_\rho^{M(\cdot, x_n)}(g(x_n) - \rho N(Sx_n, Tx_n)) \right] + \alpha_n e_n \right\| \\ &\leq \|y_{n+1} - u^*\| + \left\| (1 - \alpha_n)y_n + \alpha_n \left[x_n \right. \right. \\ &\quad \left. \left. - g(x_n) + J_\rho^{M(\cdot, x_n)}(g(x_n) - \rho N(Sx_n, Tx_n)) \right] + \alpha_n e_n - u^* \right\| \\ &\leq \|y_{n+1} - u^*\| + (1 - \alpha_n(1 - \theta))\|y_n - u^*\| + \alpha_n(1 - \theta)d_n \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$. This completes the proof. □



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents



Go Back


Close

Quit

Page 25 of 29

References

- [1] S. ADLY, Perturbed algorithm and sensitivity analysis for a general class of variational inclusions, *J. Math. Anal. Appl.*, **201** (1996), 609–630.
- [2] X.P. DING, Generalized quasi-variational-like inclusions with nonconvex functionals, *Appl. Math. Comput.*, **122** (2001), 267–282.
- [3] X.P. DING AND C.L. LUO, Perturbed proximal point algorithms for generalized quasi-variational-like inclusions, *J. Comput. Appl. Math.*, **113** (2000), 153–165.
- [4] F. GIANNESSI AND A. MAUGERI, *Variational Inequalities and Network Equilibrium Problems*, Plenum, New York, 1995.
- [5] A. HASSOUNI AND A. MOUDAFI, A perturbed algorithm for variational inequalities, *J. Math. Anal. Appl.*, **185** (1994), 706–712.
- [6] N.J. HUANG, Generalized nonlinear variational inclusions with noncompact valued mapping, *Appl. Math. Lett.*, **9**(3) (1996), 25–29.
- [7] N.J. HUANG, On the generalized implicit quasivariational inequalities, *J. Math. Anal. Appl.*, **216** (1997), 197–210.
- [8] N.J. HUANG, A new completely general class of variational inclusions with noncompact valued mappings, *Computers Math. Appl.*, **35**(10) (1998), 9–14.
- [9] N.J. HUANG AND C.X. DENG, Auxiliary principle and iterative algorithms for generalized set-valued strongly nonlinear mixed variational-like inequalities, *J. Math. Anal. Appl.*, **256** (2001), 345–359.



Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings

Mao-Ming Jin

Title Page

Contents




Go Back

Close

Quit

Page 26 of 29

- [10] N.J. HUANG AND Y.P. FANG, A new class of general variational inclusions involving maximal η -monotone mappings, *Publ. Math. Debrecen*, **62** (2003).
- [11] M.M. JIN, Generalized nonlinear Implicit quasivariational inclusions with relaxed monotone mappings, *J. Adv. Nonlinear Var. Inequal.*, **7**(2) (2004), 173–181.
- [12] M.M. JIN, Sensitivity analysis for strongly nonlinear quasivariational inclusions involving generalized m -accretive mappings, *J. Nonlinear Anal. Forum*, **8**(1) (2003), 93–99.
- [13] M.M. JIN AND Q.K. LIU, Nonlinear quasi-variational inclusions involving generalized m -accretive mappings, *Nonlinear Funct. Anal. Appl.*, (in press).
- [14] K.R. KAZMI, Mann and Ishikawa type perturbed iterative algorithms for generalized quasivariational inclusions, *J. Math. Anal. Appl.*, **209** (1997), 572–584.
- [15] C.H. LEE, Q.H. ANSARI AND J.C. YAO, A perturbed algorithm for strongly nonlinear variational-like inclusions, *Bull. Austral. Math. Soc.*, **62** (2000), 417–426.
- [16] L.S. LIU, Ishikawa and Mann iterative proces with errors for nonlinear strongly accretive mappings in Banach spaces, *J. Math. Anal. Appl.*, **194** (1995), 114–125.
- [17] S.B. NADLER, Multivalued contraction mappings, *Pacific J. Math.*, **30** (1969), 475–488.



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents




Go Back

Close

Quit

Page 27 of 29

- [18] M.A. NOOR, Set-valued quasi-variational inequalities, *J. Comput Appl Math.*, **7** (2000), 101–113.
- [19] M.A. NOOR, Three-step iterative algorithms for multivalued quasi-variational inclusions, *J. Math. Anal. Appl.*, **255** (2001), 589–604.
- [20] M.A. NOOR, Two-step approximation schemes for multivalued quasi-variational inclusions, *Nonlinear Funct Anal Appl.*, **7**(1) (2002), 1–14.
- [21] M.A. NOOR, Multivalued quasi-variational inclusions and implicit resolvent equations, *Nonlinear Anal TMA.* **48**(2) (2002), 159–174.
- [22] M.A. NOOR, Operator-splitting methods for general mixed variational inequalities, *J. Inequal. Pure and Appl. Math.*, **3**(5) (2002), Art. 67. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=219>].
- [23] P.D. PANAGIOTOPOULOS AND G.E. STAVROULAKIS, New types of variational principles based on the notion of quasidifferentiability, *Acta Mech.*, **94** (1992), 171–194.
- [24] A.H. SIDDIQI AND Q.H. ANSARI, General strongly nonlinear variational inequalities, *J. Math. Anal. Appl.*, **116** (1992), 386–392.
- [25] A.H. SIDDIQI AND Q.H. ANSARI, Strongly nonlinear quasivariational inequalities, *J. Math. Anal. Appl.*, **149** (1990), 444–450.
- [26] GEORGE X.Z. YUAN, *KKM Theory and Applications in Nonlinear Analysis*, Marcel Dekker, New York, 1999.



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents




Go Back

Close

Quit

Page 28 of 29

- [27] J.X. ZHOU AND G. CHEN, Diagonal convexity conditions for problems in convex analysis and quasivariational inequalities, *J. Math. Anal. Appl.*, **132** (1988), 213–225.



**Generalized Nonlinear Mixed
Quasi-Variational Inequalities
Involving Maximal η -Monotone
Mappings**

Mao-Ming Jin

Title Page

Contents



Go Back

Close

Quit

Page 29 of 29