

HILBERT-PACHPATTE TYPE INEQUALITIES FROM BONSALL'S FORM OF HILBERT'S INEQUALITY

JOSIP PEČARIĆ

Faculty of Textile Technology
University of Zagreb
Pierottijeva 6, 10000 Zagreb, Croatia

EEmail: pecaric@hazu.hr

IVAN PERIĆ

Faculty of Food Technology and Biotechnology
University of Zagreb
Pierottijeva 6, 10000 Zagreb, Croatia

EEmail: iperic@pbf.hr

PREDRAG VUKOVIĆ

Faculty of Teacher Education
University of Zagreb, Ante Starčevića 55
40000 Čakovec, Croatia

EEmail: predrag.vukovic@vus-ck.hr

Received: 19 June, 2007

Accepted: 09 October, 2007

Communicated by: **B. Yang**

2000 AMS Sub. Class.: 26D15.

Key words: Inequalities, Hilbert's inequality, sequences and functions, homogeneous kernels, conjugate and non-conjugate exponents, the Beta function.

Abstract: The main objective of this paper is to deduce Hilbert-Pachpatte type inequalities using Bonsall's form of Hilbert's and Hardy-Hilbert's inequalities, both in discrete and continuous case.



**Hilbert-Pachpatte Type
Inequalities**

Josip Pečarić, Ivan Perić
and Predrag Vuković

vol. 9, iss. 1, art. 9, 2008

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 1 of 27

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Contents

1	Introduction	3
2	Integral Case	8
3	Discrete Case	17
4	Non-conjugate Exponents	23



**Hilbert-Pachpatte Type
Inequalities**

Josip Pečarić, Ivan Perić
and Predrag Vuković

vol. 9, iss. 1, art. 9, 2008

Title Page

Contents



Page 2 of 27

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



1. Introduction

An interesting feature of one of the forms of Hilbert-Pachpatte type inequalities, is that it controls the size (in the sense of L^p or l^p spaces) of the modified Hilbert transform of a function or of a series with the size of its derivative or its backward differences, respectively. We start with the following results of Zhongxue Lü from [9], for both continuous and discrete cases. For a sequence $a : \mathbb{N}_0 \rightarrow \mathbb{R}$, the sequence $\nabla a : \mathbb{N} \rightarrow \mathbb{R}$ is defined by $\nabla a(n) = a(n) - a(n-1)$. For a function $u : (0, \infty) \rightarrow \mathbb{R}$, u' denotes the usual derivative of u .

Theorem A. Let $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $s > 2 - \min\{p, q\}$, and $f(x), g(y)$ be real-valued continuous functions defined on $[0, \infty)$, respectively, and let $f(0) = g(0) = 0$, and

$$0 < \int_0^\infty \int_0^x x^{1-s} |f'(\tau)|^p d\tau dx < \infty, \quad 0 < \int_0^\infty \int_0^y y^{1-s} |g'(\delta)|^q d\delta dy < \infty,$$

then

$$(1.1) \quad \int_0^\infty \int_0^\infty \frac{|f(x)||g(y)|}{(qx^{p-1} + py^{q-1})(x+y)^s} dx dy \\ \leq \frac{B\left(\frac{q+s-2}{q}, \frac{p+s-2}{p}\right)}{pq} \cdot \left(\int_0^\infty \int_0^x x^{1-s} |f'(\tau)|^p d\tau dx\right)^{\frac{1}{p}} \\ \times \left(\int_0^\infty \int_0^y y^{1-s} |g'(\delta)|^q d\delta dy\right)^{\frac{1}{q}}.$$

Theorem B. Let $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $s > 2 - \min\{p, q\}$, and $\{a(m)\}$ and $\{b(n)\}$ be

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 3 of 27


[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 4 of 27

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

two sequences of real numbers where $m, n \in \mathbb{N}_0$, and $a(0) = b(0) = 0$, and

$$0 < \sum_{m=1}^{\infty} \sum_{\tau=1}^m m^{1-s} |\nabla a(\tau)|^p < \infty, \quad 0 < \sum_{n=1}^{\infty} \sum_{\delta=1}^n n^{1-s} |\nabla b(\delta)|^q < \infty,$$

then


$$(1.2) \quad \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{|a_m| |b_n|}{(qm^{p-1} + pn^{q-1})(m+n)^s} \\ \leq \frac{B\left(\frac{q+s-2}{q}, \frac{p+s-2}{p}\right)}{pq} \cdot \left(\sum_{m=1}^{\infty} \sum_{\tau=1}^m m^{1-s} |\nabla a(\tau)|^p \right)^{\frac{1}{p}} \\ \times \left(\sum_{n=1}^{\infty} \sum_{\delta=1}^n n^{1-s} |\nabla b(\delta)|^q \right)^{\frac{1}{q}}.$$

Note that the condition $s > 2 - \min\{p, q\}$ from Theorem B is not sufficient. Namely, the author of the proof of Theorem B used the following result

$$(1.3) \quad \sum_{n=1}^{\infty} \frac{1}{(m+n)^s} \left(\frac{m}{n}\right)^{\frac{2-s}{q}} < B\left(\frac{q+s-2}{q}, \frac{p+s-2}{p}\right) m^{1-s},$$

for $m \in \{1, 2, \dots\}$ and $s > 2 - \min\{p, q\}$. For $p = q = 2$, $s = 18$ and $m = 1$, the left-hand side of (1.3) is greater than the right-hand side of (1.3). Therefore, we refer to a paper of Krnić and Pečarić, [4], where the next inequality is given:

$$(1.4) \quad \sum_{n=1}^{\infty} \frac{1}{(m+n)^s} \frac{m^{\alpha_1}}{n^{\alpha_2}} < m^{1-s+\alpha_1-\alpha_2} B(1-\alpha_2, s+\alpha_2-1),$$



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 5 of 27

[Go Back](#)

[Full Screen](#)

[Close](#)

where $0 < s \leq 14$, $1 - s < \alpha_2 < 1$ for $s \leq 2$ and $-1 \leq \alpha_2 < 1$ for $s > 2$. By using this result, here we shall obtain a generalization of Theorem B but with the condition $2 - \min\{p, q\} < s \leq 2 + \min\{p, q\}$. Also, the following result is given in [9]:

$$(1.5) \quad \sum_{n=1}^{\infty} \frac{1}{m^s + n^s} \left(\frac{m}{n}\right)^{\frac{2-s}{q}} < \frac{1}{s} B\left(\frac{q+s-2}{sq}, \frac{p+s-2}{sp}\right) m^{1-s},$$

for $m \in \{1, 2, \dots\}$ and $s > 2 - \min\{p, q\}$. Similarly as before, for $p = q = 2$, $s = 6$ and $m = 1$, the left-hand side of (1.5) is greater than the right-hand side of (1.5). The case of nontrivial weights is essential in Theorem A and Theorem B, since for $s = 1$ only the trivial functions and sequences satisfy the assumptions.

In 1951, Bonsall established the following conditions for non-conjugate exponents (see [1]). Let p and q be real parameters, such that

$$(1.6) \quad p > 1, q > 1, \frac{1}{p} + \frac{1}{q} \geq 1,$$

and let p' and q' respectively be their conjugate exponents, that is, $\frac{1}{p} + \frac{1}{p'} = 1$ and $\frac{1}{q} + \frac{1}{q'} = 1$. Further, define

$$(1.7) \quad \lambda = \frac{1}{p'} + \frac{1}{q'}$$

and note that $0 < \lambda \leq 1$ for all p and q as in (1.6). In particular, $\lambda = 1$ holds if and only if $q = p'$, that is, only when p and q are mutually conjugate. Otherwise, we have $0 < \lambda < 1$, and in such cases p and q will be referred to as non-conjugate exponents. Also, in this paper we shall obtain some generalizations of (1.1). It will be done in simpler way than in [9]. Our results will be based on the following results of Pečarić et al., [2], for the non-conjugate and conjugate exponents.



Theorem C. Let p, p', q, q' and λ be as in (1.6) and (1.7). If K, φ, ψ, f and g are non-negative measurable functions, then the following inequalities hold and are equivalent

$$(1.8) \quad \int_{\Omega^2} K^\lambda(x, y) f(x) g(y) dx dy \leq \left(\int_{\Omega} (\varphi F f)^p(x) dx \right)^{\frac{1}{p}} \left(\int_{\Omega} (\psi G g)^q(y) dy \right)^{\frac{1}{q}}$$

and

$$(1.9) \quad \left(\int_{\Omega} \left(\frac{1}{\psi G(y)} \int_{\Omega} K^\lambda(x, y) f(x) dx \right)^{q'} dy \right)^{\frac{1}{q'}} \leq \left(\int_{\Omega} (\varphi F f)^p(x) dx \right)^{\frac{1}{p}},$$

where the functions F, G are defined by

$$F(x) = \left(\int_{\Omega} \frac{K(x, y)}{\psi^{q'}(y)} dy \right)^{\frac{1}{q'}} \quad \text{and} \quad G(y) = \left(\int_{\Omega} \frac{K(x, y)}{\varphi^{p'}(x)} dx \right)^{\frac{1}{p'}}.$$

The next inequalities from [5] can be seen as a special case of (1.8) and (1.9) respectively for the conjugate exponents:

$$(1.10) \quad \int_{\Omega^2} K(x, y) f(x) g(y) dx dy \leq \left(\int_{\Omega} \varphi^p(x) F(x) f^p(x) dx \right)^{\frac{1}{p}} \left(\int_{\Omega} \psi^q(y) G(y) g^q(y) dy \right)^{\frac{1}{q}}$$

and

$$(1.11) \quad \int_{\Omega} G^{1-p}(y) \psi^{-p}(y) \left(\int_{\Omega} K(x, y) f(x) dx \right)^p dy \leq \int_{\Omega} \varphi^p(x) F(x) f^p(x) dx,$$

[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 6 of 27


[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



Title Page

Contents



Page 7 of 27

Go Back

Full Screen

Close

where

$$(1.12) \quad F(x) = \int_{\Omega} \frac{K(x, y)}{\psi^p(y)} dy \quad \text{and} \quad G(y) = \int_{\Omega} \frac{K(x, y)}{\varphi^q(x)} dx.$$

In particular, inequalities (1.10) and (1.11) are equivalent.


On the other hand, here we also refer to a paper of Brnetić et al., [8], where a general Hilbert-type inequality was obtained for $n \geq 2$ conjugate exponents, that is, real parameters $p_1, \dots, p_n > 1$, such that $\sum_{i=1}^n \frac{1}{p_i} = 1$. Namely, we let $K : \Omega^n \rightarrow \mathbb{R}$ and $\phi_{ij} : \Omega \rightarrow \mathbb{R}$, $i, j = 1, \dots, n$, be non-negative measurable functions. If $\prod_{i,j=1}^n \phi_{ij}(x_j) = 1$, then the inequality

$$(1.13) \quad \int_{\Omega^n} K(x_1, \dots, x_n) \prod_{i=1}^n f_i(x_i) dx_1 \dots dx_n \leq \prod_{i=1}^n \left(\int_{\Omega} F_i(x_i) (\phi_{ii} f_i)^{p_i}(x_i) dx_i \right)^{\frac{1}{p_i}},$$

holds for all non-negative measurable functions $f_1, \dots, f_n : \Omega \rightarrow \mathbb{R}$, where

$$(1.14) \quad F_i(x_i) = \int_{\Omega^{n-1}} K(x_1, \dots, x_n) \prod_{j=1, j \neq i}^n \phi_{ij}^{p_i}(x_j) dx_1 \dots dx_{i-1} dx_{i+1} \dots dx_n,$$

for $i = 1, \dots, n$.



[Title Page](#)

[Contents](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

Page 8 of 27

[Go Back](#)

[Full Screen](#)

[Close](#)

2. Integral Case

In this section we shall state our main results. We suppose that all integrals converge and shall omit these types of conditions. Thus, we have the following.

Theorem 2.1. *Let $\frac{1}{p} + \frac{1}{q} = 1$ with $p > 1$. If $K(x, y)$, $\varphi(x)$, $\psi(y)$ are non-negative functions and $f(x)$, $g(y)$ are absolutely continuous functions such that $f(0) = g(0) = 0$, then the following inequalities hold*

$$\begin{aligned}
 (2.1) \quad & \int_0^\infty \int_0^\infty \frac{K(x, y)|f(x)||g(y)|}{qx^{p-1} + py^{q-1}} dx dy \\
 & \leq \int_0^\infty \int_0^\infty K(x, y)|f(x)||g(y)| d\left(x^{\frac{1}{p}}\right) d\left(y^{\frac{1}{q}}\right) \\
 & \leq \frac{1}{pq} \left(\int_0^\infty \int_0^x \varphi^p(x) F(x) |f'(\tau)|^p d\tau dx \right)^{\frac{1}{p}} \\
 & \quad \times \left(\int_0^\infty \int_0^y \psi^q(y) G(y) |g'(\delta)|^q d\delta dy \right)^{\frac{1}{q}},
 \end{aligned}$$

and

$$\begin{aligned}
 (2.2) \quad & \int_0^\infty G^{1-p}(y) \psi^{-p}(y) \left(\int_0^\infty K(x, y) |f(x)| d\left(x^{\frac{1}{p}}\right) \right)^p dy \\
 & \leq \frac{1}{p^p} \int_0^\infty \int_0^x \varphi^p(x) F(x) |f'(\tau)|^p d\tau dx,
 \end{aligned}$$

where $F(x)$ and $G(y)$ are defined as in (1.12).



Proof. By using Hölder's inequality, (see also [9]), we have

$$(2.3) \quad |f(x)| |g(y)| \leq x^{\frac{1}{q}} y^{\frac{1}{p}} \left(\int_0^x |f'(\tau)|^p d\tau \right)^{\frac{1}{p}} \left(\int_0^y |g'(\delta)|^q d\delta \right)^{\frac{1}{q}}.$$

From (2.3) and using the elementary inequality

$$(2.4) \quad xy \leq \frac{x^p}{p} + \frac{y^q}{q}, \quad x \geq 0, \quad y \geq 0, \quad \frac{1}{p} + \frac{1}{q} = 1, \quad p > 1,$$

we observe that

$$(2.5) \quad \frac{pq|f(x)| |g(y)|}{qx^{p-1} + py^{q-1}} \leq \frac{|f(x)| |g(y)|}{x^{\frac{1}{q}} y^{\frac{1}{p}}} \leq \left(\int_0^x |f'(\tau)|^p d\tau \right)^{\frac{1}{p}} \left(\int_0^y |g'(\delta)|^q d\delta \right)^{\frac{1}{q}}$$

and therefore


$$(2.6) \quad \begin{aligned} pq \int_0^\infty \int_0^\infty \frac{K(x, y) |f(x)| |g(y)|}{qx^{p-1} + py^{q-1}} dx dy \\ \leq \int_0^\infty \int_0^\infty \frac{K(x, y)}{x^{\frac{1}{q}} y^{\frac{1}{p}}} |f(x)| |g(y)| dx dy \\ \leq \int_0^\infty \int_0^\infty K(x, y) \left(\int_0^x |f'(\tau)|^p d\tau \right)^{\frac{1}{p}} \left(\int_0^y |g'(\delta)|^q d\delta \right)^{\frac{1}{q}} dx dy. \end{aligned}$$

Applying the substitutions

$$f_1(x) = \left(\int_0^x |f'(\tau)|^p d\tau \right)^{\frac{1}{p}}, \quad g_1(y) = \left(\int_0^y |g'(\delta)|^q d\delta \right)^{\frac{1}{q}}$$

Title Page

Contents



Page 9 of 27

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 10 of 27

[Go Back](#)

[Full Screen](#)

[Close](#)

and (1.10), we obtain

$$\begin{aligned} (2.7) \quad & \int_0^\infty \int_0^\infty K(x, y) f_1(x) g_1(y) dx dy \\ & \leq \left(\int_0^\infty \varphi^p(x) F(x) f_1^p(x) dx \right)^{\frac{1}{p}} \left(\int_0^\infty \psi^q(y) G(y) g_1^q(y) dy \right)^{\frac{1}{q}} \\ & = \left(\int_0^\infty \int_0^x \varphi^p(x) F(x) |f'(\tau)|^p d\tau dx \right)^{\frac{1}{p}} \\ & \quad \times \left(\int_0^\infty \int_0^y \psi^q(y) G(y) |g'(\delta)|^q d\delta dy \right)^{\frac{1}{q}}. \end{aligned}$$

By using (2.6) and (2.7) we obtain (2.1). The second inequality (2.2) can be proved by applying (1.11) and the inequality


$$|f(x)| \leq x^{\frac{1}{q}} \left(\int_0^x |f'(t)|^p dt \right)^{\frac{1}{p}}.$$

□

Now we can apply our main result to non-negative homogeneous functions. Recall that for a homogeneous function of degree $-s$, $s > 0$, the equality $K(tx, ty) = t^{-s} K(x, y)$ is satisfied. Further, we define

$$k(\alpha) := \int_0^\infty K(1, u) u^{-\alpha} du$$

and suppose that $k(\alpha) < \infty$ for $1 - s < \alpha < 1$. To prove first application of our main results we need the following lemma.



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 11 of 27

[Go Back](#)

[Full Screen](#)

[Close](#)

Lemma 2.2. *If $s > 0$, $1 - s < \alpha < 1$ and $K(x, y)$ is a non-negative homogeneous function of degree $-s$, then*

$$(2.8) \quad \int_0^\infty K(x, y) \left(\frac{x}{y}\right)^\alpha dy = x^{1-s} k(\alpha).$$

Proof. By using the substitution $u = \frac{y}{x}$ and the fact that $K(x, y)$ is homogeneous function, the equation (2.8) follows easily. \square

Corollary 2.3. *Let $s > 0$, $\frac{1}{p} + \frac{1}{q} = 1$ with $p > 1$. If $f(x)$, $g(y)$ are absolutely continuous functions such that $f(0) = g(0) = 0$, and $K(x, y)$ is a non-negative symmetrical and homogeneous function of degree $-s$, then the following inequalities hold*

$$(2.9) \quad \int_0^\infty \int_0^\infty \frac{K(x, y) |f(x)| |g(y)|}{qx^{p-1} + py^{q-1}} dx dy$$

$$\leq \int_0^\infty \int_0^\infty K(x, y) |f(x)| |g(y)| d\left(x^{\frac{1}{p}}\right) d\left(y^{\frac{1}{q}}\right)$$

$$\leq \frac{L}{pq} \left(\int_0^\infty \int_0^x x^{1-s+p(A_1-A_2)} |f'(\tau)|^p d\tau dx \right)^{\frac{1}{p}}$$

$$\times \left(\int_0^\infty \int_0^y y^{1-s+q(A_2-A_1)} |g'(\delta)|^q d\delta dy \right)^{\frac{1}{q}}$$

and

$$(2.10) \quad \int_0^\infty y^{(p-1)(s-1)+p(A_1-A_2)} \left(\int_0^\infty K(x, y) |f(x)| d\left(x^{\frac{1}{p}}\right) \right)^p dy$$

$$\leq \left(\frac{L}{p}\right)^p \int_0^\infty \int_0^x x^{1-s+p(A_1-A_2)} |f'(\tau)|^p d\tau dx,$$



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 12 of 27

[Go Back](#)

[Full Screen](#)

[Close](#)

where $A_1 \in (\frac{1-s}{q}, \frac{1}{q})$, $A_2 \in (\frac{1-s}{p}, \frac{1}{p})$ and $L = k(pA_2)^{\frac{1}{p}}k(qA_1)^{\frac{1}{q}}$.

Proof. Let $F(x)$, $G(y)$ be the functions defined as in (1.12). Setting $\varphi(x) = x^{A_1}$ and $\psi(y) = y^{A_2}$, by Lemma 2.2 we obtain

$$\begin{aligned}(2.11) \quad & \int_0^\infty \int_0^x \varphi^p(x) F(x) |f'|^p d\tau dx \\ &= \int_0^\infty \int_0^x |f'(\tau)|^p \left(\int_0^\infty K(x, y) \left(\frac{x}{y}\right)^{pA_2} dy \right) x^{p(A_1-A_2)} d\tau dx \\ &= k(pA_2) \int_0^\infty \int_0^x x^{1-s+p(A_1-A_2)} |f'(\tau)|^p d\tau dx,\end{aligned}$$


and similarly

$$\begin{aligned}(2.12) \quad & \int_0^\infty \int_0^y \psi^q(y) G(y) |g'(\delta)|^q d\delta dy \\ &= k(qA_1) \int_0^\infty \int_0^y y^{1-s+q(A_2-A_1)} |g'(\delta)|^q d\delta dy.\end{aligned}$$

From (2.1), (2.11) and (2.12), we get (2.9). Similarly, the inequality (2.10) follows from (2.2). \square


We proceed with some special homogeneous functions. First, by putting $K(x, y) = \frac{1}{(x+y)^s}$ in Corollary 2.3, we get the following.

Corollary 2.4. *Let $s > 0$, $\frac{1}{p} + \frac{1}{q} = 1$ with $p > 1$. If $f(x)$, $g(y)$ are absolutely continuous functions such that $f(0) = g(0) = 0$, then the following inequalities*



[Title Page](#)

[Contents](#)



Page 13 of 27

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

hold

$$\begin{aligned} & \int_0^\infty \int_0^\infty \frac{|f(x)| |g(y)|}{(qx^{p-1} + py^{q-1})(x+y)^s} dx dy \\ & \leq \int_0^\infty \int_0^\infty \frac{|f(x)| |g(y)|}{(x+y)^s} d\left(x^{\frac{1}{p}}\right) d\left(y^{\frac{1}{q}}\right) \\ & \leq \frac{L_1}{pq} \left(\int_0^\infty \int_0^x x^{1-s+p(A_1-A_2)} |f'(\tau)|^p d\tau dx \right)^{\frac{1}{p}} \\ & \quad \times \left(\int_0^\infty \int_0^y y^{1-s+q(A_2-A_1)} |g'(\delta)|^q d\delta dy \right)^{\frac{1}{q}} \end{aligned}$$

and

$$\begin{aligned} & \int_0^\infty y^{(p-1)(s-1)+p(A_1-A_2)} \left(\int_0^\infty \frac{|f(x)|}{(x+y)^s} d\left(x^{\frac{1}{p}}\right) \right)^p dy \\ & \leq \left(\frac{L_1}{p} \right)^p \int_0^\infty \int_0^x x^{1-s+p(A_1-A_2)} |f'(\tau)|^p d\tau dx, \end{aligned}$$

where $A_1 \in (\frac{1-s}{q}, \frac{1}{q})$, $A_2 \in (\frac{1-s}{p}, \frac{1}{p})$ and

$$L_1 = [B(1 - pA_2, pA_2 + s - 1)]^{\frac{1}{p}} [B(1 - qA_1, qA_1 + s - 1)]^{\frac{1}{q}}.$$



Remark 1. By putting $A_1 = A_2 = \frac{2-s}{pq}$ in Corollary 2.4, with the condition $s > 2 - \min\{p, q\}$, we obtain Theorem A from the introduction.

Since the function $K(x, y) = \frac{\ln \frac{y}{x}}{y-x}$ is symmetrical and homogeneous of degree -1 , by using Corollary 2.3 we obtain:



Title Page

Contents



Page 14 of 27

Go Back

Full Screen

Close

Corollary 2.5. Let $\frac{1}{p} + \frac{1}{q} = 1$ with $p > 1$. If $f(x)$, $g(y)$ are absolutely continuous functions such that $f(0) = g(0) = 0$, then the following inequalities hold

$$\begin{aligned} & \int_0^\infty \int_0^\infty \frac{\ln \frac{y}{x} |f(x)| |g(y)|}{(qx^{p-1} + py^{q-1})(y-x)} dx dy \\ & \leq \int_0^\infty \int_0^\infty \frac{\ln \frac{y}{x} |f(x)| |g(y)|}{y-x} d\left(x^{\frac{1}{p}}\right) d\left(y^{\frac{1}{q}}\right) \\ & \leq \frac{L_2}{pq} \left(\int_0^\infty \int_0^x x^{p(A_1-A_2)} |f'(\tau)|^p d\tau dx \right)^{\frac{1}{p}} \left(\int_0^\infty \int_0^y y^{q(A_2-A_1)} |g'(\delta)|^q d\delta dy \right)^{\frac{1}{q}} \end{aligned}$$

and

$$\int_0^\infty y^{p(A_1-A_2)} \left(\int_0^\infty \frac{|f(x)| \ln \frac{y}{x}}{y-x} d\left(x^{\frac{1}{p}}\right) \right)^p dy \leq \left(\frac{L_2}{p} \right)^p \int_0^\infty \int_0^x x^{p(A_1-A_2)} |f'(\tau)|^p d\tau dx,$$

where $A_1 \in (0, \frac{1}{q})$, $A_2 \in (0, \frac{1}{p})$ and

$$L_2 = \pi^2 (\sin pA_2\pi)^{-\frac{2}{p}} (\sin qA_1\pi)^{-\frac{2}{q}}.$$


Similarly, for the symmetrical homogeneous function of degree $-s$, $K(x, y) = \frac{1}{\max\{x, y\}^s}$, we have:

Corollary 2.6. Let $s > 0$, $\frac{1}{p} + \frac{1}{q} = 1$ with $p > 1$. If $f(x)$, $g(y)$ are absolutely continuous functions such that $f(0) = g(0) = 0$, then the following inequalities



Title Page

Contents



Page 15 of 27

Go Back

Full Screen

Close

hold

$$\begin{aligned} & \int_0^\infty \int_0^\infty \frac{|f(x)| |g(y)|}{(qx^{p-1} + py^{q-1}) \max\{x, y\}^s} dx dy \\ & \leq \int_0^\infty \int_0^\infty \frac{|f(x)| |g(y)|}{\max\{x, y\}^s} d\left(x^{\frac{1}{p}}\right) d\left(y^{\frac{1}{q}}\right) \\ & \leq \frac{L_3}{pq} \left(\int_0^\infty \int_0^x x^{1-s+p(A_1-A_2)} |f'(\tau)|^p d\tau dx \right)^{\frac{1}{p}} \\ & \quad \times \left(\int_0^\infty \int_0^y y^{1-s+q(A_2-A_1)} |g'(\delta)|^q d\delta dy \right)^{\frac{1}{q}} \end{aligned}$$


and

$$\begin{aligned} & \int_0^\infty y^{(p-1)(s-1)+p(A_1-A_2)} \left(\int_0^\infty \frac{|f(x)|}{\max\{x, y\}^s} d\left(x^{\frac{1}{p}}\right) \right)^p dy \\ & \leq \left(\frac{L_3}{p} \right)^p \int_0^\infty \int_0^x x^{1-s+p(A_1-A_2)} |f'(\tau)|^p d\tau dx, \end{aligned}$$

where $A_1 \in \left(\frac{1-s}{q}, \frac{1}{q}\right)$, $A_2 \in \left(\frac{1-s}{p}, \frac{1}{p}\right)$ and $L_3 = k(pA_2)^{\frac{1}{p}} k(qA_1)^{\frac{1}{q}}$, where $k(\alpha) = \frac{s}{(1-\alpha)(s+\alpha-1)}$.

At the end of this section we give a generalization of the inequality (2.1) from Theorem 2.1. In the proof we used a general Hilbert-type inequality (1.13) of Brnetić et al., [8].

Theorem 2.7. Let $n \in \mathbb{N}$, $n \geq 2$, $\sum_{i=1}^n \frac{1}{p_i} = 1$ with $p_i > 1$, $i = 1, \dots, n$. Let q_i , α_i , $i = 1, \dots, n$, are defined with $\frac{1}{q_i} = 1 - \frac{1}{p_i}$ and $\alpha_i = \prod_{j=1, j \neq i}^n p_j$. If $K(x_1, \dots, x_n)$, $\phi_{ij}(x_j)$, $i, j = 1, \dots, n$, are non-negative functions such that $\prod_{i,j=1}^n \phi_{ij}(x_j) = 1$,



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 16 of 27

Go Back


Full Screen

Close

and $f_i(x_i)$, $i = 1, \dots, n$, are absolutely continuous functions such that $f_i(0) = 0$, $i = 1, \dots, n$, then the following inequality holds

$$\begin{aligned} & \int_0^\infty \dots \int_0^\infty \frac{K(x_1, \dots, x_n) \prod_{i=1}^n |f_i(x_i)|}{\sum_{i=1}^n \alpha_i x_i^{p_i-1}} dx_1 \dots dx_n \\ & \leq \int_0^\infty \dots \int_0^\infty K(x_1, \dots, x_n) \prod_{i=1}^n |f_i(x_i)| d\left(x_1^{\frac{1}{p_1}}\right) \dots d\left(x_n^{\frac{1}{p_n}}\right) \\ & \leq \frac{1}{p_1 \dots p_n} \prod_{i=1}^n \left(\int_0^\infty \int_0^{x_i} \phi_{ii}^{p_i}(x_i) F_i(x_i) |f_i'(\tau_i)|^{p_i} d\tau_i dx_i \right)^{\frac{1}{p_i}}, \end{aligned}$$

where $F_i(x_i)$ are defined as in (1.14) for $i = 1, \dots, n$.



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 17 of 27

[Go Back](#)

[Full Screen](#)

[Close](#)

3. Discrete Case

We also give results for the discrete case. For that, we apply the following result from [5].

Theorem 3.1. *If $\{a(m)\}$ and $\{b(n)\}$ are non-negative real sequences, $K(x, y)$ is non-negative homogeneous function of degree $-s$ strictly decreasing in both parameters x and y , $\frac{1}{p} + \frac{1}{q} = 1$, $p > 1$, $A, B, \alpha, \beta > 0$, then the following inequalities hold and are equivalent*

$$(3.1) \quad \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} K(Am^{\alpha}, Bn^{\beta}) a_m b_n$$

$$< N \left(\sum_{m=1}^{\infty} m^{\alpha(1-s) + \alpha p(A_1 - A_2) + (p-1)(1-\alpha)} a_m^p \right)^{\frac{1}{p}}$$

$$\cdot \left(\sum_{n=1}^{\infty} n^{\beta(1-s) + \beta q(A_2 - A_1) + (q-1)(1-\beta)} b_n^q \right)^{\frac{1}{q}},$$


and

$$(3.2) \quad \sum_{n=1}^{\infty} n^{\beta(s-1)(p-1) + p\beta(A_1 - A_2) + \beta - 1} \left(\sum_{m=1}^{\infty} K(Am^{\alpha}, Bn^{\beta}) a_m \right)^p$$

$$< N^p \sum_{m=1}^{\infty} m^{\alpha(1-s) + \alpha p(A_1 - A_2) + (p-1)(1-\alpha)} a_m^p,$$

where $A_1 \in (\max\{\frac{1-s}{q}, \frac{\alpha-1}{\alpha q}\}, \frac{1}{q})$, $A_2 \in (\max\{\frac{1-s}{p}, \frac{\beta-1}{\beta p}\}, \frac{1}{p})$ and

$$(3.3) \quad N = \alpha^{-\frac{1}{q}} \beta^{-\frac{1}{p}} A^{\frac{2-s}{p} + A_1 - A_2 - 1} B^{\frac{2-s}{q} + A_2 - A_1 - 1} k(pA_2)^{\frac{1}{p}} k(2-s-qA_1)^{\frac{1}{q}}.$$



Title Page

Contents



Page 18 of 27

Go Back

Full Screen

Close

Applying Theorem 3.1, we obtain the following.


Corollary 3.2. Let $s > 0$, $\frac{1}{p} + \frac{1}{q} = 1$ with $p > 1$. Let $\{a(m)\}$ and $\{b(n)\}$ be two sequences of real numbers where $m, n \in \mathbb{N}_0$, and $a(0) = b(0) = 0$. If $K(x, y)$ is a non-negative homogeneous function of degree $-s$ strictly decreasing in both parameters x and y , $A, B, \alpha, \beta > 0$, then the following inequalities hold

$$\begin{aligned}
 (3.4) \quad & \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{K(Am^{\alpha}, Bn^{\beta})|a_m||b_n|}{qm^{p-1} + pn^{q-1}} \\
 & \leq \frac{1}{pq} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{K(Am^{\alpha}, Bn^{\beta})|a_m||b_n|}{m^{\frac{1}{q}}n^{\frac{1}{p}}} \\
 & < \frac{N}{pq} \left(\sum_{m=1}^{\infty} \sum_{\tau=1}^m m^{\alpha(1-s)+\alpha p(A_1-A_2)+(p-1)(1-\alpha)} |\nabla a(\tau)|^p \right)^{\frac{1}{p}} \\
 & \quad \cdot \left(\sum_{n=1}^{\infty} \sum_{\delta=1}^n n^{\beta(1-s)+\beta q(A_2-A_1)+(q-1)(1-\beta)} |\nabla b(\delta)|^q \right)^{\frac{1}{q}},
 \end{aligned}$$

and


$$\begin{aligned}
 (3.5) \quad & \sum_{n=1}^{\infty} n^{\beta(s-1)(p-1)+p\beta(A_1-A_2)+\beta-1} \left(\sum_{m=1}^{\infty} K(Am^{\alpha}, Bn^{\beta}) \frac{|a_m|}{m^{\frac{1}{q}}} \right)^p \\
 & < N^p \sum_{m=1}^{\infty} \sum_{\tau=1}^m m^{\alpha(1-s)+\alpha p(A_1-A_2)+(p-1)(1-\alpha)} |\nabla a(\tau)|^p,
 \end{aligned}$$

where $A_1 \in \left(\max \left\{ \frac{1-s}{q}, \frac{\alpha-1}{\alpha q} \right\}, \frac{1}{q} \right)$, $A_2 \in \left(\max \left\{ \frac{1-s}{p}, \frac{\beta-1}{\beta p} \right\}, \frac{1}{p} \right)$ and the constant N is defined as in (3.3).



Title Page

Contents



Page 19 of 27

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Proof. By using Hölder's inequality, (see also [9]), we have

$$(3.6) \quad |a_m| |b_n| \leq m^{\frac{1}{q}} n^{\frac{1}{p}} \left(\sum_{\tau=1}^m |\nabla a(\tau)|^p \right)^{\frac{1}{p}} \left(\sum_{\delta=1}^n |\nabla b(\delta)|^q \right)^{\frac{1}{q}}.$$

From (2.4) and (3.6), we get

$$(3.7) \quad \frac{pq|a_m| |b_n|}{qm^{p-1} + pn^{q-1}} \leq \frac{|a_m| |b_n|}{m^{\frac{1}{q}} n^{\frac{1}{p}}} \leq \left(\sum_{\tau=1}^m |\nabla a(\tau)|^p \right)^{\frac{1}{p}} \left(\sum_{\delta=1}^n |\nabla b(\delta)|^q \right)^{\frac{1}{q}},$$

and therefore

$$(3.8) \quad pq \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{K(Am^{\alpha}, Bn^{\beta}) |a_m| |b_n|}{qm^{p-1} + pn^{q-1}} \\ \leq \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{K(Am^{\alpha}, Bn^{\beta}) |a_m| |b_n|}{m^{\frac{1}{q}} n^{\frac{1}{p}}} \\ \leq \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} K(Am^{\alpha}, Bn^{\beta}) \left(\sum_{\tau=1}^m |\nabla a(\tau)|^p \right)^{\frac{1}{p}} \left(\sum_{\delta=1}^n |\nabla b(\delta)|^q \right)^{\frac{1}{q}}.$$

Applying the substitutions $\tilde{a}_m = \left(\sum_{\tau=1}^m |\nabla a(\tau)|^p \right)^{\frac{1}{p}}$, $\tilde{b}_n = \left(\sum_{\delta=1}^n |\nabla b(\delta)|^q \right)^{\frac{1}{q}}$ and (3.1), we have

$$(3.9) \quad \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} K(Am^{\alpha}, Bn^{\beta}) \tilde{a}_m \tilde{b}_n$$



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 20 of 27

[Go Back](#)

[Full Screen](#)

[Close](#)

$$\begin{aligned}
 &< N \left(\sum_{m=1}^{\infty} m^{\alpha(1-s)+\alpha p(A_1-A_2)+(p-1)(1-\alpha)} \tilde{a}_m^p \right)^{\frac{1}{p}} \\
 &\cdot \left(\sum_{n=1}^{\infty} n^{\beta(1-s)+\beta q(A_2-A_1)+(q-1)(1-\beta)} \tilde{b}_n^q \right)^{\frac{1}{q}}, \\
 &= \left(\sum_{m=1}^{\infty} \sum_{\tau=1}^m m^{\alpha(1-s)+\alpha p(A_1-A_2)+(p-1)(1-\alpha)} |\nabla a(\tau)|^p \right)^{\frac{1}{p}} \\
 &\cdot \left(\sum_{n=1}^{\infty} \sum_{\delta=1}^n n^{\beta(1-s)+\beta q(A_2-A_1)+(q-1)(1-\beta)} |\nabla b(\delta)|^q \right)^{\frac{1}{q}},
 \end{aligned}$$


where $A_1 \in (\max\{\frac{1-s}{q}, \frac{\alpha-1}{\alpha q}\}, \frac{1}{q})$, $A_2 \in (\max\{\frac{1-s}{p}, \frac{\beta-1}{\beta p}\}, \frac{1}{p})$ and the constant N is defined as in (3.3). Now, by applying (3.8) and (3.9) we obtain (3.4). The second inequality (3.5) can be proved by using (3.2) and the inequality

$$|a_m| \leq m^{\frac{1}{q}} \left(\sum_{\tau=1}^m |\nabla a(\tau)|^p \right)^{\frac{1}{p}}.$$

□


Remark 2. If the function $K(x, y)$ from the previous corollary is symmetrical, then $k(2-s-qA_1) = k(qA_1)$. So, if $K(x, y) = \frac{1}{(x+y)^s}$, then we can put $A_1 = A_2 = \frac{2-s}{pq}$, $A = B = \alpha = \beta = 1$ in Corollary 3.2 and obtain Theorem B from the introduction but with the condition $2 - \min\{p, q\} < s < 2$.

By using (1.4), see [4], we will obtain a larger interval for the parameter s . More precisely, we have:



Title Page

Contents



Page 21 of 27

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

Corollary 3.3. Let $\frac{1}{p} + \frac{1}{q} = 1$ with $p > 1$ and $2 - \min\{p, q\} < s \leq 2 + \min\{p, q\}$. Let $\{a(m)\}$ and $\{b(n)\}$ be two sequences of real numbers where $m, n \in \mathbb{N}_0$, and $a(0) = b(0) = 0$. Then the following inequalities hold

$$(3.10) \quad \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{|a_m| |b_n|}{(qm^{p-1} + pn^{q-1})(m+n)^s} \\ \leq \frac{1}{pq} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{|a_m| |b_n|}{m^{\frac{1}{q}} n^{\frac{1}{p}} (m+n)^s} \\ < \frac{N_1}{pq} \left(\sum_{m=1}^{\infty} \sum_{\tau=1}^m m^{1-s} |\nabla a(\tau)|^p \right)^{\frac{1}{p}} \cdot \left(\sum_{n=1}^{\infty} \sum_{\delta=1}^n n^{1-s} |\nabla b(\delta)|^q \right)^{\frac{1}{q}},$$

and

$$(3.11) \quad \sum_{n=1}^{\infty} n^{(s-1)(p-1)} \left(\sum_{m=1}^{\infty} \frac{|a_m|}{m^{\frac{1}{q}} (m+n)^s} \right)^p < N_1^p \sum_{m=1}^{\infty} \sum_{\tau=1}^m m^{1-s} |\nabla a(\tau)|^p,$$

where $N_1 = B\left(\frac{s+q-2}{q}, \frac{s+p-2}{p}\right)$.

Proof. As in the proof of Corollary 3.2, by using Hölder's inequality we obtain

$$\sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{|a_m| |b_n|}{(qm^{p-1} + pn^{q-1})(m+n)^s} \\ \leq \frac{1}{pq} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{|a_m| |b_n|}{m^{\frac{1}{q}} n^{\frac{1}{p}} (m+n)^s} \\ \leq \frac{1}{pq} \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{1}{(m+n)^s} \left(\sum_{\tau=1}^m |\nabla a(\tau)|^p \right)^{\frac{1}{p}} \left(\sum_{\delta=1}^n |\nabla b(\delta)|^q \right)^{\frac{1}{q}} \left(\frac{m}{n} \right)^{\frac{2-s}{pq}} \left(\frac{n}{m} \right)^{\frac{2-s}{pq}}$$



Title Page

Contents

◀◀ ▶▶

◀ ▶

Page 22 of 27

Go Back

Full Screen

Close

$$\leq \frac{1}{pq} \left(\sum_{m=1}^{\infty} \sum_{\tau=1}^m \left(\sum_{n=1}^{\infty} \frac{1}{(m+n)^s} \left(\frac{m}{n} \right)^{\frac{2-s}{q}} \right) |\nabla a(\tau)|^p \right)^{\frac{1}{p}} \\ \cdot \left(\sum_{n=1}^{\infty} \sum_{\delta=1}^n \left(\sum_{m=1}^{\infty} \frac{1}{(m+n)^s} \left(\frac{n}{m} \right)^{\frac{2-s}{p}} \right) |\nabla b(\delta)|^q \right)^{\frac{1}{q}}.$$

Now, the inequality (3.10) follows from (1.4). Let us show that the inequality (3.11) is valid. For this purpose we use the following inequality from [4]

$$(3.12) \quad \sum_{n=1}^{\infty} n^{(s-1)(p-1)} \left(\sum_{m=1}^{\infty} \frac{a_m}{(m+n)^s} \right)^p < L_1 \sum_{m=1}^{\infty} m^{1-s} a_m^p,$$

where $2 - \min\{p, q\} < s \leq 2 + \min\{p, q\}$ and $L_1 = B\left(\frac{s+p-2}{p}, \frac{s+q-2}{q}\right)$. Setting

$$a_m = \left(\sum_{\tau=1}^m |\nabla a(\tau)|^p \right)^{\frac{1}{p}}$$

in (3.12) and using


$$|a_m| \leq m^{\frac{1}{q}} \left(\sum_{\tau=1}^m |\nabla a(\tau)|^p \right)^{\frac{1}{p}},$$

the inequality (3.11) follows easily. □



[Title Page](#)

[Contents](#)



Page 23 of 27

[Go Back](#)

[Full Screen](#)

[Close](#)

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

4. Non-conjugate Exponents

Let p , p' , q , q' and λ be as in (1.6) and (1.7). To obtain an analogous result for the case of non-conjugate exponents, we introduce real parameters r' , r such that $p \leq r' \leq q'$ and $\frac{1}{r'} + \frac{1}{r} = 1$. For example, we can define $\frac{1}{r'} = \frac{1}{q'} + \frac{1-\lambda}{2}$ or $r' = (2-\lambda)p$.

It is easy to see that

$$(4.1) \quad x^{\frac{1}{p'}} y^{\frac{1}{q'}} \leq \frac{1}{rr'} \left(rx^{\frac{r'}{p'}} + r'y^{\frac{r}{q'}} \right), \quad x \geq 0, \quad y \geq 0,$$

and


$$(4.2) \quad |f(x)| |g(y)| \leq x^{\frac{1}{p'}} y^{\frac{1}{q'}} \left(\int_0^x |f'(\tau)|^p d\tau \right)^{\frac{1}{p}} \left(\int_0^y |g'(\delta)|^q d\delta \right)^{\frac{1}{q}},$$

hold, where $f(x)$, $g(y)$ are absolutely continuous functions on $(0, \infty)$.

Applying Theorem C, (4.1) and (4.2) in the same way as in the proof of Theorem 2.1, we obtain the following result for non-conjugate exponents.

Theorem 4.1. *Let p , p' , q , q' and λ be as in (1.6) and (1.7). Let r' , r be real parameters such that $p \leq r' \leq q'$ and $\frac{1}{r'} + \frac{1}{r} = 1$. If $K(x, y)$, $\varphi(x)$, $\psi(y)$ are non-negative functions and $f(x)$, $g(y)$ are absolutely continuous functions such that $f(0) = g(0) = 0$, then the following inequalities hold*

$$(4.3) \quad \int_0^\infty \int_0^\infty \frac{K^\lambda(x, y) |f(x)| |g(y)|}{rx^{\frac{r'}{p'}} + r'y^{\frac{r}{q'}}} dx dy \\ \leq \frac{pq}{rr'} \int_0^\infty \int_0^\infty K^\lambda(x, y) |f(x)| |g(y)| d \left(x^{\frac{1}{p'}} \right) d \left(y^{\frac{1}{q'}} \right)$$



[Title Page](#)

[Contents](#)

◀◀ ▶▶

◀ ▶

Page 24 of 27

[Go Back](#)

[Full Screen](#)

[Close](#)

$$\leq \frac{1}{rr'} \left(\int_0^\infty \int_0^x (\varphi F)^p(x) |f'(\tau)|^p d\tau dx \right)^{\frac{1}{p}} \\ \times \left(\int_0^\infty \int_0^y (\psi G)^q(y) |g'(\delta)|^q d\delta dy \right)^{\frac{1}{q}},$$

and


$$(4.4) \quad \left(\int_0^\infty \left(\frac{1}{\psi G(y)} \int_0^\infty K^\lambda(x, y) |f(x)| d(x^{\frac{1}{p}}) \right)^{q'} dy \right)^{\frac{1}{q'}} \\ \leq \frac{1}{p} \left(\int_0^\infty \int_0^x (\varphi F)^p(x) |f'(\tau)|^p d\tau dx \right)^{\frac{1}{p}},$$

where $F(x)$ and $G(y)$ are defined as in Theorem C.

Obviously, Theorem 4.1 is the generalization of Theorem 2.1. Namely, if $\lambda = 1$, $r' = p$ and $r = q$, then the inequalities (4.3) and (4.4) become respectively the inequalities (2.1) and (2.2). If $K(x, y)$ is a non-negative symmetrical and homogeneous function of degree $-s$, $s > 0$, then we obtain:



Corollary 4.2. *Let $s > 0$, p, p', q, q' and λ be as in (1.6) and (1.7). If $f(x), g(y)$ are absolutely continuous functions such that $f(0) = g(0) = 0$, and $K(x, y)$ is a non-negative symmetrical and homogeneous function of degree $-s$, then the following inequalities hold*

$$(4.5) \quad \int_0^\infty \int_0^\infty \frac{K^\lambda(x, y) |f(x)| |g(y)|}{qx^{(p-1)(2-\lambda)} + py^{(q-1)(2-\lambda)}} dx dy$$



Title Page

Contents



Page 25 of 27

Go Back

Full Screen

Close

$$\begin{aligned} &\leq \frac{1}{2-\lambda} \int_0^\infty \int_0^\infty K^\lambda(x, y) |f(x)| |g(y)| d\left(x^{\frac{1}{p}}\right) d\left(y^{\frac{1}{q}}\right) \\ &\leq \frac{M}{pq(2-\lambda)} \left(\int_0^\infty \int_0^x x^{\frac{p}{q'}(1-s)+p(A_1-A_2)} |f'(\tau)|^p d\tau dx \right)^{\frac{1}{p}} \\ &\quad \cdot \left(\int_0^\infty \int_0^y y^{\frac{q}{p'}(1-s)+q(A_2-A_1)} |g'(\delta)|^q d\delta dy \right)^{\frac{1}{q}} \end{aligned}$$

and

$$(4.6) \quad \left(\int_0^\infty y^{\frac{q'}{p'}(s-1)+q'(A_1-A_2)} \left(\int_0^\infty K^\lambda(x, y) |f(x)| d\left(x^{\frac{1}{p}}\right) \right)^{q'} dy \right)^{\frac{1}{q'}} \\ \leq \frac{M}{p} \left(\int_0^\infty \int_0^x x^{\frac{p}{q'}(1-s)+p(A_1-A_2)} |f'(\tau)|^p d\tau dx \right)^{\frac{1}{p}},$$

where $A_1 \in \left(\frac{1-s}{p'}, \frac{1}{p'}\right)$, $A_2 \in \left(\frac{1-s}{q'}, \frac{1}{q'}\right)$ and $M = k(p'A_1)^{\frac{1}{p'}} k(q'A_2)^{\frac{1}{q'}}$.

Proof. The proof follows directly from Theorem 4.1 setting $r' = (2-\lambda)p$, $r = (2-\lambda)q$, $\varphi(x) = x^{A_1}$ and $\psi(y) = y^{A_2}$ in the inequalities (4.3) and (4.4). Namely, if $F(x)$ and $G(y)$ are the functions defined by

$$F(x) = \left(\int_0^\infty \frac{K(x, y)}{\psi^{q'}(y)} dy \right)^{\frac{1}{q'}} \quad \text{and} \quad G(y) = \left(\int_0^\infty \frac{K(x, y)}{\varphi^{p'}(x)} dx \right)^{\frac{1}{p'}},$$



then applying Lemma 2.2 we have

$$\begin{aligned}(4.7) \quad (\varphi F)^p(x) &= x^{pA_1} \left(\int_0^\infty K(x, y) y^{-q'A_2} dy \right)^{\frac{p}{q}} \\ &= x^{pA_1 - pA_2} \left(\int_0^\infty K(x, y) \left(\frac{x}{y} \right)^{q'A_2} dy \right)^{\frac{p}{q}} \\ &= x^{\frac{p}{q'}(1-s) + p(A_1 - A_2)} k^{\frac{p}{q'}}(q'A_2),\end{aligned}$$

and similarly

$$(4.8) \quad (\psi G)^q(y) = y^{\frac{q}{p'}(1-s) + q(A_2 - A_1)} k^{\frac{q}{p'}}(p'A_1).$$


Now, by using (4.3), (4.7) and (4.8) we obtain (4.5).

The second inequality (4.6) follows directly from (4.4). \square

Remark 3. Setting $K(x, y) = \frac{1}{(x+y)^s}$ in Corollary 4.2 we obtain that the constant M is equal to $M = B(1 - p'A_1, p'A_1 + s - 1)^{\frac{1}{p'}} B(1 - q'A_2, q'A_2 + s - 1)^{\frac{1}{q'}}$.

Title Page

Contents



Page 26 of 27

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756

References

- [1] F.F. BONSALL, Inequalities with non-conjugate parameters, *Quart. J. Math. Oxford Ser. (2)*, **2** (1951), 135–150.
- [2] I. BRNETIĆ, M. KRNIĆ AND J. PEČARIĆ, Multiple Hilbert's and Hardy-Hilbert's integral inequality with non-conjugate parameters, *Bull. Austral. Math. Soc.*, **71** (2005), 447–457.
- [3] G.H. HARDY, J.E. LITTLEWOOD AND G. PÓLYA, *Inequalities*, 2nd Edition, Cambridge University Press, Cambridge, 1967.
- [4] M. KRNIĆ AND J. PEČARIĆ, Extension of Hilbert's inequality, *J. Math. Anal. Appl.*, **324**(1) (2006), 150–160.
- [5] M. KRNIĆ AND J. PEČARIĆ, General Hilbert's and Hardy's Inequalities, *Math. Inequal. Appl.*, **8**(1) (2005), 29–52.
- [6] D.S. MITRINOVIĆ, J.E. PEČARIĆ AND A.M. FINK, *Classical and New Inequalities in Analysis*, Kluwer Academic Publishers, Dordrecht/Boston/London, 1993.
- [7] B.G. PACHPATTE, On some new inequalities similar to Hilbert's inequality, *J. Math. Anal. Appl.*, **226** (1998), 166–179.
- [8] B. YANG, I. BRNETIĆ, M. KRNIĆ AND J. PEČARIĆ, Generalization of Hilbert and Hardy-Hilbert integral inequalities, *Math. Inequal. Appl.*, **8**(2) (2005), 259–272.
- [9] ZHONGXUE LÜ, Some new inequalities similar to Hilbert-Pachpatte type inequalities, *J. Inequal. Pure Appl. Math.*, **4**(2) (2003), Art. 33. [ONLINE: <http://jipam.vu.edu.au/article.php?sid=271>].




Hilbert-Pachpatte Type
Inequalities

Josip Pečarić, Ivan Perić
and Predrag Vuković

vol. 9, iss. 1, art. 9, 2008

Title Page

Contents



Page 27 of 27

Go Back

Full Screen

Close

journal of **inequalities**
in pure and applied
mathematics

issn: 1443-5756