


## APPLICATIONS OF DIFFERENTIAL SUBORDINATION TO CERTAIN SUBCLASSES OF MEROMORPHICALLY MULTIVALENT FUNCTIONS

H.M. SRIVASTAVA AND J. PATEL

Department of Mathematics and Statistics  
University of Victoria  
Victoria, British Columbia V8W 3P4  
Canada.  
*EMail:* [harimsri@math.uvic.ca](mailto:harimsri@math.uvic.ca)

Department of Mathematics  
Utkal University  
Vani Vihar, Bhubaneswar 751004  
Orissa, India.  
*EMail:* [jpatelmath@sify.com](mailto:jpatelmath@sify.com)



---

volume 6, issue 3, article 88,  
2005.

*Received 18 April, 2005;  
accepted 05 July, 2005.*

*Communicated by: Th.M. Rassias*

---

Abstract

Contents



Home Page

Go Back

Close

Quit

## Abstract

By making use of the principle of differential subordination, the authors investigate several inclusion relationships and other interesting properties of certain subclasses of meromorphically multivalent functions which are defined here by means of a linear operator. They also indicate relevant connections of the various results presented in this paper with those obtained in earlier works.


*2000 Mathematics Subject Classification:* Primary 30C45; Secondary 30D30, 33C20.

*Key words:* Meromorphic functions, Differential subordination, Hadamard product (or convolution), Multivalent functions, Linear operator, Hypergeometric function.

The present investigation was supported, in part, by the *Natural Sciences and Engineering Research Council of Canada* under Grant OGP0007353 and, in part, by the *University Grants Commission of India* under its DRS Financial Assistance Program.

## Contents

1	Introduction and Definitions .....	3
2	Preliminary Lemmas .....	7
3	The Main Subordination Theorems and The Associated Functional Inequalities .....	10
	References	



---

An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

---

Title Page

Contents



Go Back

Close

Quit

Page 2 of 31

# 1. Introduction and Definitions

For any integer  $m > -p$ , let  $\Sigma_{p,m}$  denote the class of all meromorphic functions  $f(z)$  normalized by

$$(1.1) \quad f(z) = z^{-p} + \sum_{k=m}^{\infty} a_k z^k \quad (p \in \mathbb{N} := \{1, 2, 3, \dots\}),$$

which are analytic and  $p$ -valent in the punctured unit disk

$$\mathbb{U}^* = \{z : z \in \mathbb{C} \quad \text{and} \quad 0 < |z| < 1\} = \mathbb{U} \setminus \{0\}.$$

For convenience, we write

$$\Sigma_{p,-p+1} = \Sigma_p.$$

If  $f(z)$  and  $g(z)$  are analytic in  $\mathbb{U}$ , we say that  $f(z)$  is subordinate to  $g(z)$ , written symbolically as follows:

$$f \prec g \quad \text{in } \mathbb{U} \quad \text{or} \quad f(z) \prec g(z) \quad (z \in \mathbb{U}),$$

if there exists a Schwarz function  $w(z)$ , which (by definition) is analytic in  $\mathbb{U}$  with

$$w(0) = 0 \quad \text{and} \quad |w(z)| < 1 \quad (z \in \mathbb{U})$$

such that

$$f(z) = g(w(z)) \quad (z \in \mathbb{U}).$$

Indeed it is known that

$$f(z) \prec g(z) \quad (z \in \mathbb{U}) \implies f(0) = g(0) \quad \text{and} \quad f(\mathbb{U}) \subset g(\mathbb{U}).$$



---

An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

---

Title Page

Contents



Go Back

Close

Quit

Page 3 of 31

In particular, if the function  $g(z)$  is univalent in  $\mathbb{U}$ , we have the following equivalence (cf., e.g., [5]; see also [6, p. 4]):

$$f(z) \prec g(z) \quad (z \in \mathbb{U}) \quad \iff \quad f(0) = g(0) \quad \text{and} \quad f(\mathbb{U}) \subset g(\mathbb{U}).$$

For functions  $f(z) \in \Sigma_{p,m}$ , given by (1.1), and  $g(z) \in \Sigma_{p,m}$  defined by

$$(1.2) \quad g(z) = z^{-p} + \sum_{k=m}^{\infty} b_k z^k \quad (m > -p; p \in \mathbb{N}),$$

we define the Hadamard product (or convolution) of  $f(z)$  and  $g(z)$  by


$$(1.3) \quad (f \star g)(z) := z^{-p} + \sum_{k=m}^{\infty} a_k b_k z^k =: (g \star f)(z) \\ (m > -p; p \in \mathbb{N}; z \in \mathbb{U}).$$

Following the recent work of Liu and Srivastava [3], for a function  $f(z)$  in the class  $\Sigma_{p,m}$ , given by (1.1), we now define a linear operator  $D^n$  by

$$D^0 f(z) = f(z), \\ D^1 f(z) = z^{-p} + \sum_{k=m}^{\infty} (p+k+1) a_k z^k = \frac{(z^{p+1} f(z))'}{z^p},$$

and (in general)

$$(1.4) \quad D^n f(z) = D(D^{n-1} f(z)) = z^{-p} + \sum_{k=m}^{\infty} (p+k+1)^n a_k z^k \\ = \frac{(z^{p+1} D^{n-1} f(z))'}{z^p} \quad (n \in \mathbb{N}).$$



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 4 of 31

It is easily verified from (1.4) that

$$(1.5) \quad z(D^n f(z))' = D^{n+1} f(z) - (p+1)D^n f(z) \\ \left( f \in \Sigma_{p,m}; n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\} \right).$$

The case  $m = 0$  of the linear operator  $D^n$  was introduced recently by Liu and Srivastava [3], who investigated (among other things) several inclusion relationships involving various subclasses of meromorphically  $p$ -valent functions, which they defined by means of the linear operator  $D^n$  (see also [2]). A special case of the linear operator  $D^n$  for  $p = 1$  and  $m = 0$  was considered earlier by Uralegaddi and Somanatha [13]. Aouf and Hossen [1] also obtained several results involving the operator  $D^n$  for  $m = 0$  and  $p \in \mathbb{N}$ .


Making use of the principle of differential subordination as well as the linear operator  $D^n$ , we now introduce a subclass of the function class  $\Sigma_{p,m}$  as follows.

**Definition.** For fixed parameters  $A$  and  $B$  ( $-1 \leq B < A \leq 1$ ), we say that a function  $f(z) \in \Sigma_{p,m}$  is in the class  $\Sigma_{p,m}^n(A, B)$ , if it satisfies the following subordination condition:

$$(1.6) \quad -\frac{z^{p+1}(D^n f(z))'}{p} \prec \frac{1 + Az}{1 + Bz} \quad (n \in \mathbb{N}_0; z \in \mathbb{U}).$$

In view of the definition of differential subordination, (1.6) is equivalent to the following condition:

$$\left| \frac{z^{p+1}(D^n f(z))' + p}{Bz^{p+1}(D^n f(z))' + pA} \right| < 1 \quad (z \in \mathbb{U}).$$




---

An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

---

Title Page

Contents



Go Back

Close

Quit

Page 5 of 31

For convenience, we write

$$\Sigma_{p,m}^n \left( 1 - \frac{2\alpha}{p}, -1 \right) = \Sigma_{p,m}^n(\alpha),$$

where  $\Sigma_{p,m}^n(\alpha)$  denotes the class of functions in  $\Sigma_{p,m}$  satisfying the following inequality:


$$\Re \left( -z^{p+1} (D^n f(z))' \right) > \alpha \quad (0 \leq \alpha < p; z \in \mathbb{U}).$$

In particular, we have

$$\Sigma_{p,0}^n(A, B) = \mathcal{R}_{n,p}(A, B),$$

where  $\mathcal{R}_{n,p}(A, B)$  is the function class introduced and studied by Liu and Srivastava [3]. The function class  $\mathcal{H}(p; A, B)$ , considered by Mogra [7], happens to be a *further* special case of the Liu-Srivastava class  $\mathcal{R}_{n,p}(A, B)$  when  $n = 0$ .

In the present paper, we derive several inclusion relationships for the function class  $\Sigma_{p,m}^n(A, B)$  and investigate various other properties of functions belonging to the class  $\Sigma_{p,m}^n(A, B)$ , which we have defined here by means of the linear operator  $D^n$ . These include (for example) some mapping properties involving the linear operator  $D^n$ . Relevant connections of the results presented in this paper with those obtained in earlier works are also pointed out.




---

**An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem**

H.M. Srivastava and J. Patel

---

Title Page

Contents



Go Back

Close

Quit

Page 6 of 31

## 2. Preliminary Lemmas

In proving our main results, we need each of the following lemmas.

**Lemma 1 (Miller and Mocanu [5]; see also [6]).** *Let the function  $h(z)$  be analytic and convex (univalent) in  $\mathbb{U}$  with  $h(0) = 1$ . Suppose also that the function  $\phi(z)$  given by*

$$(2.1) \quad \phi(z) = 1 + c_{p+m}z^{p+m} + c_{p+m+1}z^{p+m+1} + \dots$$

is analytic in  $\mathbb{U}$ . If

$$(2.2) \quad \phi(z) + \frac{z\phi'(z)}{\gamma} \prec h(z) \quad (\Re(\gamma) \geq 0; \gamma \neq 0; z \in \mathbb{U}),$$

then

$$\phi(z) \prec \psi(z) = \frac{\gamma}{p+m} z^{-\frac{\gamma}{p+m}} \int_0^z t^{\frac{\gamma}{p+m}-1} h(t) dt \prec h(z) \quad (z \in \mathbb{U}),$$


and  $\psi(z)$  is the best dominant of (2.2).

With a view to stating a well-known result (Lemma 2 below), we denote by  $\mathcal{P}(\gamma)$  the class of functions  $\varphi(z)$  given by

$$(2.3) \quad \varphi(z) = 1 + b_1z + b_2z^2 + \dots,$$

which are analytic in  $\mathbb{U}$  and satisfy the following inequality:

$$\Re(\varphi(z)) > \gamma \quad (0 \leq \gamma < 1; z \in \mathbb{U}).$$



---

An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

---

Title Page

Contents



Go Back

Close

Quit

Page 7 of 31

**Lemma 2 (cf., e.g., Pashkouleva [9]).** Let the function  $\varphi(z)$ , given by (2.3), be in the class  $\mathcal{P}(\gamma)$ . Then

$$\Re\{\varphi(z)\} \geq 2\gamma - 1 + \frac{2(1 - \gamma)}{1 + |z|} \quad (0 \leq \gamma < 1; z \in \mathbb{U}).$$

**Lemma 3 (see [12]).** For  $0 \leq \gamma_1, \gamma_2 < 1$ ,

$$\mathcal{P}(\gamma_1) \star \mathcal{P}(\gamma_2) \subset \mathcal{P}(\gamma_3) \quad (\gamma_3 := 1 - 2(1 - \gamma_1)(1 - \gamma_2)).$$

The result is the best possible.

For real or complex numbers  $a, b$ , and  $c$  ( $c \notin \mathbb{Z}_0^- := \{0, -1, -2, \dots\}$ ), the Gauss hypergeometric function is defined by


$${}_2F_1(a, b; c; z) = 1 + \frac{ab}{c} \cdot \frac{z}{1!} + \frac{a(a+1)b(b+1)}{c(c+1)} \cdot \frac{z^2}{2!} + \dots$$

We note that the above series converges absolutely for  $z \in \mathbb{U}$  and hence represents an analytic function in  $\mathbb{U}$  (see, for details, [14, Chapter 14]).

Each of the identities (asserted by Lemma 4 below) is well-known (cf., e.g., [14, Chapter 14]).

**Lemma 4.** For real or complex parameters  $a, b$ , and  $c$  ( $c \notin \mathbb{Z}_0^-$ ),

$$(2.4) \quad \int_0^1 t^{b-1}(1-t)^{c-b-1}(1-zt)^{-a} dt = \frac{\Gamma(b)\Gamma(c-b)}{\Gamma(c)} {}_2F_1(a, b; c; z) \quad (\Re(c) > \Re(b) > 0);$$



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 8 of 31

$$(2.5) \quad {}_2F_1(a, b; c; z) = (1 - z)^{-a} {}_2F_1\left(a, c - b; c; \frac{z}{z - 1}\right);$$

$$(2.6) \quad {}_2F_1(a, b; c; z) = {}_2F_1(a, b - 1; c; z) + \frac{az}{c} {}_2F_1(a + 1, b; c + 1; z);$$


$$(2.7) \quad {}_2F_1\left(a, b; \frac{a + b + 1}{2}; \frac{1}{2}\right) = \frac{\sqrt{\pi} \Gamma\left(\frac{a + b + 1}{2}\right)}{\Gamma\left(\frac{a + 1}{2}\right) \Gamma\left(\frac{b + 1}{2}\right)}.$$

We now recall a result due to Singh and Singh [11] as Lemma 5 below.

**Lemma 5.** *Let  $\Phi(z)$  be analytic in  $\mathbb{U}$  with*

$$\Phi(0) = 1 \quad \text{and} \quad \Re(\Phi(z)) > \frac{1}{2} \quad (z \in \mathbb{U}).$$

*Then, for any function  $F(z)$  analytic in  $\mathbb{U}$ ,  $(\Phi \star F)(\mathbb{U})$  is contained in the convex hull of  $F(\mathbb{U})$ .*



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 9 of 31

### 3. The Main Subordination Theorems and The Associated Functional Inequalities

Unless otherwise mentioned, we shall assume *throughout the sequel* that  $m$  is an integer greater than  $-p$ , and that

$$-1 \leq B < A \leq 1, \lambda > 0, n \in \mathbb{N}_0, \text{ and } p \in \mathbb{N}.$$

**Theorem 1.** Let the function  $f(z)$  defined by (1.1) satisfy the following subordination condition:

$$-\frac{(1-\lambda)z^{p+1}(D^n f(z))' + \lambda z^{p+1}(D^{n+1} f(z))'}{p} \prec \frac{1+Az}{1+Bz} \quad (z \in \mathbb{U}).$$

Then


$$(3.1) \quad -\frac{z^{p+1}(D^n f(z))'}{p} \prec Q(z) \prec \frac{1+Az}{1+Bz} \quad (z \in \mathbb{U}),$$

where the function  $Q(z)$  given by

$$Q(z) = \begin{cases} \frac{A}{B} + \left(1 - \frac{A}{B}\right) (1+Bz)^{-1} {}_2F_1\left(1, 1; \frac{1}{\lambda(p+m)} + 1; \frac{Bz}{1+Bz}\right) & (B \neq 0) \\ 1 + \frac{A}{\lambda(p+m)+1} z & (B = 0) \end{cases}$$

is the best dominant of (3.1). Furthermore,

$$(3.2) \quad \Re\left(-\frac{z^{p+1}(D^n f(z))'}{p}\right) > \rho \quad (z \in \mathbb{U}),$$



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 10 of 31

where

$$\rho = \begin{cases} \frac{A}{B} + (1 - \frac{A}{B})(1 - B)^{-1} {}_2F_1\left(1, 1; \frac{1}{\lambda(p+m)} + 1; \frac{B}{B-1}\right) & (B \neq 0) \\ 1 - \frac{A}{\lambda(p+m)+1} & (B = 0). \end{cases}$$

The inequality in (3.2) is the best possible.

*Proof.* Consider the function  $\phi(z)$  defined by


$$(3.3) \quad \phi(z) = -\frac{z^{p+1}(D^n f(z))'}{p} \quad (z \in \mathbb{U}).$$

Then  $\phi(z)$  is of the form (2.1) and is analytic in  $\mathbb{U}$ . Applying the identity (1.5) in (3.3) and differentiating the resulting equation with respect to  $z$ , we get

$$\begin{aligned} -\frac{(1 - \lambda)z^{p+1}(D^n f(z))' + \lambda z^{p+1}(D^{n+1} f(z))'}{p} \\ = \phi(z) + \lambda z \phi'(z) \prec \frac{1 + Az}{1 + Bz} \quad (z \in \mathbb{U}). \end{aligned}$$

Now, by using Lemma 1 for  $\gamma = 1/\lambda$ , we deduce that

$$\begin{aligned} -\frac{z^{p+1}(D^n f(z))'}{p} \prec Q(z) \\ = \frac{1}{\lambda(p+m)} z^{-\{1/\lambda(p+m)\}} \int_0^z t^{\{1/\lambda(p+m)\}-1} \left( \frac{1 + At}{1 + Bt} \right) dt \end{aligned}$$




---

An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

---

Title Page

Contents




Go Back

Close

Quit

Page 11 of 31



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 12 of 31

$$= \begin{cases} \frac{A}{B} + \left(1 - \frac{A}{B}\right) (1 + Bz)^{-1} {}_2F_1\left(1, 1; \frac{1}{\lambda(p+m)} + 1; \frac{Bz}{1+Bz}\right) & (B \neq 0) \\ 1 + \frac{A}{\lambda(p+m)+1} z & (B = 0), \end{cases}$$

by change of variables followed by the use of the identities (2.4), (2.5), and (2.6) (with  $b = 1$  and  $c = a + 1$ ). This proves the assertion (3.1) of Theorem 1.

Next, in order to prove the assertion (3.2) of Theorem 1, it suffices to show that

$$(3.4) \quad \inf_{|z| < 1} \{\Re(Q(z))\} = Q(-1).$$

Indeed, for  $|z| \leq r < 1$ ,

$$\Re\left(\frac{1 + Az}{1 + Bz}\right) \geq \frac{1 - Ar}{1 - Br} \quad (|z| \leq r < 1).$$

Upon setting

$$\mathcal{G}(s, z) = \frac{1 + Asz}{1 + Bsz} \quad \text{and} \\ d\nu(s) = \frac{1}{\lambda(p+m)} s^{\{1/\lambda(p+m)\}-1} ds \quad (0 \leq s \leq 1),$$

which is a positive measure on the closed interval  $[0, 1]$ , we get

$$Q(z) = \int_0^1 \mathcal{G}(s, z) d\nu(s),$$

so that

$$\Re(Q(z)) \geq \int_0^1 \left( \frac{1 - Asr}{1 - Bsr} \right) d\nu(s) = Q(-r) \quad (|z| \leq r < 1).$$

Letting  $r \rightarrow 1-$  in the above inequality, we obtain the assertion (3.2) of Theorem 1.

Finally, the estimate in (3.2) is the best possible as the function  $Q(z)$  is the best dominant of (3.1).  $\square$

For  $\lambda = 1$  and  $m = 0$ , Theorem 1 yields the following result which improves the corresponding work of Liu and Srivastava [3, Theorem 1].

**Corollary 1.** *The following inclusion property holds true for the function class  $\mathcal{R}_{n,p}(A, B)$ :*

$$\mathcal{R}_{n+1,p}(A, B) \subset \mathcal{R}_{n,p}(1 - 2\varrho, -1) \subset \mathcal{R}_{n,p}(A, B),$$

where


$$\varrho = \begin{cases} \frac{A}{B} + (1 - \frac{A}{B})(1 - B)^{-1} {}_2F_1\left(1, 1; \frac{1}{p} + 1; \frac{B}{B-1}\right) & (B \neq 0) \\ 1 - \frac{A}{p+1} & (B = 0). \end{cases}$$

The result is the best possible.

Putting

$$A = 1 - \frac{2\alpha}{p}, \quad B = -1, \quad \lambda = 1, \quad n = 0, \quad \text{and} \quad m = -p + 2$$

in Theorem 1, we get Corollary 2 below.



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 13 of 31

**Corollary 2.** If  $f(z) \in \Sigma_{p,-p+2}$  satisfies the following inequality:

$$\Re\left(-z^{p+1}\{(p+2)f'(z) + zf''(z)\}\right) > \alpha \quad (0 \leq \alpha < p; z \in \mathbb{U}),$$

then

$$\Re(-z^{p+1}f'(z)) > \alpha + (p - \alpha)\left(\frac{\pi}{2} - 1\right) \quad (z \in \mathbb{U}).$$

The result is the best possible.

**Remark 1.** From Corollary 2, we note that, if  $f(z) \in \Sigma_{p,-p+2}$  satisfies the following inequality:

$$\Re\left(-z^{p+1}\{(p+2)f'(z) + zf''(z)\}\right) > -\frac{p(\pi - 2)}{4 - \pi} \quad (z \in \mathbb{U}),$$

then

$$\Re(-z^{p+1}f'(z)) > 0 \quad (z \in \mathbb{U}).$$

This result is the best possible.

The result (asserted by Remark 1 above) was also obtained by Pap [8].


**Theorem 2.** If  $f(z) \in \Sigma_{p,m}^n(\alpha)$  ( $0 \leq \alpha < p$ ), then

$$(3.5) \quad \Re\left(-z^{p+1}\left\{(1-\lambda)(D^n f(z))' + \lambda(D^{n+1}f(z))'\right\}\right) > \alpha \quad (|z| < R),$$

where

$$R = \left(\sqrt{1 + \lambda^2(p+m)^2} - \lambda(p+m)\right)^{\frac{1}{p+m}}.$$

The result is the best possible.



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 14 of 31

*Proof.* We begin by writing

$$(3.6) \quad -z^{p+1}(D^n f(z))' = \alpha + (p - \alpha)u(z) \quad (z \in \mathbb{U}).$$

Then, clearly,  $u(z)$  is of the form (2.1), is analytic in  $\mathbb{U}$ , and has a positive real part in  $\mathbb{U}$ . Making use of the identity (1.5) in (3.6) and differentiating the resulting equation with respect to  $z$ , we observe that

$$(3.7) \quad -\frac{z^{p+1}\left((1 - \lambda)(D^n f(z))' + \lambda(D^{n+1}f(z))'\right) + \alpha}{p - \alpha} = u(z) + \lambda zu'(z).$$

Now, by applying the following estimate [4]:

$$\frac{|zu'(z)|}{\Re\{u(z)\}} \leq \frac{2(p + m)r^{p+m}}{1 - r^{2(p+m)}} \quad (|z| = r < 1)$$

in (3.7), we get

$$(3.8) \quad \Re\left(-\frac{z^{p+1}\left((1 - \lambda)(D^n f(z))' + \lambda(D^{n+1}f(z))'\right) + \alpha}{p - \alpha}\right) \geq \Re(u(z)) \cdot \left(1 - \frac{2\lambda(p + m)r^{p+m}}{1 - r^{2(p+m)}}\right).$$

It is easily seen that the right-hand side of (3.8) is positive, provided that  $r < R$ , where  $R$  is given as in Theorem 2. This proves the assertion (3.5) of Theorem 2.



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 15 of 31

In order to show that the bound  $R$  is the best possible, we consider the function  $f(z) \in \Sigma_{p,m}$  defined by

$$-z^{p+1}(D^n f(z))' = \alpha + (p - \alpha) \frac{1 + z^{p+m}}{1 - z^{p+m}} \quad (0 \leq \alpha < p; z \in \mathbb{U}).$$

Noting that

$$\begin{aligned} & - \frac{z^{p+1} \left( (1 - \lambda)(D^n f(z))' + \lambda(D^{n+1} f(z))' \right) + \alpha}{p - \alpha} \\ & = \frac{1 - z^{2(p+m)} + 2\lambda(p + m)z^{p+m}}{(1 - z)^{2(p+m)}} = 0 \end{aligned}$$

for

$$z = R \cdot \exp\left(\frac{i\pi}{p + m}\right),$$


we complete the proof of Theorem 2. □

Putting  $\lambda = 1$  in Theorem 2, we deduce the following result.

**Corollary 3.** *If  $f(z) \in \Sigma_{p,m}^n(\alpha)$  ( $0 \leq \alpha < p$ ), then  $f(z) \in \Sigma_{p,m}^{n+1}(\alpha)$  for  $|z| < \tilde{R}$ , where*

$$\tilde{R} = \left( \sqrt{1 + (p + m)^2} - (p + m) \right)^{\frac{1}{p+m}}.$$

*The result is the best possible.*



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 16 of 31

**Theorem 3.** Let  $f(z) \in \Sigma_{p,m}^n(A, B)$  and let

$$(3.9) \quad \mathcal{F}_{\delta,p}(f)(z) = \frac{\delta}{z^{\delta+p}} \int_0^z t^{\delta+p-1} f(t) dt \quad (\delta > 0; z \in \mathbb{U}^*).$$

Then

$$(3.10) \quad -\frac{z^{p+1} (D^n \mathcal{F}_{\delta,p}(f)(z))'}{p} \prec \Theta(z) \prec \frac{1 + Az}{1 + Bz} \quad (z \in \mathbb{U}),$$

where the function  $\Theta(z)$  given by

$$\Theta(z) = \begin{cases} \frac{A}{B} + \left(1 - \frac{A}{B}\right) (1 + Bz)^{-1} {}_2F_1\left(1, 1; \frac{\delta}{p+m} + 1; \frac{Bz}{1+Bz}\right) & (B \neq 0) \\ 1 + \frac{A\delta}{\delta+p+m} z & (B = 0) \end{cases}$$


is the best dominant of (3.10). Furthermore,

$$(3.11) \quad \Re\left(-\frac{z^{p+1} (D^n \mathcal{F}_{\delta,p}(f)(z))'}{p}\right) > \varkappa \quad (z \in \mathbb{U}),$$

where

$$\varkappa = \begin{cases} \frac{A}{B} + \left(1 - \frac{A}{B}\right) (1 - B)^{-1} {}_2F_1\left(1, 1; \frac{\delta}{p+m} + 1; \frac{B}{B-1}\right) & (B \neq 0) \\ 1 - \frac{A\delta}{\delta+p+m} & (B = 0). \end{cases}$$

The result is the best possible.



**An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem**

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 17 of 31

*Proof.* Setting

$$(3.12) \quad \phi(z) = -\frac{z^{p+1}(D^n \mathcal{F}_{\delta,p}(f)(z))'}{p} \quad (z \in \mathbb{U}),$$

we note that  $\phi(z)$  is of the form (2.1) and is analytic in  $\mathbb{U}$ . Using the following operator identity:

$$(3.13) \quad z(D^n \mathcal{F}_{\delta,p}(f)(z))' = \delta D^n f(z) - (\delta + p) D^n \mathcal{F}_{\delta,p}(f)(z)$$

in (3.12), and differentiating the resulting equation with respect to  $z$ , we find that

$$-\frac{z^{p+1}(D^n f(z))'}{p} = \phi(z) + \frac{z\phi'(z)}{\delta} \prec \frac{1 + Az}{1 + Bz} \quad (z \in \mathbb{U}).$$

Now the remaining part of Theorem 3 follows by employing the techniques that we used in proving Theorem 1 above.  $\square$

Setting  $m = 0$  in Theorem 3, we obtain the following result which improves the corresponding work of Liu and Srivastava [3, Theorem 2].

**Corollary 4.** *If  $\delta > 0$  and  $f(z) \in \mathcal{R}_{n,p}(A, B)$ , then*

$$\mathcal{F}_{\delta,p}(f)(z) \in \mathcal{R}_{n,p}(1 - 2\xi, -1) \subset \mathcal{R}_{n,p}(A, B),$$

where

$$\xi = \begin{cases} \frac{A}{B} + \left(1 - \frac{A}{B}\right) (1 - B)^{-1} {}_2F_1\left(1, 1; \frac{\delta}{p} + 1; \frac{B}{B-1}\right) & (B \neq 0) \\ 1 - \frac{A\delta}{\delta+p} & (B = 0). \end{cases}$$

*The result is the best possible.*



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 18 of 31

**Remark 2.** By observing that

$$(3.14) \quad z^{p+1} (D^n \mathcal{F}_{\delta,p}(f)(z))' = \frac{\delta}{z^\delta} \int_0^z t^{\delta+p} (D^n f(t))' dt$$

$$(f \in \Sigma_{p,m}; z \in \mathbb{U}),$$

Corollary 4 can be restated as follows.

If  $\delta > 0$  and  $f(z) \in \mathcal{R}_{n,p}(A, B)$ , then

$$\Re \left( -\frac{\delta}{pz^\delta} \int_0^z t^{\delta+p} (D^n f(t))' dt \right) > \xi \quad (z \in \mathbb{U}),$$

where  $\xi$  is given as in Corollary 4.

In view of (3.14), Theorem 3 for

$$A = 1 - \frac{2\alpha}{p}, \quad B = -1, \quad \text{and} \quad n = 0$$

yields

**Corollary 5.** If  $\delta > 0$  and if  $f(z) \in \Sigma_{p,m}$  satisfies the following inequality:


$$\Re \left( -z^{p+1} f'(z) \right) > \alpha \quad (0 \leq \alpha < p; z \in \mathbb{U}),$$

then

$$\Re \left( -\frac{\delta}{z^\delta} \int_0^z t^{\delta+p} f'(t) dt \right)$$

$$> \alpha + (p - \alpha) \left[ {}_2F_1 \left( 1, 1; \frac{\delta}{p+m} + 1; \frac{1}{2} \right) - 1 \right] \quad (z \in \mathbb{U}).$$

The result is the best possible.



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 19 of 31

**Theorem 4.** Let  $f(z) \in \Sigma_{p,m}$ . Suppose also that  $g(z) \in \Sigma_{p,m}$  satisfies the following inequality:

$$\Re(z^p D^n g(z)) > 0 \quad (z \in \mathbb{U}).$$

If

$$\left| \frac{D^n f(z)}{D^n g(z)} - 1 \right| < 1 \quad (n \in \mathbb{N}_0; z \in \mathbb{U}),$$

then

$$\Re \left( -\frac{z(D^n f(z))'}{D^n f(z)} \right) > 0 \quad (|z| < R_0),$$

where

$$R_0 = \frac{\sqrt{9(p+m)^2 + 4p(2p+m)} - 3(p+m)}{2(2p+m)}.$$

*Proof.* Letting

$$(3.15) \quad w(z) = \frac{D^n f(z)}{D^n g(z)} - 1 = \kappa_{p+m} z^{p+m} + \kappa_{p+m+1} z^{p+m+1} + \dots,$$

we note that  $w(z)$  is analytic in  $\mathbb{U}$ , with

$$w(0) = 0 \quad \text{and} \quad |w(z)| \leq |z|^{p+m} \quad (z \in \mathbb{U}).$$

Then, by applying the familiar Schwarz lemma, we get

$$w(z) = z^{p+m} \Psi(z),$$



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 20 of 31

where the function  $\Psi(z)$  is analytic in  $\mathbb{U}$  and

$$|\Psi(z)| \leq 1 \quad (z \in \mathbb{U}).$$

Therefore, (3.15) leads us to

$$(3.16) \quad D^n f(z) = D^n g(z)(1 + z^{p+m}\Psi(z)) \quad (z \in \mathbb{U}).$$

Making use of logarithmic differentiation in (3.16), we obtain

$$(3.17) \quad \frac{z(D^n f(z))'}{D^n f(z)} = \frac{z(D^n g(z))'}{D^n g(z)} + \frac{z^{p+m}\{(p+m)\Psi(z) + z\Psi'(z)\}}{1 + z^{p+m}\Psi(z)}.$$

Setting  $\phi(z) = z^p D^n g(z)$ , we see that the function  $\phi(z)$  is of the form (2.1), is analytic in  $\mathbb{U}$ ,


$$\Re(\phi(z)) > 0 \quad (z \in \mathbb{U}),$$

and

$$\frac{z(D^n g(z))'}{D^n g(z)} = \frac{z\phi'(z)}{\phi(z)} - p,$$

so that we find from (3.17) that

$$(3.18) \quad \Re\left(-\frac{z(D^n f(z))'}{D^n f(z)}\right) \geq p - \left| \frac{z\phi'(z)}{\phi(z)} \right| - \left| \frac{z^{p+m}\{(p+m)\Psi(z) + z\Psi'(z)\}}{1 + z^{p+m}\Psi(z)} \right| \quad (z \in \mathbb{U}).$$



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 21 of 31

Now, by using the following known estimates [10] (see also [4]):

$$\left| \frac{\phi'(z)}{\phi(z)} \right| \leq \frac{2(p+m)r^{p+m-1}}{1-r^{2(p+m)}} \quad \text{and}$$

$$\left| \frac{(p+m)\Psi(z) + z\Psi'(z)}{1+z^{p+m}\Psi(z)} \right| \leq \frac{(p+m)}{1-r^{p+m}} \quad (|z| = r < 1)$$

in (3.18), we obtain

$$\Re \left( -\frac{z(D^n f(z))'}{D^n f(z)} \right) \geq \frac{p - 3(p+m)r^{p+m} - (2p+m)r^{2(p+m)}}{1-r^{2(p+m)}} \quad (|z| = r < 1),$$

which is certainly positive, provided that  $r < R_0$ ,  $R_0$  being given as in Theorem 4.  $\square$

**Theorem 5.** Let  $-1 \leq B_j < A_j \leq 1$  ( $j = 1, 2$ ). If each of the functions  $f_j(z) \in \Sigma_p$  satisfies the following subordination condition:


$$(3.19) \quad (1-\lambda)z^p D^n f_j(z) + \lambda z^p D^{n+1} f_j(z) \prec \frac{1+A_j z}{1+B_j z} \quad (j = 1, 2; z \in \mathbb{U}),$$

then

$$(3.20) \quad (1-\lambda)z^p D^n H(z) + \lambda z^p D^{n+1} H(z) \prec \frac{1+(1-2\eta)z}{1-z} \quad (z \in \mathbb{U}),$$

where

$$H(z) = D^n(f_1 \star f_2)(z)$$



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 22 of 31

and

$$\eta = 1 - \frac{4(A_1 - B_1)(A_2 - B_2)}{(1 - B_1)(1 - B_2)} \left[ 1 - \frac{1}{2} {}_2F_1 \left( 1, 1; \frac{1}{\lambda} + 1; \frac{1}{2} \right) \right].$$

The result is the best possible when  $B_1 = B_2 = -1$ .

*Proof.* Suppose that each of the functions  $f_j(z) \in \Sigma_p$  ( $j = 1, 2$ ) satisfies the condition (3.19). Then, by letting

$$(3.21) \quad \varphi_j(z) = (1 - \lambda)z^p D^n f_j(z) + \lambda z^p D^{n+1} f_j(z) \quad (j = 1, 2),$$

we have

$$\varphi_j(z) \in \mathcal{P}(\gamma_j) \quad \left( \gamma_j = \frac{1 - A_j}{1 - B_j}; j = 1, 2 \right).$$

By making use of the operator identity (1.5) in (3.21), we observe that


$$D^n f_j(z) = \frac{1}{\lambda} z^{-p-(1/\lambda)} \int_0^z t^{(1/\lambda)-1} \varphi_j(t) dt \quad (j = 1, 2),$$

which, in view of the definition of  $H(z)$  given already with (3.20), yields

$$(3.22) \quad D^n H(z) = \frac{1}{\lambda} z^{-p-(1/\lambda)} \int_0^z t^{(1/\lambda)-1} \varphi_0(t) dt,$$

where, for convenience,

$$(3.23) \quad \begin{aligned} \varphi_0(z) &= (1 - \lambda)z^p D^n H(z) + \lambda z^p D^{n+1} H(z) \\ &= \frac{1}{\lambda} z^{-(1/\lambda)} \int_0^z t^{(1/\lambda)-1} (\varphi_1 \star \varphi_2)(t) dt. \end{aligned}$$



**An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem**

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 23 of 31

Since  $\varphi_1(z) \in \mathcal{P}(\gamma_1)$  and  $\varphi_2(z) \in \mathcal{P}(\gamma_2)$ , it follows from Lemma 3 that


$$(3.24) \quad (\varphi_1 \star \varphi_2)(z) \in \mathcal{P}(\gamma_3) \quad (\gamma_3 = 1 - 2(1 - \gamma_1)(1 - \gamma_2)).$$

Now, by using (3.24) in (3.23) and then appealing to Lemma 2 and Lemma 4, we get

$$\begin{aligned} \Re\{\varphi_0(z)\} &= \frac{1}{\lambda} \int_0^1 u^{(1/\lambda)-1} \Re\{(\varphi_1 \star \varphi_2)\}(uz) du \\ &\geq \frac{1}{\lambda} \int_0^1 u^{(1/\lambda)-1} \left( 2\gamma_3 - 1 + \frac{2(1 - \gamma_3)}{1 + u|z|} \right) du \\ &> \frac{1}{\lambda} \int_0^1 u^{(1/\lambda)-1} \left( 2\gamma_3 - 1 + \frac{2(1 - \gamma_3)}{1 + u} \right) du \\ &= 1 - \frac{4(A_1 - B_1)(A_2 - B_2)}{(1 - B_1)(1 - B_2)} \left( 1 - \frac{1}{\lambda} \int_0^1 u^{(1/\lambda)-1} (1 + u)^{-1} du \right) \\ &= 1 - \frac{4(A_1 - B_1)(A_2 - B_2)}{(1 - B_1)(1 - B_2)} \left[ 1 - \frac{1}{2} {}_2F_1 \left( 1, 1; \frac{1}{\lambda} + 1; \frac{1}{2} \right) \right] \\ &= \eta \quad (z \in \mathbb{U}). \end{aligned}$$

When  $B_1 = B_2 = -1$ , we consider the functions  $f_j(z) \in \Sigma_p$  ( $j = 1, 2$ ), which satisfy the hypothesis (3.19) of Theorem 5 and are defined by

$$D^n f_j(z) = \frac{1}{\lambda} z^{-(1/\lambda)} \int_0^z t^{(1/\lambda)-1} \left( \frac{1 + A_j t}{1 - z} \right) dt \quad (j = 1, 2).$$



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 24 of 31

Thus it follows from (3.23) and Lemma 4 that

$$\begin{aligned} \varphi_0(z) &= \frac{1}{\lambda} \int_0^1 u^{(1/\lambda)-1} \left( 1 - (1 + A_1)(1 + A_2) + \frac{(1 + A_1)(1 + A_2)}{1 - uz} \right) du \\ &= 1 - (1 + A_1)(1 + A_2) \\ &\quad + (1 + A_1)(1 + A_2)(1 - z)^{-1} {}_2F_1 \left( 1, 1; \frac{1}{\lambda} + 1; \frac{z}{z - 1} \right) \\ &\longrightarrow 1 - (1 + A_1)(1 + A_2) \\ &\quad + \frac{1}{2}(1 + A_1)(1 + A_2) {}_2F_1 \left( 1, 1; \frac{1}{\lambda} + 1; \frac{1}{2} \right) \quad \text{as } z \rightarrow -1, \end{aligned}$$

which evidently completes the proof of Theorem 5.  $\square$

By setting

$$A_j = 1 - 2\alpha_j, \quad B_j = -1 \quad (j = 1, 2), \quad \text{and} \quad n = 0$$

in Theorem 5, we obtain the following result which refines the work of Yang [15, Theorem 4].

**Corollary 6.** *If the functions  $f_j(z) \in \Sigma_p$  ( $j = 1, 2$ ) satisfy the following inequality:*

$$(3.25) \quad \Re \left( (1 + \lambda p) z^p f_j(z) + \lambda z^{p+1} f'_j(z) \right) > \alpha_j \\ (0 \leq \alpha_j < 1; \quad j = 1, 2; \quad z \in \mathbb{U}),$$



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 25 of 31

then

$$\Re\left((1 + \lambda p)z^p(f_1 \star f_2)(z) + \lambda z^{p+1}(f_1 \star f_2)'(z)\right) > \eta_0 \quad (z \in \mathbb{U}),$$

where

$$\eta_0 = 1 - 4(1 - \alpha_1)(1 - \alpha_2) \left[1 - \frac{1}{2} {}_2F_1\left(1, 1; \frac{1}{\lambda} + 1; \frac{1}{2}\right)\right].$$

The result is the best possible.

**Theorem 6.** If  $f(z) \in \Sigma_{p,m}$  satisfies the following subordination condition:

$$(1 - \lambda)z^p D^n f(z) + \lambda z^p D^{n+1} f(z) \prec \frac{1 + Az}{1 + Bz} \quad (z \in \mathbb{U}),$$

then

$$\Re\left((z^p D^n f(z))^{1/q}\right) > \rho^{1/q} \quad (q \in \mathbb{N}; z \in \mathbb{U}),$$


where  $\rho$  is given as in Theorem 1. The result is the best possible.

*Proof.* Defining the function  $\phi(z)$  by

$$(3.26) \quad \phi(z) = z^p D^n f(z) \quad (f \in \Sigma_{p,m}; z \in \mathbb{U}),$$

we see that the function  $\phi(z)$  is of the form (2.1) and is analytic in  $\mathbb{U}$ . Using the identity (1.5) in (3.26) and differentiating the resulting equation with respect to  $z$ , we find that

$$(1 - \lambda)z^p D^n f(z) + \lambda z^p D^{n+1} f(z) = \phi(z) + \lambda z \phi'(z) \prec \frac{1 + Az}{1 + Bz} \quad (z \in \mathbb{U}).$$



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 26 of 31

Now, by following the lines of proof of Theorem 1 *mutatis mutandis*, and using the elementary inequality:

$$\Re(w^{1/q}) \geq (\Re(w))^{1/q} \quad (\Re(w) > 0; q \in \mathbb{N}),$$

we arrive at the result asserted by Theorem 6. □

Upon setting

$$A = \left[ {}_2F_1 \left( 1, 1; \frac{1}{\lambda(p+m)} + 1; \frac{1}{2} \right) - 1 \right] \cdot \left[ 2 - {}_2F_1 \left( 1, 1; \frac{1}{\lambda(p+m)} + 1; \frac{1}{2} \right) \right]^{-1},$$

$$B = -1, \quad n = 0, \quad \text{and} \quad q = 1$$

in Theorem 6, we deduce Corollary 7 below.


**Corollary 7.** *If  $f(z) \in \Sigma_{p,m}$  satisfies the following inequality:*

$$(3.27) \quad \Re((1 + \lambda p)z^p f(z) + \lambda z^{p+1} f'(z)) > \frac{3 - 2 {}_2F_1 \left( 1, 1; \frac{1}{\lambda(p+m)} + 1; \frac{1}{2} \right)}{2 \left[ 2 - {}_2F_1 \left( 1, 1; \frac{1}{\lambda(p+m)} + 1; \frac{1}{2} \right) \right]} \quad (z \in \mathbb{U}),$$

then

$$\Re(z^p f(z)) > \frac{1}{2} \quad (z \in \mathbb{U}).$$

*The result is the best possible.*



**An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem**

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 27 of 31

From Corollary 6 and Theorem 6 (for  $m = -p + 1$ ,  $A = 1 - 2\eta_0$ ,  $B = -1$ , and  $q = 1$ ), we deduce the following result.

**Corollary 8.** *If the functions  $f_j(z) \in \Sigma_p$  ( $j = 1, 2$ ) satisfy the inequality (3.25), then*

$$\Re\left(z^p(f_1 \star f_2)(z)\right) > \eta_0 + (1 - \eta_0) \left[ {}_2F_1\left(1, 1; \frac{1}{\lambda} + 1; \frac{1}{2}\right) - 1 \right] \quad (z \in \mathbb{U}),$$

where  $\eta_0$  is given as in Corollary 6. The result is the best possible.

**Theorem 7.** *Let  $f(z) \in \Sigma_{p,m}^n(A, B)$  and let  $g(z) \in \Sigma_{p,m}$  satisfy the following inequality:*

$$\Re(z^p g(z)) > \frac{1}{2} \quad (z \in \mathbb{U}).$$

Then

$$(f \star g)(z) \in \Sigma_{p,m}^n(A, B).$$

*Proof.* We have


$$-\frac{z^{p+1}(D^n(f \star g)(z))'}{p} = -\frac{z^{p+1}(D^n f(z))'}{p} \star z^p g(z) \quad (z \in \mathbb{U}).$$

Since

$$\Re(z^p g(z)) > \frac{1}{2} \quad (z \in \mathbb{U})$$

and the function

$$\frac{1 + Az}{1 + Bz}$$



An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

Title Page

Contents



Go Back

Close

Quit

Page 28 of 31

is convex (univalent) in  $\mathbb{U}$ , it follows from (1.6) and Lemma 5 that


$$(f \star g)(z) \in \Sigma_{p,m}^n(A, B).$$

This completes the proof of Theorem 7. □

In view of Corollary 7 and Theorem 7, we have Corollary 9 below.

**Corollary 9.** *If  $f(z) \in \Sigma_{p,m}^n(A, B)$  and the function  $g(z) \in \Sigma_{p,m}$  satisfies the inequality (3.27), then*

$$(f \star g)(z) \in \Sigma_{p,m}^n(A, B).$$



---

An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

---

Title Page

Contents



Go Back


Close

Quit

Page 29 of 31

## References

- [1] M.K. AOUF AND H.M. HOSSEN, New criteria for meromorphic  $p$ -valent starlike functions, *Tsukuba J. Math.*, **17** (1993), 481–486.
- [2] J.-L. LIU AND H.M. SRIVASTAVA, Classes of meromorphically multivalent functions associated with the generalized hypergeometric function, *Math. Comput. Modelling*, **39** (2004), 21–34.
- [3] J.-L. LIU AND H.M. SRIVASTAVA, Subclasses of meromorphically multivalent functions associated with a certain linear operator, *Math. Comput. Modelling*, **39** (2004), 35–44.
- [4] T.H. MacGREGOR, Radius of univalence of certain analytic functions, *Proc. Amer. Math. Soc.*, **14** (1963), 514–520.
- [5] S.S. MILLER AND P.T. MOCANU, Differential subordinations and univalent functions, *Michigan Math. J.*, **28** (1981), 157–171.
- [6] S.S. MILLER AND P.T. MOCANU, *Differential Subordinations: Theory and Applications*, Series on Monographs and Textbooks in Pure and Applied Mathematics, Vol. 225, Marcel Dekker, New York and Basel, 2000.
- [7] M.L. MOGRA, Meromorphic multivalent functions with positive coefficients. I and II, *Math. Japon.*, **35** (1990), 1–11 and 1089–1098.
- [8] M. PAP, On certain subclasses of meromorphic  $m$ -valent close-to-convex functions, *Pure Math. Appl.*, **9** (1998), 155–163.



---

An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

---

Title Page

Contents




Go Back

Close

Quit

Page 30 of 31



---

An Inequality of Ostrowski Type  
via Pompeiu's Mean Value  
Theorem

H.M. Srivastava and J. Patel

---

- [9] D.Ž. PASHKOULEVA, The starlikeness and spiral-convexity of certain subclasses of analytic functions, in : H.M. Srivastava and S. Owa (Editors), *Current Topics in Analytic Function Theory*, pp. 266–273, World Scientific Publishing Company, Singapore, New Jersey, London and Hong Kong, 1992.
- [10] J. PATEL, Radii of  $\gamma$ -spirallikeness of certain analytic functions, *J. Math. Phys. Sci.*, **27** (1993), 321–334.
- [11] R. SINGH AND S. SINGH, Convolution properties of a class of starlike functions, *Proc. Amer. Math. Soc.*, **106** (1989), 145–152.
- [12] J. STANKIEWICZ AND Z. STANKIEWICZ, Some applications of the Hadamard convolution in the theory of functions, *Ann. Univ. Mariae Curie-Skłodowska Sect. A*, **40** (1986), 251–265.
- [13] B.A. URALEGADDI AND C. SOMANATHA, New criteria for meromorphic starlike univalent functions, *Bull. Austral. Math. Soc.*, **43** (1991), 137–140.
- [14] E.T. WHITTAKER AND G.N. WATSON, *A Course on Modern Analysis: An Introduction to the General Theory of Infinite Processes and of Analytic Functions; With an Account of the Principal Transcendental Functions*, Fourth Edition (Reprinted), Cambridge University Press, Cambridge, 1927.
- [15] D.-G. YANG, Certain convolution operators for meromorphic functions, *Southeast Asian Bull. Math.*, **25** (2001), 175–186.

Title Page

Contents



Go Back

Close

Quit

Page 31 of 31