



Dickson Polynomials, Chebyshev Polynomials, and Some Conjectures of Jeffery

Stefano Barbero
Department of Mathematics
University of Turin
via Carlo Alberto 10
10123 Turin
Italy
stefano.barbero@unito.it

Abstract

By using Dickson polynomials in several variables and Chebyshev polynomials of the second kind, we derive the explicit expression of the entries in the array defining the sequence [A185905](#). As a result, we obtain a straightforward proof of some conjectures of Jeffery concerning this sequence and other related ones.

1 Proof of main conjectures

As we can find in the *Encyclopedia of Integer Sequences* [1], the sequence [A185095](#) is a rectangular array read by antidiagonals, where the n -th row has generating function

$$F_n(z) = \frac{\sum_{r=0}^n (n+1-r)(-1)^r \binom{2(n+1)-r}{r} z^r}{\sum_{r=0}^{n+1} (-1)^r \binom{2(n+1)-r}{r} z^r} \quad (1)$$

for $n = 0, 1, 2, \dots$. In this section we find the explicit expression for the entries of this rectangular array and we determine the ordinary generating functions for the columns.

Let us consider the Dickson polynomials of the first kind in several indeterminates, as defined in the book of Lidl, Mullen, and Turnwald [2].

Definition 1. The Dickson polynomials of the first kind in n indeterminates of total degree k

$$D_k^{(i)}(x_1, \dots, x_n, a), \quad i = 1, \dots, n$$

satisfy the functional equations

$$D_k^{(i)}(x_1, \dots, x_n, a) = S_i(u_1^k, \dots, u_{n+1}^k), \quad i = 1, \dots, n \quad (2)$$

where $x_i = S_i(u_1, \dots, u_{n+1})$, $S_i(y_1, \dots, y_{n+1})$ is the i -th symmetric function of y_1, \dots, y_{n+1} , and $u_1 \cdot \dots \cdot u_{n+1} = a$.

In particular when $i = 1$ we have $D_k^{(1)}(x_1, \dots, x_n, a) = S_1(u_1^k, \dots, u_{n+1}^k) = \sum_{j=1}^{n+1} u_j^k$. Moreover if we pose $x_0 = 1$ and $x_{n+1} = a$, the Dickson polynomials of the first kind $D_k^{(1)}(x_1, \dots, x_n, a)$ have the following generating function (see Lidl, Mullen, and Turnwald [2, Lemma 2.23]):

$$\sum_{k=0}^{+\infty} D_k^{(1)}(x_1, \dots, x_n, a) z^k = \frac{\sum_{r=0}^n (n+1-r) (-1)^r x_r z^r}{\sum_{r=0}^{n+1} (-1)^r x_r z^r}, \quad k \geq 0. \quad (3)$$

Proposition 2. The k -th entry in the n -th row $R_{n,k}$, $k, n \geq 0$, of the rectangular array which defines the sequence [A185095](#), corresponds to $D_k^{(1)}(x_1, \dots, x_n, 1)$ where $x_r = \binom{2(n+1)-r}{r}$ for $r = 0, \dots, n+1$. Furthermore we have that

$$R_{n,k} = 2^{2k} \sum_{j=1}^{n+1} \cos^{2k} \left(\frac{j\pi}{2n+3} \right). \quad (4)$$

Proof. After a comparison between the two generating functions (1) and (3), we immediately find that $R_{n,k} = D_k^{(1)}(x_1, \dots, x_n, 1)$. Moreover, by definition (1), the role played by symmetric functions when $x_r = \binom{2(n+1)-r}{r}$ for $r = 0 \dots, n+1$, allow us to prove that

$$R_{n,k} = D_k^{(1)}(x_1, \dots, x_n, 1) = \sum_{j=1}^{n+1} \alpha_j^k,$$

where α_j , for $j = 1, \dots, n+1$ are the zeros of the polynomial

$$P_n(x) = \sum_{j=0}^{n+1} (-1)^j \binom{2(n+1)-j}{j} x^{n+1-j}. \quad (5)$$

Now let us recall the definition of the Chebyshev polynomials of the second kind $U_h \left(\frac{x}{2} \right)$ (see for references the books of Rivlin [3] and of Mason and Hascomb [4]):

$$U_h \left(\frac{x}{2} \right) = \sum_{j=0}^{\lfloor \frac{h}{2} \rfloor} (-1)^j \binom{h-j}{j} x^{h-2j}. \quad (6)$$

It is clear that $P_n(x) = U_{2n+2}\left(\frac{\sqrt{x}}{2}\right)$. So α_j is a zero of $P_n(x)$ if and only if $\frac{\sqrt{\alpha_j}}{2}$ is a positive zero for $U_{2n+2}(x)$. Since the zeros of $U_{2n+2}(x)$ are $x_j = \cos\left(\frac{j\pi}{2n+3}\right)$, $j = 1, \dots, 2n+3$, a simple trigonometric consideration allow us to find the positive ones when $j = 1, \dots, n+1$, showing that $\alpha_j = 2^2 \cos^2\left(\frac{j\pi}{2n+3}\right)$. The thesis immediately follows. \square

Corollary 3. *The entry $R_{n,k}$ for $n \geq 1$ has the alternative expression*

$$R_{n,k} = \binom{2k}{k}n + \frac{3}{2} \binom{2k}{k} - 2^{2k-1}, \quad k \geq 1 \quad (7)$$

and $R_{n,0} = n + 1$, $R_{0,k} = 1$.

Proof. Obviously if $k = 0$ we obtain $R_{n,0} = n + 1$ and if $n = 0$ we have $R_{0,k} = 1$. When $k \geq 1$, by (4) and thanks to the summation formulas (see Gradshteyn and Ryzhik [5]):

$$\cos^{2k}(x) = \frac{1}{2^{2k}} \left\{ \binom{2k}{k} + 2 \sum_{h=0}^{k-1} \binom{2k}{h} \cos(2(k-h)x) \right\}$$

and

$$\sum_{j=0}^n \cos(jx) = \frac{1}{2} \left[1 + \frac{\sin\left(\frac{(n+1)x}{2}\right)}{\sin\left(\frac{x}{2}\right)} \right],$$

we get

$$\begin{aligned} R_{n,k} &= \sum_{j=1}^{n+1} \left(\binom{2k}{k} + 2 \sum_{h=0}^{k-1} \binom{2k}{h} \cos\left(2(k-h)\frac{j\pi}{2n+3}\right) \right) = \\ &= \binom{2k}{k}(n+1) + 2 \sum_{h=0}^{k-1} \binom{2k}{h} \left(\sum_{j=0}^{n+1} \cos\left(2(k-h)\frac{j\pi}{2n+3}\right) - 1 \right) = \\ &= \binom{2k}{k}(n+1) + 2 \sum_{h=0}^{k-1} \binom{2k}{h} \left(\frac{1}{2} \left[1 + \frac{\sin\left(\frac{(2n+3)2(k-h)\pi}{2(2n+3)}\right)}{\sin\left(\frac{(k-h)\pi}{2n+3}\right)} \right] - 1 \right) = \\ &= \binom{2k}{k}(n+1) - \sum_{h=0}^{k-1} \binom{2k}{h}. \end{aligned}$$

Finally, by using the following well known identity

$$\sum_{h=0}^{k-1} \binom{2k}{h} = 2^{2k-1} - \frac{1}{2} \binom{2k}{k},$$

we easily have the thesis. \square

Considering the rectangular array in the definition of sequence [A 185095](#) we can also study its columns, finding their generating functions. Thanks to the previous result we have the following

Proposition 4. *The ordinary generating function $G_k(z)$ for the k -th column of the rectangular array in the definition of [A 185095](#) is*

$$G_k(z) = \frac{(2^{2k-1} - \frac{3}{2}\binom{2k}{k} + 1)z^2 + (\frac{5}{2}\binom{2k}{k} - 2^{2k-1} - 2)z + 1}{(1-z)^2}, \quad k \geq 0. \quad (8)$$

Proof. Using the alternative expression determined in Corollary 7, and the well-known formulas $\sum_{n=0}^{+\infty} z^n = \frac{1}{1-z}$ and $\sum_{n=0}^{+\infty} (n+1)z^n = \frac{1}{(1-z)^2}$, we can compute the ordinary generating function $G_k(z)$ as follows:

$$\begin{aligned} G_k(z) &= \sum_{n=0}^{+\infty} R_{n,k} z^n = 1 + \binom{2k}{k} \left(\sum_{n=0}^{+\infty} (n+1)z^n - 1 \right) - \left(2^{2k-1} - \frac{1}{2}\binom{2k}{k} \right) \left(\sum_{n=0}^{+\infty} z^n - 1 \right) = \\ &= 1 + \binom{2k}{k} \frac{(2z - z^2)}{(1-z)^2} - \frac{z}{1-z} \left(2^{2k-1} - \frac{1}{2}\binom{2k}{k} \right) = \\ &= \frac{(2^{2k-1} - \frac{3}{2}\binom{2k}{k} + 1)z^2 + (\frac{5}{2}\binom{2k}{k} - 2^{2k-1} - 2)z + 1}{(1-z)^2}. \end{aligned}$$

□

2 Concluding remarks

In this section we discuss some consequences of the previous results concerning the sequence [A185905](#). We show that the proof of the remaining conjectures and the relations with other sequences are immediate. First of all we observe that the k -th column of the array defining the sequence [A185905](#) satisfies the recurrence relation

$$R_{n+1,k} = 2R_{n,k} - R_{n-1,k}, \quad \text{for } n \geq 1,$$

which is a straightforward consequence of (7). The rectangular array which generates [A186740](#) clearly is the transpose of the rectangular array that we have considered. In fact, if we start by numbering by 0 the columns of the rectangular array which defines [A186740](#), we find that the entry in the k -th row and n -th column corresponds to $R_{n,k}$ by definition. Moreover by (7) and using our numeration for the columns we have

$R_{n,0} = n + 1, n \geq 0$: the column 0 corresponds to the natural numbers [A000027](#) ;

$R_{n,1} = 2n + 1, n \geq 0$: the column 1 is the sequence of odd integers [A005408](#) ;

$R_{n,2} = 6n + 1, n \geq 1$: the column 2 is [A016921](#) ;

$R_{n,3} = 20n - 2, n \geq 1$: the column 3 is [A114698](#) ;

$R_{n,4} = 70n - 93, n \geq 1$: the column 4 is [A114646](#) ,

and so on .

Finally another interesting consequence of the formula for $R_{n,k}$ concerns the sequence [A198632](#). By considering the array $w(h, 2k)$, as defined in the page related to [A198632](#) (see OEIS [1]), we have immediately $w(2(n+1), 2k) = 2R_{n,k}$. In fact

$$w(h, l) = \text{Tr}(J_h^l) = \sum_{j=1}^h \lambda_j^l,$$

where $\lambda_j, j = 1, \dots, h$, are the eigenvalues of the adjacency matrix J_h , a Jacobi matrix, whose characteristic polynomial is $U_h(\frac{x}{2})$. Thus

$$w(2(n+1), 2k) = \sum_{j=1}^{2n+2} \left(2 \cos \left(\frac{j\pi}{2n+3} \right) \right)^{2k} = 2^{2k+1} \sum_{j=1}^{n+1} \left(\cos \left(\frac{j\pi}{2n+3} \right) \right)^{2k} = 2R_{n,k}.$$

3 Acknowledgements

In this paper we point out some relations between many sequences and we think that a lot of interesting formulas will also rise, considering the role played by Dickson and Chebyshev polynomials, their zeros and the summation identities involved. The author sincerely thanks L. Edson Jeffery for all his interesting work and for his invitation to try a proof of his conjectures.

References

- [1] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, <http://oeis.org> .
- [2] R. Lidl, G. L. Mullen, and G. Turnwald, *Dickson Polynomials*, John Wiley and Sons, 1993.
- [3] T. J. Rivlin, *The Chebyshev Polynomials*, John Wiley and Sons, 1974.
- [4] J. C. Mason and D. C. Handscomb, *Chebyshev Polynomials*, Chapman and Hall, 2002.
- [5] I. S. Gradshteyn and I. M. Ryzhik, *Tables of Integrals, Series and Products*, Academic Press, 2007.

2010 *Mathematics Subject Classification*: Primary 11C08; Secondary 11B83.

Keywords: Dickson polynomial, Chebyshev polynomial.

(Concerned with sequences [A000027](#), [A005408](#), [A016921](#), [A114646](#), [A114698](#), [A114968](#), [A185095](#), [A185905](#), [A186740](#), [A198632](#), and [A198636](#).)

Received December 7 2013; revised version received January 25 2014. Published in *Journal of Integer Sequences*, February 16 2014.

Return to [Journal of Integer Sequences home page](#).