



Integer Compositions and Higher-Order Conjugation

Augustine O. Munagi
School of Mathematics
University of the Witwatersrand
Wits 2050, Johannesburg
South Africa
augustine.munagi@wits.ac.za

Abstract

We consider the classical MacMahon conjugation of compositions or ordered partitions of positive integers. Using both algebraic and graphical methods we provide a natural extension of the standard conjugation of a composition to higher orders. The higher-order conjugates of a composition are obtained by varying the increments used in standard conjugation to turn strings of ones into larger summands and vice versa. It turns out that every nontrivial composition has an integral conjugation order beyond which it is not conjugable. We also discuss recursive conjugation and provide enumeration formulas and combinatorial identities between different classes of compositions.

1 Introduction

A composition of a positive integer n is an ordered partition of n or a sequence of positive integers that sum to n . The terms of the sequence are called parts and n is the *weight* of the composition. For example 4 has eight compositions namely

$$(4), (3, 1), (1, 3), (2, 2), (2, 1, 1), (1, 2, 1), (1, 1, 2), (1, 1, 1, 1).$$

It is well known that there are $c(n) = 2^{n-1}$ compositions of n , and $c(n, k) = \binom{n-1}{k-1}$ compositions of n with exactly k parts which are also called k -compositions (see for example [2, 9]).

The *conjugate* of a composition may be obtained by drawing its *zig-zag graph*. Here each part is represented by a row of dots such that the first dot on a row is aligned with the last dot on the previous row. For example, the zig-zag graph of $(5, 3, 1, 2, 2)$ is shown in Figure 1.



Figure 1: Zig-zag graph of $(5, 3, 1, 2, 2)$

The conjugate is the composition corresponding to the columns of zig-zag graphs, from left to right. Thus the conjugate of $(5, 3, 1, 2, 2)$, from the Figure 1, is $(1, 1, 1, 1, 2, 1, 3, 2, 1)$. The conjugate of a composition C will be denoted by C' .

This is an expository paper aimed at providing a natural extension of the definition of the standard conjugation of a composition using algebraic and graphical methods. This will be followed by a brief discussion of how the resulting viewpoint refines or enlarges some familiar topics in the theory.

Percy Alexander MacMahon (1854–1929) introduced the zig-zag graph in [14]. Subsequently, in the classic text *Combinatory Analysis*, he contrasted the zig-zag graph with the partition Ferrers graph and remarked that the latter was rigidly fixed between two perpendicular axes [12, Sec. IV, p. 153]. However, he had previously defined conjugation using the *Line Graph* [12, 13] which provided an intuitive complementary relationship between a pair of mutually conjugate compositions. MacMahon also explained a diagram-free short-cut for writing the conjugates directly (now called the ‘DD’ method) [13].

In recent years conjugate compositions have played a pivotal role in the discovery of composition analogues of classical partition identities while affording natural insights into difficult proofs, see for example [8, 15, 16, 17]. The contemporary text [9] provides a wide-ranging coverage of the subject.

Agarwal [1] introduced n -color compositions which are extended compositions in which a part of size m may come in m different colors. Subsequently, Brian Hopkins [11] devised a MacMahon-type zig-zag graph for n -color compositions by replacing the original zig-zag graph dots with unit squares and then placing dots in some of the squares to indicate colors of part sizes. However, the graphical conjugation operation proposed by Hopkins failed to yield conjugates to many n -color compositions. Several other generalizations of compositions have appeared in the literature in the form of weighted compositions [6, 7], locally restricted compositions [3, 4] and compositions whose parts belong to a prescribed integer set [9, 10], and so forth. But none of these species of compositions emerged with specialized conjugation considerations.

Two further methods of obtaining the conjugate composition are outlined in Section 2 together with other essential preliminaries. In Section 3 we introduce the concept of higher-

The respective conjugates of the four types of compositions are given according to the rule below.

- (i') $C' = (a_1 + 1, 1^{b_1-2}, a_2 + 2, 1^{b_2-2}, \dots, 1^{b_r-2}, a_{r+1} + 1)$;
- (ii') $C' = (a_1 + 1, 1^{b_1-2}, a_2 + 2, 1^{b_2-2}, \dots, a_r + 2, 1^{b_r-1})$;
- (iii') $C' = (1^{b_1-1}, a_1 + 2, 1^{b_2-2}, a_2 + 2, \dots, 1^{b_r-2}, a_r + 1)$;
- (iv') $C' = (1^{b_1-1}, a_1 + 2, 1^{b_2-2}, a_2 + 2, \dots, a_{r-1} + 2, 1^{b_r-1})$.

The DD technique shows that C and C' have the same number of symbolic parts.

For example, the conjugate of $(5, 3, 1, 2, 2) = (5, 1^0, 3, 1, 2, 1^0, 2)$ is given by

$$(1^4, 2, 1, 3, 1^0, 2, 1).$$

In the sequel compositions of the form (1^n) or (n) will be called *trivial*. Thus a *nontrivial* composition contains 1 and a big part or at least two big parts.

The four types of compositions in (i) to (iv) will be referred to as $1c1$, $1c2$, $2c1$ and $2c2$, compositions, respectively. Thus a $1c1$ composition has the first and last parts equal to 1, a $1c2$ composition has a first part 1 and a last part > 1 , and so forth. Their conjugates in (i') to (iv') are $2c2$, $2c1$, $1c2$ and $1c1$ compositions. We will mostly use $1c2$ or $2c1$ compositions for demonstrations because they possess the features of all four types. Note that a $1c2$ or $2c1$ composition is never trivial.

A nontrivial composition in symbolic form consists of an alternation of big parts and strings of 1's. Initial and final strings of 1's or big parts are *boundary* objects while others are *interior* or *intermediate* objects.

A composition C is deemed to be empty (or non-existent), denoted by $C = \emptyset$, if a defining inequality is violated. For example, if C is a $1c2$ composition (see (ii) above), then $C = \emptyset$ if $b_i < 2$ for some i or $a_1 < 1$ or $a_i < 0$, $i = 2, 3, \dots, r$.

3 Higher-order conjugation

Consider the general $1c2$ composition

$$C = (1^{a_1}, b_1, 1^{a_2}, b_2, \dots, 1^{a_r}, b_r), \quad a_1 > 0, a_i \geq 0, i > 1, b_i > 1, \forall i.$$

As already seen, the conjugate using the DD conjugation rule, is

$$C' = (a_1 + 1, 1^{b_1-2}, a_2 + 2, 1^{b_2-2}, \dots, a_r + 2, 1^{b_r-1}).$$

This is the classical conjugation, and we say that it is of order 1. Notice that this process involves subtracting 1 from a boundary big part, and subtracting $2 = 2 \times 1$ from an intermediate big part to make 1's (and correspondingly adding 1 to a boundary string of 1's, and adding 2 to an intermediate string of 1's, to make a big part).

For example, $C = (1^3, 6, 4, 1^2, 5) = (1^3, 6, 1^0, 4, 1^2, 5)$ has the conjugate of order 1 given by

$$C' = (3 + 1, 1^{6-2}, 0 + 2, 1^{4-2}, 2 + 2, 1^{5-1}) = (4, 1^4, 2, 1^2, 4, 1^4).$$

On the other hand, C has the order-2 conjugate

$$C'' = (3 + 2, 1^{6-4}, 0 + 4, 1^{4-4}, 2 + 4, 1^{5-2}) = (5, 1^2, 4, 1^0, 6, 1^3).$$

In the second conjugation, we subtracted 2 from a boundary big part, and subtracted $4 = 2 \times 2$ from an intermediate big part, to make 1's.

However, $C''' = \emptyset$: interior big parts of size at least 6 are required. This may be extended to higher orders.

Definition 1 (Conjugate of Order t). Let t be a positive integer. A nontrivial composition C possesses a conjugate of order t if every boundary big part of C is greater than t and every interior big part is greater than or equal to $2t$.

Let C be a composition that possesses a conjugate of order t . Then to obtain the conjugate of order t , we subtract t from each boundary big part and subtract $2t$ from every intermediate big part to make 1's, and correspondingly add t to a boundary string of 1's, and add $2t$ to an intermediate string of 1's, to make a big part.

Thus a $1c2$ composition C that possesses a conjugate of order $t > 0$ has the form

$$C = (1^{a_1}, b_1, 1^{a_2}, \dots, 1^{a_r}, b_r),$$

$$a_1 \geq 1, a_i \geq 0, i > 1, b_i \geq 2t, 1 \leq i \leq r - 1, b_r > t. \quad (1)$$

The conjugate of order t is a composition E given by

$$E = (a_1 + t, 1^{b_1-2t}, a_2 + 2t, 1^{b_2-2t}, \dots, a_r + 2t, 1^{b_r-t}). \quad (2)$$

Notice that E is a $2c1$ composition as expected. It is easily checked that the parts of E satisfy $a_1 + t > t$, $b_i - 2t \geq 0$ and $b_r - t > 0$. So the parts of E fulfill the same properties as the parts of C . Therefore E in turn possesses a conjugate of order t . It may be verified that the conjugate of E , of order t , is C .

Definition 2. Given a positive integer t , a composition is t -conjugable if it possesses a conjugate of order t . A t -conjugable composition C is also said to possess a t -conjugate, denoted successively, for $t = 1, 2, 3, \dots$, by $C', C'', C''', \dots, C^{(t)}, C^{(t+1)}, \dots$.

The trivial compositions (1^n) and (n) possess conjugates of arbitrary order. So they are conventionally m -conjugable for all positive integers m .

$$(1^n)^{(m)} = (n) \quad \text{and} \quad (n)^{(m)} = (1^n), \quad m > 0.$$

The following characterization is equivalent to the first part of Definition 1.

By this definition every nontrivial composition C of $n \geq 3$ satisfies

$$1 \leq \text{ht}(C) \leq n - 2, \quad n > 2, \quad (3)$$

where equality holds on the right for the compositions $(1, n - 1), (n - 1, 1)$. There is no nontrivial composition of n with height greater than $n - 2$.

If $\text{ht}(C) = t > 1$, then C is $(t - 1)$ -conjugable. If $\text{ht}(C) = t$, then $C^{(t+1)} = \emptyset$.

Theorem 5. *A nontrivial t -conjugable composition $C = (c_1, c_2, \dots, c_k)$ has conjugate height t if and only if either of the following conditions hold:*

(i) *Every $c_i > 1$, $2 \leq i \leq k - 1$ satisfies $c_i \geq 2t$ and $t + 1 \in \{c_1, c_k\}$.*

(ii) *$c_i \in \{2t, 2t + 1\}$ for some $i \neq 1, k$, and $c_1, c_k \geq t + 1$.*

Proof. We use the standard 1c2 composition $C = (1^{a_1}, b_1, 1^{a_2}, b_2, \dots, 1^{a_r}, b_r)$. Thus $c_k = b_r$, and any $c_i > 1$ corresponds to b_i for some $i = 1, \dots, r - 1$.

Assume $\text{ht}(C) = t$. Then $C^{(t)}$ exists but not $C^{(t+1)}$. Consider

$$C^{(t)} = (a_1 + t, 1^{b_1 - 2t}, \dots, a_r + 2t, 1^{b_r - t}) \neq \emptyset, \quad (4)$$

$$C^{(t+1)} = (a_1 + t + 1, 1^{b_1 - 2t - 2}, \dots, a_r + 2t + 2, 1^{b_r - t - 1}) = \emptyset. \quad (5)$$

Since $C^{(t+1)} = \emptyset$, Equations (4) and (5) imply that b_r may satisfy $b_r - t > 0$ and $b_r - t - 1 < 1$, which imply $1 \leq b_r - t \leq 1$ or $b_r = t + 1$ (where we have used Theorem 3).

Alternatively, we might have $b_i - 2t \geq 0$ and $b_i - 2t - 2 < 0$ for some $i \in \{1, \dots, r - 1\}$. This gives $0 \leq b_i - 2t \leq 1$ or $2t \leq b_i \leq 2t + 1$. So conditions (i) and (ii) hold.

Conversely (i) implies that $b_r = t + 1$. But from (5) $C^{(t+1)} = \emptyset$ since its last string of 1's will be $1^{b_r - t - 1} = 1^0$. Condition (ii) implies that $b_i \in \{2t, 2t + 1\}$, $i \neq r$. Then $1^{b_i - 2t - 2} = 1^u$, $u < 0$; thus $C^{(t+1)} = \emptyset$. So $\text{ht}(C) = t$ and the assertion is proved. \square

The set of compositions with height $\geq t$ is precisely the set of t -conjugable compositions: $C^{(t)} \neq \emptyset \iff \text{ht}(C) \geq t$.

Let $G_t(n)$ be the set of compositions of n with conjugate height t and let $C_t(n)$ be the set of t -conjugable compositions of n with $g_t(n) = |G_t(n)|$ and $c_t(n) = |C_t(n)|$. Then we have $g_1(n) \geq g_2(n) \geq \dots \geq g_{n-2}(n) \geq g_x(n)$, $x \geq n - 1$, even though $G_i(n) \cap G_j(n) = \emptyset$, $i \neq j$.

However, $c_1(n) \geq c_2(n) \geq \dots \geq c_{n-2}(n) \geq c_x(n)$ with $C_1(n) \supseteq C_2(n) \supseteq \dots \supseteq C_{n-2}(n) \supseteq C_x(n)$. Hence $c_1(n) = c(n) = 2^{n-1}$, and

$$C_t(n) = \bigcup_{i \geq t} G_i(n) \quad \text{and} \quad G_t(n) = C_t(n) \setminus C_{t+1}(n). \quad (6)$$

For example, $g_1(6) = 22$, $g_2(6) = 4$, $g_3(6) = 2$, $g_4(6) = 2$ and $g_x(6) = 2$, $x \geq 5$, where the enumerated sets are shown below.

$G_1(6)$: (2, 4), (4, 2), (1, 2, 3), (1, 3, 2), (2, 1, 3), (2, 2, 2), (2, 3, 1), (3, 1, 2), (3, 2, 1),
(1, 1, 2, 2), (1, 1, 3, 1), (1, 2, 1, 2), (1, 2, 2, 1), (1, 3, 1, 1), (2, 1, 1, 2), (2, 1, 2, 1),
(2, 2, 1, 1), (1, 1, 1, 1, 2), (1, 1, 1, 2, 1), (1, 1, 2, 1, 1), (1, 2, 1, 1, 1), (2, 1, 1, 1, 1);

$G_2(6)$: (3, 3), (1, 4, 1), (1, 1, 1, 3), (3, 1, 1, 1);

$G_3(6)$: (1, 1, 4), (4, 1, 1);

$G_4(6)$: (1, 5), (5, 1);

$G_x(6)$: (6), (1, 1, 1, 1, 1, 1).

If C and E are compositions of n with $\text{ht}(C) = \text{ht}(E) = t$, then $\text{ht}(C^{(t)}) \neq \text{ht}(E^{(t)})$ in general. For example consider the 2-conjugates of $C = (1^4, 5, 1^8, 4, 1^6)$ and $E = (1^4, 5, 1^5, 7, 1^6)$. Then $C'' = (6, 1, 12, 1^0, 8)$ satisfies $\text{ht}(C'') = 5$ while $E'' = (6, 1, 9, 1^3, 8)$ gives $\text{ht}(E'') = 4$.

When $C^{(t)} \neq \emptyset$, it is clear that $\text{ht}(C) \leq \text{ht}(C^{(t)})$; but when do we have $\text{ht}(C) = \text{ht}(C^{(t)})$?

The following criterion is obtained by applying Theorem 5 to $C^{(t)}$ with height m and the fact that the big parts of $C^{(t)}$ are determined by the strings of 1's in C .

Corollary 6. *Let C be a t -conjugable composition with height m ($m \geq t$). Then $C^{(t)}$ has height m if and only if*

(i) *every boundary string 1^a in C satisfies $a = m - t + 1$ and every interior string 1^a satisfies $a \geq 2(m - t)$, or*

(ii) *every boundary string 1^a in C satisfies $a \geq m - t + 1$ and every interior string 1^a satisfies $a \in \{2(m - t), 2(m - t) + 1\}$,*

In particular, when $m = t$, we obtain the next result.

Corollary 7. *Let C be a composition with height t . Then $C^{(t)}$ has height t if and only if C contains an isolated 1 or a pair of consecutive big parts (i.e., C is not reciprocal (see definition in Section 5)).*

Proposition 8. *If C is a composition with height $t \geq s > 0$, then $\text{ht}((C^{(t)})^{(s)}) = s$.*

In particular $\text{ht}((C^{(t)})') = 1$.

Proof. Assume that $\text{ht}(C) = t$. Then for some interior part $b \in \{2t, 2t + 1\}$, the operations $C \rightarrow C^{(t)} \rightarrow (C^{(t)})^{(s)}$ induce $b \rightarrow 1^{b-2t} \rightarrow b-2t+2s$ which imply that $b-2t+2s \in \{2s, 2s+1\}$. Alternatively for a boundary big part $b = t + 1$ we have $b \rightarrow 1^{b-t} \rightarrow b - t + s$ which imply that $b - t + s = s + 1$. \square

5 Reciprocal compositions and recursive conjugation

Let $C = (\dots, 1^{a_u}, b_v, 1^{a_{u+1}}, b_{v+1}, \dots)$ be a composition of n such that $a_i, b_i \geq 2$ for all i . We define the *reciprocal* of C , denoted by $r(C)$, to be the composition obtained by replacing each string 1^{a_i} with a_i and each big part b_i with 1^{b_i} , i.e.,

$$r(C) = (\dots, a_u, 1^{b_v}, a_{u+1}, 1^{b_{v+1}}, \dots), \quad a_i, b_i > 1.$$

Note that $r(C) = \emptyset$ if $a_i < 2$ for some i . Thus the reciprocal operation is an involution on the set of compositions with no adjacent big parts in which every string of 1's occurs with multiplicity ≥ 2 . If $r(C) \neq \emptyset$, we call C a *reciprocal composition*.

For example, $r((1^2, 4, 1^2, 3, 1^3)) = (2, 1^4, 2, 1^3, 3)$, and $r((1^2, 4, 1)) = \emptyset$.

Let $C = (1^{a_1}, b_1, \dots, 1^{a_s}, b_s)$. Then observe from (1) that when $t = 0$ in $C^{(t)}$ we obtain $r(C)$:

$$\begin{aligned} (C^{(t)})|_{t=0} &= C^{(0)} = (a_1 + 0, 1^{b_1-2(0)}, a_2 + 2(0), \dots, a_s + 2(0), 1^{b_s-0}) \\ &= (a_1, 1^{b_1}, a_2, \dots, a_s, 1^{b_s}) \\ &= r(C). \end{aligned}$$

However, the resulting expression for $r(C)$ may not be defined.

Thus it is intuitive to define $C^{(0)}$ as $r(C)$, so that when $r(C) \neq \emptyset$ we have

$$r(r(C)) = r(C^{(0)}) = (C^{(0)})^{(0)} = C.$$

Lemma 9. *If C is t -conjugable, then $C^{(j)}$ is reciprocal, where $j = 1, 2, \dots, t-1$:*

$$C^{(t)} \neq \emptyset \implies r(C^{(j)}) \neq \emptyset, \quad 1 \leq j < t. \quad (7)$$

This follows from the defining inequalities for $C^{(t)}$ in (1): each string of 1's in $C^{(j)}$ has the form 1^{b_i-2j} , $i < r$, or 1^{b_r-j} , and either exponent is at least 2 because $b_i \geq 2t$ and $b_r \geq t+1$ with $j < t$.

Equation (7) implies

$$\text{ht}(C) > t \iff r(C^{(t)}) \neq \emptyset.$$

Proposition 10. *Let C be a composition. Then*

$$\text{ht}(C) \geq t \text{ and } \text{ht}(r(C)) \geq u \implies \text{ht}(C^{(t)}) \geq t + u.$$

In particular $\text{ht}(C) \geq t$ and $r(C) \neq \emptyset \iff \text{ht}(C^{(t)}) > t$.

Proof. We have that $r(C) \neq \emptyset$ and $\text{ht}(C) \geq t \iff$ every 1^a has $a > 1 \iff$ every boundary big part b in $C^{(t)}$ satisfies $b > t+1$ and every interior big part b satisfies $b > 2t+1 \iff \text{ht}(C^{(t)}) > t$. \square

Note that $(C^{(t)})' \neq C^{(t+1)}$. The correct relation is

$$(C^{(t)})' = r(C^{(t-1)}), \quad t > 0.$$

This may be extended as follows:

Theorem 11. *We have*

$$(C^{(t)})^{(u)} = r(C^{(t-u)}), \quad 0 \leq u \leq t. \quad (8)$$

Note that $u = 0$ and $u = t$ require $C^{(t)}$ and C to be reciprocal respectively.

Proof. Compute directly:

$$C = (1^{a_1}, b_1, 1^{a_2}, b_2, \dots, 1^{a_s}, b_s).$$

$$C^{(t)} = (a_1 + t, 1^{b_1-2t}, a_2 + 2t, 1^{b_2-2t}, \dots, a_s + 2t, 1^{b_s-t}).$$

$$(C^{(t)})^{(u)} = (1^{a_1+t-u}, b_1 - 2t + 2u, 1^{a_2+2t-2u}, \dots, b_s - t + u) = r(C^{(t-u)}). \quad \square$$

The following restatement provides a recursive procedure for computing $C^{(t)}$.

Corollary 12. *We have*

$$C^{(t)} = r(C^{(t-u)})^{(u)}, \quad 0 \leq u \leq t. \quad (9)$$

Note that because of (7), Equation (8) or (9) holds whenever $C^{(t)}$ exists. The process can be iterated. Let $t = t_1 + \dots + t_s$, $t_i > 0$, $\forall i$. Then writing $r(C^{(t_i)})^{t_j}$ instead of $(r(C^{(t_i)}))^{t_j}$ we have

$$C^{(t)} = r(r(\dots(r(C^{(t_1)})^{(t_2)} \dots)^{(t_{s-1})})^{(t_s)}),$$

where the reciprocal operation is applied $s - 1$ times.

For example we obtain the 5-conjugate of $C = (1^3, 15, 1, 7)$ using the decomposition $5 = 2 + 1 + 2$:

$$\begin{aligned} C^{(5)} &= r(r(C'')')'' \\ &= r(r((5, 1^{11}, 5, 1^5))')'' \\ &= r((1^5, 11, 1^5, 5)')'' \\ &= r((6, 1^9, 7, 1^4))'' \\ &= (1^6, 9, 1^7, 4)'' \\ &= (8, 1^5, 11, 1^2). \end{aligned}$$

If $C^{(t+1)} \neq \emptyset$, then (9) implies $(C^{(t+1)})^{(t)} = r(C')$, $\forall t \geq 0$. But when $r(C) \neq \emptyset$, we have

$$(C^{(t)})^{(t+1)} = r(C)'. \quad (10)$$

Indeed since $r(C) \neq \emptyset$ the following computation is justified by Proposition 10 when

$C = (1^{a_1}, b_1, 1^{a_2}, \dots, 1^{a_r}, b_r)$ with $\text{ht}(C^{(t)}) > t$:

$$\begin{aligned} r(C)' &= ((a_1, 1^{b_1}, a_2, \dots, a_r, 1^{b_r}))' \\ &= (1^{a_1-1}, b_1 + 2, 1^{a_2-2}, \dots, 1^{a_r-2}, b_r + 1) \\ &= ((a_1 + t, 1^{b_1-2t}, a_2 + 2t, \dots, a_r + 2t, 1^{b_r-t}))^{(t+1)} \\ &= (C^{(t)})^{(t+1)}. \end{aligned}$$

Thus using (10) the reciprocal of a t -conjugable reciprocal composition has the following expression.

$$r(C) = ((C^{(t)})^{(t+1)})'.$$

In particular the following relation holds for every reciprocal composition C .

$$r(C) = ((C')'').$$

However, $r(C)' = (C')'' \iff \text{ht}(C') > 1$. Hence the next result follows.

Proposition 13. *A composition C is reciprocal if and only if $\text{ht}(C') \geq 2$. Hence the number of reciprocal compositions of n is equal to the number of 2-conjugable compositions of n (cf. see (18)).*

6 Zig-zag graphs and higher-order conjugates

Given a t -conjugable composition C we show how $C^{(t)}$ is obtained using zig-zag graphs. The zig-zag graph of C will be denoted by $\text{ZG}(C)$.

We emphasize that the zig-zag graph is primarily structured for the classical 1-conjugation. So $C^{(t)}$ may only be obtained indirectly through $\text{ZG}(C)$ when $t > 1$.

Generally $C^{(t)}$ is obtained from $\text{ZG}(C)$ by reading off the composition corresponding to the columns of the zig-zag graph of a certain (transitional) composition $X_t(C)$ satisfying $C^{(t)} = X_t(C)'$, where $\text{ZG}(X_t(C))$ is visually compatible with $\text{ZG}(C)$.

It is expected that $X_1(C) = C$. In general we set $X_t(C) = r(C^{(t-1)})$ because $C^{(t)} = r(C^{(t-1)})'$ from (9). The visual connection between $\text{ZG}(C)$ and $\text{ZG}(r(C^{(t-1)}))$ is evident from the following procedure.

Algorithm to obtain $\text{ZG}(r(C^{(t-1)}))$ from $\text{ZG}(C)$.

- (i) Draw $\text{ZG}(C)$ depicting the dots as open circles;
- (ii) Shade the first or last $b - t + 1$ circles if the first or last row contains $b > 1$ circles, and shade the middle $b - 2(t - 1)$ circles in every other row containing $b > 1$ circles.
- (iii) Straighten each ‘zig-zag’ string of unshaded circles and align them vertically so that each string of shaded circles is in a different row from an unshaded circle.

The figure obtained in step (iii) is $ZG(r(C^{(t-1)}))$, that is, $ZG((C^{(t)})')$.

This procedure is reversible as one may verify by switching the roles of C and $C^{(t)}$; note that $r((C^{(t)})^{t-1}) = r(r(C')) = C'$.

We illustrate the process with $C = (1^2, 8, 1, 6, 5)$. Assume that $t = 3$, that is, we wish to obtain C''' using $ZG(C)$. Then the first diagram in Figure 2 shows $ZG(C)$ after the execution of step (ii) of the algorithm; the second diagram shows $ZG(r(C''))$.

Now the conjugate C''' is given by the composition corresponding to the columns of $ZG(r(C''))$, that is, $C''' = (5, 1^2, 7, 6, 1^2)$.

Note that $r(C'') = r((1^2, 8, 1, 6, 5)'') = r((4, 1^4, 5, 1^2, 4, 1^3)) = (1^4, 4, 1^5, 2, 1^4, 3) = (C''')'$.




Figure 2: Zig-zag graphs of $C = (1^2, 8, 1, 6, 5)$ and $r(C'')$

7 Enumeration

Let $c_t(n, k)$ denote the number of t -conjugable k -compositions of n . Thus $c_t(n) = \sum_k c_t(n, k)$.

Note that if $t > 1$ and $n - t + 1 \leq k \leq n - 1$, then $c_t(n, k) = 0$ since the longest composition after (1^n) is $(1, \dots, 1, t + 1)$ or $(t + 1, 1, \dots, 1)$ and any composition of the form $(1, \dots, 1, x)$, $2 \leq x \leq t$ is not t -conjugable.

7.1 Formulas for the number of conjugable compositions

Following MacMahon [12, 13] we demonstrate how to use the line graph to obtain the formula for $c_t(n, k)$ for various types of compositions.

Let $c_t(n, k \mid P) :=$ number of t -conjugable k -compositions of n that satisfy condition P . For example, consider the function $c_t(n, k \mid \text{parts} > 1)$. Then an enumerated composition

has the form

$$C = (b_1, b_2, \dots, b_{k-1}, b_k), b_1, b_k \geq t + 1, b_i \geq 2t, 2 \leq i \leq k - 1.$$

The line graph of C contains $n - 2t - 2 \geq 0$ central lines as shown in (11).

$$\begin{array}{c} \overbrace{\text{---} \cdots \text{---}}^{n-2t-1 \text{ gaps}} \\ \underbrace{\text{---} \cdots \text{---}}_{t+1 \text{ lines}} \quad \underbrace{\text{---} \cdots \text{---}}_{n-2t-2 \text{ lines}} \quad \underbrace{\text{---} \cdots \text{---}}_{t+1 \text{ lines}} \end{array} \quad (11)$$

In order to specify C we create $k - 2$ adjacent strings of lines in-between the $n - 2t - 1$ central gaps such that the length of each string is at least $2t$. To achieve this we apply the known formula $w(n, k, v) = \binom{n-(v-1)(k-1)}{k}$ which is the number of subsets $\{n_1, \dots, n_k\} \subseteq \{1, \dots, n\}$ satisfying $|n_i - n_j| \geq v, i \neq j$ [5].

Hence the desired number of compositions is given by $w(n - 2t - 1, k - 1, 2t)$, that is,

$$c_t(n, k \mid \text{parts} > 1) = \binom{n - 2t - 1 - (2t - 1)(k - 2)}{k - 1}. \quad (12)$$

A similar method may be applied to the function $c_t(n, k \mid 2c2 \text{ and } s \text{ ones})$, that is, the number of t -conjugable $2c2$ compositions of n into k parts containing s interior 1's, $s \geq 0$. Thus $c_t(n, k \mid 2c2 \text{ and } 0 \text{ ones}) = c_t(n, k \mid \text{parts} > 1)$.

Using (11) we find that $c_t(n, k \mid 2c2 \text{ and } s \text{ ones}) = \binom{k-2}{s} w(n - s - 2t - 1, k - s - 1, 2t)$, that is,

$$c_t(n, k \mid 2c2 \text{ and } s \text{ ones}) = \binom{k - 2}{s} \binom{n - s - 2t - 1 - (2t - 1)(k - s - 2)}{k - s - 1},$$

where $n \geq 2t + 2 + s + 2t(k - 1 - s), k > 1$.

Further variations of the shape of C and the formulas are clearly possible.

For a general formula we need the generating function for $c_t(n, k)$.

Theorem 14. *Let $n \geq k > 1$ be integers. Then for all integers $t > 0$,*

$$\sum_{n=k}^{\infty} c_t(n, k) x^n = \frac{(x - x^2 + x^{t+1})^2 (x - x^2 + x^{2t})^{k-2}}{(1 - x)^k}.$$

In particular,

$$\sum_{n=k}^{\infty} c_1(n, k) x^n = \frac{x^k}{(1 - x)^k} = \sum_{n=k}^{\infty} \binom{n - 1}{k - 1} x^n.$$

Proof. We use the following decomposition of an enumerated composition C , where $k > 1$.

$$\begin{aligned} C = & (\text{a composition into one part} \in \{1, t + 1, t + 2, \dots\}) \\ & \times (\text{a } (k - 2)\text{-composition using the parts } 1, 2t, 2t + 1, \dots) \\ & \times (\text{a composition into one part} \in \{1, t + 1, t + 2, \dots\}). \end{aligned}$$

Hence the generating function is given by

$$(x + x^{t+1} + x^{t+2} + \dots)(x + x^{2t} + x^{2t+1} + \dots)^{k-2}(x + x^{t+1} + x^{t+2} + \dots),$$

That is,

$$\sum_{n=k}^{\infty} c_t(n, k)x^n = \left(x + \frac{x^{t+1}}{1-x}\right)^2 \left(x + \frac{x^{2t}}{1-x}\right)^{k-2}$$

which simplifies to the stated result. \square

Now if $\frac{(x - x^2 + x^{2t})^{k-2}}{(1-x)^k} = \sum_{n \geq k} V(n, k)x^n$, then it is a routine process to show that

$$V_t(n, k) = \sum_{r=0}^{n-k+2} \sum_{j=0}^{n-k+2} (-1)^{r+j} \binom{k-2}{r} \binom{n-j+1}{k-1} \binom{k-2-r}{j-r(2t-1)}. \quad (13)$$

Therefore the full generating function is given by

$$\begin{aligned} \sum_{n=k}^{\infty} c_t(n, k)x^n &= (x - x^2 + x^{t+1})^2 \sum_{n=k}^{\infty} V_t(n, k)x^n. \\ &= (x^2 - 2x^3 + x^4 + 2x^{t+2} - 2x^{t+3} + x^{2t+2}) \sum_{n=k}^{\infty} V_t(n, k)x^n. \end{aligned}$$

This gives the following formula.

Theorem 15. *We have*

$$\begin{aligned} c_t(n, k) &= V_t(n-2, k) - 2V_t(n-3, k) + V_t(n-4, k) \\ &\quad + 2V_t(n-t-2, k) - 2V_t(n-t-3, k) + V_t(n-2t-2, k), \end{aligned}$$

where $V_t(n, k)$ is defined in (13).

In particular since $c_1(n, k) = V_1(n-2, k)$ we equate this with a known formula to get the identity:

$$\binom{n-1}{k-1} = \sum_{r=0}^{n-k} \sum_{j=0}^{n-k} (-1)^{r+j} \binom{k-2}{r} \binom{n-j-1}{k-1} \binom{k-2-r}{j-r}.$$

The generating function for $c_t(n) = \sum_k c_t(n, k)$ is given by

$$\begin{aligned} \sum_{n=1}^{\infty} c_t(n)x^n &= (x + x^2 + \dots) + \sum_{k=2}^{\infty} \frac{(x - x^2 + x^{t+1})^2 (x - x^2 + x^{2t})^{k-2}}{(1-x)^k} \\ &= \frac{x + x^2 - x^{t+1}}{1-x-x^t}. \end{aligned}$$

Theorem 16. *We have*

$$\sum_{n=1}^{\infty} c_t(n)x^n = \frac{x(1+x-x^t)}{1-x-x^t}.$$

In order to obtain an explicit formula for $c_t(n)$ we first consider the function $A(n, t)$ given by

$$\frac{x}{1-x-x^t} = \sum_{n=1}^{\infty} A(n, t)x^n. \quad (14)$$

When $t = 1$ and $t = 2$ one immediately recognizes the generating functions of two famous enumerators in combinatorial mathematics, namely, $A(n, 1) = 2^{n-1}$, the number of all subsets of $\{1, \dots, n-1\}$ or the number of all compositions of n ; and $A(n, 2) = F_n$, the n th Fibonacci number ($F_1 = F_2 = 1$, $F_n = F_{n-1} + F_{n-2}$, $n > 2$).

For the general coefficient one can show that

$$A(n, t) = \sum_{i=0}^{\lfloor \frac{n-1}{t} \rfloor} \binom{n-1-(t-1)i}{i}, \quad n > 0, \quad (15)$$

which is a generalized ‘Narayana’s cows’ sequence [18, [A000930](#)].

Since $\sum_{n \geq 1} c_t(n)x^n = (1+x-x^t) \sum_{n \geq 1} A(n, t)x^n$, we obtain

$$c_t(n) = A(n, t) + A(n-1, t) - A(n-t, t). \quad (16)$$

The formula (16) may be simplified using (15) and the Pascal triangle recurrence.

Theorem 17. *Given any positive integers n, t with $1 \leq t \leq n-2$ we have*

$$c_t(n) = 2 \sum_{i=0}^{\lfloor \frac{n-2}{t} \rfloor} \binom{n-2-(t-1)i}{i}, \quad (17)$$

where $c_t(1) = 1$, $c_t(n) = 2$, $t \geq n-1$.

In particular we see that $c_1(n) = 2 \cdot 2^{n-2} = 2^{n-1}$, and

$$c_2(n) = 2 \sum_{i=0}^{\lfloor \frac{n-2}{2} \rfloor} \binom{n-2-i}{i} = 2F_{n-1}, \quad n > 1. \quad (18)$$

7.1.1 Recurrence relations

Theorem 16 translates into the following recurrence relation but we also give a direct combinatorial proof.

Theorem 18. *Given positive integers n, t with $n \geq 2t + 2$, then*

$$c_t(n) = c_t(n-1) + c_t(n-t),$$

$$c_t(1) = 1, c_t(n) = 2, 1 \leq n \leq t+1; c_t(n) = 2(n-t), t+2 \leq n \leq 2t+1.$$

Proof. Let $H_t(n)$ be the set enumerated by $c_t(n)$. A composition $C = (u_1, u_2, \dots)$ in $H_t(n)$, falls under one of the four categories:

1. $u_1 = 1$,
2. $u_1 = t + 1$,
3. $t + 2 \leq u_1 \leq 2t$, $t > 1$, and
4. $u_1 \geq 2t + 1$:

Delete $u_1 = 1$ from C in (1) or subtract 1 from u_1 in (3) to obtain a member of $H_t(n-1)$. Subtract t from u_1 in (2) or (4) to obtain a member of $H_t(n-t)$.

Conversely, if $(e_1, e_2, \dots) \in H_t(n-1)$ satisfies $e_1 = 1$ or $e_1 \geq 2t$, then $C = (1, e_1, e_2, \dots) \in H_t(n)$, otherwise $C = (1 + e_1, e_2, \dots) \in H_t(n)$. If $(e_1, e_2, \dots) \in H_t(n-t)$, then $(t + e_1, e_2, \dots) \in H_t(n)$.

The compositions (1^n) and (n) are t -conjugable for any integer $t > 0$. So by convention we set $c_1(1) = 1 = c_t(1)$ and $c_t(n) = 2, 1 \leq n \leq t+1$. Then for $t+2 \leq n \leq 2t+1$ we see that $c_t(n) = c_t(n-1) + 2$, where C is obtained by adding 1 to the big part in a composition enumerated by $c_t(n-1)$ or by inserting 1 into (1^{n-1}) or by using a member of $\{(1^{n-t-1}, t+1), (t+1, 1^{n-t-1})\}$. Then the recurrence $c_t(n) = c_t(n-1) + 2$, with $c_t(t+1) = 2$, has the solution $c_t(n) = 2(n-t)$. \square

Another recurrence is given in terms of the number of compositions with last part 1. Denote the number of compositions in $H_t(n)$ with last part x by $c_t(n)_x$. Then

$$c_t(n) = 2c_t(n)_1, c_t(t)_1 = 1.$$

Indeed we have $c_t(n) = c_t(n)_1 + c_t(n)_{>1}$. But since every composition enumerated by $c_t(n)_{>1}$ has a conjugate with last part 1, we see that $c_t(n)_{>1} = c_t(n)_1$. Hence the assertion follows.

7.2 Number of compositions with a fixed conjugate height

Let $g_t(n, k)$ denote the number of k -compositions of n with conjugate height exactly t so that $g_t(n) = \sum_k g_t(n, k)$.

The formulas for $g_t(n, k)$ and $g_t(n)$ are derived from those of $c_t(n, k)$ and $c_t(n)$ (cf. Equation (6)).

$$g_t(n, k) = c_t(n, k) - c_{t+1}(n, k), \quad n \geq t > 1,$$

$$g_t(n) = c_t(n) - c_{t+1}(n), \quad n \geq t > 1. \tag{19}$$

From Equations (19) and (17) we obtain

Corollary 19. *Given any positive integers n, t , then*

$$g_t(n) = 2 \sum_{i=0}^{\lfloor \frac{n-2}{t} \rfloor} \binom{n-2-(t-1)i}{i} - 2 \sum_{i=0}^{\lfloor \frac{n-2}{t+1} \rfloor} \binom{n-2-ti}{i}.$$

Note that $g_t(n) > 0$ only for $n \geq t+2$ in view of (3). The case $t = 1$ gives

$$g_1(n) = 2^{n-1} - 2F_{n-1}, \quad n > 1.$$

The following result is a consequence of the generating function of $c_t(n)$.

Corollary 20. *We have*

$$\sum_{n=1}^{\infty} g_t(n)x^n = \frac{2x^{t+2}(1-x)}{(1-x-x^t)(1-x-x^{t+1})}.$$

This generating function implies the following recurrence relation.

Corollary 21. *Given integers n, t with $t > 1$, $n \geq 2t+2$, we have*

$$\begin{aligned} g_t(n) &= 2g_t(n-1) - g_t(n-2) + g_t(n-t) - g_t(n-t-2) - g_t(n-2t-1), \\ g_t(n) &= 0, \text{ for } 1 \leq n \leq t+1, \quad g_1(4) = 4 \text{ and } g_t(n) = 2, \text{ for } t+2 \leq n \leq 2t+1. \end{aligned}$$

An open problem is to find a direct combinatorial proof of Corollary 21.

Figure 3 shows some values of the enumeration functions $c_t(n)$ and $g_t(n)$ for small n . The second columns of the tables give the sequences $c_1(n) = 2^{n-1}$ and $g_1(n) = 2^{n-1} - 2F_{n-1}$, $n > 0$, respectively.

We remark that the binomial coefficient sequences $\binom{n}{k}$, $n \geq k \geq 0$, in Sloane [18] are reproduced here in the form of the enumeration function $c_1(n, k)$ (see Theorem 14).

The 2-conjugable compositions of n are enumerated by the double Fibonacci numbers $2F_n$ (A055389). The sequences $\frac{1}{2}c_t(n)$, $t \geq 1$ occur in [18] as a family of sequences that satisfy the recurrence of Theorem 18: [A000045](#), [A000079](#), [A000930](#), [A003269](#), [A003520](#), [A005708](#), [A005709](#), [A005710](#), [A005711](#).

The functions $c_t(n, 2)$, $t = 1, 2, 3$, $n > t$ agree with [A233583](#) for suitable sets of initial values, while $c_2(n, 3)$, $n > 7$ coincides with [A145018](#). On the other hand, $\frac{1}{2}g_t(n)$ is listed only for $t = 1$, namely [A027934](#).

Lastly we note that the following sequences are given in [18] as coefficients in the expansions of the rational function $\frac{(1-x)}{1-x-x^t}$, $t > 3$ (cf. Equation (14)): [A017898](#), [A017899](#), [A017900](#), [A017901](#), [A017902](#), [A017903](#), [A017904](#).

		$c_t(n)$							
	t	1	2	3	4	5	6	7	8
1	1	1	1	1	1	1	1	1	1
2	2	2	2	2	2	2	2	2	2
3	4	2	2	2	2	2	2	2	2
4	8	4	2	2	2	2	2	2	2
5	16	6	4	2	2	2	2	2	2
6	32	10	6	4	2	2	2	2	2
7	64	16	8	6	4	2	2	2	2
8	128	26	12	8	6	4	2	2	2
9	256	42	18	10	8	6	4	2	2
10	512	68	26	14	10	8	6	4	4

		$g_t(n)$							
	t	1	2	3	4	5	6	7	8
1	1	0	0	0	0	0	0	0	0
2	0	0	0	0	0	0	0	0	0
3	2	0	0	0	0	0	0	0	0
4	4	2	0	0	0	0	0	0	0
5	10	2	2	0	0	0	0	0	0
6	22	4	2	2	0	0	0	0	0
7	48	8	2	2	2	0	0	0	0
8	102	14	4	2	2	2	0	0	0
9	214	24	8	2	2	2	2	0	0
10	444	42	12	4	2	2	2	2	2

Figure 3: The numbers $c_t(n)$ and $g_t(n)$ for $1 \leq n \leq 10$ and $1 \leq t \leq 8$

8 Combinatorial identities

Call a composition s -separated if every pair of consecutive big parts is separated by s ones, i.e., C has the form $C = (\dots, b_i, 1^s, b_{i+1}, 1^s, b_{i+2}, 1^s, \dots)$, $b_j > 1, \forall j$.

Then the following result is a consequence of conjugation.

Lemma 22. *The number of t -conjugable $2c2$ compositions of n that are s -separated is equal to the number of t -conjugable $1c1$ compositions of n using only the parts 1 and $2t + s$:*

$$c_t(n \mid 2c2, s\text{-separated}) = c_t(n \mid 1c1, \text{parts } 1, 2t + s).$$

For a generating function proof we have

$$\begin{aligned} \sum_{n=2}^{\infty} c_t(n, k \mid 1c1, \text{parts } 1, 2t + s) x^n &= (x)(x + x^{2t+s})^{k-2}(x), \quad k \geq 2; \\ \sum_{n=r}^{\infty} c_t(n \mid 2c2, s\text{-separated}, r \text{ parts } > 1) x^n, \quad r \geq 2 \\ &= (x^{t+1} + x^{t+2} + \dots)(x^{2t} + x^{2t+1} + \dots)^{r-2}(x^{t+1} + x^{t+2} + \dots)(x^s)^{r-1} \\ &= \left(\frac{x^{t+1}}{1-x}\right)^2 \left(\frac{x^{2t}}{1-x}\right)^{r-2} x^{s(r-1)}. \end{aligned}$$

The lemma then follows from the easily verified identity

$$\sum_{k \geq 2} x^2 (x + x^{2t+s})^{k-2} = \frac{x^2}{1-x} + \sum_{r=2}^{\infty} \left(\frac{x^{t+1}}{1-x}\right)^2 \left(\frac{x^{2t}}{1-x}\right)^{r-2} x^{s(r-1)} = \frac{x^2}{1-x-x^{2t+s}},$$

where $(x^2 + x^3 + \dots) = x^2/(1 - x)$ is the generating function for 1-part $2c2$ compositions.

Since a positive integer may have different compositions of the form $(2t, s)$ for different choices of t and s , the next assertion follows from Lemma 22.

Theorem 23. *Let $2t + s = 2h + u$ for positive integers t, s, h, u . Then*

$$c_t(n \mid 2c2, s\text{-separated}) = c_h(n \mid 2c2, u\text{-separated}), \quad \forall n > 1.$$

For example, since $5 = 2 \cdot 1 + 3 = 2 \cdot 2 + 1$, we have $c_1(n \mid 2c2, 3\text{-separated}) = c_2(n \mid 2c2, 1\text{-separated})$ for all $n > 1$. Thus when $n = 18$, some (bijective) pairs of enumerated objects follow:

$$c_1(18 \mid 2c2, 3\text{-separated}) : (18), (4, 1^3, 11), (3, 1^3, 7, 1^3, 2), (2, 1^3, 3, 1^3, 2, 1^3, 2), \dots;$$

$$c_2(18 \mid 2c2, 1\text{-separated}) : (18), (5, 1, 12), (4, 1, 9, 1, 3), (3, 1, 5, 1, 4, 1, 3), \dots$$

The case $s = 0$ in Lemma 22 leads to a 5-way combinatorial identity. We provide only bijective proofs below.

Theorem 24. *Let t be a positive integer. Then the following enumeration functions are equal.*

$$(i) \quad c_t(n \mid \text{no } 1\text{'s});$$

$$(ii) \quad c_t(n - 2 \mid \text{parts } 1, 2t);$$

$$(iii) \quad c_{2t}(n \mid \text{first part } 1);$$

$$(iv) \quad c_t(n - 1 \mid \text{parts} \equiv 1 \pmod{2t});$$

$$(v) \quad c_t(n + 2t - 2 \mid \text{parts} \geq 2t).$$

The common value, from Equation (12), is given by

$$\sum_k \binom{n - 2t - 1 - (2t - 1)(k - 2)}{k - 1}.$$

Proof. We will replace the lower-case letter c in each enumeration function with the upper-case C to denote the corresponding enumerated set.

(i) \iff (ii). There is an obvious bijection

$$\alpha : C_t(n - 2 \mid \text{parts } 1, 2t) \rightarrow C_t(n \mid 1c1, \text{parts } 1, 2t), \quad (20)$$

where $\alpha((c_1, \dots, c_k)) = (1, c_1, \dots, c_k, 1)$. Thus (i) \iff (ii) is essentially a restatement of Lemma 22 with $s = 0$.

(ii) \iff (iii). By (20) and Lemma 22 it will suffice to establish the bijection

$$\varphi : C_t(n \mid 1c1, \text{parts } 1, 2t) \rightarrow C_{2t}(n \mid \text{1st part } 1).$$

Each $U \in C_t(n \mid 1c1, \text{parts } 1, 2t)$ has the symbolic form

$$U = (1^{a_1}, 2t, 1^{a_2}, 2t, \dots, 1^{a_r}, 2t, 1^{a_{r+1}}), \quad a_1, a_{r+1} \geq 1, \quad a_i \geq 0, \quad 1 < i < r + 1.$$

Now define φ as follows.

If r is even, then $\varphi(U) = (1^{a_1}, 4t + a_2, 1^{a_3}, 4t + a_4, 1^{a_5}, \dots, 4t + a_r, 1^{a_{r+1}})$, giving a 1c1 member of $C_{2t}(n \mid \text{1st part } 1)$.

If r is odd, then $\varphi(U) = (1^{a_1}, 4t + a_2, 1^{a_3}, 4t + a_4, 1^{a_5}, \dots, 4t + a_{r-1}, 1^{a_r}, 2t + a_{r+1})$, giving a 1c2 member of $C_{2t}(n \mid \text{1st part } 1)$. Clearly φ is one-to-one.

Conversely, since each interior big part b of any $E \in C_{2t}(n \mid \text{1st part } 1)$ satisfies $b \geq 4t$, it splits into three symbolic parts $(2t, 1^a, 2t)$, $a \geq 0$. Correspondingly a boundary big part b of E satisfies $b \geq 2t + 1$ and splits into $(2t, 1^a)$, $a > 0$. So E can be transformed into a unique member of $C_t(n \mid 1c1, \text{parts } 1, 2t)$. Hence the existence of φ^{-1} is guaranteed.

(ii) \iff (iv). An integer of the form $2tj + 1$, $j > 0$ has a composition of the form $(1, 2t, \dots, 2t)$. If we apply this transformation to each big part of $C \in C_t(n - 1 \mid \text{parts } \equiv 1 \pmod{2t})$, and delete the first part ($= 1$), we obtain a unique member of $C_t(n - 2 \mid \text{parts } 1, 2t)$.

(i) \iff (v). Since any $(c_1, \dots, c_k) \in C_t(n \mid \text{no } 1\text{'s})$ satisfies $c_1, c_k > t$ and $c_i \geq 2t$, $1 < i < k$, it follows that $(c_1 + t - 1, c_2, \dots, c_{k-1}, c_k + t - 1) \in C_t(n + 2t - 2 \mid \text{parts } \geq 2t)$.

□

9 Acknowledgment

The author is grateful to the editor and the referee for valuable comments which facilitated the revision of this paper. This work is based on research supported in part by the National Research Foundation of South Africa (Grant number 115024).

References

- [1] A. K. Agarwal, n -color compositions, *Indian J. Pure Appl. Math.* **31** (2000), 1421–1427.
- [2] G. E. Andrews, *The Theory of Partitions*, Cambridge University Press, 1998.
- [3] E. A. Bender and E. R. Canfield, Locally restricted compositions I. Restricted adjacent differences, *Electron. J. Combin.* **12** (2005), Paper R57.

- [4] E. A. Bender and E. R. Canfield, Locally restricted compositions II. General restrictions and infinite matrices, *Electron. J. Combin.* **16** (2009), Paper R108.
- [5] C. Chuan-Chong and K. Khee-Meng, *Principles and Techniques in Combinatorics*, World Scientific, 1992.
- [6] S. Eger, Restricted weighted integer compositions and extended binomial coefficients, *J. Integer Seq.* **16** (2013), [Article 13.1.3](#).
- [7] S. Eger, Some Elementary Congruences for the number of weighted integer compositions, *J. Integer Seq.* **18** (2015), [Article 15.4.1](#).
- [8] Y. Guo, Inverse-conjugate compositions with odd parts, *Ars Combinatoria*, **135** (2017), 93–102.
- [9] S. Heubach and T. Mansour, *Combinatorics of Compositions and Words*, Discrete Mathematics and its Applications, CRC Press, 2010.
- [10] S. Heubach and T. Mansour, Compositions of n with parts in a set, *Congressus Numerantium* **168** (2004), 127–143.
- [11] B. Hopkins, Spotted tilings and n -color compositions, *Integers* **12B** (2012), Paper A6.
- [12] P. A. MacMahon, *Combinatory Analysis*, Volume 1, Cambridge University Press, 1915
- [13] P. A. MacMahon, Memoir on the theory of the compositions of numbers, *Philos. Trans. Roy. Soc. London Ser. A*, **184** (1893), 835–901.
- [14] P. A. MacMahon, Second memoir on the theory of the compositions of numbers, *Philos. Trans. Roy. Soc. London Ser. A*, **207** (1908), 65–134.
- [15] A. O. Munagi, Euler-type identities for integer compositions via zig-zag graphs, *Integers* **12** (2012), Paper A60.
- [16] A. O. Munagi, Primary classes of compositions of numbers, *Ann. Math. Inform.* **41** (2013), 193–204.
- [17] A. O. Munagi, Zig-zag graphs and partition identities of A. K. Agarwal, *Ann. Comb.* **19** (2015), 557–566.
- [18] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>.

2000 *Mathematics Subject Classification*: Primary: 05A17 Secondary: 11P81, 05A15, 05A19.

Keywords: composition, zig-zag graph, line graph, direct detection, conjugate, height, reciprocal.

(Concerned with sequences: [A000045](#), [A000079](#), [A000930](#), [A003269](#), [A003520](#), [A005708](#), [A005709](#), [A005710](#), [A005711](#), [A017898](#), [A017899](#), [A017900](#), [A017901](#), [A017902](#), [A017903](#), [A017904](#), [A027934](#), [A055389](#), [A145018](#), and [A233583](#).)

Received May 29 2018; revised versions received August 30 2018; August 31 2018. Published in *Journal of Integer Sequences*, November 25 2018.

Return to [Journal of Integer Sequences home page](#).