

INCLUSION THEOREMS FOR THE SPACES \mathcal{F}_α

Stojan Duborija

Abstract. We give a new proof of one known inclusion theorem for the space \mathcal{F}_0 that enables us to extend this theorem from the unit disc in \mathbf{C} to the unit ball in \mathbf{C}^n , $n > 1$. We also improve an inclusion relation between Bergman spaces and the spaces \mathcal{F}_α , $\alpha > 0$.

1. Introduction

Let Δ denote the unit disc in \mathbf{C} with boundary T . We denote by m the Lebesgue measure on Δ and by σ the normalized Lebesgue measure on T .

Let \mathcal{M} denote the set of complex valued Borel measures on T . For each $\alpha \geq 0$ let $\mathcal{F}_\alpha = \mathcal{F}_\alpha(\Delta)$ denote the family of functions on Δ having the property that there exists a measure $\mu \in \mathcal{M}$ such that

$$f(z) = \int_T K_\alpha(z\bar{\xi}) d\mu(\xi), \quad z \in \Delta, \quad (1.1)$$

where for $\alpha > 0$, $K_\alpha(w) = (1-w)^{-\alpha}$, $w \in \Delta$, and $K_0(w) = 1 + \log \frac{1}{1-w}$, $w \in \Delta$. In (1.1) and throughout this paper each logarithm means the principal branch. The family \mathcal{F}_α is a Banach space with respect to the norm defined by $\|f\|_{\mathcal{F}_\alpha} = \inf \|\mu\|$, where μ varies over all measures in \mathcal{M} for which (1.1) holds and where $\|\mu\|$ denotes the total variation norm of μ .

A function f holomorphic in Δ (abbreviated $f \in H(\Delta)$) is said to belong to the Hardy space H^p , $0 < p < \infty$, if $\|f\|_{H^p} = \sup_{0 < r < 1} M_p(r, f) < \infty$ and to the weighted Bergman space $A^{p,\alpha}$, $0 < p < \infty$, $\alpha > -1$, if

$$\|f\|_{A^{p,\alpha}}^p = \int_0^1 (1-r)^\alpha M_p^p(r, f) dr < \infty.$$

Here $M_p(r, f)$ is the $L^p(\{z \in \Delta : |z| = r\})$ “norm” of f .

Let $g(z) = \sum_{k=0}^\infty a_k z^k$ be holomorphic in Δ . We define the multiplier transformation $D^\beta g$ of g , where β is a real number, by

$$D^\beta g(z) = \sum_{k=0}^\infty (k+1)^\beta a_k z^k.$$

AMS Subject Classification: 30 E 20, 30 D 50, 30 D 55

Keywords and phrases: Fractional Cauchy integral, Hardy space, BMOA, Bergman space.

A function $f \in H(\Delta)$ is said to belong to the space H_s^p if $\|D^s f\|_{H^p} < \infty$ and to the space $A_s^{p,\alpha}$ if $\|D^s f\|_{A^{p,\alpha}} < \infty$.

For an arc $I \subset T$ and a function $g \in H^1$ let $MO(g, I)$ denote the mean oscillation of g over I , i.e.

$$MO(g, I) = \frac{1}{\sigma(I)} \int_I \left| g(\xi) - \frac{1}{\sigma(I)} \int_I g(\eta) d\sigma(\eta) \right| d\sigma(\xi),$$

and let, as usual, $BMOA = BMOA(\Delta)$ be the space of functions $g \in H^1$ such that $\|g\|_{BMOA} = \sup MO(g, I) < \infty$, where the supremum runs over all arcs $I \subset T$.

We are now ready to state our first result

THEOREM 1. $\mathcal{F}_0(\Delta) \subset BMOA(\Delta)$.

Theorem is known (see [5]). The argument given in [5] is limited to the one dimensional case. Our proof is different and may be easily extended to the corresponding spaces $\mathcal{F}_0(B^n)$ and $BMOA(B^n)$ of holomorphic functions on the unit ball B^n in \mathbf{C}^n , $n > 1$. Thus we have

THEOREM 2. $\mathcal{F}_0(B^n) \subset BMOA(B^n)$.

In [6] it is shown that if $\alpha > 0$, then $A_1^{1,\alpha-1} \subset \mathcal{F}_\alpha$ (Lemma 1 and Lemma 2, pp. 159–160). Since $A_1^{1,\alpha-1} \subset H_{1-\alpha}^1$ and $H_{1-\alpha}^1 \setminus A_1^{1,\alpha-1}$ is a non-empty set, the following theorem is an improvement of this inclusion.

THEOREM 3. Let $\alpha > 0$. Then $H_{1-\alpha}^1 \subset \mathcal{F}_\alpha$.

2. Proof of Theorem 1.

Let $f \in \mathcal{F}_0$. Then there exists $\mu \in \mathcal{M}$ such that

$$f(z) = f(0) + \int_T \log \frac{1}{1 - z\xi} d\mu(\xi). \quad (2.1)$$

To show that $f \in BMOA$ it suffices to show that

$$\sup_{z \in \Delta} \int_{\Delta} \frac{1 - |z|^2}{|1 - z\bar{w}|^2} |f'(w)|^2 (1 - |w|^2) dm(w) < \infty,$$

(see [4], p. 240). Using (2.1) we find that

$$\int_{\Delta} \frac{1 - |z|^2}{|1 - z\bar{w}|^2} |f'(w)|^2 (1 - |w|^2) dm(w) \leq C \int_T d|\mu|(\xi) \int_{\Delta} \frac{(1 - |z|^2)(1 - |w|^2) dm(w)}{|1 - z\bar{w}|^2 |1 - \bar{w}\xi|^2}.$$

Since $|\mu|(T) < \infty$, it is sufficient to show that

$$\int_{\Delta} \frac{(1 - |w|^2) dm(w)}{|1 - z\bar{w}|^2 |1 - \bar{w}\xi|^2} \leq \frac{C}{1 - |z|^2}, \quad \text{for all } z \in \Delta \text{ and } \xi \in T. \quad (2.2)$$

In this note we follow the custom of using letters C, C_1, C_2, \dots , to stand for positive constants which change their values from one occurrence to another while remaining independent of the important variables.

Given $z \in \Delta$ and $\xi \in T$ we consider the following partition of Δ :

$$\begin{aligned}\Omega_1 &= \{ w \in \Delta : |1 - z\bar{w}| \leq \frac{1}{2}|1 - z\bar{\xi}| \}, \\ \Omega_2 &= \{ w \in \Delta : |1 - w\bar{\xi}| \leq \frac{1}{2}|1 - z\bar{\xi}| \}, \\ \Omega_3 &= \{ w \in \Delta : \frac{1}{2}|1 - z\bar{\xi}| < |1 - z\bar{w}| \leq |1 - w\bar{\xi}| \}, \\ \Omega_4 &= \{ w \in \Delta : \frac{1}{2}|1 - z\bar{\xi}| < |1 - w\bar{\xi}| \leq |1 - w\bar{z}| \}.\end{aligned}$$

With this notation we have

$$\begin{aligned}|1 - \bar{w}z| &\leq C_1|1 - z\bar{\xi}| \leq C_2|1 - \bar{w}\xi|, & w \in \Omega_1, \\ |1 - \bar{w}\xi| &\leq C_1|1 - z\bar{\xi}| \leq C_2|1 - z\bar{w}|, & w \in \Omega_2, \\ |1 - z\bar{\xi}| &\leq C_1|1 - z\bar{w}| \leq C_2|1 - \bar{w}\xi|, & w \in \Omega_3, \\ |1 - z\bar{\xi}| &\leq C_1|1 - \bar{w}\xi| \leq C_2|1 - \bar{w}z|, & w \in \Omega_4.\end{aligned}$$

Using this we find

$$\begin{aligned}\int_{\Omega_1} \frac{(1 - |w|^2) dm(w)}{|1 - z\bar{w}|^2 |1 - \xi\bar{w}|^2} &\leq C \int_{\Delta} \frac{(1 - |w|^2) dm(w)}{|1 - z\bar{w}|^4} \\ &\leq \int_0^1 (1 - \rho) d\rho \int_T \frac{|d\eta|}{|1 - z\rho\bar{\eta}|^4} \leq C \int_0^1 \frac{(1 - \rho) d\rho}{(1 - \rho|z|)^3} \leq \frac{C}{1 - |z|},\end{aligned}\quad (2.3)$$

$$\begin{aligned}\int_{\Omega_2} \frac{(1 - |w|^2) dm(w)}{|1 - z\bar{w}|^2 |1 - \xi\bar{w}|^2} &\leq C \int_{\Omega_2} \frac{(1 - |w|^2) dm(w)}{(|1 - z\bar{\xi}| + |1 - w\bar{\xi}|)^2 |1 - \bar{w}\xi|^2} \\ &\leq C \int_{\Delta} \frac{(1 - |w|^2) dm(w)}{(|1 - z\bar{\xi}| + (1 - |w|))^2 |1 - \bar{w}\xi|^2} \leq C \int_0^1 \frac{d\rho}{(|1 - z\bar{\xi}| + 1 - \rho)^2} \\ &\leq \frac{C}{|1 - z\bar{\xi}|} \int_0^\infty \frac{dx}{(1 + x)^2} \leq \frac{C}{|1 - z\bar{\xi}|} \leq \frac{C}{1 - |z|^2},\end{aligned}\quad (2.4)$$

$$\int_{\Omega_3} \frac{(1 - |w|^2) dm(w)}{|1 - z\bar{w}|^2 |1 - \xi\bar{w}|^2} \leq C \int_{\Omega_3} \frac{(1 - |w|^2) dm(w)}{|1 - z\bar{w}|^4} \leq \frac{C}{1 - |z|^2},\quad (2.5)$$

$$\begin{aligned}\int_{\Omega_4} \frac{(1 - |w|^2) dm(w)}{|1 - z\bar{w}|^2 |1 - \xi\bar{w}|^2} &\leq C \int_{\Omega_4} \frac{(1 - |w|^2) dm(w)}{(|1 - z\bar{\xi}| + |1 - \bar{w}\xi|)^2 |1 - \bar{w}\xi|^2} \\ &\leq C \int_{\Delta} \frac{(1 - |w|^2) dm(w)}{(|1 - z\bar{\xi}| + (1 - |w|))^2 |1 - \bar{w}\xi|^2} \leq \frac{C}{|1 - z\bar{\xi}|} \leq \frac{C}{1 - |z|^2}.\end{aligned}\quad (2.6)$$

Now (2.2) follows from (2.3), (2.4), (2.5) and (2.6). This finishes the proof of Theorem 1. ■

3. Proof of Theorem 3.

Let α be a positive real numbers. Define the function H_α by

$$H_\alpha(z) = \sum_{n=0}^{\infty} (n+1)^{\alpha-1} z^n, \quad \text{for } z \in \Delta.$$

Let \mathcal{H}_α denote the family of functions $f(z) = \sum_{k=0}^{\infty} a_k z^k$, $z \in \Delta$, having the property that there exists $\mu \in \mathcal{M}$ such that

$$a_n = (n+1)^{\alpha-1} \int_T \bar{\xi}^n d\mu(\xi), \quad n = 0, 1, 2, \dots \quad (3.1)$$

Let $\|f\|_{\mathcal{H}_\alpha} = \inf \|\mu\|$, where μ varies over all members of \mathcal{M} for which (3.1) holds. Then \mathcal{H}_α is a Banach space.

For the proof of Theorem 3 the following lemma is needed.

LEMMA 3.1. *If $\alpha > 0$ then $f \in \mathcal{F}_\alpha$ if and only if $f \in \mathcal{H}_\alpha$. There is a positive constant C depending only on α such that if $f \in \mathcal{F}_\alpha$ then*

$$C^{-1} \|f\|_{\mathcal{F}_\alpha} \leq \|f\|_{\mathcal{H}_\alpha} \leq C \|f\|_{\mathcal{F}_\alpha}.$$

Proof. Suppose that $f \in \mathcal{F}_\alpha$ and $f(z) = \sum_{n=0}^{\infty} a_n z^n$ for $z \in \Delta$. Then there exists $\mu \in \mathcal{M}$ such that

$$a_n = A_n(\alpha) \int_T \bar{\xi}^n d\mu(\xi), \quad n = 0, 1, 2, \dots, \quad (3.2)$$

where $A_n(\alpha) = \frac{\Gamma(n+\alpha)}{\Gamma(\alpha)\Gamma(n+1)}$, $n = 0, 1, 2, \dots$. From this it follows that

$$A_n(\alpha) = (n+1)^{\alpha-1} \left(\frac{1}{\Gamma(\alpha)} + B_n(\alpha) \right), \quad \text{for } n = 1, 2, \dots, \quad (3.3)$$

and there is a positive constant $B(\alpha)$ such that $|B_n(\alpha)| \leq B(\alpha)/n$, for $n = 1, 2, \dots$. For $n = 1, 2, \dots$ let

$$c_n(\alpha) = B_n(\alpha) \int_T \bar{\xi}^n d\mu(\xi)$$

and define the function g by $g(z) = \sum_{n=1}^{\infty} c_n(\alpha) z^n$, for $z \in \Delta$. Since $|c_n(\alpha)| \leq |B_n(\alpha)| \|\mu\| \leq \frac{B(\alpha)}{n} \|\mu\|$, $g \in H^2$. Therefore $g \in \mathcal{F}_1$ (see [2]) and hence there exists $\nu \in \mathcal{M}$ such that

$$g(z) = \int_T \frac{d\nu(\xi)}{1-z\xi}.$$

This implies that

$$c_n(\alpha) = \int_T \bar{\xi}^n d\nu(\xi), \quad \text{for } n=1, 2, \dots$$

Thus

$$a_n = (n+1)^{\alpha-1} \left(\frac{1}{\Gamma(\alpha)} \int_T \bar{\xi}^n d\mu(\xi) + \int_T \bar{\xi}^n d\nu(\xi) \right), \quad \text{for } n = 1, 2, \dots$$

Let $\lambda = \frac{1}{\Gamma(\alpha)}\mu + \nu + b\sigma$, where $b = \mu(T)(\frac{-1}{\Gamma(\alpha)} + 1) - \nu(T)$. Then

$$a_n = (n+1)^{\alpha-1} \int_T \bar{\xi}^n d\lambda(\xi), \quad n = 0, 1, 2, \dots$$

Since $\lambda \in \mathcal{M}$, $f \in \mathcal{H}_\alpha$.

The argument given above shows that

$$\|f\|_{\mathcal{H}_\alpha} \leq \frac{1}{\Gamma(\alpha)} \|\mu\| + \|\nu\| + |b| \leq C(\|\mu\| + \|\nu\|) \leq C\|\mu\|.$$

Here we have used again that $H^2 \subset H^1 \subset \mathcal{F}_1$. This inequality holds for every $\mu \in \mathcal{M}$ for which (3.2) holds. Hence $\|f\|_{\mathcal{H}_\alpha} \leq C\|f\|_{\mathcal{F}_\alpha}$.

The same argument shows that if $f \in \mathcal{H}_\alpha$ then $f \in \mathcal{F}_\alpha$ and $\|f\|_{\mathcal{F}_\alpha} \leq C\|f\|_{\mathcal{H}_\alpha}$. Instead of (3.3) it should use the relation

$$(n+1)^{\alpha-1} = A_n(\alpha)[\Gamma(\alpha) + D_n(\alpha)], \quad \text{for } n = 1, 2, \dots, \quad (3.4)$$

and $|D_n(\alpha)| \leq \frac{D(\alpha)}{n}$, for some positive constant $D(\alpha)$. For (3.3) and (3.4), see [3] and [7]. ■

Proof of Theorem 3. Let $f \in H_{1-\alpha}^1$ and $f(z) = \sum_{k=0}^n a_k z^k$, $z \in \Delta$. Then $D^{1-\alpha}f \in H^1$. Since $H^1 \in \mathcal{F}_1$ we have $D^{1-\alpha}f \in \mathcal{F}_1$. Hence there exists a measure $\mu \in \mathcal{M}$ such that

$$(n+1)^{1-\alpha} a_n = \int_T \bar{\xi}^n d\mu(\xi), \quad n = 0, 1, 2, \dots,$$

or equivalently

$$a_n = (n+1)^{\alpha-1} \int_T \bar{\xi}^n d\mu(\xi), \quad n = 0, 1, 2, \dots$$

Therefore, $f \in \mathcal{H}_\alpha$ and by Lemma 3.1 we have $f \in \mathcal{F}_\alpha$. Thus, $H_{1-\alpha}^1 \subset \mathcal{F}_\alpha$. ■

REFERENCES

- [1] Baernstein, A. II, *Analytic functions of bounded mean oscillation*, in: *Aspects of Contemporary Complex Analysis*, Academic Press, New York 1981, pp. 3–36.
- [2] Duren, P. L., *Theory of H^p Spaces*, Academic Press, New York 1970.
- [3] Erdelyi, A., *Higher Transcendental Functions, I*, McGraw Hill, 1953.
- [4] Garnett, J., *Bounded Analytic Functions*, Academic Press, New York 1981.
- [5] Hallenbeck, D. J., Samotij, K., *On Cauchy integrals of logarithmic potentials and their multipliers*, J. Math. Anal. Appl. **194** (1993), 614–634.
- [6] Hallenbeck D. J., MacGregor, T. H., Samotij, K., *Cauchy transforms, inner functions and multipliers*, Proc. London Math. Soc. **72** (1996), 157–187.
- [7] Hibscheiller, R., MacGregor, T. H., *Bounded analytic families of Cauchy-Stieltjes integrals*, Rocky Mountain J. Math. **23** (1993), 187–202.
- [8] MacGregor, T. H., *Analytic and univalent functions with integral representations involving complex measures*, Indiana Univ. Math. J. **36** (1987), 109–130.

(received 20.01.2000.)

Prirodno-matematički fakultet, Cetinjski put b.b., 81000 Podgorica, Yugoslavia