

## **Brunnian planar braids and simplicial groups**

**Valeriy G. Bardakov, Pravin Kumar  
and Mahender Singh**

**ABSTRACT.** Twin groups are planar analogues of Artin braid groups and play a crucial role in the Alexander-Markov correspondence for the isotopy classes of immersed circles on the 2-sphere without triple and higher intersections. These groups admit diagrammatic representations, leading to maps obtained by the addition and deletion of strands. This paper explores Brunnian twin groups, which are subgroups of twin groups composed of twins that become trivial when any of their strands are deleted. We establish that Brunnian twin groups consisting of more than two strands are free groups. Furthermore, we provide a necessary and sufficient condition for a Brunnian doodle on the 2-sphere to be the closure of a Brunnian twin. Additionally, we delve into two generalizations of Brunnian twins, namely,  $k$ -decomposable twins and Cohen twins, and prove some structural results about these groups. We also investigate a simplicial structure on pure twin groups that admits a simplicial homomorphism from Milnor's construction of the simplicial 2-sphere. This gives a possibility to provide a combinatorial description of homotopy groups of the 2-sphere in terms of pure twins.

### CONTENTS

1. Introduction	1236
2. Bi- $\Delta$ -set structure on twin and pure twin groups	1237
3. Brunnian twins	1242
4. $k$ -decomposable twins and Cohen twins	1246
5. Brunnian doodles on the 2-sphere	1252
6. Simplicial structure on pure twin groups	1255
7. Declaration	1261
References	1261

---

Received February 12, 2024.

2020 *Mathematics Subject Classification.* Primary 20F55, 20F36; Secondary 18N50.

*Key words and phrases.* Brunnian twin, Cohen twin, doodle, homotopy group, Milnor construction, pure twin group, simplicial group, twin group.

## 1. Introduction

The twin group, or the planar braid group  $T_n$  on  $n \geq 2$  strands, is a right angled Coxeter group generated by  $n - 1$  involutions that admit only far commutativity relations. These groups appeared in the work of Khovanov [19] on real  $K(\pi, 1)$  subspace arrangements and were further investigated in [17]. Twin groups have a geometrical interpretation similar to the one for Artin braid groups [17, 19]. We fix parallel lines  $y = 0$  and  $y = 1$  on the plane  $\mathbb{R}^2$  with  $n$  marked points on each line. Consider the set of configurations of  $n$  strands in the strip  $\mathbb{R} \times [0, 1]$  connecting the  $n$  marked points on the line  $y = 1$  to those on the line  $y = 0$  such that each strand is monotonic and no three strands have a point in common. Two such configurations are equivalent if one can be deformed into the other by a homotopy of strands, keeping the end points fixed throughout the homotopy. Such an equivalence class is called a *twin*. Placing one twin on top of another and rescaling the interval turns the set of all twins on  $n$  strands into a group isomorphic to  $T_n$ . The generators  $t_i$  of  $T_n$  can be geometrically represented by configurations as shown in Figure 1.




FIGURE 1. The generator  $t_i$  of  $T_n$ .

Analogous to classical knot theory, it is evident that the closure of a twin gives a *doodle* on the 2-sphere. In general, a doodle on a closed surface is a collection of finitely many piecewise-linear closed curves without triple intersections. These objects first appeared in the work of Fenn and Taylor [13]. In [17], Khovanov proved that every oriented doodle on the 2-sphere is the closure of a twin. A Markov theorem for doodles on the 2-sphere has been established by Gotin [16], although the idea has been implicit in [18]. These constructions have been generalised by Bartholomew-Fenn-Kamada-Kamada [5], where they consider a collection of immersed circles in closed oriented surfaces of arbitrary genus.

The pure twin group, denoted as  $PT_n$ , is defined as the kernel of the natural homomorphism from the twin group  $T_n$  to the symmetric group  $S_n$ , which maps the twin  $t_i$  to the transposition  $(i, i + 1)$ . A nice topological interpretation of  $PT_n$  is known due to Khovanov. Consider the space

$$X_n = \mathbb{R}^n \setminus \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_i = x_j = x_k \text{ for } i \neq j \neq k \neq i\},$$

which is the complement of triple diagonals  $x_i = x_j = x_k$ . In [19], Khovanov proved that the fundamental group  $\pi_1(X_n)$  is isomorphic to  $PT_n$ . Prior to this,

Björner and Welker [9] had investigated the cohomology of these spaces, establishing that each  $H^i(X_n, \mathbb{Z})$  is free.

Simplicial structures on braid groups are connected with homotopy groups of some manifolds [6, 8, 20, 23, 27]. Notably, they provide a description of elements in homotopy groups of the 2-sphere in terms of Brunnian braids [6], with a generalization to higher dimensional spheres [27]. A set of generators for the Brunnian braid group of a surface other than the 2-sphere and the projective plane has been provided in [2]. Furthermore, Brunnian subgroups of mapping class groups have been considered in [7]. In this paper, we explore simplicial structures on pure twin groups. The geometrical interpretation of elements in twin groups allows us to define face and degeneracy maps obtained by the deletion and addition of strands, thereby transforming the family of pure twin groups into a simplicial group. We adopt the approach introduced by Cohen and Wu in [11] for Artin pure braid groups.

The paper is organised as follows. In Section 2, we prove that the natural maps of deletion and addition of strands turn the sequence  $\{T_n\}_{n \geq 1}$  into a bi- $\Delta$ -set, whereas the sequence  $\{PT_n\}_{n \geq 1}$  is turned into a bi- $\Delta$ -group (Proposition 2.4). In Section 3, we investigate Brunnian twins, which are twins that become trivial when any one of their strands is removed. We prove that the group  $\text{Brun}(T_n)$  of Brunnian twins on  $n$  strands is free for  $n \geq 3$  (Proposition 3.9), and give an infinite free generating set for  $\text{Brun}(T_4)$  (Theorem 3.5). In Section 4, we consider two generalisations of Brunnian twins, namely,  $k$ -decomposable twins and Cohen twins. A twin is  $k$ -decomposable if it becomes trivial after removing any  $k$  of its strands. We give a complete description of  $k$ -decomposable twins on  $n \geq 4$  strands (Proposition 4.4). A twin on  $n$  strands is said to be Cohen if the twins obtained by removing any one of its strands are all the same. We give a characterisation for a twin to be Cohen (Theorem 4.10). In Section 5, we consider Brunnian doodles on the 2-sphere, and prove that an  $m$ -component Brunnian doodle on the 2-sphere is the closure of a Brunnian twin if and only if its twin index is  $m$  (Theorem 5.4). In Section 6, we observe that pure twin groups admit the structure of a simplicial group  $SPT_*$ . We relate it with the well-known Milnor's construction for simplicial spheres by establishing a homomorphism  $\Theta : F[S^2]_* \rightarrow SPT_*$  of simplicial groups. We also identify some low degree terms of the image of  $\Theta$  as free groups (Theorem 6.6). A complete description of the image of  $\Theta$  gives a possibility to provide a combinatorial description of homotopy groups of the 2-sphere in terms of pure twins.

## 2. Bi- $\Delta$ -set structure on twin and pure twin groups

For  $n \geq 2$ , the *twin group*  $T_n$  on  $n$  strands is generated by  $\{t_1, \dots, t_{n-1}\}$  and it is defined by the following relations:

$$t_i^2 = 1 \quad \text{for } 1 \leq i \leq n-1$$

and

$$t_i t_j = t_j t_i \quad \text{for } |i - j| \geq 2.$$

Clearly, each  $T_n$  is a right angled Coxeter group. Further, there is a surjective homomorphism  $\nu : T_n \rightarrow S_n$ , that sends the generator  $t_i$  to the transposition  $\tau_i = (i, i + 1)$  in the symmetric group  $S_n$ . It's kernel, denoted as  $PT_n$ , is called the pure twin group. It is not difficult to see that  $PT_2$  is trivial and  $PT_3$  is the infinite cyclic group generated by the pure twin  $(t_1 t_2)^3$  [1]. Figure 2 represents the pure twin  $(t_1 t_2)^3$ .




FIGURE 2. The pure twin  $(t_1 t_2)^3$ .

Let us consider the following definitions [33, p.16].

**Definition 2.1.** A sequence of sets  $\{G_n\}_{n \geq 0}$  is called a  $\Delta$ -set if there are maps  $d_i : G_n \rightarrow G_{n-1}$  for each  $0 \leq i \leq n$  such that

$$d_j d_i = d_i d_{j+1} \quad (2.1)$$

for all  $j \geq i$ . The maps  $d_i$  are called face maps. If each  $G_n$  is a group and each face map is a group homomorphism, then  $\{G_n\}_{n \geq 0}$  is called a  $\Delta$ -group.

**Definition 2.2.** A sequence of sets  $\{G_n\}_{n \geq 0}$  is called a bi- $\Delta$ -set if there are face maps  $d_i : G_n \rightarrow G_{n-1}$  and coface maps  $d^i : G_{n-1} \rightarrow G_n$  for each  $0 \leq i \leq n$  such that the following identities hold:

- (1)  $d_j d_i = d_i d_{j+1}$  for  $j \geq i$ ,
- (2)  $d^j d^i = d^{i+1} d^j$  for  $j \leq i$ ,
- (3)  $d_j d^i = d^{i-1} d_j$  for  $j < i$ ,
- (4)  $d_j d^i = \text{id}$  for  $j = i$ ,
- (5)  $d_j d^i = d^i d_{j-1}$  for  $j > i$ .

Moreover, if each  $G_n$  is a group and each face and coface map is a group homomorphism, then  $\{G_n\}_{n \geq 0}$  is called a bi- $\Delta$ -group.

We define a bi- $\Delta$ -set structure on twin groups that would induce a bi- $\Delta$ -group structure on pure twin groups. For geometrical reasons, we take  $G_n = T_{n+1}$  or

$PT_{n+1}$  for each  $n \geq 0$ . For each  $0 \leq i \leq n$ , define the map

$$d_i : T_{n+1} \rightarrow T_n$$

that deletes the  $(i + 1)$ -th strand from the diagram of a twin on  $n + 1$  strands. Note that,  $d_i$  is not a group homomorphism, but it satisfies

$$d_i(uw) = d_i(u)d_{\nu(u)(i+1)-1}(w) \tag{2.2}$$

for all  $u, w \in T_{n+1}$ , where  $\nu : T_{n+1} \rightarrow S_{n+1}$  is the natural surjection. On the other hand, we have  $d_i(PT_{n+1}) \subseteq PT_n$  for each  $0 \leq i \leq n$ . Further, it follows from (2.2) that  $d_i : PT_{n+1} \rightarrow PT_n$  is a surjective group homomorphism for each  $0 \leq i \leq n$ .

**Remark 2.3.** The homomorphism  $d_i : PT_{n+1} \rightarrow PT_n$  has an alternative interpretation. Consider the space

$$X_n = \mathbb{R}^n \setminus \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_i = x_j = x_k, \quad i \neq j \neq k \neq i\},$$

which is the complement of triple diagonals  $x_i = x_j = x_k$  in  $\mathbb{R}^n$ . For each  $1 \leq i \leq n + 1$ , let

$$(p_i)_\# : \pi_1(X_{n+1}) \rightarrow \pi_1(X_n)$$

be the group homomorphism induced by the coordinate projection  $p_i : X_{n+1} \rightarrow X_n$ , where

$$p_i(x_1, \dots, x_{n+1}) = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_{n+1}).$$

By [19, Proposition 3.1], we identify  $\pi_1(X_n)$  with  $PT_n$ , and observe that  $(p_i)_\# = d_{i-1}$  for each  $1 \leq i \leq n + 1$ .




FIGURE 3. The geometrical interpretation of a coface map.

In analogy with [33, Example 1.2.8], for each  $0 \leq i \leq n$ , we define  $d^i : T_n \rightarrow T_{n+1}$  on generators by

$$d^i(t_j) = \begin{cases} t_j & \text{for } j < i, \\ t_{i+1}t_it_{i+1} & \text{for } j = i, \\ t_{j+1} & \text{for } j > i. \end{cases} \tag{2.3}$$

A direct calculation shows that each  $d^i$  satisfy the defining relations in  $T_n$ , and hence is a group homomorphism. Geometrically, the coface maps  $d^0$  and  $d^n$  simply insert a trivial strand on the left and on the right of the diagram of a twin, respectively. Further, for  $1 \leq i \leq n-1$ , the map  $d^i$  inserts a trivial strand between the  $i$ -th and the  $(i+1)$ -th strands of the diagram of the twin generator  $t_i$  such this new strand passes from the right of the crossing corresponding to  $t_i$ . For other twin generators  $t_j$  with  $j \neq i$ , the map  $d^i$  simply inserts a trivial strand between the  $i$ -th and the  $(i+1)$ -th strands of  $t_j$ . See Figure 3 for an illustration.

**Proposition 2.4.** *Consider the sequence of groups  $\{T_n\}_{n \geq 1}$ . For each  $0 \leq i \leq n$ , let  $d_i : T_{n+1} \rightarrow T_n$  be the map satisfying (2.2) and  $d^i : T_n \rightarrow T_{n+1}$  be the homomorphism defined by (2.3). Then  $\{T_n, d_i, d^i\}_{n \geq 1}$  is a bi- $\Delta$ -set and  $\{PT_n, d_i, d^i\}_{n \geq 1}$  is a bi- $\Delta$ -group.*

**Proof.** For each  $0 \leq i \leq n$ , the map  $d_i : T_{n+1} \rightarrow T_n$  clearly satisfies (2.1). Let  $d^i : T_n \rightarrow T_{n+1}$  be the homomorphism defined by (2.3). A direct computation yields

$$d^j d^i(t_k) = d^{i+1} d^j(t_k) = \begin{cases} t_k & \text{for } k < j \leq i, \\ t_{k+2} t_{k+1} t_k t_{k+1} t_{k+2} & \text{for } j = k = i, \\ t_{k+1} t_k t_{k+1} & \text{for } j = k < i, \\ t_{k+2} t_{k+1} t_{k+2} & \text{for } j < k = i, \\ t_{k+1} & \text{for } j < k < i, \\ t_{k+2} & \text{for } j \leq i < k, \end{cases}$$

for all  $j \leq i$ . This proves the identity (2). The identities (3)-(5) follows from the geometrical interpretation of  $d_i$  and  $d^i$ . Hence,  $\{T_n, d_i, d^i\}_{n \geq 1}$  is a bi- $\Delta$ -set.

We already noticed that, for each  $0 \leq i \leq n$ ,  $d_i(PT_{n+1}) \subseteq PT_n$  and  $d_i : PT_{n+1} \rightarrow PT_n$  is a group homomorphism. The inclusion  $d^i(PT_n) \subseteq PT_{n+1}$  follows from the geometrical interpretation of the map  $d^i$ . Alternatively, for each  $n \geq 1$ , let  $\eta^i : S_n \rightarrow S_{n+1}$  be the map defined by

$$\eta^i(\tau_j) = \begin{cases} \tau_j & \text{for } j < i, \\ \tau_{i+1} \tau_i \tau_{i+1} & \text{for } j = i, \\ \tau_{j+1} & \text{for } j > i. \end{cases}$$

As with the case of  $d^i$ , each  $\eta^i$  satisfies the far commutativity and involutory relations of generators of  $S_n$ . For braid relations, we see that

$$\begin{aligned} \eta^i(\tau_k) \eta^i(\tau_{k+1}) \eta^i(\tau_k) &= \begin{cases} \tau_k \tau_{k+1} \tau_k & \text{for } k+1 < i, \\ \tau_k \tau_{k+1} \tau_{k+2} \tau_{k+1} \tau_k & \text{for } i = k, k+1, \\ \tau_{k+1} \tau_{k+2} \tau_{k+1} & \text{for } k > i, \end{cases} \\ &= \eta^i(\tau_{k+1}) \eta^i(\tau_k) \eta^i(\tau_{k+1}), \end{aligned}$$

and hence each  $\eta^i$  is a group homomorphism. Then the inclusion  $d^i(PT_n) \subseteq PT_{n+1}$  also follows from the commutativity of the following diagram

$$\begin{array}{ccc} T_n & \xrightarrow{d^i} & T_{n+1} \\ \downarrow \nu & & \downarrow \nu \\ S_n & \xrightarrow{\eta^i} & S_{n+1}. \end{array}$$

Finally, we prove that each  $d^i$  is group homomorphism at the level of twin groups itself. Clearly,  $(d^i(t_k))^2 = 1$  for all  $i$  and  $k$ . Further, for  $k < \ell$  with  $|k - \ell| \geq 2$ , we have

$$\begin{aligned} d^i(t_k)d^i(t_\ell) &= \begin{cases} t_k t_\ell & \text{for } \ell < i, \\ t_k t_{\ell+1} t_\ell t_{\ell+1} & \text{for } \ell = i, \\ t_k t_{\ell+1} & \text{for } k < i < \ell, \\ t_{k+1} t_k t_{k+1} t_{\ell+1} & \text{for } k = i, \\ t_{k+1} t_{\ell+1} & \text{for } i < k, \end{cases} \\ &= d^i(t_\ell)d^i(t_k). \end{aligned}$$

This proves that  $\{PT_n, d_i, d^i\}_{n \geq 1}$  is a bi- $\Delta$ -group.  $\square$

**Remark 2.5.** For each  $0 \leq i \leq n$ , we can also define the coface maps  $d^i : T_n \rightarrow T_{n+1}$  by

$$d^i(t_j) = \begin{cases} t_j & \text{for } j < i, \\ t_i t_{i+1} t_i & \text{for } j = i, \\ t_{j+1} & \text{for } j > i. \end{cases}$$

It can be verified that the analogue of Proposition 2.4 holds with these coface maps.

We now use the bi- $\Delta$ -set structure on  $\{T_n\}_{n \geq 1}$  to give a new presentation for  $T_{n+1}$ . We use the coface maps  $d^i$  as defined in Proposition 2.4.

**Proposition 2.6.** *Let  $q_k := d^n d^{n-1} \cdots d^k(t_k)$  for  $1 \leq k \leq n-1$  and  $q_n := d^{n-2}(t_{n-1})$ . Then  $T_{n+1}$  admits a presentation with generating set  $\{q_1, \dots, q_n\}$  and the following defining relations:*

- (1)  $q_i^2 = 1$  for all  $i$ ,
- (2)  $[q_{i+1} q_i q_{i+1}, q_n] = 1$  for  $i < n-1$ ,
- (3)  $[q_{i+1} q_i q_{i+1}, q_{j+1} q_j q_{j+1}]$  for  $|i-j| > 2$  and  $i, j \leq n-1$ .

**Proof.** Using the simplicial identity  $d^j d^i = d^{i+1} d^j$  for  $j \leq i$ , we can assume that  $i_{n-1} > i_{n-2} > \cdots > i_1$  in the composite map  $d^{i_{n-1}} d^{i_{n-2}} \cdots d^{i_1}$ . We see that

$$\begin{aligned} q_k &= d^{n-1} d^{n-2} \cdots d^k(t_k) \\ &= t_n t_{n-1} \cdots t_{k+1} t_k t_{k+1} \cdots t_{n-1} t_n \end{aligned}$$

for  $1 \leq k \leq n-1$  and

$$q_n = d^{n-2}(t_{n-1}) = t_n.$$

A direct check gives  $t_k = q_{k+1}q_kq_{k+1}$  for each  $1 \leq k \leq n-1$ , and hence  $\{q_1, \dots, q_n\}$  generates  $T_{n+1}$ . Further, the defining relations of  $T_{n+1}$  in terms of the Coxeter generating set  $\{t_1, \dots, t_n\}$  gives the defining relations for the new generating set as follows:

- (1)  $q_i^2 = 1$  for all  $i$ ,
- (2)  $[q_{i+1}q_iq_{i+1}, q_n] = 1$  for  $i < n-1$ ,
- (3)  $[q_{i+1}q_iq_{i+1}, q_{j+1}q_jq_{j+1}]$  for  $|i-j| > 2$  and  $i, j \leq n-1$ .

□

### 3. Brunnian twins

In the influential work [6], a connection has been established between certain quotients of the Brunnian braid groups of the 2-sphere and its higher homotopy groups.

**Definition 3.1.** *A pure twin is said to be Brunnian if it becomes trivial after removing any one of its strands.*

Let  $\text{Brun}(T_n)$  denote the set of all Brunnian twins on  $n$  strands.

**Proposition 3.2.**  *$\text{Brun}(T_n)$  is a normal subgroup of  $PT_n$ .*

**Proof.** For each  $0 \leq i \leq n-1$ , let  $d_i : PT_n \rightarrow PT_{n-1}$  be the face map of Proposition 2.4. Since each  $d_i$  is a group homomorphism and

$$\text{Brun}(T_n) = \bigcap_{i=0}^{n-1} \ker(d_i),$$

it follows that  $\text{Brun}(T_n)$  is a normal subgroup of  $T_n$ . □

Next, we attempt to understand the groups of Brunnian twins.

**Proposition 3.3.** *For  $n \geq 4$ ,  $\text{Brun}(T_n)$  does not contain any element from  $\{(t_i t_{i+1})^3 \mid 1 \leq i \leq n-1\}$ .*

**Proof.** For  $n \geq 4$ , removing a trivial strand from  $(t_i t_{i+1})^3$  gives a non-trivial twin, and hence the assertion follows. □

**Proposition 3.4.**  *$\text{Brun}(T_3) \cong PT_3 \cong \mathbb{Z}$ .*

**Proof.** We already have  $\text{Brun}(T_3) \subseteq PT_3$ . By [1, Theorem 2], we know that  $PT_3$  is the infinite cyclic group generated by  $(t_1 t_2)^3$  (see Figure 2), and clearly  $(t_1 t_2)^3 \in \text{Brun}(T_3)$ . □

In contrast, it is proved in [22] that  $\text{Brun}(B_3)$  is the commutator subgroup of the Artin pure braid group  $P_3$ .

**Theorem 3.5.**  *$\text{Brun}(T_4)$  is a free group of infinite rank.*



**Proof.** By [1, Theorem 2],  $PT_4$  is a free group of rank 7 generated by

$$\begin{aligned} x_1 &= (t_1 t_2)^3, & x_2 &= ((t_1 t_2)^3)^{t_3}, & x_3 &= ((t_1 t_2)^3)^{t_3 t_2}, & x_4 &= ((t_1 t_2)^3)^{t_3 t_2 t_1}, \\ x_5 &= (t_2 t_3)^3, & x_6 &= ((t_2 t_3)^3)^{t_1}, & x_7 &= ((t_2 t_3)^3)^{t_1 t_2}. \end{aligned}$$

Denote the generator  $(t_1 t_2)^3$  of  $PT_3$  by  $y$ . Direct computations show that images of  $x_i$ 's under the face maps  $d_i$ 's are as follows:

$$\begin{aligned} d_0(x_1) &= d_0(x_2) = d_0(x_3) = d_0(x_6) = d_0(x_7) = 1, & d_0(x_4) &= d_0(x_5) = y, \\ d_1(x_1) &= d_1(x_2) = d_1(x_4) = d_1(x_5) = d_1(x_7) = 1, & d_1(x_3) &= d_1(x_6) = y, \\ d_2(x_1) &= d_2(x_3) = d_2(x_4) = d_2(x_5) = d_2(x_6) = 1, & d_2(x_2) &= d_2(x_7) = y, \\ d_3(x_2) &= d_3(x_3) = d_3(x_4) = d_3(x_5) = d_3(x_6) = d_3(x_7) = 1, & d_3(x_1) &= y. \end{aligned}$$

For each generator  $x_i$ , let  $\log_i(w)$  denote the sum of the powers of  $x_i$  in the word  $w$ . Then, it follows that

$$\begin{aligned} \ker(d_0) &= \{w \in PT_4 \mid \log_4(w) + \log_5(w) = 0\}, \\ \ker(d_1) &= \{w \in PT_4 \mid \log_3(w) + \log_6(w) = 0\}, \\ \ker(d_2) &= \{w \in PT_4 \mid \log_2(w) + \log_7(w) = 0\}, \\ \ker(d_3) &= \{w \in PT_4 \mid \log_1(w) = 0\}, \end{aligned}$$

and hence

$$\begin{aligned} &\text{Brun}(T_4) \\ &= \bigcap_{i=0}^3 \ker(d_i) \\ &= \left\{ w \in PT_4 \mid \log_4(w) + \log_5(w) = 0, \log_3(w) + \log_6(w) = 0, \right. \\ &\quad \left. \log_2(w) + \log_7(w) = 0, \log_1(w) = 0 \right\}. \end{aligned}$$

Clearly,  $\text{Brun}(T_4)$  is free being a subgroup of the free group  $PT_4$ . We now find an infinite free basis for  $\text{Brun}(T_4)$ . It follows from the preceding description of  $\text{Brun}(T_4)$  that the commutator subgroup of  $PT_4$  is contained in  $\text{Brun}(T_4)$ . In fact, the containment is strict since  $x_4 x_5^{-1} \in \text{Brun}(T_4)$ , but  $x_4 x_5^{-1} \notin PT'_n$ . Thus,  $PT_4/\text{Brun}(T_4)$  is a non-trivial abelian group. Let  $q : PT_4 \rightarrow PT_4/\text{Brun}(T_4)$  be the quotient map with  $q(x_i) = y_i$  for  $1 \leq i \leq 7$ . Since  $x_2 x_7^{-1}, x_3 x_6^{-1}, x_4 x_5^{-1} \in \text{Brun}(T_4)$ , the group  $PT_4/\text{Brun}(T_4)$  is generated by the set  $\{y_1, y_2, y_3, y_4\}$ . Note that  $x_i x_j^{-1} \notin \text{Brun}(T_4)$  for all  $i \neq j \in \{1, 2, 3, 4\}$  and  $x_i^k \notin \text{Brun}(T_4)$  for  $k > 0$ . Thus, by the fundamental theorem for finitely generated abelian groups,  $PT_4/\text{Brun}(T_4)$  is a free abelian group of rank 4.

Consider the short exact sequence

$$1 \rightarrow \text{Brun}(T_4) \rightarrow PT_4 \rightarrow \mathbb{Z}^4 \rightarrow 1.$$

We fix a Schreier system  $\{x_1^{k_1} x_2^{k_2} x_3^{k_3} x_4^{k_4} \mid k_1, k_2, k_3, k_4 \in \mathbb{Z}\}$  of coset representatives of  $\text{Brun}(T_4)$  in  $PT_4$ . This gives a free basis for  $\text{Brun}(T_4)$  consisting of

elements of the form

$$\begin{aligned} &x_1^{k_1} x_2^{k_2} x_3^{k_3} x_4^{k_4} x_1(x_1^{k_1+1} x_2^{k_2} x_3^{k_3} x_4^{k_4})^{-1}, && x_1^{k_1} x_2^{k_2} x_3^{k_3} x_4^{k_4} x_2(x_1^{k_1} x_2^{k_2+1} x_3^{k_3} x_4^{k_4})^{-1}, \\ &x_1^{k_1} x_2^{k_2} x_3^{k_3} x_4^{k_4} x_3(x_1^{k_1} x_2^{k_2} x_3^{k_3+1} x_4^{k_4})^{-1}, && x_1^{k_1} x_2^{k_2} x_3^{k_3} x_4^{k_4} x_4(x_1^{k_1} x_2^{k_2} x_3^{k_3} x_4^{k_4+1})^{-1}, \\ &x_1^{k_1} x_2^{k_2} x_3^{k_3} x_4^{k_4} x_5(x_1^{k_1} x_2^{k_2} x_3^{k_3} x_4^{k_4+1})^{-1}, && x_1^{k_1} x_2^{k_2} x_3^{k_3} x_4^{k_4} x_6(x_1^{k_1} x_2^{k_2} x_3^{k_3+1} x_4^{k_4})^{-1}, \\ &x_1^{k_1} x_2^{k_2} x_3^{k_3} x_4^{k_4} x_7(x_1^{k_1} x_2^{k_2+1} x_3^{k_3} x_4^{k_4})^{-1}, \end{aligned}$$

for  $k_1, k_2, k_3, k_4 \in \mathbb{Z}$ . This completes the proof. □

**Remark 3.6.**  $PT_n/\text{Brun}(T_n)$  is non-abelian for  $n \geq 5$  since  $PT'_n \not\subseteq \text{Brun}(T_n)$  for  $n \geq 5$ . For example, if  $w = [(t_1 t_2)^3, (t_2 t_3)^3]$ , then  $d_4(w) \neq 1$ .

**Proposition 3.7.**  $PT_n/\text{Brun}(T_n)$  is torsion free for each  $n \geq 4$  and

$$PT_4/\text{Brun}(T_4) \cong \mathbb{Z}^4.$$

**Proof.** The homomorphisms  $d_i : PT_n \rightarrow PT_{n-1}$  induce an injective homomorphism

$$PT_n/\text{Brun}(T_n) \hookrightarrow \underbrace{PT_{n-1} \times \cdots \times PT_{n-1}}_{n \text{ times}}.$$

By [1, Theorem 3],  $PT_n$  is torsion-free for  $n \geq 3$ . Hence,  $PT_{n-1} \times \cdots \times PT_{n-1}$  is torsion free, and therefore  $PT_n/\text{Brun}(T_n)$  is so. The second assertion is proven in the proof of Theorem 3.5. □

**Problem 3.8.** Describe the structure of the group  $PT_n/\text{Brun}(T_n)$  for  $n \geq 5$ .

Recall from [1, 30] that the virtual twin group  $VT_n$  on  $n \geq 2$  strands is the group generated by  $\{t_1, \dots, t_{n-1}, \rho_1, \dots, \rho_{n-1}\}$  and having the following defining relations:

$$\begin{aligned} t_i^2 &= 1 \quad \text{for } 1 \leq i \leq n-1, \\ t_i t_j &= t_j t_i \quad \text{for } |i-j| \geq 2, \\ \rho_i^2 &= 1 \quad \text{for } 1 \leq i \leq n-1, \\ \rho_i \rho_j &= \rho_j \rho_i \quad \text{for } |i-j| \geq 2, \\ \rho_i \rho_{i+1} \rho_i &= \rho_{i+1} \rho_i \rho_{i+1} \quad \text{for } 1 \leq i \leq n-2, \\ \rho_i t_j &= t_j \rho_i \quad \text{for } |i-j| \geq 2, \\ \rho_i \rho_{i+1} t_i &= t_{i+1} \rho_i \rho_{i+1} \quad \text{for } 1 \leq i \leq n-2. \end{aligned}$$

The group  $VT_n$  plays the role of virtual braid groups in the Alexander-Markov correspondence for the planar analogue of virtual knot theory. There is a surjective homomorphism  $\mu : VT_n \rightarrow S_n$  given by

$$\mu(t_i) = \mu(\rho_i) = (i, i+1)$$

for all  $1 \leq i \leq n-1$ . The kernel  $PVT_n$  of this surjection is called the *pure virtual twin group* on  $n$  strands.

For each  $n \geq 2$ , we have surjective homomorphisms  $d_{n-1} : PT_n \rightarrow PT_{n-1}$  and  $\bar{d}_{n-1} : PVT_n \rightarrow PVT_{n-1}$  that delete the  $n$ -th strand from the diagram of a pure twin and pure virtual twin. In the reverse directions, we have homomorphisms  $d^{n-1} : PT_{n-1} \rightarrow PT_n$  and  $\bar{d}^{n-1} : PVT_{n-1} \rightarrow PVT_n$  that add a trivial strand to the right side of the diagram. Further, we have  $d_{n-1} d^{n-1} = \text{id}_{PT_{n-1}}$  and  $\bar{d}_{n-1} \bar{d}^{n-1} = \text{id}_{PVT_{n-1}}$ . Setting  $U_n = \ker(d_{n-1})$  and  $V_n = \ker(\bar{d}_{n-1})$ , we have split short exact sequences

$$1 \rightarrow U_n \rightarrow PT_n \rightarrow PT_{n-1} \rightarrow 1$$

and

$$1 \rightarrow V_n \rightarrow PVT_n \rightarrow PVT_{n-1} \rightarrow 1.$$

In other words,  $PT_n \cong U_n \rtimes PT_{n-1}$  and  $PVT_n \cong V_n \rtimes PVT_{n-1}$ .

**Proposition 3.9.** *Brun( $T_n$ ) is free for all  $n \geq 3$ .*

**Proof.** The map  $t_i \mapsto t_i$  gives an embedding of  $T_n$  into  $PVT_n$  [31, Corollary 3.5]. Restricting to  $PT_n$ , this gives an inclusion  $\psi_n : PT_n \rightarrow PVT_n$  such that the following diagram commutes

$$\begin{array}{ccc} PT_n & \xrightarrow{d_{n-1}} & PT_{n-1} \\ \downarrow \psi_n & & \downarrow \psi_{n-1} \\ PVT_n & \xrightarrow{\bar{d}_{n-1}} & PVT_{n-1}. \end{array}$$

This gives  $U_n \cong \psi_n(U_n) = \psi_n(\ker(d_{n-1})) \leq \ker(\bar{d}_{n-1}) = V_n$ . Since  $V_n$  is free for  $n \geq 2$  [30, Theorem 4.1], it follows that  $U_n$  is also free. Note that the subgroup  $U_i = \ker(d_i)$  is conjugate to  $U_n$  by the element  $t_{n-1}t_{n-2} \cdots t_{i+1}$ . Thus,  $U_i$  is free group for each  $1 \leq i \leq n$ , and hence  $\text{Brun}(T_n) = \bigcap_{i=1}^n U_i$  is a free group. □

At this juncture, the ensuing problem naturally arises.

**Problem 3.10.** *Determine a free generating set for  $\text{Brun}(T_n)$  for  $n \geq 5$ .*

We conclude the section with a consequence of the Decomposition Theorem for bi- $\Delta$ -groups in our setting [33, Proposition 1.2.9].

**Proposition 3.11.** *The pure twin group  $PT_{n+1}$  is the iterated semi-direct product of subgroups*

$$\{d^k d^{i_{k-1}} \cdots d^{i_1}(\text{Brun}(T_{n-k+1})) \mid 0 \leq i_1 < i_2 < \cdots < i_k \leq n \text{ and } 0 \leq k \leq n\}$$

*with the lexicographic order on the indexing set*

$$\{(i_k, i_{k-1}, \dots, i_1, \underbrace{i_0, i_0, \dots, i_0}_{n-k \text{ times}}) \mid 0 \leq i_1 < i_2 < \cdots < i_k \leq n \text{ and } 0 \leq k \leq n\}$$

*from the left, where  $i_0$  is the blank symbol considered smaller than all other indices.*

**Example 3.12.** For  $n = 3$ , we have

$$\begin{aligned} d^0(PT_3) &= \langle (t_2 t_3)^3 \rangle, & d^1(PT_3) &= \langle (t_2 t_1 t_2 t_3)^3 \rangle, \\ d^2(PT_3) &= \langle (t_1 t_3 t_2 t_3)^3 \rangle, & d^3(PT_3) &= \langle (t_1 t_2)^3 \rangle. \end{aligned}$$




FIGURE 4. Images of  $(t_1 t_2)^3$  under the coface maps  $d^0, d^1, d^2$  and  $d^3$ .

See Figure 4. Observing the proof of [33, Proposition 1.2.9], we get

$$PT_4 = \ker(d_3) \rtimes \langle (t_1 t_2)^3 \rangle,$$

where  $\ker(d_3)$  is normal in  $PT_4$  and  $\langle (t_1 t_2)^3 \rangle$  acts on  $\ker(d_3)$  via conjugation. At the second stage, we obtain

$$\ker(d_3) = (\ker(d_2) \cap \ker(d_3)) \rtimes \langle (t_1 t_3 t_2 t_3)^3 \rangle,$$

where  $\ker(d_2) \cap \ker(d_3)$  is normal in  $\ker(d_3)$  and the subgroup

$$\langle (t_1 t_3 t_2 t_3)^3 \rangle = \langle d^2((t_1 t_2)^3) \rangle \leq \ker(d_3)$$

acts on  $\ker(d_2) \cap \ker(d_3)$  via conjugation. At the third stage, we get

$$\ker(d_2) \cap \ker(d_3) = (\ker(d_1) \cap \ker(d_2) \cap \ker(d_3)) \rtimes \langle (t_2 t_1 t_2 t_3)^3 \rangle,$$

where  $\ker(d_1) \cap \ker(d_2) \cap \ker(d_3)$  is normal in  $\ker(d_2) \cap \ker(d_3)$  and the subgroup  $\langle (t_2 t_1 t_2 t_3)^3 \rangle = \langle d^1((t_1 t_2)^3) \rangle \leq \ker(d_2) \cap \ker(d_3)$  acts on  $\ker(d_1) \cap \ker(d_2) \cap \ker(d_3)$  via conjugation. Finally, we have

$$\ker(d_1) \cap \ker(d_2) \cap \ker(d_3) = \text{Brun}(T_4) \rtimes \langle (t_2 t_3)^3 \rangle,$$

where  $\text{Brun}(T_4)$  is normal and the subgroup  $\langle (t_2 t_3)^3 \rangle = \langle d^0((t_1 t_2)^3) \rangle$  acts on  $\text{Brun}(T_4)$  via conjugation. Thus, we obtain the following decomposition of  $PT_4$  as an iterated semi-direct product

$$PT_4 = (((\text{Brun}(T_4) \rtimes \langle (t_2 t_3)^3 \rangle) \rtimes \langle (t_2 t_1 t_2 t_3)^3 \rangle) \rtimes \langle (t_1 t_3 t_2 t_3)^3 \rangle) \rtimes \langle (t_1 t_2)^3 \rangle.$$

Similarly, there are 16 non-trivial terms in the decomposition of  $PT_5$  with the leftmost term being the Brunnian subgroup  $\text{Brun}(T_5)$ .

#### 4. $k$ -decomposable twins and Cohen twins

In this section, we consider two generalisations of Brunnian twins.




FIGURE 5. Converting a pure twin into a Brunnian twin.

**4.1.  $k$ -decomposable twins.** We begin with the following definition.

**Definition 4.1.** A pure twin on  $n$  strands is said to be  $k$ -decomposable if it becomes trivial after removing any  $k$  of its strands.

Clearly, a 1-decomposable twin is simply a Brunnian twin. Further, the set of all  $k$ -decomposable twins on  $n$  strands forms a normal subgroup of  $PT_n$  and we denote this subgroup by  $D_{k,n}$ . For  $w \in PT_n$  and  $1 \leq i < j < k \leq n$ , let  $w_{i,j,k}$  be the pure twin obtained from  $w$  by deleting all the strands except those indexed  $i, j, k$ . We can still view each  $w_{i,j,k}$  as an element of  $PT_n$  by adding trivial  $(n - 3)$  strands on its right. See Figure 5 for an example for  $n = 4$ . Using ideas from [21], we prove the following result.

**Proposition 4.2.** For  $n \geq 4$ ,

$$D_{n-3,n} = \left\{ w \prod_{1 \leq i < j < k \leq n} (w_{i,j,k}^{-1})^{c_{i,j,k}} \mid w \in PT_n \right\},$$

where  $c_{i,j,k} \in T_n$  is a coset representative of the permutation in  $T_n/PT_n \cong S_n$  which takes  $i, j, k$  to  $1, 2, 3$ , respectively, and fix everything else.

**Proof.** In view of Proposition 3.4, we have  $w_{i,j,k} \in \text{Brun}(T_3)$ . A direct check shows that for any  $w \in PT_n$ , the pure twin

$$w \prod_{1 \leq i < j < k \leq n} (w_{i,j,k}^{-1})^{c_{i,j,k}}$$

is a  $(n - 3)$ -decomposable twin on  $n$  strands. Note that the map  $\phi : PT_n \rightarrow D_{n-3,n}$  given by

$$\phi(w) = w \prod_{1 \leq i < j < k \leq n} (w_{i,j,k}^{-1})^{c_{i,j,k}}$$

is a retraction, that is, the restriction of  $\phi$  on  $D_{n-3,n}$  is the identity map. Hence, it follows that each element of  $D_{n-3,n}$  arises in this fashion.  $\square$

**Corollary 4.3.**  $\text{Brun}(T_4) = \{w w_{1,2,3}^{-1} (w_{1,2,4}^{-1})^{t_3} (w_{1,3,4}^{-1})^{t_2 t_3} (w_{2,3,4}^{-1})^{t_1 t_2 t_3} \mid w \in PT_4\}$ .

Next, we describe a process of constructing  $D_{k-1,n}$  from  $D_{k,n}$ . Let  $w \in D_{k,n}$  and  $1 \leq i_1 < i_2 \cdots < i_{n-k+1} \leq n$ . Let  $w_{i_1, i_2, \dots, i_{n-k+1}}$  be the pure twin obtained from  $w$  by removing the  $k - 1$  strands except those indexed  $i_1, i_2, \dots, i_{n-k+1}$ . Since  $w \in D_{k,n}$ , we have  $w_{i_1, i_2, \dots, i_{n-k+1}} \in \text{Brun}(T_{n-k+1})$ . The following result can be proved along the lines of Proposition 4.2.

**Proposition 4.4.** For  $n \geq 4$ ,

$$D_{k-1,n} = \{w \prod_{1 \leq i_1 < i_2 \cdots < i_{n-k+1} \leq n} (w_{i_1, i_2, \dots, i_{n-k+1}}^{-1})^{c_{i_1, i_2, \dots, i_{n-k+1}}} \mid w \in D_{k,n}\},$$

where  $c_{i_1, i_2, \dots, i_{n-k+1}} \in T_n$  is a coset representative of the permutation in  $T_n/PT_n \cong S_n$  which takes  $i_1, i_2, \dots, i_{n-k+1}$  to  $1, 2, \dots, n - k + 1$ , respectively, and fix everything else.

Beginning with  $PT_n = D_{n-2,n} = D_{n-1,n}$  and iterating the procedure of constructing  $D_{k-1,n}$  from  $D_{k,n}$ , we can construct all Brunnian twins on  $n$  strands.

**4.2. Cohen twins.** Next, we consider another generalisation of Brunnian twins motivated by an idea due to Fred Cohen [10], and developed further for surface braid groups in [4]. Recall that, for  $0 \leq i \leq n - 1$ , the face map  $d_i : T_n \rightarrow T_{n-1}$  deletes the  $(i + 1)$ -st strand from the diagram of a twin. Although  $d_i$  is not a group homomorphism, it satisfies

$$d_i(uw) = d_i(u) d_{\nu(u)(i+1)-1}(w), \tag{4.1}$$

where  $\nu : T_{n+1} \rightarrow S_{n+1}$  is the natural surjection. For an arbitrary  $u \in T_{n-1}$ , we ask whether there exists  $w \in T_n$  which is a solution of the system of equations

$$\begin{cases} d_0(w) = u, \\ d_1(w) = u, \\ \vdots \\ d_{n-1}(w) = u. \end{cases} \tag{4.2}$$

Taking  $u = 1$  amounts to  $w \in T_n$  being a Brunnian twin.




FIGURE 6. Elements  $\delta_3, \delta_4$  and  $\gamma_3, \gamma_4$ .

**Definition 4.5.** A twin  $w \in T_n$  is called a Cohen twin if  $d_0(w) = d_1(w) = \dots = d_{n-1}(w)$ .

For  $n \geq 2$ , let us set

$$CT_n = \{w \in T_n \mid d_0(w) = d_1(w) = \dots = d_{n-1}(w)\}.$$

In other words, a twin on  $n$  strands lie in  $CT_n$  if it gives the same twin on  $(n - 1)$  strands after removing any one of its strands. For example, the twin

$$\delta_n := (t_1 t_2 \dots t_{n-1})(t_1 t_2 \dots t_{n-2}) \dots (t_1 t_2) t_1$$

lies in  $CT_n$  for all  $n \geq 2$  and  $d_0(\delta_n) = \delta_{n-1}$  (see Figure 6). Similarly, we define

$$CPT_n = CT_n \cap PT_n = \{w \in PT_n \mid d_0(w) = d_1(w) = \dots = d_{n-1}(w)\}.$$

We refer to elements of  $CPT_n$  as pure Cohen twins. For instance, the pure twin

$$\gamma_n := (t_1 t_2 \dots t_{n-1})^n$$

lies in  $CPT_n$  for all  $n \geq 2$  and  $d_0(\gamma_n) = \gamma_{n-1}$  (see Figure 6).

If  $\phi, \psi : G \rightarrow H$  are group homomorphisms, then their equalizer is the subgroup of  $G$  given by

$$\{g \in G \mid \phi(g) = \psi(g)\}.$$

Hence,  $CPT_n$  is a subgroup of  $PT_n$  being the equalizer of group homomorphisms  $d_0, d_1, \dots, d_{n-1} : PT_n \rightarrow PT_{n-1}$ .

**Proposition 4.6.** The following assertions hold:

- (1) For each  $0 \leq i \leq n - 1$ ,  $d_i(CPT_n) \subseteq CPT_{n-1}$  and the map  $d_0 = d_1 = \dots = d_{n-1} : CPT_n \rightarrow CPT_{n-1}$  is a group homomorphism.
- (2) The set  $CT_n$  is a subgroup of  $T_n$ . Moreover, for each  $0 \leq i \leq n - 1$ ,  $d_i(CT_n) \subseteq CT_{n-1}$  and the map  $d_0 = d_1 = \dots = d_{n-1} : CT_n \rightarrow CT_{n-1}$  is a group homomorphism.

**Proof.** Let  $w \in CPT_n$  and  $0 \leq i \leq n - 1$ . Then, using (2.1), we obtain

$$d_j(d_i(w)) = d_j(d_0(w)) = d_0(d_{j+1}(w)) = d_0(d_i(w)) \tag{4.3}$$

for each  $0 \leq j \leq n - 2$ , and hence  $d_i(CPT_n) \subseteq CPT_{n-1}$ . That  $d_0 = d_1 = \dots = d_{n-1} : CPT_n \rightarrow CPT_{n-1}$  is a group homomorphism follows from Proposition 2.4.

For the second assertion, let  $u, w \in CT_n$ . By (4.1), we have

$$d_i(uw) = d_i(u)d_{\nu(u)(i+1)-1}(w) = d_0(u)d_{\nu(u)(1)-1}(w) = d_0(uw) \tag{4.4}$$

for each  $0 \leq i \leq n - 1$ , and hence  $uw \in CT_n$ . Further, the equation

$$1 = d_i(u^{-1}u) = d_i(u^{-1})d_{\nu(u^{-1})(i+1)-1}(u) = d_i(u^{-1})d_0(u),$$

gives

$$d_i(u^{-1}) = (d_0(u))^{-1}$$

for each  $0 \leq i \leq n - 1$ , and hence  $CT_n$  is a subgroup of  $T_n$ . The proof of  $d_i(CT_n) \subseteq CT_{n-1}$  follows from 4.3. Finally, (4.4) also shows that  $d_0 = d_1 = \dots = d_{n-1} : CT_n \rightarrow CT_{n-1}$  is a group homomorphism.  $\square$

**Proposition 4.7.**  $CPT_n$  is an index two subgroup of  $CT_n$  for  $n \geq 3$ .

**Proof.** The topological interpretation of elements of  $T_n$  can be applied to elements of  $S_n$  as well by allowing triple intersection points. Thus, for each  $0 \leq i \leq n - 1$ , there is a map  $\bar{d}_i : S_n \rightarrow S_{n-1}$  (thought of as deleting the  $(i + 1)$ -st strand) such the following diagram commutes

$$\begin{array}{ccccc} PT_n & \hookrightarrow & T_n & \xrightarrow{\nu_n} \twoheadrightarrow & S_n \\ \downarrow d_i & & \downarrow d_i & & \downarrow \bar{d}_i \\ PT_{n-1} & \hookrightarrow & T_{n-1} & \xrightarrow{\nu_{n-1}} \twoheadrightarrow & S_{n-1}. \end{array}$$

Set  $CS_n := \nu_n(CT_n)$  for each  $n \geq 2$ . Note that  $CS_2 = \nu_2(T_2) = S_2 \cong \mathbb{Z}_2$ . The commutativity of the preceding diagram shows that every  $\tau \in CS_n$  satisfy  $\bar{d}_0(\tau) = \bar{d}_1(\tau) = \dots = \bar{d}_{n-1}(\tau)$ . By Proposition 4.6(2), we have  $d_0(CT_n) \subseteq CT_{n-1}$ . The commutativity of the preceding diagram implies that  $\bar{d}_0(CS_n) = \bar{d}_0\nu_n(CT_n) = \nu_{n-1}d_0(CT_n) \subseteq \nu_{n-1}(CT_{n-1}) = CS_{n-1}$ . Thus, for  $n \geq 3$ , the restriction of the map  $\bar{d}_0 : S_n \rightarrow S_{n-1}$  induces a map  $\bar{d}_0 : CS_n \rightarrow CS_{n-1}$  such that  $\ker(\bar{d}_0) = \bigcap_{i=0}^{n-1} \ker(\bar{d}_i)$ . Direct computation gives  $\ker(\bar{d}_0) = 1$ , and hence the map  $\bar{d}_0 \dots \bar{d}_0 : CS_n \rightarrow CS_2$  is injective. Since  $\nu_n(\delta_n) \neq 1$ , we have  $CT_n/CPT_n \cong CS_n \cong \mathbb{Z}_2$ , and the proof is complete.  $\square$

The following result follows along the lines of [32, Lemma 2.10].



**Proposition 4.8.** *For each  $1 \leq k \leq n - 1$ , the map*

$$\underbrace{d_0 \cdots d_0}_{(n-k) \text{ times}} : CPT_n \rightarrow CPT_k$$

*is surjective. In particular,  $d_0 : CPT_n \rightarrow CPT_{n-1}$  is surjective for  $n \geq 2$ .*

**Proof.** Let us set  $d_{n-k,n} = \underbrace{d_0 \cdots d_0}_{(n-k) \text{ times}}$ . We use induction on  $k$ . Clearly, for  $k = 1$ ,

the map  $d_{n-1,n} : CPT_n \rightarrow CPT_1$  is surjective. Assume that  $d_{n-k+1,n}$  is surjective with  $k > 1$ , and let  $w \in CPT_k$ .

Case 1: Suppose that  $w \in \ker(d_0 : CPT_k \rightarrow CPT_{k-1})$ . Then consider the element

$$w_{k,n} = \prod_{0 \leq i_1 < i_2 < \cdots < i_{n-k} \leq n-1} d^{i_{n-k}} d^{i_{n-k-1}} \cdots d^{i_1}(w)$$

of  $PT_n$  with lexicographic order on the indices from the right. Since  $w \in \ker(d_0 : CPT_k \rightarrow CPT_{k-1})$ , a straightforward computation shows that  $w_{k,n} \in CPT_n$  and  $d_{n-k,n}(w_{k,n}) = w$ . For instance, taking  $n = 4$  and  $k = 1$ , we have

$$w_{1,4} = \prod_{0 \leq i_1 < i_2 < i_3 \leq 3} d^{i_3} d^{i_2} d^{i_1}(w)$$

with lexicographic order from the right. Note that  $(i_1, i_2, i_3)$  all lie in the set  $\{(0, 1, 2), (0, 1, 3), (0, 2, 3), (1, 2, 3)\}$  and

$$w_{1,4} = d^2 d^1 d^0(w) d^3 d^1 d^0(w) d^3 d^2 d^0(w) d^3 d^2 d^1(w).$$

Direct computations give

$$\begin{aligned} d_0(w_{1,4}) &= d^1 d^0(w) d^2 d^0(w) d^2 d^1(w) d^2 d^1 d^0(d_0(w)), \\ d_1(w_{1,4}) &= d^1 d^0(w) d^2 d^0(w) d^2 d^1 d^0(d_0(w)) d^2 d^1(w), \\ d_2(w_{1,4}) &= d^1 d^0(w) d^2 d^1 d^0(d_0(w)) d^2 d^0(w) d^2 d^1(w), \\ d_3(w_{1,4}) &= d^2 d^1 d^0(d_0(w)) d^1 d^0(w) d^2 d^0(w) d^2 d^1(w). \end{aligned}$$

Since  $w \in \ker(d_0 : CPT_k \rightarrow CPT_{k-1})$ ,  $d^2 d^1 d^0(d_0(w)) = 1$ , and hence  $w_{1,4} \in CPT_4$ .

Case 2: Now, suppose that  $1 \neq \delta = d_0(w) \in CPT_{k-1}$ . By induction hypothesis, there exists  $\gamma \in CPT_n$  such that  $d_{n-k+1,n}(\gamma) = d_0(d_{n-k,n}(\gamma)) = \delta$ . Note that

$$w d_{n-k,n}(\gamma)^{-1} \in \ker(d_0 : CPT_k \rightarrow CPT_{k-1}).$$

Thus, by Case 1, there exists  $\lambda \in CPT_n$  such that

$$d_{n-k,n}(\lambda) = w d_{n-k,n}(\gamma)^{-1},$$

and hence  $d_{n-k,n}(\lambda\gamma) = w$ . This proves that the map  $d_{n-k,n}$  is surjective.  $\square$

**Proposition 4.9.** *The map  $d_0 : CT_n \rightarrow CT_{n-1}$  is surjective for each  $n \geq 2$ .*

**Proof.** In view of Proposition 4.7, we can write  $CT_{n-1} = CPT_{n-1} \cup \delta_{n-1} CPT_{n-1}$ . Let us take  $w \in CT_{n-1}$ . If  $w \in CPT_{n-1}$ , then by Proposition 4.8, there exists an  $u \in CPT_n$  such that  $d_0(u) = w$ . If  $w \in \delta_{n-1} CPT_{n-1}$ , then again by Proposition 4.8, there exists  $v \in CPT_n$ , such that  $d_0(v) = \delta_{n-1}^{-1} w$ , and hence  $d_0(\delta_n v) = w$ . This complete the proof.  $\square$

Thus, we obtain the following short exact sequences

$$1 \rightarrow \text{Brun}(T_n) \rightarrow CT_n \rightarrow CT_{n-1} \rightarrow 1$$

and

$$1 \rightarrow \text{Brun}(T_n) \rightarrow CPT_n \rightarrow CPT_{n-1} \rightarrow 1.$$

Observe that  $CPT_2 = \text{Brun}(T_2) = PT_2 = 1$  and  $CPT_3 = \text{Brun}(T_3) = PT_3 = \langle (t_1 t_2)^3 \rangle \cong \mathbb{Z}$ . Thus, the preceding exact sequence gives  $CPT_4 = \text{Brun}(T_4) \rtimes \langle (t_1 t_2)^3 \rangle$ .

**Theorem 4.10.** *For each  $u \in PT_{n-1}$  or  $u \in T_{n-1}$ , the system of equations*

$$\begin{cases} d_0(w) = u, \\ d_1(w) = u, \\ \vdots \\ d_{n-1}(w) = u, \end{cases} \quad (4.5)$$

*has a solution if and only if  $u$  satisfies the condition*

$$d_0(u) = d_1(u) = \cdots = d_{n-2}(u).$$

**Proof.** Let  $u \in PT_{n-1}$  such that the system of equations (4.2) has a solution. Then there exists  $w \in PT_n$  such that  $d_0(w) = \cdots = d_{n-1}(w) = u$ . It follows from Proposition 4.6 that  $u \in CPT_{n-1}$ , and hence  $d_0(u) = \cdots = d_{n-2}(u)$ . Conversely, suppose that  $d_0(u) = \cdots = d_{n-2}(u)$ , that is,  $u \in CPT_{n-1}$ . By Proposition 4.8,  $d_0 : CPT_n \rightarrow CPT_{n-1}$  is surjective, and hence there exists  $w \in CPT_n$  which is a solution to (4.2). The proof for the case when  $u \in T_{n-1}$  is similar.  $\square$

## 5. Brunnian doodles on the 2-sphere

Note that the closure of a Brunnian braid is a Brunnian link. The converse is not true and there exist Brunnian links that cannot be obtained as the closure of Brunnian braids (see [12]). The same scenario occurs with doodles on the 2-sphere. Consider the Brunnian doodle on the 2-sphere as shown in Figure 7. We will justify in Remark 5.6 that this Brunnian doodle cannot be realised as the closure of a Brunnian twin.

**Definition 5.1.** *A doodle diagram on the 2-sphere is called minimal if it has no monogons and bigons.*




FIGURE 7. A Brunnian doodle which is not the closure of a Brunnian twin.




FIGURE 8. Transformation of doodle diagrams.




FIGURE 9. Reidemeister moves.

**Theorem 5.2.** [17, Theorem 2.2] *Any doodle has a unique (up to the transformation shown in Figure 8) minimal doodle diagram with a minimal number of intersection points. Further, this minimal doodle diagram can be constructed from any other doodle diagram by applying Reidemeister moves R1 and R2 (see Figure 9) that reduce the number of intersection points.*

For a given reduced word  $w = t_{i_1} \dots t_{i_k} \in T_n$ , let  $\ell(w) = k$  be the length of  $w$ . For each  $1 \leq i \leq n - 1$ , if  $\log_i(w)$  denote the number of  $t_i$ 's present in the expression  $w$ , then

$$\ell(w) = \sum_{i=1}^{n-1} \log_i(w).$$

A cyclic permutation of a word  $w = t_{i_1} \dots t_{i_k} \in T_n$  (not necessarily reduced) is a word  $w' = t_{i_r} t_{i_{r+1}} \dots t_{i_k} t_{i_1} t_{i_2} \dots t_{i_{r-1}}$  for some  $1 \leq r \leq k$ . It is easy to see that  $w$  and  $w'$  are conjugate to each other in  $T_n$ , in fact,

$$w' = (t_{i_1} t_{i_2} \dots t_{i_{r-1}})^{-1} w (t_{i_1} t_{i_2} \dots t_{i_{r-1}}).$$

A word  $w$  is called *cyclically reduced* if each cyclic permutation of  $w$  is reduced. Clearly, a cyclically reduced word is reduced.

**Lemma 5.3.** *Let  $w \in PT_n$  be a pure twin. Then the following assertions hold:*

- (1) *If  $\ell(w)$  is minimal among all the elements in the conjugacy class of  $w$ , then the closure of  $w$  is a minimal doodle diagram.*
- (2) *The closure of  $w$  is an  $n$ -component trivial doodle if and only if  $w$  is a trivial twin.*

**Proof.** It follows from [29, Corollary 2.4] that each word in  $T_n$  is conjugate to some cyclically reduced word. Since  $\ell(w)$  is minimal among all the elements in the conjugacy class of  $w$ , it follows that  $w$  is a cyclically reduced word. Hence, the closure of  $w$  has no bigons. Since  $w$  is pure twin, its closure has no monogons, and hence the diagram is minimal.

By Markov Theorem for doodles on the 2-sphere [16, Theorem 4.1], conjugate twins have the same closure. Thus, we can assume that  $\ell(w)$  is minimal among all the elements in the conjugacy class of  $w$ . It follows from assertion (1) that the closure of  $w$  is a minimal doodle diagram. Note that the number of double points in the closure of the twin  $w$  equals  $\ell(w)$ , and hence  $\ell(w) = 0$ . But, this implies that  $w$  is trivial twin. The converse implication in assertion (2) is obvious.  $\square$

Let  $\widehat{w}$  denote the closure of a twin  $w$  on the 2-sphere. By [17, Theorem 2.1], every oriented doodle on the 2-sphere is the closure of a twin. The *twin index*  $I(D)$  of a doodle  $D$  on the 2-sphere is the minimal  $n$  such that there is a twin  $w \in T_n$  whose closure is equivalent to  $D$ .

**Theorem 5.4.** *An  $m$ -component Brunnian doodle  $D$  on the 2-sphere is the closure of a Brunnian twin if and only if  $I(D) = m$ .*

**Proof.** If  $u$  is a Brunnian twin on  $m$  strands, then its closure on the 2-sphere is a Brunnian doodle on  $m$  components with  $I(\widehat{u}) = m$ . Conversely, if  $D$  is a Brunnian doodle on  $m$  components and  $I(D) = m$ , then there exist  $w \in PT_m$  such that  $\widehat{w} = D$ . Removing any strand from  $w$  corresponds to removing a component from  $D$ . Thus,  $\widehat{d_i(w)}$  is a trivial doodle for each  $i$ . By Lemma 5.3,  $d_i(w) = 1$  for each  $i$ , and hence  $w$  is a Brunnian twin.  $\square$

**Remark 5.5.** An analogue of Theorem 5.4 for Brunnian links in  $S^3$  is proved in [24, Theorem 2.2].

**Remark 5.6.** The Brunnian doodle in Figure 7 cannot have twin index 3, since the closure of a pure twin on 3 strands is a minimal doodle diagram with the number of crossings being a multiple of 6. Hence, this Brunnian doodle cannot be realised as the closure of a Brunnian twin.

## 6. Simplicial structure on pure twin groups

In this section, we discuss simplicial structures on twin and pure twin groups and relate them with Milnor's construction for simplicial spheres.

**6.1. Simplicial sets and simplicial groups.** We recall some basic definitions and constructions [25, 26].

**Definition 6.1.** A sequence of sets  $X_* = \{X_n\}_{n \geq 0}$  is called a simplicial set if there are face maps

$$d_i : X_n \longrightarrow X_{n-1} \text{ for } 0 \leq i \leq n$$

and degeneracy maps

$$s_i : X_n \longrightarrow X_{n+1} \text{ for } 0 \leq i \leq n,$$

which satisfy the following simplicial identities:

- (1)  $d_i d_j = d_{j-1} d_i$  if  $i < j$ ,
- (2)  $s_i s_j = s_{j+1} s_i$  if  $i \leq j$ ,
- (3)  $d_i s_j = s_{j-1} d_i$  if  $i < j$ ,
- (4)  $d_j s_j = \text{id} = d_{j+1} s_j$ ,
- (5)  $d_i s_j = s_j d_{i-1}$  if  $i > j + 1$ .

We view  $X_n$  geometrically as the set of  $n$ -simplices including all possible degenerate simplices. Here, a simplex  $x$  is *degenerate* if  $x = s_i(y)$  for some simplex  $y$  and degeneracy operator  $s_i$ , otherwise  $x$  is *non-degenerate*. A simplicial set  $X_*$  is *pointed* if we fix a basepoint  $\star \in X_0$  that creates one and only one degenerate  $n$ -simplex in each  $X_n$  by applying iterated degeneracy operations on it. A *simplicial group* is a simplicial set  $X_*$  such that each  $X_n$  is a group and all face and degeneracy maps are group homomorphisms.

**Remark 6.2.** In the context of braid-type groups (for example, braid group  $B_n$ , virtual braid group  $VB_n$ , welded braid group  $WB_n$ , etc.), the maps  $d_i$  usually represents deleting of the  $(i + 1)$ -th strand and  $s_i$  represents doubling of the  $(i + 1)$ -th strand.

**Remark 6.3.** Note that the defining identities of a bi- $\Delta$ -set and that of a simplicial set are similar. The only differences are that we don't have  $d_{j+1} s_j = \text{id}$  for bi- $\Delta$ -sets, and when viewed as maps from  $X_{n-1} \rightarrow X_n$ , the number of degeneracy maps is one less than the number of coface maps. We have used the bi- $\Delta$ -set structure at three instances in the preceding sections. The first instance of usage of a bi- $\Delta$ -set is Proposition 2.4, though its arguments can be modified to adapt to a simplicial set structure. The second instance is the proof of Proposition 4.8, where we defined the element  $w_{k,n}$  and showed that  $w_{k,n} \in CPT_n$ . In the latter case, a simplicial structure would not be helpful. Finally, using the Decomposition Theorem for bi- $\Delta$ -groups, we have given a decomposition of pure twin groups in Proposition 3.11 with Brunnian subgroups as constituents.

Let  $G_* = \{G_n\}_{n \geq 0}$  be a simplicial group. The group of *Moore  $n$ -cycles*  $Z_n(G_*) \leq G_n$  is defined by

$$Z_n(G_*) = \bigcap_{i=0}^n \text{Ker}(d_i : G_n \rightarrow G_{n-1})$$

and the group of *Moore  $n$ -boundaries*  $B_n(G_*) \leq G_n$  is defined by

$$B_n(G_*) = d_0 \left( \bigcap_{i=1}^{n+1} \text{Ker}(d_i : G_{n+1} \rightarrow G_n) \right).$$

Simplicial identities guarantees that  $B_n(G_*)$  is a (normal) subgroup of  $Z_n(G_*)$  (see [6, Proposition 4.1.3] or [14, Example 7.7]). The  $n$ -th *Moore homotopy group*  $\pi_n(G_*)$  of  $G_*$  is defined by

$$\pi_n(G_*) = Z_n(G_*)/B_n(G_*).$$

It is a classical result due to Moore [28] that  $\pi_n(G_*) \cong \pi_n(|G_*|)$ , where  $|G_*|$  is the geometric realisation of  $G_*$ . A simplicial group  $G_*$  is called *contractible* if  $\pi_n(G_*) = 1$  for all  $n > 0$ .

Milnor's  $F[K]$  construction is the adjoint functor to the forgetful functor from the category of pointed simplicial groups to the category of pointed simplicial sets. For a given pointed simplicial set  $K_* = \{K_n, \star\}_{n \geq 0}$ , Milnor's  $F[K]$  construction is the simplicial group with  $F[K]_n = F(K_n \setminus \star)$ , the free group on  $K_n \setminus \star$ , with the face and the degeneracy maps induced from the face and degeneracy maps of  $K_*$ . It is well-known from [26] that there is weak homotopy equivalence

$$|F[K]_*| \simeq \Omega\Sigma|K_*|, \quad (6.1)$$

where  $|X_*|$  denotes the geometric realisation of a simplicial set  $X_*$ . Here,  $\Omega Z$  is the loop space of all based loops in a pointed topological space  $Z$  and  $\Sigma Z$  is the reduced suspension of  $Z$ .

Consider the pointed simplicial 2-sphere  $S^2 = \Delta[2]/\partial\Delta[2]$  with

$$S_0^2 = \{\star\}, S_1^2 = \{\star\}, S_2^2 = \{\star, \sigma\}, S_3^2 = \{\star, s_0(\sigma), s_1(\sigma), s_2(\sigma)\}, \dots,$$

$$S_n^2 = \{\star, x_{ij} \mid 0 \leq i < j \leq n-1\}, \dots$$

where  $\sigma = (0, 1, 2)$  is the non-degenerate 2-simplex and

$$x_{ij} = s_{n-1} \dots s_{j+1} \hat{s}_j s_{j-1} \dots s_{i+1} \hat{s}_i s_{i-1} \dots s_0(\sigma)$$

with  $\widehat{s}_k$  meaning that the degeneracy map  $s_k$  is omitted. Then  $F[S^2]$  construction has the following terms:

$$\begin{aligned} F[S^2]_0 &= 1, \\ F[S^2]_1 &= 1, \\ F[S^2]_2 &= F(\sigma), \\ F[S^2]_3 &= F(s_0(\sigma), s_1(\sigma), s_2(\sigma)), \\ F[S^2]_4 &= F(s_1s_0(\sigma), s_2s_0(\sigma), s_3s_0(\sigma), s_2s_1(\sigma), s_3s_1(\sigma), s_3s_2(\sigma)), \\ &\vdots \\ F[S^2]_n &= F(x_{ij}; 0 \leq i < j \leq n-1), \\ &\vdots \end{aligned}$$

For each  $n \geq 2$ , the group  $F[S^2]_n$  is a free group of rank  $n(n-1)/2$ . In this construction of the simplicial 2-sphere, it is convenient to present the degeneracy map  $s_i$  as a doubling of the  $(i+1)$ -th component and the face map  $d_i$  as deletion of the  $(i+1)$ -th component. For example,

$$\begin{aligned} s_0(\sigma) &= (0, 0, 1, 2), & s_1(\sigma) &= (0, 1, 1, 2), & s_2(\sigma) &= (0, 1, 2, 2), \\ s_1s_0(\sigma) &= (0, 0, 0, 1, 2), & s_2s_0(\sigma) &= (0, 0, 1, 1, 2), & s_3s_0(\sigma) &= (0, 0, 1, 2, 2), \\ s_2s_1(\sigma) &= (0, 1, 1, 1, 2), & s_3s_1(\sigma) &= (0, 1, 1, 2, 2), & s_3s_2(\sigma) &= (0, 1, 2, 2, 2). \end{aligned}$$

The face and degeneracy maps are determined with respect to the standard simplicial identities for simplicial groups. For example, the first non-trivial face maps  $d_i : F[S^2]_3 \rightarrow F[S^2]_2$  are given by

$$\begin{aligned} d_0 &: s_0(\sigma) \mapsto \sigma, & s_1(\sigma) &\mapsto \star, & s_2(\sigma) &\mapsto \star, \\ d_1 &: s_0(\sigma) \mapsto \sigma, & s_1(\sigma) &\mapsto \sigma, & s_2(\sigma) &\mapsto \star, \\ d_2 &: s_0(\sigma) \mapsto \star, & s_1(\sigma) &\mapsto \sigma, & s_2(\sigma) &\mapsto \sigma, \\ d_3 &: s_0(\sigma) \mapsto \star, & s_1(\sigma) &\mapsto \star, & s_2(\sigma) &\mapsto \sigma. \end{aligned}$$

Milnor's construction gives a possibility to define the homotopy groups  $\pi_n(S^3)$  combinatorially, in terms of free groups. By (6.1), the geometric realisation of  $F[S^2]_*$  is weakly homotopically equivalent to the loop space  $\Omega S^3$ . Thus, the homotopy groups of  $S^3$  are isomorphic to the Moore homotopy groups of  $F[S^2]$ , that is,

$$\pi_{n+1}(S^3) \cong Z_n(F[S^2]_*)/B_n(F[S^2]_*). \quad (6.2)$$

**6.2. Simplicial pure twin group.** By [1, Theorem 2], we have

$$PT_3 = \langle (t_1t_2)^3 \rangle \cong \mathbb{Z}, \quad PT_4 \cong F_7,$$

where  $F_7$  is the free group on the elements

$$x_1 = (t_1t_2)^3, \quad x_2 = ((t_1t_2)^3)^{t_3}, \quad x_3 = ((t_1t_2)^3)^{t_3t_2}, \quad x_4 = ((t_1t_2)^3)^{t_3t_2t_1},$$

$$x_5 = (t_2 t_3)^3, \quad x_6 = ((t_2 t_3)^3)^{t_1}, \quad x_7 = ((t_2 t_3)^3)^{t_1 t_2}.$$

Let  $SPT_* = \{SPT_n\}_{n \geq 0}$ , where  $SPT_n = PT_{n+1}$  for each  $n \geq 0$ . Following the methodology of [11], consider the sequence of groups

$$\dots \begin{array}{c} \xrightarrow{\dots} \\ \xleftrightarrow{\dots} \\ \xleftarrow{\dots} \end{array} PT_4 \begin{array}{c} \xrightarrow{\dots} \\ \xleftrightarrow{\dots} \\ \xleftarrow{\dots} \end{array} PT_3 \begin{array}{c} \xrightarrow{\dots} \\ \xleftrightarrow{\dots} \\ \xleftarrow{\dots} \end{array} PT_2 \begin{array}{c} \xrightarrow{\dots} \\ \xleftrightarrow{\dots} \\ \xleftarrow{\dots} \end{array} PT_1$$

with face and degeneracy homomorphisms

$$d_i : SPT_n = PT_{n+1} \rightarrow SPT_{n-1} = PT_n, \\ s_i : SPT_n = PT_{n+1} \rightarrow SPT_{n+1} = PT_{n+2},$$

where the face map  $d_i$  is the deleting of the  $(i + 1)$ -th strand and the degeneracy map  $s_i$  is the doubling of the  $(i + 1)$ -th strand for each  $0 \leq i \leq n$ . For example, we prove in the proof of Proposition 3.5 that  $d_3 : PT_4 \rightarrow PT_3$  is given by  $d_3(x_1) = y$  and

$$d_3(x_2) = d_3(x_3) = d_3(x_4) = d_3(x_5) = d_3(x_6) = d_3(x_7) = 1,$$

where  $y = (t_1 t_2)^3 \in PT_3$ . As in the classical case it is not difficult to prove the following result, whose proof is adapted from [3, Proposition 3.1].

**Proposition 6.4.** *SPT\* is a contractible simplicial group.*

**Proof.** Let  $x \in Z_n(SPT_*)$  be a Moore  $n$ -cycle, that is,  $x \in SPT_n$  and  $d_i(x) = 1$  for all  $0 \leq i \leq n$ . Note that  $SPT_*$  admits an additional degeneracy map  $\iota_{n+1} : SPT_n \rightarrow SPT_{n+1}$ , which adds a trivial strand on the left of the diagram of the twin. If we set  $y = \iota_{n+1}(x) \in SPT_{n+1}$ , then we see that  $d_j(y) = 1$  for all  $1 \leq j \leq n + 1$  and  $d_0(y) = x$ . Thus,  $x \in B_n(SPT_*)$  is a Moore  $n$ -boundary, and hence  $\pi_n(SPT_*) = 1$  for all  $n$ . □

We write  $U_{n,i} := \text{Ker}(d_i : PT_n \rightarrow PT_{n-1})$  for each  $0 \leq i \leq n - 1$ . Then, we have the following short exact sequence

$$1 \longrightarrow U_{n,i} \longrightarrow PT_n \xrightarrow{d_i} PT_{n-1} \longrightarrow 1$$

with the splitting given by  $d^i : PT_{n-1} \rightarrow PT_n$  as defined in Proposition 2.4. This gives a semi-direct product decomposition  $PT_n = U_{n,i} \rtimes PT_{n-1}$ . Clearly,  $U_{3,0} = U_{3,1} = U_{3,2} = PT_3$ . The following problem seems interesting.

**Problem 6.5.** *Find presentations of  $U_{n,i}$  for  $n \geq 4$ .*

We construct a simplicial subgroup  $K_*$  of  $SPT_*$ , which would be the image of the simplicial sphere  $S^2$  under a simplicial map. We set  $K_0 = K_1 = 1$ ,  $K_2 = SPT_2 = \langle c_{0,1;2} \rangle$ , the infinite cyclic group generated by  $c_{0,1;2} = (t_1 t_2)^3$ , and

$$K_3 = \langle c_{1,2;3} = s_0(c_{0,1;2}), c_{0,2;3} = s_1(c_{0,1;2}), c_{0,1;3} = s_2(c_{0,1;2}) \rangle.$$

In general, we define

$$K_n = \langle c_{i,j;n} = s_{n-1} \dots s_{j+1} \widehat{s}_j s_{j-1} \dots s_{i+1} \widehat{s}_i s_{i-1} \dots s_0(c_{0,1;2}) \mid 0 \leq i < j \leq n - 1 \rangle,$$



the subgroup of  $SPT_n$  generated by  $n(n-1)/2$  elements. It follows from the simplicial identities that  $d_i(c_{i,j;n}) \in K_{n-1}$  and  $s_j(c_{i,j;n}) \in K_{n+1}$  for each generator  $c_{i,j;n}$  of  $K_n$  and all  $d_i, s_j$ . Thus, for each  $n \geq 0$ , restriction of face maps  $d_i : SPT_n \rightarrow SPT_{n-1}$  gives face maps  $d_i : K_n \rightarrow K_{n-1}$ . Similarly, restriction of degeneracy maps  $s_i : SPT_n \rightarrow SPT_{n+1}$  induce degeneracy maps  $s_i : K_n \rightarrow K_{n+1}$ , turning  $K_* = \{K_n\}_{n \geq 0}$  into a simplicial subgroup of  $SPT_*$ .

**Theorem 6.6.**  $K_3 \cong F[S^2]_3$  and  $K_4 \cong F[S^2]_4$ .

**Proof.** Using the geometrical interpretation of  $c_{0,1;2}$  (see Figure 2) and degeneracy maps  $s_i$ , we write the generators of  $K_3$  in terms of the generators of  $PT_4$  as follows:

$$\begin{aligned} c_{1,2;3} &= (t_2 t_1 t_3 t_2)(t_1 t_2 t_3)(t_1 t_2 t_3) = ((t_1 t_2)^3)^{t_3 t_2} (t_2 t_3)^3 = x_3 x_5, \\ c_{0,2;3} &= (t_1 t_2 t_3)(t_2 t_1 t_3 t_2)(t_1 t_2 t_3) = ((t_2 t_3)^3)^{t_1} ((t_1 t_2)^3)^{t_3} = x_6 x_2, \\ c_{0,1;3} &= (t_1 t_2 t_3)(t_1 t_2 t_3)(t_2 t_1 t_3 t_2) = (t_1 t_2)^3 ((t_2 t_3)^3)^{t_1 t_2} = x_1 x_7. \end{aligned}$$

Since  $PT_4$  is a free group of rank 7, it follows that  $K_3$  is a free group of rank 3, and hence  $K_3 \cong F[S^2]_3$ . It is known from [15, Section 5] that  $SPT_4 = PT_5$  is free group of rank 31, but [15] does not give any free generating set for  $PT_5$ . However, using [1, Theorem 4], we obtain a generating set for  $PT_5$  of cardinality 43. By removing the redundant generators, we obtain the following minimal generating set for  $PT_5$ :

$$\begin{aligned} a_1 &= (t_1 t_2)^3 & a_{17} &= ((t_2 t_3)^3)^{t_4} \\ a_2 &= ((t_1 t_2)^3)^{t_3} & a_{18} &= ((t_2 t_3)^3)^{t_1 t_2 t_4 t_3 t_2 t_1} \\ a_3 &= ((t_1 t_2)^3)^{t_3 t_2} & a_{19} &= ((t_2 t_3)^3)^{t_1 t_2 t_4 t_3 t_2} \\ a_4 &= ((t_1 t_2)^3)^{t_3 t_2 t_1} & a_{20} &= ((t_2 t_3)^3)^{t_1 t_2 t_4 t_3} \\ a_5 &= ((t_1 t_2)^3)^{t_3 t_2 t_1 t_4 t_3 t_2} & a_{21} &= ((t_2 t_3)^3)^{t_1 t_2 t_4} \\ a_6 &= ((t_1 t_2)^3)^{t_3 t_2 t_1 t_4 t_3} & a_{22} &= ((t_2 t_3)^3)^{t_1 t_4 t_3 t_2 t_1} \\ a_7 &= ((t_1 t_2)^3)^{t_3 t_2 t_1 t_4} & a_{23} &= ((t_2 t_3)^3)^{t_1 t_4 t_3 t_2} \\ a_8 &= ((t_1 t_2)^3)^{t_3 t_2 t_4 t_3} & a_{24} &= ((t_2 t_3)^3)^{t_1 t_4 t_3} \\ a_9 &= ((t_1 t_2)^3)^{t_3 t_4 t_3 t_2} & a_{25} &= ((t_2 t_3)^3)^{t_1 t_4} \\ a_{10} &= ((t_1 t_2)^3)^{t_3 t_4} & a_{26} &= ((t_3 t_4)^3) \\ a_{11} &= (t_2 t_3)^3 & a_{27} &= ((t_3 t_4)^3)^{t_2 t_1 t_3 t_2} \\ a_{12} &= ((t_2 t_3)^3)^{t_1} & a_{28} &= ((t_3 t_4)^3)^{t_2 t_1 t_3} \\ a_{13} &= ((t_2 t_3)^3)^{t_1 t_2} & a_{29} &= ((t_3 t_4)^3)^{t_2 t_1} \\ a_{14} &= ((t_2 t_3)^3)^{t_4 t_3 t_2 t_1} & a_{30} &= ((t_3 t_4)^3)^{t_2 t_3} \\ a_{15} &= ((t_2 t_3)^3)^{t_4 t_3 t_2} & a_{31} &= ((t_3 t_4)^3)^{t_2} \\ a_{16} &= ((t_2 t_3)^3)^{t_4 t_3} & & \end{aligned}$$

By definition, we have

$$K_4 = \langle s_1 s_0(c_{0,1;2}), s_2 s_0(c_{0,1;2}), s_3 s_0(c_{0,1;2}), s_2 s_1(c_{0,1;2}), s_3 s_1(c_{0,1;2}), s_3 s_2(c_{0,1;2}) \rangle.$$

Direct calculation gives

$$\begin{aligned} s_1(x_3 x_5) &= a_8 a_{16} a_{26}, & s_2(x_3 x_5) &= a_{23} a_9 a_{31} a_{17}, \\ s_3(x_3 x_5) &= a_3 a_{19} a_{11} a_{30}, & s_2(x_6 x_2) &= a_{29} a_{25} a_{10}, \\ s_3(x_6 x_2) &= a_{12} a_{28} a_2 a_{20}, & s_3(x_1 x_7) &= a_1 a_{13} a_{27}. \end{aligned}$$

Thus,  $K_4$  is free of rank 6, and hence  $K_4 \cong F[S^2]_4$ .  $\square$

**Problem 6.7.** Determine a presentation of  $K_n$  for  $n \geq 4$ .

We consider  $c_{0,1;2}$  as a 2-simplex in the simplicial group  $SPT_*$ . Since

$$d_0(c_{0,1;2}) = d_1(c_{0,1;2}) = d_2(c_{0,1;2}) = 1,$$

there is a (unique) simplicial map

$$\theta : S^2 \rightarrow SPT_*$$

such that  $\theta(\sigma) = c_{0,1;2}$ , where  $\sigma = (0, 1, 2)$  is the non-degenerate 2-simplex of the simplicial 2-sphere  $S^2$ . By Milnor's construction, the simplicial map  $\theta$  extends uniquely to a simplicial homomorphism

$$\Theta : F[S^2]_* \longrightarrow SPT_*.$$

We note that  $K_* = \Theta(F[S^2]_*)$  and it is the smallest simplicial subgroup of  $SPT_*$  containing  $c_{0,1;2}$ . Further, by Proposition 6.6,

$$\Theta_n : F[S^2]_n \longrightarrow SPT_n$$

is injective for  $n \leq 4$ . If each  $\Theta_n : F[S^2]_n \rightarrow SPT_n$  is injective, then by (6.2), we have

$$\pi_{n+1}(S^3) \cong Z_n(F[S^2]_*)/B_n(F[S^2]_*) \cong Z_n(K_*)/B_n(K_*) \cong \pi_n(K_*).$$

Thus, if  $\Theta$  is injective, then we can describe  $\pi_{n+1}(S^3)$  as a quotient of a subgroup of  $PT_{n+1}$ . For instance, the generator of  $\pi_3(S^3) \cong \mathbb{Z}$  can be represented by the pure twin  $(t_1 t_2)^3$ .

It appears that the following statement holds.

**Conjecture 6.8.**  $\Theta : F[S^2]_* \longrightarrow K_*$  is an isomorphism.

**Acknowledgement.** The authors are grateful to the anonymous referee for many valuable suggestions. Valeriy Bardakov is supported by the state contract of the Sobolev Institute of Mathematics, SB RAS (No. I.1.5, Project FWNF-2022-0009). Pravin Kumar is supported by the PMRF fellowship at IISER Mohali. Mahender Singh is supported by the Swarna Jayanti Fellowship grants DST/SJF/MSA-02/2018-19 and SB/SJF/2019-20/04.

## 7. Declaration

The authors declare that there is no data associated to this paper and that there are no conflicts of interests.

## References

- [1] BARDAKOV, VALERIY; SINGH, MAHENDER.; VESNIN, ANDREI. Structural aspects of twin and pure twin groups. *Geom. Dedicata* **203** (2019), 135–154. [MR4027588](#), [Zbl 1454.20080](#), [arXiv:1811.04020](#), doi: [10.1007/s10711-019-00429-1](#). [1238](#), [1242](#), [1243](#), [1244](#), [1257](#), [1259](#)
- [2] BARDAKOV, VALERY G.; MIKHAILOV, ROMAN; VERSHININ, VLADIMIR V.; WU, JIE. Brunnian braids on surfaces. *Algebr. Geom. Topol.* **12** (2012), no. 3, 1607–1648. [MR2966697](#), [Zbl 1270.20036](#), [arXiv:0909.3387](#), doi: [10.2140/agt.2012.12.1607](#). [1237](#)
- [3] BARDAKOV, VALERY G.; WU, JIE. Lifting theorem for the virtual pure braid groups. Preprint, 2023. [arXiv:2002.08686](#). [1258](#)
- [4] BARDAKOV, V. G.; VERSHININ, VLADIMIR V.; WU, JIE. On Cohen braids. *Proc. Steklov Inst. Math.* **286** (2014), no. 1, 16–32. [MR3482589](#), [Zbl 1319.20033](#), [arXiv:1312.2924](#), doi: [10.1134/S0081543814060029](#). [1248](#)
- [5] BARTHOLOMEW, ANDREW; FENN, ROGER; KAMADA, NAOKO; KAMADA, SEIICHI. Doodles on surfaces. *J. Knot Theory Ramifications* **27** (2018), no. 12, 1850071, 26 pp. [MR3876348](#), [Zbl 1403.57003](#), [arXiv:1612.08473](#), doi: [10.1142/S0218216518500712](#). [1236](#)
- [6] BERRICK, A. J.; COHEN, FREDERICK R.; WONG, YAN L.; WU, JIE. Configurations, braids, and homotopy groups. *J. Amer. Math. Soc.* **19** (2006), no. 2, 265–326. [MR2188127](#), [Zbl 1188.55007](#), doi: [10.1090/S0894-0347-05-00507-2](#). [1237](#), [1242](#), [1256](#)
- [7] BERRICK, A. J.; HANBURY, ELIZABETH; WU, JIE. Brunnian subgroups of mapping class groups and braid groups, *Proc. Lond. Math. Soc.* (3) **107** (2013), no. 4, 875–906. [MR3108834](#), [Zbl 1282.57002](#), doi: [10.1112/plms/pds096](#). [1237](#)
- [8] BERRICK, A. J.; HANBURY, ELIZABETH; WU, JIE. Delta-structures on mapping class groups and braid groups. *Trans. Amer. Math. Soc.* **366** (2014), no. 4, 1879–1903. [MR3152716](#), [Zbl 1434.20022](#), doi: [10.1090/S0002-9947-2013-05889-8](#). [1237](#)
- [9] BJÖRNER, ANDERS; WELKER, VOLKMAR. The homology of “ $k$ -equal” manifolds and related partition lattices. *Adv. Math.* **110** (1995), no. 2, 277–313. [MR1317619](#), [Zbl 0845.57020](#), doi: [10.1006/aima.1995.1012](#). [1237](#)
- [10] COHEN, FREDERICK R. On combinatorial group theory in homotopy. *Homotopy theory and its applications* (Cocoyoc, 1993), 57–63, *Contemp. Math.* **188**, Amer. Math. Soc., Providence, RI, 1995. ISBN:0-8218-0305-0. [MR1349129](#), [Zbl 0849.55015](#). [1248](#)
- [11] COHEN, FREDERICK R.; WU, JIE. Artin’s braid groups, free groups, and the loop space of the 2-sphere. *Q. J. Math.* **62** (2011), no. 4, 891–921. [MR2853222](#), [Zbl 1252.55005](#), [arXiv:math/0409307](#), doi: [10.1093/qmath/haq010](#). [1237](#), [1258](#)
- [12] DUZHIN, FEDOR; WONG, SHAO MING Z. On two constructions of Brunnian links. *J. Knot Theory Ramifications* **23** (2014), no. 3, 1420002, 6 pp. [MR3200492](#), [Zbl 1294.57005](#), doi: [10.1142/S0218216514200028](#). [1252](#)
- [13] FENN, ROGER; TAYLOR, PAUL. Introducing doodles. *Topology of low-dimensional manifolds* (Proc. Second Sussex Conf., Chelwood Gate, 1977), pp. 37–43. *Lecture Notes in Math.* **722**. Springer, Berlin, 1979. ISBN:3-540-09506-3. [MR0547452](#), [Zbl 0409.57003](#). [1236](#)
- [14] FRIEDMAN, GREG. Survey article: An elementary illustrated introduction to simplicial sets. *Rocky Mountain J. Math.* **42** (2012), no.2, 353–423. [MR2915498](#), [Zbl 1248.55001](#), [arXiv:0809.4221](#), doi: [10.1216/RMJ-2012-42-2-353](#). [1256](#)
- [15] GONZÁLEZ, JESÚS; LEÓN-MEDINA, JOSÉ L.; ROQUE-MÁRQUEZ, CHRISTOPHER. Linear motion planning with controlled collisions and pure planar braids. *Homology Homotopy Appl.* **23** (2021), no. 1, 275–296. [MR4170471](#), [Zbl 1455.55001](#), [arXiv:1902.06190](#), doi: [10.4310/HHA.2021.v23.n1.a15](#). [1259](#)

- [16] GOTIN, KONSTANTIN. Markov theorem for doodles on two-sphere. Preprint, 2018. [arXiv:1807.05337](https://arxiv.org/abs/1807.05337). [1236](#), [1254](#)
- [17] KHOVANOV, MIKHAIL. Doodle groups. *Trans. Amer. Math. Soc.* **349** (1997), no. 6, 2297–2315. [MR1370644](#), [Zbl 0869.57001](#), doi: [10.1090/S0002-9947-97-01706-6](https://doi.org/10.1090/S0002-9947-97-01706-6). [1236](#), [1253](#), [1254](#)
- [18] KHOVANOV, MIKHAIL. New geometrical constructions in low-dimensional topology. Preprint, 1990. [1236](#)
- [19] KHOVANOV, MIKHAIL. Real  $K(\pi, 1)$  arrangements from finite root systems. *Math. Res. Lett.* **3** (1996), no. 2, 261–274. [MR1386845](#), [Zbl 0861.55010](#), doi: [10.4310/MRL.1996.v3.n2.a11](https://doi.org/10.4310/MRL.1996.v3.n2.a11). [1236](#), [1239](#)
- [20] LEI, FENGCHUN; LI, FENGLING; WU, JIE. On simplicial resolutions of framed links. *Trans. Amer. Math. Soc.* **366** (2014), no. 6, 3075–3093. [MR3180740](#), [Zbl 1297.57017](#), doi: [10.1090/S0002-9947-2013-05957-0](https://doi.org/10.1090/S0002-9947-2013-05957-0). [1237](#)
- [21] LEVINSON, HENRY W. Decomposable braids and linkages. *Trans. Amer. Math. Soc.* **178** (1973), 111–126. [MR324684](#), [Zbl 0275.55002](#), doi: [10.2307/1996692](https://doi.org/10.2307/1996692). [1247](#)
- [22] LEVINSON, HENRY W. Decomposable braids as subgroups of braid groups. *Trans. Amer. Math. Soc.* **202** (1975), 51–55. [MR362287](#), [Zbl 0305.55002](#) doi: [10.2307/1997296](https://doi.org/10.2307/1997296). [1242](#)
- [23] LI, JINGYAN; WU, JIE. Artin braid groups and homotopy groups. *Proc. Lond. Math. Soc.* (3) **99** (2009), no. 3, 521–556. [MR2551462](#), [Zbl 1241.20042](#), doi: [10.1112/plms/pdp005](https://doi.org/10.1112/plms/pdp005). [1237](#)
- [24] MAES, ARNAUD; CERF, CORINNE. A family of Brunnian links based on Edwards’ construction of Venn diagrams. *J. Knot Theory Ramifications* **10** (2001), no. 1, 97–107. [MR1822143](#), [Zbl 0998.57020](#), doi: [10.1142/S0218216501000743](https://doi.org/10.1142/S0218216501000743). [1254](#)
- [25] MAY, J. P. Simplicial objects in algebraic topology. Van Nostrand Mathematical Studies, 11. D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1967. vi+161 pp. [MR0222892](#), [Zbl 0165.26004](#). [1255](#)
- [26] MILNOR, JOHN W.; ADAMS, JOHN F. On the construction FK. In: Algebraic topology: a student’s guide by J.F. Adams, pp. 119–136. London Mathematical Society Lecture Note Series, 4. Cambridge Univ. Press. 1972. [MR0445484](#), [Zbl 0234.55002](#), doi: [10.1017/CBO9780511662584.011](https://doi.org/10.1017/CBO9780511662584.011). [1255](#), [1256](#)
- [27] MIKHAILOV, ROMAN V.; WU, JIE. Combinatorial group theory and the homotopy groups of finite complexes. *Geom. Topol.* **17** (2013), 235–272. [MR3035327](#), [Zbl 1270.55011](#), doi: [10.2140/gt.2013.17.235](https://doi.org/10.2140/gt.2013.17.235). [1237](#)
- [28] MOORE, J. C. Homotopie des complexes monoïdaux. In: Séminaire Henri Cartan de l’Ecole Normale Supérieure, 1954/1955. Algèbres d’Eilenberg-MacLane et homotopie. *Secrétariat Mathématique, 11 rue Pierre Curie, Paris*, 1956. iii+234 pp. [MR0087935](#), [Zbl 0067.15601](#), [http://www.numdam.org/item/SHC\\_1954-1955\\_\\_7\\_2\\_A8\\_0/](http://www.numdam.org/item/SHC_1954-1955__7_2_A8_0/). [1256](#)
- [29] NAIK, TUSHAR K.; NANDA, NEHA; SINGH, MAHENDER. Conjugacy classes and automorphisms of twin groups. *Forum Math.* **32** (2020), no. 5, 1095–1108. [MR4145210](#), [Zbl 1530.20086](#), [arXiv:1906.06723](https://arxiv.org/abs/1906.06723), doi: [10.1515/forum-2019-0321](https://doi.org/10.1515/forum-2019-0321). [1254](#)
- [30] NAIK, TUSHAR K.; NANDA, NEHA; SINGH, MAHENDER. Structure and automorphisms of pure virtual twin groups. *Monatsh. Math.* **202** (2023), no. 3, 555–582. [MR4651964](#), [Zbl 7750833](#), [arXiv:2008.10035](https://arxiv.org/abs/2008.10035), doi: [10.1007/s00605-023-01851-0](https://doi.org/10.1007/s00605-023-01851-0). [1244](#), [1245](#)
- [31] NAIK, TUSHAR K.; NANDA, NEHA; SINGH, MAHENDER. Virtual planar braid groups and permutations. *J. Group Theory* **27** (2024), no. 3, 443–483. [MR4739232](#), [Zbl 1537.20087](#), [arXiv:2109.13035](https://arxiv.org/abs/2109.13035), doi: [10.1515/jgth-2023-0010](https://doi.org/10.1515/jgth-2023-0010). [1245](#)
- [32] WU, JIE. Homotopy theory of the suspensions of the projective plane. *Mem. Amer. Math. Soc.* **162** (2003), no. 769, x+130 pp. [MR1955357](#), [Zbl 1022.55009](#), doi: [10.1090/memo/0769](https://doi.org/10.1090/memo/0769). [1250](#)
- [33] WU, JIE. On maps from loop suspensions to loop spaces and the shuffle relations on the Cohen groups. *Mem. Amer. Math. Soc.* **180** (2006), no. 851, vi+64 pp. [MR2203532](#), [Zbl 1098.55006](#), doi: [10.1090/memo/0851](https://doi.org/10.1090/memo/0851). [1238](#), [1239](#), [1245](#), [1246](#)

(Valeriy G. Bardakov) SOBOLEV INSTITUTE OF MATHEMATICS, ACAD. KOPTYUG AVE. 4, 630090 NOVOSIBIRSK, RUSSIA; NOVOSIBIRSK STATE AGRARIAN UNIVERSITY, DOBROLYUBOVA STR., 160, 630039 NOVOSIBIRSK, RUSSIA; REGIONAL SCIENTIFIC AND EDUCATIONAL MATHEMATICAL CENTER OF TOMSK STATE UNIVERSITY, LENIN AVE. 36, 634009 TOMSK, RUSSIA.

[bardakov@math.nsc.ru](mailto:bardakov@math.nsc.ru)

(Pravin Kumar) DEPARTMENT OF MATHEMATICAL SCIENCES, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH (IISER) MOHALI, SECTOR 81, SAS NAGAR, P O MANAULI, PUNJAB 140306, INDIA

[pravin444enaj@gmail.com](mailto:pravin444enaj@gmail.com)

(Mahender Singh) DEPARTMENT OF MATHEMATICAL SCIENCES, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH (IISER) MOHALI, SECTOR 81, SAS NAGAR, P O MANAULI, PUNJAB 140306, INDIA

[mahender@iisermohali.ac.in](mailto:mahender@iisermohali.ac.in)

This paper is available via <http://nyjm.albany.edu/j/2024/30-55.html>.