

Preimages question for surjective endomorphisms on $(\mathbb{P}^1)^n$

Xiao Zhong

ABSTRACT. Let K be a number field and let $f : (\mathbb{P}^1)^n \rightarrow (\mathbb{P}^1)^n$ be a dominant endomorphism defined over K . We show that if V is an f -invariant subvariety (that is, $f(V) = V$) then there is a positive integer s_0 such that

$$(f^{-s-1}(V) \setminus f^{-s}(V))(K) = \emptyset$$

for every integer $s \geq s_0$, answering the Preimages Question of Matsuzawa, Meng, Shibata, and Zhang in the case of $(\mathbb{P}^1)^n$.

CONTENTS

1. Introduction	633
2. Proof of the main theorem	635
Acknowledgments	648
References	648

1. Introduction

Let X be a projective variety and let $f : X \rightarrow X$ be a surjective self-map such that both X and f are defined over a number field K . To study the dynamics of (X, f) , it is important to identify the closed subvarieties of $Y \subseteq X$ that are invariant under f ; i.e., subvarieties with $f(Y) \subseteq Y$. For an invariant subvariety Y for the map f , it is natural to study its preimages under iterates of f . An important principle within arithmetic dynamics is that the underlying geometric structure should exert significant influence on the arithmetic structure, which gives rise to the expectation that the tower of K -points:

$$Y(K) \subseteq (f^{-1}(Y))(K) \subseteq (f^{-2}(Y))(K) \subseteq \dots$$

should eventually stabilize. This expectation has been made precise in the form of the Preimages Question of Matsuzawa, Meng, Shibata, and Zhang [MMS23, Question 8.4(1)].

Received February 15, 2024.

2020 *Mathematics Subject Classification.* 37P55, 14G05.

Key words and phrases. Preimages Question, invariant subvarieties, arithmetic dynamics, rational points on varieties.

The author was supported in part by NSERC grant RGPIN-2022-02951.

There are recent works dealing with some special cases of the Preimages Question. Notably, it has been solved with affirmative answer when X is a smooth variety with non-negative Kodaira dimension and f is étale [BMS23, Theorem 1.2]. Additionally, if we consider the case when $f = (g, g) : X \times X \rightarrow X \times X$ is a diagonal map, with $g : X \rightarrow X$ is a surjective morphism, then the diagonal subvariety $\Delta \subseteq X \times X$ is an invariant subvariety under f , and in this special case the Preimages Question becomes a cancellation problem, which asks whether there exists a natural number s with the property that, for all $x, y \in X(K)$, if $g^n(x) = g^n(y)$ for some natural number n then we must in fact have $g^s(x) = g^s(y)$. This special form is a dynamical cancellation problem considered in the work of Bell, Matsuzawa, and Satriano [BMS23], and this question is again answered affirmatively when X is a curve [BMS23, Theorem 1.3] by applying p -adic uniformization techniques of Rivera-Letelier [Riv03]. Considerably more is known for polynomial maps on \mathbb{P}^1 , and a more general dynamical cancellation results allowing multiple polynomial self-maps on \mathbb{P}^1 is also proved along these lines in [BMS23, Theorem 1.7] and [Zho23, Theorem 1.2].

Our main result is to give a positive solution to the Preimages Question for surjective endomorphisms on $(\mathbb{P}^1)^n$. It is well-known that a surjective endomorphism of $(\mathbb{P}^1)^n$ has some iterate that becomes a split rational map (f_1, \dots, f_n) . Since answering the Preimages question for an iterate of a self-map gives one the answer for the original map, our main theorem, stated below, solves the Preimages Question for surjective endomorphisms on $(\mathbb{P}^1)^n$.

Theorem 1.1. *Let K be a number field, let $n \geq 1$, and let $f = (f_1, \dots, f_n) : (\mathbb{P}_K^1)^n \rightarrow (\mathbb{P}_K^1)^n$ be a split rational map defined over a number field K with at least one f_i of degree greater than 1. If $V \subseteq (\mathbb{P}^1)^n$ is a subvariety defined over K that is invariant under f then there exists a non-negative integer s_0 such that*

$$(f^{-s-1}(V) \setminus f^{-s}(V))(K) = \emptyset$$

for all $s \geq s_0$.

It is also well known that Theorem 1.1 holds when $n = 1$.

Remark 1.2. This paper works over a number field K for the reader's convenience, but the argument also works if we let K instead be a finitely generated extension of \mathbb{Q} . Proposition 2.4 uses the fact that the set of roots of unity in the union of all finite field extensions of \mathbb{Q} of bounded degree is finite and it remains true that the set of roots of unity inside the union of field extensions of bounded degree over K is finite when K is a finitely generated extension over \mathbb{Q} . Proposition 2.5 uses [BMS23, Theorem 2.3] whose proof is based on embedding K into a finite extension of \mathbb{Q}_p for a suitable prime p , which works for any finitely generated field extension of \mathbb{Q} as well (see [BGT16, Proposition 2.5.3.1]). Everything else goes through directly without additional changes when considering K as a finitely generated field extension of \mathbb{Q} .

2. Proof of the main theorem

In this section we give the proof of the main result. Our first result shows that we can reduce to the case when our maps all have degree ≥ 2 .

Lemma 2.1. *Let $n \geq 2$ be a natural number and let $f = (f_1, \dots, f_n) : (\mathbb{P}^1)^n \rightarrow (\mathbb{P}^1)^n$ be a split rational map defined over a algebraically closed characteristic zero field and suppose that there exists a positive integer $k \in \{1, \dots, n-1\}$ such that $\deg(f_i) > 1$ when $1 \leq i \leq k$ and $\deg(f_i) = 1$ when $i > k$. If V is an irreducible subvariety of $(\mathbb{P}^1)^n$ that is invariant under f then there exist subvarieties $V_1 \subseteq (\mathbb{P}^1)^k$ and $V_2 \subseteq (\mathbb{P}^1)^{n-k}$ such that $V = V_1 \times V_2$ and such that V_1 is invariant under $g_1 = (f_1, \dots, f_k)$ and V_2 is invariant under $g_2 = (f_{k+1}, \dots, f_n)$.*

Proof. The proof is similar to [Xie23, Proposition 3.14]. Let d denote the dimension of V and let $\pi_1 : (\mathbb{P}^1)^n \rightarrow (\mathbb{P}^1)^k$ and $\pi_2 : (\mathbb{P}^1)^n \rightarrow (\mathbb{P}^1)^{n-k}$ be respectively the projection onto the first k factors and the projection onto the last $n-k$ factors.

Let $V_i = \pi_i(V)$ and $d_i = \dim(V_i)$ for $i = 1, 2$. Since $V \subseteq V_1 \times V_2$ we have $d \leq d_1 + d_2$ and $d = d_1 + d_2$ if and only if $V = V_1 \times V_2$.

Thus we may assume without loss of generality that $d < d_1 + d_2$. We take α_1 to be numerical class of the ample line bundle $\pi_1^* \mathcal{O}_{(\mathbb{P}^1)^k}(1, 1, \dots, 1)$ and α_2 to be the numerical class of the ample line bundle $\pi_2^* \mathcal{O}_{(\mathbb{P}^1)^{n-k}}(1, 1, \dots, 1)$. Furthermore, we let β_j denote the numerical class of the line bundle $p_j^* \mathcal{O}_{\mathbb{P}^1}(1)$ for each $j \in \{1, 2, \dots, n\}$ and $p_j : (\mathbb{P}^1)^n \rightarrow \mathbb{P}^1$ the projection onto the j -th coordinate. Then

$$\alpha_1 = \beta_1 + \dots + \beta_k, \quad (2.1)$$

$$\alpha_2 = \beta_{k+1} + \dots + \beta_n. \quad (2.2)$$

Notice that for any $j \in \{0, 1, \dots, d\}$, $\alpha_1^j \cdot \alpha_2^{d-j} \cdot V \geq 0$ and it is positive if $j = d_1$ or $d - j = d_2$. Let's denote $I_1 = \{1, 2, \dots, k\}$ and $I_2 = \{k+1, \dots, n\}$ from now on.

Since there are only finitely many collections of indices (allowing repetition) of size $d - d_2$ which is inside I_1 and also only finitely many collections of indices (allowing repetition) of size d_2 which is inside I_2 , there exists a $\{r_1, r_2, \dots, r_{d-d_2}\} \subseteq I_1$, and a $\{e_1, e_2, \dots, e_{d_2}\} \subseteq I_2$ such that

$$V \cdot \prod_{t=1}^{d-d_2} \beta_{r_t} \prod_{l=1}^{d_2} \beta_{e_l} > 0,$$

and $C = \prod_{t=1}^{d-d_2} c_{r_t}$, where $c_i = \deg(f_i)$ for $i \in I$, is the maximum in the set

$$\left\{ \prod_{i \in I'} c_i : V \cdot \prod_{i \in I'} \beta_i \prod_{v \in J'} \beta_v > 0, I' \subseteq I_1, J' \subseteq I_2, |I'| = d - d_2, |J'| = d_2 \right\}.$$

We let V' be an irreducible subvariety of $V \cap \bigcap_{t=1}^{d-d_2} \beta_{r_t}$ of dimension d_2 and here we abuse notation to view β_{r_t} 's as some suitable hypersurfaces in the numerical classes $p_{r_t}^* \mathcal{O}_{\mathbb{P}^1}(1)$'s. Then noticed that $\dim(\pi_1(V')) = d_1 - (d - d_2)$.

We have

$$V' \cdot (\alpha_1)^{d_1-(d-d_2)} \cdot (\alpha_2)^{d-d_1} > 0.$$

This implies that there exists collections of indices

$$\{j_1 = r_1, j_2 = r_2, \dots, j_{d-d_2} = r_{d-d_2}, j_{d-d_2+1}, \dots, j_{d_1}\} \subseteq I_1$$

and

$$\{u_1, u_2, \dots, u_{d-d_1}\} \subseteq I_2$$

such that

$$V \cdot \prod_{t=1}^{d_1} \beta_{j_t} \prod_{l=1}^{d-d_1} \beta_{u_l} > 0. \quad (2.3)$$

Notice that

$$\prod_{t=1}^{d_1} c_{j_t} > C \quad (2.4)$$

by construction and Equation (2.3) implies that

$$V \cdot \alpha_1^{d_1} \cdot \alpha_2^{d-d_1} > 0.$$

Let $u_1, u_2 \in \mathbb{R}$, we have

$$\begin{aligned} (\deg(f|_V))(V \cdot (u_1\alpha_1 + u_2\alpha_2)^d) &= f_*(V) \cdot ((u_1\alpha_1 + u_2\alpha_2)^d) \\ &= V \cdot (u_1(g_1)^*(\alpha_1) + u_2(g_2)^*(\alpha_2))^d \\ &= V \cdot \left(u_1 \sum_{i=1}^k c_i \beta_i + u_2 \sum_{i=k+1}^n c_i \beta_i \right)^d. \end{aligned}$$

Now we compare the coefficients of the $u_1^{d_1} u_2^{d-d_1}$ and $u_1^{d-d_2} u_2^{d_2}$ terms and we obtain that for each positive integer m :

$$\deg(f^m|_V) \alpha_1^{d_1} \cdot \alpha_2^{d-d_1} \cdot V = \left(\sum_{i=1}^k c_i^m \beta_i \right)^{d_1} \cdot \left(\sum_{i=k+1}^n c_i^m \beta_i \right)^{d-d_1} \cdot V,$$

and

$$\deg(f^m|_V) \alpha_1^{d-d_2} \cdot \alpha_2^{d_2} \cdot V = \left(\sum_{i=1}^k c_i^m \beta_i \right)^{d-d_2} \cdot \left(\sum_{i=k+1}^n c_i^m \beta_i \right)^{d_2} \cdot V.$$

Therefore

$$\deg(f^m|_V) = \left(\sum_{i=1}^k c_i^m \beta_i \right)^{d_1} \cdot \alpha_2^{d-d_1} \cdot V / (\alpha_1^{d_1} \cdot \alpha_2^{d-d_1} \cdot V) \quad (2.5)$$

and

$$\deg(f^m|_V) = \left(\sum_{i=1}^k c_i^m \beta_i \right)^{d-d_2} \cdot \alpha_2^{d_2} \cdot V / (\alpha_1^{d-d_2} \cdot \alpha_2^{d_2} \cdot V), \quad (2.6)$$

since $c_i = 1$ when $i \in I_2$,

$$\alpha_1^{d_1} \cdot \alpha_2^{d-d_1} \cdot V > 0,$$

and

$$\alpha_1^{d-d_2} \cdot \alpha_2^{d_2} \cdot V > 0.$$

Now, by (2.4) and Equation (2.5), we have

$$\deg(f^m|_V) = aM^m + o(M^m)$$

for some positive integer a and M , such that $M > C \geq \prod_{i=1}^{d-d_2} c_{l_i}$ for all $\{l_1, \dots, l_{d-d_2}\} \subseteq I_1$ and $\{v_1, \dots, v_{d_2}\} \subseteq I_2$ with the property

$$\prod_{t=1}^{d-d_2} \beta_{l_t} \prod_{s=1}^{d_2} \beta_{v_s} \cdot V > 0.$$

On the other hand, Equation (2.6) gives that $\deg(f^m|_V) = \mathcal{O}(C^m)$. This is a contradiction and so the result follows. \square

Lemma 2.1 allows us to reduce to the case that all maps f_i in the statement of Theorem 1.1 have degree at least two when proving the result, and we now consider this restricted case. We begin with a result in which the maps are special.

Definition 2.2. We define the Chebyshev polynomial of degree r to be the unique polynomial T_r of degree r such that

$$T_r((x + x^{-1})/2) = (x^r + x^{-r})/2$$

for any $x \in \mathbb{C}^*$.

Remark 2.3. We introduce the notation here to avoid confusion in the later discussion: Let $n, D \in \mathbb{N}$ and K be a field. For a quasi-projective variety $W \subseteq (\mathbb{P}_K^1)^n$ defining over K and a set L which is a union of the finite field extensions of K of degree not exceeding D , which is not necessarily a field or a ring, we let $W(L)$ denote the largest subset of points in $W(\overline{K})$ such that for every $i \in \{1, 2, \dots, n\}$,

$$\pi_i(W(L)) \subseteq \bigcup_{K': K' \subseteq L} \mathbb{P}_{K'}^1,$$

where π_i is the projection to the i -th factor and K' ranges in the set of field extensions of K contained in L as a set. We call this set of points $W(L)$ the L -points of W .

More concretely, after fixing a coordinate of W defined over K , we have $W(L)$ is the set of points $(x_1, x_2, \dots, x_n) \in W \subseteq (\mathbb{P}_K^1)^n$ such that $x_i \in L \cup \{\infty\}$, for each $i \in \{1, 2, \dots, n\}$.

Moreover, we let L^* denote the set $L \setminus \{0\}$.

Proposition 2.4. *Let K be a number field, let n and D be positive integers that are at least 2, and let K' be the union of all the finite field extensions of K of degree not greater than D . Suppose that $f = (f_1, \dots, f_n) : (\mathbb{P}_K^1)^n \rightarrow (\mathbb{P}_K^1)^n$ is a split rational map defined over K in which the f_i 's all have the same degree $d \geq 2$ and each f_i is conjugate to either $x^{\pm d}$ or $\pm T_d$ for $i \in \{1, 2, \dots, n\}$. If $V \subset (\mathbb{P}^1)^n$ is an irreducible f -invariant hypersurface defined over K , then there exists a non-negative integer s_0 such that*

$$(f^{-s-1}(V) \setminus f^{-s}(V))(K') = \emptyset$$

for all $s \geq s_0$.

Proof. We first enlarge our number field K to a larger number field so that each f_i can be conjugated to either $x^{\pm d}$ or to $\pm T_d$ by a map $v_i \in \mathrm{PGL}_2(K)$ and $\{i, e^{\pi i/4}\} \subseteq K$. Notice that it is enough to prove the statement in this larger field we henceforth assume that K is a number field satisfying this condition. We can then perform a change of coordinates and may assume without the loss of the generality that f is of the form $f = (f_1(x_1), \dots, f_n(x_n)) = (x_1^{\pm d}, x_2^{\pm d}, \dots, x_k^{\pm d}, \pm T_d(x_{k+1}), \dots, \pm T_d(x_n))$ for some $k \in \{0, \dots, n\}$.

Let

$$\mu = (x_1, \dots, x_k, (x_{k+1} + x_{k+1}^{-1})/2, \dots, (x_n + x_n^{-1})/2) : (\mathbb{C}^*)^n \rightarrow (\mathbb{P}^1)^n,$$

and

$$G' = (x_1^{\pm d}, \dots, x_k^{\pm d}, \pm x_{k+1}^d, \dots, \pm x_n^d) : (\mathbb{C}^*)^n \rightarrow (\mathbb{C}^*)^n$$

such that $\mu \circ G' = f \circ \mu$. Then take

$$\mu' = (x_1, \dots, x_k, \xi_{k+1} x_{k+1}, \dots, \xi_n x_n)$$

and

$$G = (x_1^{\pm d}, \dots, x_k^{\pm d}, x_{k+1}^d, \dots, x_n^d),$$

where ξ_{k+1}, \dots, ξ_n are some roots of unity such that $\mu' \circ G = G' \circ \mu'$. Enlarge K so that $\{\xi_{k+1}, \dots, \xi_n\} \subseteq K$ and abuse notation from now on to let μ be $\mu \circ \mu'$. So we have

$$\mu \circ G = f \circ \mu.$$

Notice that further abusing notations by replacing G by G^2 and f by f^2 , we may let

$$G = (x_1^d, \dots, x_k^d, x_{k+1}^d, \dots, x_n^d)$$

and

$$\mu \circ G = f \circ \mu.$$

Now we prove the statement by induction on the dimension n . If $n = 1$, then the statement is certainly true as V is a just a finite set of points. Now, we suppose it is true when $n \leq n_0$ for some positive integer n_0 and we want to show it is true when $n = n_0 + 1$. From now on, let $n = n_0 + 1$. Let W be the preimage of V under μ , which is a subvariety in $(\mathbb{C}^*)^n$. Since $f(V) = V$, we have

$$G(W) \subseteq W.$$

Thus, there exists a subvariety $W' \subseteq W$ such that $G(W') = W'$, after replacing f and G by a suitable iterate, we have $W \subseteq G^{-1}(W')$. Therefore, by [Hin88, Lemma 10], we have W' is a finite union of translation of algebraic subgroups of $(\mathbb{C}^*)^n$. Notice that $(\xi_i x_i + \xi_i^{-1} x_i^{-1})/2$ maps $\{0, \infty\}$ to $\{\infty\}$, for each $i \in \{k+1, \dots, n\}$. Thus, we have

$$V' = V \setminus \mu(W) = \left(\bigcup_{i=1}^n (V \cap H_i) \right) \cup \bigcup_{i=1}^k (V \cap P_i),$$

where H_i is the hypersurface in $(\mathbb{P}^1)^n$ defined by having the i -th coordinate equals to ∞ and P_i is the hypersurface with the i -th coordinate equals to 0. We write $V' = \bigcup_{j=1}^{k_0} V_j$ for some positive integer k_0 , where V_j 's are irreducible components of $V \cap H_i$ or $V \cap P_i$ for some $i \in \{1, 2, \dots, n\}$ and $l \in \{1, 2, \dots, k\}$. Also, $f(V') = V'$ because $f(V) = V$, $f(V') \subseteq V'$ and $f(\mu(W)) \subseteq \mu(W)$ as power maps take 0 or ∞ to themselves and Chebyshev polynomials take ∞ to itself.

After replacing f by some iteration we may assume each V_j is invariant under f . Notice that every irreducible component of $V \cap H_i$ or $V \cap P_i$ has dimension at least $n-2$, if not empty, for each $i \in \{1, 2, \dots, n\}$ and $l \in \{1, 2, \dots, k\}$. Thus, after some reorderings of the coordinates, each V_j projects onto either an irreducible hypersurface in $(\mathbb{P}^1)^{n-1}$ or $(\mathbb{P}^1)^{n-1}$ itself, denoted as V'_j in both case, and projects to either 0 or ∞ in the remaining \mathbb{P}^1 factor. If $V'_j = (\mathbb{P}^1)^{n-1}$, then its preimages under f restricted to $(\mathbb{P}^1)^{n-1}$ is itself and there is nothing to show, so from now on we assume V'_j is an irreducible hypersurface. Also, V'_j is invariant under f'_j , the restriction of f to the corresponding $(\mathbb{P}^1)^{n-1}$ factors. Now the induction hypothesis tells us that there exists non-negative integer s'_j such that

$$((f'_j)^{-s-1}(V'_j) \setminus (f'_j)^{-s}(V'_j))(K') = \emptyset$$

for every $s \geq s'_j$. But notice that this also implies

$$(f^{-s-1}(V_j) \setminus f^{-s}(V_j))(K') = \emptyset$$

for $s \geq s'_j$, since preimages of $\{\infty, 0\}$ under x^d and preimages of ∞ under $\pm T_d$ are just themselves. Take $s'_0 = \max_{j=1}^{k_0} s'_j$.

Now we look at the preimages of V under μ . If this is empty, then V lives completely inside some H_i or P_j for some $i \in \{1, 2, \dots, n\}$ and $j \in \{1, 2, \dots, k\}$ and, using arguments above, we go to the induction steps. From now on, we assume that $W = \mu^{-1}(V)$ is not empty. Notice that $\mu^{-1}((V \setminus V')(K')) \subseteq W(L)$, where L is the union of finite field extensions of degree $D \cdot 2^n$ over K . Recall that $W \subseteq G^{-1}(W')$ and $G(W') = W'$. It is enough to show that there exists a non-negative integer s_1 such that for any $x \in (L^*)^n$, if $G^s(x) \in W'(L)$ for some $s \geq 0$, then $G^{s_1}(x) \in W'(L)$. As, if this s_1 exists and there still exists a $y \in (\mathbb{P}^1_{K'})^n$ and a $s > s_1$ such that $f^s(y) \in \mu(W)(K')$ but $f^{s_1}(y) \notin \mu(W)$, then

for any $x \in \mu^{-1}(y) \subset (L^*)^n$,

$$G^{s+1}(x) \in G(\mu^{-1}(f^s(x))) \subset G(W(L)) \subseteq W'(L).$$

This implies that $G^{s_1}(x) \in W'(L)$. But we have $G^{s_1}(x) \notin W$ from $f^{s_1}(y) \notin \mu(W)$. This is a contradiction.

Now, recall W' is a finite union of torsion translation of algebraic subgroups. It is enough to prove for each torsion translation of algebraic subgroup separately. So, without loss of generality, we assume

$$W' = V(x_1^{r_1} \dots x_n^{r_n} - \epsilon)$$

where r_1, \dots, r_n are integers that are not all zero and ϵ is an torsion element such that $\epsilon^d = \epsilon$. Enlarge K, K' and L if necessary so that $\epsilon \in K$. Then L -points in preimages of $W'(L)$ under iterates of G live in the union of $V(x_1^{r_1} \dots x_n^{r_n} - \epsilon\lambda)$, where λ ranges over elements which is d^m -torsion for some non-negative integer m . Notice that if $(x_1, \dots, x_n) \in (L^*)^n$, then $x_i \in L_i$ where L_i is a finite field extension of K such that $[L_i : K] \leq D \cdot 2^n$ for each $i \in \{1, 2, \dots, n\}$. Therefore, $\lambda \in L' = L_1 L_2 \dots L_n$ where L' is a finite field extension of K such that $[L' : K] \leq (D \cdot 2^n)^n$. Therefore, λ is a root of unity such that

$$[\mathbb{Q}(\lambda) : \mathbb{Q}] \leq (D \cdot 2^n)^n \cdot [K : \mathbb{Q}].$$

We claim that

$$M = \{\lambda \in \overline{\mathbb{Q}} : \lambda^{d^m} = 1, \text{ for some } m \in \mathbb{N}^+, [\mathbb{Q}(\lambda) : \mathbb{Q}] \leq (D \cdot 2^n)^n \cdot [K : \mathbb{Q}]\}$$

is a finite set. This is because the set of roots of unity of bounded degree is finite, which is a special case of Northcott property, and M is a subset of it. Then we can take

$$s_1 = \max_{\lambda \in M} \text{ord}(\lambda),$$

where $\text{ord}(\lambda)$ is the minimum non-negative integer m such that $\lambda^{d^m} = 1$. Then for any $y \in (L^*)^n$ such that $G^s(y) \in W'(L)$ for some $s > s_1$, we have $y \in V(x_1^{r_1} \dots x_n^{r_n} - \epsilon\lambda)$ for some $\lambda \in M$ and thus $G^{s_1}(y) \in W'(L)$. This will conclude the proof in this case.

To conclude, we take $s_0 = \max(s'_0, s_1)$. We have

$$(f^{-s-1}(V) \setminus f^{-s}(V))(K') = \emptyset$$

for any $s \geq s_0$. □

Proposition 2.5. *Let K be a number field, let n and D be positive integers that are at least 2, and let K' be the union of all the finite field extensions of K of degree not greater than D in some fixed algebraic closure of K . Suppose that $f = (f_1, \dots, f_n) : (\mathbb{P}_K^1)^n \rightarrow (\mathbb{P}_K^1)^n$ is a split rational map defined over K in which the f_i 's all have the same degree $d \geq 2$ and each f_i is a Lattès map for $i \in \{1, 2, \dots, n\}$. If $V \subset (\mathbb{P}^1)^n$ is an irreducible f -invariant hypersurface defined over K that projects dominantly onto any subset of $n - 1$ coordinate axes, then there exists a non-negative integer s_0 such that*

$$(f^{-s-1}(V) \setminus f^{-s}(V))(K') = \emptyset$$

for all $s \geq s_0$.

Proof. In this case, $f = (f_1, f_2, \dots, f_n)$ where each f_i 's is a Lattès map. We consider a diagram:

$$\begin{array}{ccc} E_1 \times E_2 \times \cdots \times E_n & \xrightarrow{G} & E_1 \times E_2 \times \cdots \times E_n \\ \pi \downarrow & & \downarrow \pi \\ (\mathbb{P}^1)^n & \xrightarrow{f} & (\mathbb{P}^1)^n \end{array} \quad (2.7)$$

where $G = (g_1, g_2, \dots, g_n)$ is also a split morphism on the abelian variety $E_1 \times E_2 \times \cdots \times E_n$ and $\pi = (\pi_1, \pi_2, \dots, \pi_n)$ is a projection map such that each g_i and π_i satisfies $f_i \circ \pi_i = \pi_i \circ g_i$ by the fact that f_i is a Lattès map. Let $W = \pi^{-1}(V)$, which is a subvariety in $E_1 \times E_2 \times \cdots \times E_n$. Since each π_i has degree bounded by 6 [Sil07, Proposition 6.37] and we can enlarge K to a larger number field so that g_i 's are defined over K , we replace K by this larger number field. It is enough to show that L -points of W are stabilized under preimages of G , for every finite field extension L of K of degree not greater than $(6^n \cdot D)^n$. We have $G(W) \subseteq W$ by construction and there exists a subvariety $W' \subseteq W$ such that, after replacing G and f by some suitable iterate, $G(W') = W'$ and $G(W) \subseteq W'$. We just need to show that there exists a non-negative integer s_0 such that for any $s \geq s_0$ and $[L : K] \leq (6^n \cdot D)^n$,

$$(G^{-s-1}(W') \setminus G^{-s}(W'))(L) = \emptyset. \quad (2.8)$$

This will imply that

$$(f^{-s-1}(V) \setminus f^{-s}(V))(K) = \emptyset, \quad (2.9)$$

since, if not, there exists $x \in (\mathbb{P}_K^1)^n$ such that $f^s(x) \in V(K)$ but $f^{s_0}(x) \notin V(K)$ for some $s > s_0$. Thus there exists a

$$y \in \pi^{-1}(x) \subset (E_1 \times E_2 \times \cdots \times E_n)(L),$$

where L is a finite field extension of K of degree not greater than $(6^n \cdot D)^n$, such that $G^{s+1}(y) \in W'(L)$, but $G^{s_0}(y) \notin W$ so is not in W' . This is a contradiction to (2.8).

Notice that it is enough to prove 2.8 for each irreducible component of W' . From now on we abuse notation and let W denote an arbitrary irreducible component of W' . To prove the statement (2.8), we first consider \tilde{W} which is a translation of W by some \tilde{K} -point in $E_1 \times E_2 \times \cdots \times E_n$ such that \tilde{W} is invariant under \tilde{G} , where \tilde{G} is a group homomorphism of the abelian variety with the property that G is the composition of \tilde{G} with a suitable translation.

Then, by [GTZ11, Theorem 3.1], \tilde{W} contains a Zariski dense set of preperiodic points of \tilde{G} .

Note that the preperiodic points of \tilde{G} are torsion points in

$$E_1 \times E_2 \times \cdots \times E_n$$

(see [GTZ11, Claim 3.2] and notice that \tilde{G} , G and f are polarizable). Therefore, we can apply the classical Manin-Mumford conjecture (proved in [Ray83]) to

get that \tilde{W} is a translation of some subabelian variety of $E_1 \times E_2 \times \cdots \times E_n$ and so is W . Thus G restricted to W is an étale morphism and therefore we apply [BMS23, Theorem 2.3] to conclude that there exists a nonnegative integer s_0 such that for any $s \geq s_0$ we have

$$(G^{-s-1}(W) \setminus G^{-s}(W))(L) = \emptyset, \quad (2.10)$$

for any finite field extension L of K of degree not greater than $(6^n \cdot D)^n$. \square

Lemma 2.6. *Let V be an irreducible hypersurface in $(\mathbb{P}^1)^n$ that projects dominantly onto every subset of $n - 1$ coordinate axes. Let $f = (f_1, \dots, f_n)$ be a split rational map with $\deg(f_i) > 1$ for all $1 \leq i \leq n$. If $f(V) = V$ then $\deg(f_1) = \deg(f_2) = \cdots = \deg(f_n)$.*

Proof. Let's consider for each $i \in \{1, 2, \dots, n\}$ the following diagram:

$$\begin{array}{ccc} V & \xrightarrow{f|_V} & V \\ \iota \downarrow & & \downarrow \iota \\ (\mathbb{P}^1)^n & \xrightarrow{f} & (\mathbb{P}^1)^n \\ \pi_i \downarrow & & \downarrow \pi_i \\ (\mathbb{P}^1)^{n-1} & \xrightarrow{g_i} & (\mathbb{P}^1)^{n-1} \end{array} \quad (2.11)$$

where $\iota : V \rightarrow (\mathbb{P}^1)^n$ is the natural embedding, π_i is the projection to the $n - 1$ coordinates excluding the i -th coordinate, and

$$g_i = (f_1, f_2, \dots, f_{i-1}, f_{i+1}, \dots, f_n),$$

i.e., we skip the i -th index and so $g_i : (\mathbb{P}^1)^{n-1} \rightarrow (\mathbb{P}^1)^{n-1}$. Since $\pi_i \circ \iota$ is a finite morphism by the assumption that V projects dominantly onto every subset of $n - 1$ coordinate axes, we have

$$\deg(f|_V) = \deg(g_i) = \prod_{j \in I_i} \deg(f_j),$$

where $I_i = \{1, 2, \dots, i - 1, i + 1, \dots, n\}$. Since this argument and equation hold for each $i \in \{1, 2, \dots, n\}$, we have

$$\prod_{j \in I_i} \deg(f_j) = \prod_{j \in I_k} \deg(f_j)$$

for any $i, k \in \{1, 2, \dots, n\}$. This implies that $\deg(f_i) = \deg(f_k)$ for any pair of i, k in $\{1, 2, \dots, n\}$. \square

Lemma 2.7. *In the case where the degrees of f_1, \dots, f_n are all at least two, it is enough to prove Theorem 1.1 with the additional assumption that V is an irreducible hypersurface of dimension not less than 1 that projects dominantly onto every subset of $n - 1$ coordinate axes in $(\mathbb{P}^1)^n$.*

Proof. The idea is the same as in [GN18, Proposition 2.1]. We assume throughout this proof that we have the hypotheses of Theorem 1.1 and that the maps f_1, \dots, f_n all have degree ≥ 2 .

We use induction on the dimension of V . If $\dim(V) = 0$, then V is a finite set of points and Theorem 1.1 is true since the preimages of K -points in V live inside a set of K -points of bounded height. Now suppose that the conclusion to Theorem 1.1 holds when $\dim(V) < D$, we prove the case that $\dim(V) = D$. Notice that it is enough to prove that the conclusion to Theorem 1.1 holds for each irreducible components of V separately after replacing f by a suitable iterate.

Thus we assume that V is irreducible. Then there exist D coordinate axes such that the projection π of V onto these axes is dominant.

Without loss of generality, we assume that they are the first D coordinates and for $j > D$ we let π_j denote the projection from V to the coordinate axes indexed by $\{1, \dots, D, j\}$.

Therefore $\pi_j(V)$ is a hypersurface in $(\mathbb{P}^1)^{D+1}$ and also

$$H_j := \pi_j(V) \times (\mathbb{P}^1)^{n-D-1}$$

is a hypersurface in $(\mathbb{P}^1)^n$. Now we claim that V is a component of $\bigcap_{j=D+1}^n H_j$. Notice that $\dim(\bigcap_{j=D+1}^n H_j) \geq D$ and $V \subset \bigcap_{j=D+1}^n H_j$. So we just need to show that $\dim(\bigcap_{j=D+1}^n H_j) = D$.

Since $\pi_j(V)$ projects dominantly onto the first D coordinate axes, there exists a Zariski open subset $U \subset (\mathbb{P}^1)^D$ such that for each

$$\alpha = (\alpha_1, \dots, \alpha_D) \in U$$

there exists a finite set $S_{\alpha,j}$ such that if $(\alpha_1, \dots, \alpha_n) \in H_j$ and $(\alpha_1, \dots, \alpha_D) \in \pi(V)$ then $\alpha_j \in S_{\alpha,j}$. This is saying for each $\alpha \in U$ there are only finitely many points in $\bigcap_{j=D+1}^n H_j$ such that the first D coordinates are equal to α . Therefore, $\dim(\bigcap_{j=D+1}^n H_j) = D$ and V is a component of $\bigcap_{j=D+1}^n H_j$. Notice that since $f(V) = V$, we also have $f(H_j) = H_j$.

We claim that if for each H_j , we have the statement of Theorem 1.1 holds, then certainly it holds for V . To prove the claim, we first notice that there exists a subvariety $H \subseteq \bigcap_{j=D+1}^n H_j$ containing V such that $f(H) = H$ and $f(\bigcap_{j=D+1}^n H_j) = H$ if we replace f by some iterate. We denote V_1, \dots, V_k as the irreducible components of H of dimension D and without loss of generality assume that $V = V_1$ and $f(V_i) = V_i$ for each $i \in \{1, \dots, k\}$ by replacing f with suitable iterates. Assume that the conclusion to Theorem 1.1 holds for each H_j . Then there exist non-negative integers s_j such that each $x \in (\mathbb{P}_K^1)^n$ satisfies $f^s(x) \in H_j(K)$ for some non-negative integer s , and we have $f^{s_j}(x) \in H_j(K)$ for each $D+1 \leq j \leq n$.

Now for each $x \in (\mathbb{P}_K^1)^n$ such that $f^s(x) \in V(K)$ for some non-negative integer s , we have certainly $f^s(x) \in H_j(K)$ for each $D+1 \leq j \leq n$. Thus letting

s'_0 denote the quantity $\max_{D+1 \leq j \leq n} \{s_j\}$, we have $f^{s'_0}(x) \in \bigcap_{j=D+1}^n H_j(K)$ and $f^{s'_0+1}(x) \in H$. If $f^{s'_0+1}(x) \notin V$ then $f^{s'_0+1}(x) \in V_i$ for some $i \in \{2, \dots, k\}$ such that $f^s(x) \in V \cap V_i$. Since $f(V) = V$ and $f(V_i) = V_i$, we have $f(V \cap V_i) \subseteq (V \cap V_i)$ and there exists $V' \subseteq V \cap V_i$ such that $f(V \cap V_i) = V'$ and $f(V') = V'$ after further replacing f by a suitable iterate. Notice that $\dim(V') < D$ and x is in the preimages of V' under f . By the induction hypothesis, there exists a positive integer s'_1 such that $(f^{-s'_1-1}(V') \setminus f^{-s'_1}(V))(K) = \emptyset$. Therefore, taking $s_0 = \max\{s'_1, s'_0 + 1\}$, we have $f^{s_0}(x) \in V(K)$. We proved that it is enough to prove that the conclusion to Theorem 1.1 holds for each H_j .

Now for each H_j , it is equivalent to prove the statement of Theorem 1.1 for $\pi_j(V) \subset (\mathbb{P}^1)^{D+1}$ which is an irreducible hypersurface invariant under $f' = (f_1, \dots, f_D)$ and projecting dominantly onto first D coordinate axes. We claim that, after reordering coordinate axes, we have $\pi_j(V) = W_j \times (\mathbb{P}^1)^{m_j}$ for some $0 \leq m_j \leq D$, where $W_j \subseteq (\mathbb{P}^1)^{D+1-m_j}$ projects dominantly onto any subset of $D - m_j$ coordinate axes. This can be proved by induction and the base case is that $D = 1$ and in this case $\pi_j(V)$ either projects dominantly onto any \mathbb{P}^1 factors or it is $W_j \times \mathbb{P}^1$, after reordering coordinate axes, where W_j is a finite set of points. Now, assuming the claim is true for $D = D_0$, $D_0 \geq 1$ is a positive integer, let's prove it for $D = D_0 + 1$. If there exists D subset of coordinate axes such that the projection, $\pi_1(\pi_j(V))$, onto those axes is not dominant and denote π_2 as the projection onto the other \mathbb{P}^1 factor, then $\dim(\pi_1(\pi_j(V))) = D - 1$ since we have

$$\dim(\pi_j(V)) \leq \dim(\pi_1(\pi_j(V))) + \dim(\pi_2(\pi_j(V))). \quad (2.12)$$

So $\dim(\pi_2(\pi_j(V))) = 1$, which implies $\pi_2(\pi_j(V)) = \mathbb{P}^1$. Then the equality actually holds in (2.12) and we have $\pi_j(V) = \pi_1(\pi_j(V)) \times \mathbb{P}^1$. Applying the induction hypothesis on $\pi_1(\pi_j(V))$ concludes the proof of the claim.

Recall, we have already shown that it is equivalent to prove the statement of Theorem 1.1 for $\pi_j(V)$. By the claim, proving for $\pi_j(V)$ is also equivalent to proving for W_j which is an irreducible hypersurface in $(\mathbb{P}^1)^{D+1-m_j}$ invariant under $f'' = (f_1, \dots, f_{D+1-m_j})$, after reordering coordinate axes, and projects dominantly onto any subset of $D - m_j$ coordinate axes. \square

Proposition 2.8. *Let K be a finitely generated field extension of \mathbb{Q} , D a positive integer and let L be the union of the finite extensions of K of degree less or equal to D . Let $f : \mathbb{P}^1_K \rightarrow \mathbb{P}^1_K$ be a surjective morphism of degree greater than 1 defined over K . Then there exists a positive integer N such that if $a, b \in \mathbb{P}^1(L)$ satisfies $f^n(a) = f^n(b)$ for some $n \geq 0$, then $f^N(a) = f^N(b)$.*

Proof. [BMS23, Theorem 3.1] proved this for L a number field. But the proof still works if we have L as above. The only changes to the proof that are needed is that instead of embedding L into \mathbb{Q}_p for a suitable prime p directly such that, after embedding, $f \in \mathbb{Z}_p[[x]]$, we embed K into \mathbb{Q}_p for a suitable prime p such that $f \in \mathbb{Z}_p[[x]]$. Then L is naturally embedded as a set in the union of the

finite field extensions of \mathbb{Q}_p of degree less or equal to D which is inside $\overline{\mathbb{Q}_p}$. Denote such an embedding map as ι .

Notice that the key ingredient of the proof of [BMS23, Theorem 3.1] is the p -adic uniformization which works over \mathbb{C}_p and thus it can be applied here without changes. Also, in the proof [BMS23, Theorem 3.1], N is based on the least common multiple of the orders of roots of unity in a finite extension of \mathbb{Q}_p which is a finite set. And now, we only need to take it to be depending on a positive integer M , the least common multiple of orders of roots of unity in $\overline{\mathbb{Q}_p}$ of degrees bounded by a constant depending only on D , f and K . In particular, the orders of roots of unity in $\iota(L) \subseteq \overline{\mathbb{Q}_p}$ are all dividing M . Notice that this is doable since the set of roots of unity in $\overline{\mathbb{Q}_p}$ such that their degrees over \mathbb{Q}_p are all bounded by a constant is also finite [BMS23, Proposition 3.6(3)] and $\iota(L)$ is contained in the union of the finite field extensions of \mathbb{Q}_p of degree less or equal to D . Thus, the proof follows just as in [BMS23, Theorem 3.1]. \square

Proof of Theorem 1.1 in the case when $\deg(f_1), \dots, \deg(f_n) \geq 2$. Notice that by Lemma 2.7, it is enough to prove the statement with the assumption that V is an irreducible hypersurface projecting dominantly onto any subset of $(n-1)$ coordinate axes in $(\mathbb{P}^1)^n$. Since $f(V) = V$, we have that V contains a Zariski dense set of preperiodic points of f [Fak03, Theorem 5.1]. Therefore, by [GNY18, Theorem 2.2], we have if $n > 2$ then

- (1) either f_1, \dots, \dots, f_n are all Lattès maps;
- (2) or f_i 's are all conjugated to $x^{\pm d_i}$ or $\pm T_{d_i}$, where T_{d_i} 's are Chebyshev polynomials of degree $d_i = \deg(f_i)$.

Notice that by Lemma 2.6 in both cases $d_1 = d_2 = \dots = d_n$. If $n = 2$, then by the proof of [GNY19, Theorem 1.3] we have either f_1, f_2 are both Lattès maps or neither of them is. So overall we have three separate cases to show:

- (1) $n = 2$, f_1 and f_2 are not Lattès maps;
- (2) $n > 2$, f_i 's are conjugated to either $x^{\pm d}$ or $\pm T_d$ for some $d > 1$;
- (3) f_i 's are all Lattès maps.

Case (2) is implied by Proposition 2.4 and Case (3) is implied by Proposition 2.5. We left to prove case (1): By [Pak23, Corollary 4.5] and [Pak23, Remark 4.3], we have that there exists rational functions U_1, U_2 and F defined over \overline{K} such that

$$U_1 \circ F = f_1 \circ U_1,$$

$$U_2 \circ F = f_2 \circ U_2$$

and $W_1 = (U_1, U_2)^{-1}(V)$ is a subvariety such that $(F, F)(W_1) \subseteq W_1$ and it contains an irreducible component $V' \subseteq W_1$ such that $(F, F)(V') = V'$ and $(F, F)(W_1) \subseteq V'$ after replacing F, f_1 and f_2 by some suitable iterate. Also by replacing K with some larger number field, we assume U_1, U_2 and F are defined over K . Let K' be the union of the finite field extensions of K , whose degree is bounded by $\deg(U_1)^2 = \deg(U_2)^2$ of K . It is enough to show that there exists a

positive integer s_0 such that

$$((F, F)^{-s-1}(V') \setminus (F, F)^{-s}(V'))(K') = \emptyset$$

for any integer $s \geq s_0$.

Now if F is Lattès map or is conjugate to either a power map or a Chebyshev polynomial or its negative, we obtain the result from Proposition 2.4 and Proposition 2.5 separately. So, we assume F is not a Lattès map nor conjugate to either a power map or a Chebyshev polynomial or its negative. Then by [Pak23, Theorem 4.15], we have there exists rational functions U_3, U_4, F_1 and F_2 over \mathbb{C} such that

$$U_3 \circ F_1 = F \circ U_3,$$

$$U_4 \circ F_2 = F \circ U_4$$

and $W = (U_3, U_4)^{-1}(W_1)$ is a subvariety such that $(F_1, F_2)(W) \subseteq W$ and it contains a subvariety W' such that $(F_1, F_2)(W') = W'$ after replacing F_1, F_2, f_1, f_2 and F by a suitable iterate. Furthermore, F_1 and F_2 are not generalized Lattès maps by the Theorem.

Now let $\tau_1 = U_1 \circ U_3$ and $\tau_2 = U_2 \circ U_4$ and replace K by a finite generated field extension of K such that τ_1, τ_2, F_1 and F_2 are all defined over K . After replacing F and F_1, F_2 with a suitable iterate and abusing notation to let W be some irreducible component of W' , we may assume $(F_1, F_2)(W) = W$. Let L be the union of the finite field extensions of K of degree bounded by $\deg(\tau_1)^2 = \deg(\tau_2)^2$. Similarly, it is enough to show that there exists a positive integer s_0 such that

$$((F_1, F_2)^{-s-1}(W) \setminus (F_1, F_2)^{-s}(W))(L) = \emptyset$$

for all integers $s \geq s_0$.

In this case, we use [Pak23, Theorem 1.1] and we get that there exists rational functions X_1, X_2, Y_1, Y_2 and B such that there exists some positive integer d satisfies

$$F_1^d = X_1 \circ Y_1 \tag{2.13}$$

$$F_2^d = X_2 \circ Y_2 \tag{2.14}$$

$$B^d = Y_1 \circ X_1 = Y_2 \circ X_2, \tag{2.15}$$

$W \subseteq (Y_1, Y_2)^{-1}(\Delta)$ and also $(F_1^d, F_2^d)((Y_1, Y_2)^{-1}(\Delta)) = W$.

Enlarging K by adjoining coefficients of X_1, Y_1, X_2 and Y_2 and abuse the notation to let K denote this larger field and L be the union of the finite field extensions of degree bounded by $\deg^2(\tau_1) = \deg^2(\tau_2)$ of K .

Now it is enough to show that there exists a non-negative integer s_0 such that for any $s \geq s_0$,

$$((F_1, F_2)^{(-s-1)d}(W) \setminus (F_1, F_2)^{-sd}(W))(L) = \emptyset. \tag{2.16}$$

We claim that to prove the above, it suffices to show that there exists s_0 such that for any $s \geq s_0$

$$((F_1, F_2)^{d(-s-1)}((Y_1, Y_2)^{-1}(\Delta)) \setminus (F_1, F_2)^{-ds}((Y_1, Y_2)^{-1}(\Delta)))(L) = \emptyset. \tag{2.17}$$

To see this, suppose there exists s'_0 such that for any $s \geq s'_0$ Equation (2.17) holds. If there exists a non-negative integer $s > s'_0$ and a $x \in (\mathbb{P}^1 \times \mathbb{P}^1)(L)$ such that $(F_1, F_2)^{sd}(x) \in W(L)$ but $(F_1, F_2)^{s'_0 d}(x) \notin W$, then

$$(F_1, F_2)^{s'_0 d}(x) \in (Y_1, Y_2)^{-1}(\Delta) \setminus W$$

by Equation (2.17). Thus $(F_1, F_2)^{(s'_0+1)d}(x) \in W$. Therefore, take $s_0 = s'_0 + 1$, we have for any $s \geq s_0$

$$((F_1, F_2)^{(-s-1)d}(W) \setminus (F_1, F_2)^{-sd}(W))(L) = \emptyset.$$

Thus, it is enough to show that there exists s_0 such that for any $s \geq s_0$

$$((F_1, F_2)^{d(-s-1)}((Y_1, Y_2)^{-1}(\Delta)) \setminus (F_1, F_2)^{-ds}((Y_1, Y_2)^{-1}(\Delta)))(L) = \emptyset \quad (2.18)$$

which is equivalent to, by Equation (2.13), (2.14) and (2.15),

$$((Y_1, Y_2)^{-1}((B^d, B^d)^{-s-1}(\Delta) \setminus (B^d, B^d)^{-s}(\Delta)))(L) = \emptyset. \quad (2.19)$$

Since, again, Y_1, Y_2 are defined over K , it is enough to show

$$((B^d, B^d)^{-s-1}(\Delta) \setminus (B^d, B^d)^{-s}(\Delta))(L) = \emptyset. \quad (2.20)$$

Notice that if there exists a non-negative integer s_0 such that for any non-negative integer s and any $x, y \in \mathbb{P}^1(L)$, we have

$$B^{ds}(x) = B^{ds}(y) \quad (2.21)$$

implies

$$B^{ds_0}(x) = B^{ds_0}(y), \quad (2.22)$$

then Equation (2.20) holds for $s \geq s_0$. While this is proved by Proposition 2.8. The result follows. \square

Proof of Theorem 1.1 in the general case. By replacing f by some suitable iterate, we have that each irreducible component of V is also an invariant subvariety of f . It is enough to prove the theorem for each irreducible component of V , so we assume that V is irreducible.

If there doesn't exist f_i such that $\deg(f_i) = 1$, for $i \in \{1, \dots, n\}$, then the statement has been proved. Thus we may assume that at least one f_i is an automorphism. We reorder the coordinates so that there exists a positive integer $k \in \{2, \dots, n-1\}$ such that $\deg(f_i) > 1$ when $i \leq k$ and $\deg(f_i) = 1$ when $i > k$. Now Lemma 2.1 implies that $V = V_1 \times V_2$, where $V_1 \subseteq (\mathbb{P}^1)^k$, $V_2 \subseteq (\mathbb{P}^1)^{n-k}$ such that $g_1(V_1) = V_1$ and $g_2(V_2) = V_2$, where $g_1 = (f_1, \dots, f_k)$ and $g_2 = (f_{k+1}, \dots, f_n)$. Notice that $g_2^{-1}(V_2) = V_2$ and there exists a non-negative integer s_0 such that $(g_1^{-s-1}(V_1) \setminus g_1^{-s}(V_1))(K) = \emptyset$ for all $s \geq s_0$, since we have established Theorem 1.1 in the case when the maps all have degree ≥ 2 . Thus, or any non-negative integer $s \geq s_0$,

$$\begin{aligned} & (f^{-s-1}(V) \setminus f^{-s}(V))(K) \\ &= (g_1^{-s-1}(V_1) \setminus g_1^{-s}(V_1))(K) \times (g_2^{-s-1}(V_2) \setminus g_2^{-s}(V_2))(K) = \emptyset. \end{aligned}$$

The result follows.



Acknowledgments

The author thanks the referee for the careful reviewing and helpful suggestions. The author thanks Jason Bell for helpful comments and suggestions and Tom Tucker for inspiring discussions. The author thanks Yohsuke Matzusawa and Kaoru Sano for their helpful comments on an earlier version of the draft and for sharing their recent preprint [MS23] where they proved the same result with $n = 2$ but used a different approach. The author also thanks Junyi Xie for pointing out the reference [Xie23].

References

- [BGT16] BELL, JASON P.; GHIOCA, DRAGOS; TUCKER, THOMAS J. The dynamical Mordell-Lang conjecture, *Mathematical Surveys and Monographs*, vol. 210, American Mathematical Society, Providence, RI, 2016. [MR3468757](#), [Zbl 1362.11001](#), doi: [10.1090/surv/210](#). 634
- [BMS23] BELL, JASON P.; MATSUZAWA, YOHISUKE; SATRIANO, MATTHEW. On Dynamical Cancellation, *Int. Math. Res. Not. IMRN* (2023), no. 8, 7099–7139. [MR4574395](#), [Zbl 1516.37147](#), doi: [10.1093/imrn/rnac058](#). 634, 642, 644, 645
- [Fak03] FAKHRUDDIN, NAJMUDDIN. Questions on self maps of algebraic varieties, *J. Ramanujan Math. Soc.* **18** (2003), no. 2, 109–122. [MR1995861](#), [Zbl 1053.14025](#). 645
- [GNY18] GHIOCA, DRAGOS; NGUYEN, KHOA D.; YE, HEXI. The dynamical Manin-Mumford conjecture and the dynamical Bogomolov conjecture for endomorphisms of $(\mathbb{P}^1)^n$, *Compos. Math.* **154** (2018), no. 7, 1441–1472. [MR3826461](#), [Zbl 1432.37122](#), doi: [10.1112/S0010437X18007157](#). 643, 645
- [GNY19] ———, The dynamical Manin-Mumford conjecture and the dynamical Bogomolov conjecture for split rational maps, *J. Eur. Math. Soc.* **21** (2019), no. 5, 1571–1594. [MR3941498](#), [Zbl 1426.37064](#), doi: [10.4171/JEMS/869](#). 645
- [GTZ11] GHIOCA, DRAGOS; TUCKER, THOMAS J.; ZHANG, SHOUWU. Towards a dynamical Manin-Mumford conjecture, *Int. Math. Res. Not. IMRN* (2011), no. 22, 5109–5122. [MR2854724](#), [Zbl 1267.37110](#), doi: [10.1093/imrn/rnq283](#). 641
- [Hin88] HINDRY, MARC. Autour d’une conjecture de Serge Lang, *Invent. Math.* **94** (1988), no. 3, 575–603. [MR969244](#), [Zbl 0638.14026](#), doi: [10.1007/BF01394276](#). 639
- [MMS23] MATSUZAWA, YOHISUKE; MENG, SHENG; SHIBATA, TAKAHIRO; ZHANG, DE-QI. Non-density of points of small arithmetic degrees, *J. Geom. Anal.* **33** (2023), no. 4, Paper No. 112, 41. [MR4543765](#), [Zbl 1512.37111](#), doi: [10.1007/s12220-022-01156-y](#). 633
- [MS23] MATSUZAWA, YOHISUKE; SANO, KAORU. On preimages question, (2023). [arXiv:math/2311.02906](#) 648
- [Pak23] PAKOVICH, FEDOR. Invariant curves for endomorphisms of $\mathbb{P}^1 \times \mathbb{P}^1$, *Math. Ann.* **385** (2023), no. 1-2, 259–307. [MR4542716](#), [Zbl 1521.37107](#), doi: [10.1007/s00208-021-02304-5](#). 645, 646
- [Ray83] RAYNAUD, MICHEL. Sous-variétés d’une variété abélienne et points de torsion, *Arithmetic and geometry*, Vol. I, Progr. Math., vol. 35, Birkhäuser Boston, Boston, MA, 1983, pp. 327–352. [MR717600](#), [Zbl 0581.14031](#). 641
- [Riv03] RIVERA-LETELIER, JUAN. Dynamique des fonctions rationnelles sur des corps locaux, *Astérisque* (2003), no. 287, xv, 147–230. [MR2040006](#), [Zbl 1140.37336](#). 634
- [Sil07] SILVERMAN, JOSEPH H. The arithmetic of dynamical systems, *Graduate Texts in Mathematics*, vol. 241, Springer, New York, 2007. [MR2316407](#), [Zbl 1130.37001](#). 641

- [Xie23] XIE, JUNYI. Algebraicity criteria, invariant subvarieties and transcendence problems from arithmetic dynamics, *Peking Math. J.* (2023). [MR4711366](#), [Zbl 1532.11099](#), doi: [10.1007/s42543-022-00059-9](#). 635, 648
- [Zho23] ZHONG, XIAO. Dynamical cancellation of polynomials, *Bull. London Math. Soc.* **55** (2023), no. 6, 2948–2962; [MR4698321](#), [Zbl 1537.14037](#), doi: [10.1112/blms.12903](#). 634

(Xiao Zhong) UNIVERSITY OF WATERLOO, DEPARTMENT OF PURE MATHEMATICS, WATERLOO, ONTARIO N2L 3G1, CANADA
x48zhong@uwaterloo.ca

This paper is available via <http://nyjm.albany.edu/j/2025/31-22.html>.