

## THE NEG.-PROPOSITIONAL CALCULUS

Milan Tasić

**Abstract.** Consistency and completeness are proved for an axiomatic system intended to be a formalization of propositional contradictions.

By changing in the truth tables the values of the logical operations of conjunction, disjunction and implication (but not of negation  $\sim$ ), so as to obtain new functions designated by  $\varepsilon$ ,  $v$ , and  $\rightarrow$ , respectively:

$\varepsilon$	$\top$	$\perp$
$\top$	$\perp$	$\top$
$\perp$	$\top$	$\top$

$v$	$\top$	$\perp$
$\top$	$\perp$	$\perp$
$\perp$	$\perp$	$\top$

$\rightarrow$	$\top$	$\perp$
$\top$	$\perp$	$\top$
$\perp$	$\perp$	$\perp$

and by taking an interest in formulas with the truth value  $\perp$ , we may construct an *algebra of contradictions* (identically *false* formulas)  $\mathfrak{A} = \{\langle \top, \perp \rangle; \varepsilon, v, \rightarrow, \sim\}$ , which could serve as the principal model of a propositional calculus called *neg.-propositional calculus*.

This formal system may be axiomatized with the following *axiom-schemata*:

1.  $(A \rightarrow B) \rightarrow A$
2.  $((A \rightarrow B) \rightarrow ((A \rightarrow C) \rightarrow B)) \rightarrow (C \rightarrow B)$
3.  $A \rightarrow \sim(A \varepsilon A)$
4.  $((A \varepsilon B) \rightarrow \sim B) \rightarrow \sim A$
- 5a.  $\sim A \rightarrow A \varepsilon B$
- 5b.  $\sim B \rightarrow A \varepsilon B$
6.  $((\sim C \rightarrow A v B) \rightarrow (B \rightarrow C)) \rightarrow (A \rightarrow C)$
- 7a.  $A v B \rightarrow \sim A$
- 7b.  $A v B \rightarrow \sim B$

$$8. \quad (\sim B \rightarrow (A \rightarrow B)) \rightarrow (\sim A \rightarrow B)$$

and the rule of inference

$$\langle I \rangle \quad \frac{A \rightarrow B, B}{A}$$

We use  $A, B, C, \dots, A_1, \dots$ , as schematic letters for formulas, and  $\Gamma, \Gamma_1$ , as letters for lists of formulas.

We can easily prove the following:

LEMMA 1. (i)  $A \vdash A$ ; (ii) if  $\Gamma \vdash A$ , then  $B, \Gamma \vdash A$ ; (iii) if  $C, \Gamma \vdash A$ , then  $C, \Gamma \vdash A$ ; (iv) if  $\Gamma_1, C, D, \Gamma_2 \vdash A$ , then  $\Gamma_1, D, C, \Gamma_2 \vdash A$ ; (v) if  $\Gamma \vdash A_1, \dots, \Gamma \vdash A_n$  and  $A_1, \dots, A_n \vdash B$ , then  $\Gamma \vdash B$ , or in particular: if  $\vdash A_1, \dots, \vdash A_n$ , and  $A_1, \dots, A_n \vdash B$ , then  $\vdash B$ .

THEOREM 1. If  $B_2, \dots, B_m \vdash A \rightarrow B_1$ , then  $B_1, B_2, \dots, B_m \vdash A$ .

*Proof.* By adding, according to Lemma 1, the formula  $B_1$  to the sequence  $B_2, \dots, B_m$ , by the rule of inference  $\langle I \rangle$  we obtain  $A$ .

COROLLARY 1. If  $\vdash A \rightarrow B$ , then  $B \vdash A$ .

COROLLARY 2. If  $\vdash (\dots (A \rightarrow B_1) \rightarrow \dots) \rightarrow B_m$ , then  $B_1, \dots, B_m \vdash A$ .

THEOREM 2 (Deduction theorem). If  $\Gamma, B \vdash A$ , then  $\Gamma \vdash A \rightarrow B$ .

*Proof.* By induction on the length  $k$  of the given deduction.

1° When  $k = 1$ , the formula  $A$  is either an axiom or one of formulas  $\Gamma$  or  $B$ . In the first case, such a deduction is

- |                                      |                          |
|--------------------------------------|--------------------------|
| 1. $A$                               | axiom-hypothesis         |
| 2. $(A \rightarrow B) \rightarrow A$ | axiom-schema 1           |
| 3. $A \rightarrow B$                 | $\langle I \rangle$ 1, 2 |

and analogously in the remaining two cases.

2° Assume the theorem holds for all lengths  $\leq k$ , and let the length of the given deduction be  $k + 1$ .

Besides the three cases already discussed, an additional case arises here:  $A$  is an immediate consequence, by the rule  $\langle I \rangle$ , of two preceding formulas of the sequence.

Let it be the formulas  $P$  (at the  $r$ -th,  $r \leq k$ ), and  $A \rightarrow P$  (at the  $s$ -th,  $s \leq k$ ) step. In both cases, by the hypothesis, it is possible to construct two deductions of  $P \rightarrow B$  and  $(A \rightarrow P) \rightarrow B$ . Let us add to them the following instance of axiom-schema 2:

$$((A \rightarrow B) \rightarrow ((A \rightarrow P) \rightarrow B)) \rightarrow (P \rightarrow B)$$

Then a double application of the rule of inference yields  $A \rightarrow B$ .

COROLLARY 1. *If  $B \vdash A$ , then  $\vdash A \rightarrow B$ .*

COROLLARY 2. *If  $B_1, \dots, B_{m-1}, B_m \vdash A$ , then  $\vdash (\dots(A \rightarrow B_1) \rightarrow \dots \rightarrow B_{m-1}) \rightarrow B_m$ .*

We derive the introduction and the elimination rules for connectives in the theorem after the following lemma:

LEMMA 1. (a) *If  $B \vdash A$  and  $B \vdash \sim A$ , then  $\vdash \sim B$ ; (b) if  $B \vdash A$ , then  $\sim A \vdash \sim B$ ; (c1)  $\sim \sim A \vdash A$ ; (c2)  $A \vdash \sim \sim A$ .*

THEOREM 3.

	<i>Introduction</i>	<i>Elimination</i>
$\rightarrow$	<i>If <math>\Gamma, B \vdash A</math>, then <math>\Gamma \vdash A \rightarrow B</math></i>	<i><math>A \rightarrow B, B \vdash A</math></i>
$\varepsilon$	<i><math>\sim A, \sim B \vdash A \varepsilon B</math></i>	<i><math>A \varepsilon B \vdash \sim A</math> <math>A \varepsilon B \vdash \sim B</math></i>
$\vee$	<i><math>\sim A \vdash A \vee B</math> <math>\sim B \vdash A \vee B</math></i>	<i>If <math>\Gamma, C \vdash A</math> and <math>\Gamma, C \vdash B</math>, then <math>\Gamma, A \vee B \vdash \sim C</math></i>
$\sim$	<i>If <math>\Gamma, B \vdash A</math> and <math>\Gamma, B \rightarrow \sim A</math>, then <math>\Gamma \vdash \sim B</math></i>	<i><math>\sim \sim A \vdash A</math></i>

*Proof.* The rule of  $\rightarrow$ -introduction is just the deduction theorem, and  $\rightarrow$ -elimination is our rule of inference  $\langle I \rangle$ . For the remaining rules we use the axiom-schema 4 for  $\varepsilon$  introduction, 5a and 5b for  $\varepsilon$ -elimination, 7a and 7b for  $\vee$ -introduction, 6 for  $\vee$ -elimination, Lemma 2 (a) for  $\sim$ -introduction, and finally Lemma 2 (c1) for  $\sim$ -elimination.

Can this formal system describe an intuitive domain of bivalent propositions? To answer this question, we define as follows the consistency and completeness of a formal system:

*Definition 1.* (i) The neg.-propositional calculus is *semantically consistent* if every provable formula, interpreted on the model of the calculus — the algebra  $\mathfrak{A}$  — is a contradiction.

(ii) The neg.-propositional calculus is *simply consistent* if there is an  $A$  such that  $\vdash A$  and  $\vdash \sim A$ .

(iii) The neg.-propositional calculus is *syntactically consistent* if a formula is unprovable in it.

THEOREM 4. *The neg.-propositional calculus is (semantically, simple, syntactically) consistent.*

*Proof.* On the indicated model, the truth value of all axioms is  $\perp$  and the rule of inference keeps this property (semantical consistency). The value  $\perp$  can not at the same time belong to a formula and to its negation (simple consistency), and, so, as an example of an unprovable formula we have  $A \varepsilon \sim A$  (syntactical consistency).

*Definition 2.* (i) The neg.-propositional calculus is *semantically complete* if every contradiction is provable in it.

(ii) The neg.-propositional calculus is *simply complete* if for every  $A$  either  $\vdash A$  or  $\vdash \sim A$ .

(iii) The neg.-propositional calculus is *syntactically complete* if it has the following property: if one adds to the axioms an unprovable formula, one can prove every formula.

Completeness in the first sense implies completeness in other two senses. Then we prove the following:

**THEOREM 5.** *The neg.-propositional calculus is semantically complete.*

Let us assign some  $n$ -tuple of values  $\top, \perp$ , to the propositional letters  $A_1, \dots, A_n$  that occur in the formula  $F$ . Then by  $B_1, \dots, B_n$  we denote a corrected  $n$ -tuple, where  $B_i$  ( $1 \leq i \leq n$ ) is  $A_i$  or  $\sim A_i$ , according as the assigned value is  $\perp$  or  $\top$ .

**LEMMA 3.**  $B_1, \dots, B_n \vdash F$  if  $\tau(F) = \perp$  and  $B_1, \dots, B_n \vdash \sim F$  if  $\tau(F) = \top$ .

*Proof.* Let us denote by  $d$  the degree of the formula  $F$ , i.e. the number of connectives in  $F$ .

*Basis* ( $d = 0$ ). Since  $F$  is a propositional letter  $A_i$ , for  $i \in \{1, \dots, n\}$ , we have  $A_i \vdash A_i$  if  $\tau(A_i) = \perp$ , or  $\sim A_i \vdash \sim A_i$  if  $\tau(A_i) = \top$ .

*Induction step.* If the degree of  $F$  is  $k + 1$ , then  $F$  has one of the forms: (a)  $\sim A$ , (b)  $A \rightarrow B$ , (c)  $A \vee B$  or (d)  $A \varepsilon B$  with  $A, B$  of degree  $\leq n$ .

(a) We must show: (a1)  $B_1, \dots, B_n \vdash \sim F$  if  $B_1, \dots, B_n \vdash A$  and (a2)  $B_1, \dots, B_n \vdash F$  if  $B_1, \dots, B_n \vdash \sim A$ , or (a1)  $A \vdash \sim \sim A$  and (a2)  $\sim A \vdash \sim A$ . The proof is carried out by Lemma 1.

(b) If  $F$  has the form  $A \rightarrow B$ , four subcases arise:

(b1) If  $B_1, \dots, B_n \vdash \sim A$  and  $B_1, \dots, B_n \vdash \sim B$ , then  $B_1, \dots, B_n \vdash A \rightarrow B$  (and by Lemma 1, it is sufficient to prove  $\sim A, \sim B \vdash A \rightarrow B$ ).

(b2) If  $B_1, \dots, B_n \vdash \sim A$  and  $B_1, \dots, B_n \vdash B$ , then  $B_1, \dots, B_n \vdash \sim(A \rightarrow B)$  ( $\sim A, B \vdash \sim(A \rightarrow B)$ ).

(b3) If  $B_1, \dots, B_n \vdash A$  and  $B_1, \dots, B_n \vdash \sim B$ , then  $B_1, \dots, B_n \vdash A \rightarrow B$  ( $A, \sim B \vdash A \rightarrow B$ );

(b4) If  $B_1, \dots, B_n \vdash A$  and  $B_1, \dots, B_n \vdash B$ , then  $B_1, \dots, B_n \vdash A \rightarrow B$  ( $A, B \vdash A \rightarrow B$ );

**SUBLEMMA 1.**  $A, \sim A \vdash B$ .

- |      |   |                      |
|------|---|----------------------|
| (b1) | 1. $B, \sim B \vdash A$                       | Sublemma 1           |
|      | 2. $\sim B \vdash A \rightarrow B$            | $\rightarrow$ -in. 1 |
|      | 3. $\sim A, \sim B \vdash A \rightarrow B$    | Lemma 1, 2           |
| (b2) | 1. $A \rightarrow B, B \vdash A$              | $\rightarrow$ -el.   |
|      | 2. $A \rightarrow B, B \vdash B$              | Lemma 1              |
|      | 3. $A \vee B, B \vdash \sim(A \rightarrow B)$ | $\vee$ -el. 1, 2     |
|      | 4. $\sim A \vdash A \vee B$                   | $\vee$ -in.          |
|      | 5. $\sim A, B \vdash A \vee B$                | Lemma 1, 4           |
|      | 6. $\sim A, B \vdash B$                       | Lemma 1              |
|      | 7. $\sim A, B \vdash \sim(A \rightarrow B)$   | Lemma 1, 3, 5, 6     |

For (b3) and (b4) we proceed similarly.

For (c1), (c2), (c3) we use  $\vee$ -in.

- |      |   |                    |
|------|---|--------------------|
| (c4) | 1. $A \rightarrow B, B \vdash A$              | $\rightarrow$ -el. |
|      | 2. $A \rightarrow B, B \vdash B$              | Lemma 1            |
|      | 3. $B, A \vee B \vdash \sim(A \vee B)$        | $\vee$ -el. 1, 2   |
|      | 4. $B, A \rightarrow B \vdash \sim(A \vee B)$ | Lemma 2, 3         |
|      | 5. $A, B \vdash B$                            | Lemma 1            |
|      | 6. $A, B \vdash A \rightarrow B$              | (b4)               |
|      | 7. $A, B \vdash \sim(A \vee B)$               | Lemma 1, 5, 6, 3   |

For (d1) we use  $\varepsilon$ -in.

SUBLEMMA 2.  $\sim A \vee \sim B \vdash \sim(A \varepsilon B)$ .

(In the proof we use  $\varepsilon$ -el. and  $\vee$ -el.)

- |      |  |                             |
|------|--|-----------------------------|
| (d2) | 1. $\sim B \vdash A \vee B$                          | $\vee$ -in.                 |
|      | 2. $B \vdash \sim A \vee \sim B$                     | $A \vdash A, B \vdash B, 1$ |
|      | 3. $\sim A \vee \sim B \vdash \sim(A \varepsilon B)$ | Sublemma 2                  |
|      | 4. $B \vdash \sim(A \varepsilon B)$                  | Lemma 1, 2, 3               |
|      | 5. $\sim A, B \vdash \sim(A \varepsilon B)$          | Lemma 1, 4                  |

Case (d3) is similar to (d2). For (d4) we use (d3).

LEMMA 4. *If for any of  $2^n$  possible sequences of values we always have  $B_1, \dots, B_n \vdash F$ , then*

$$A_1 \vee \sim A_1, \dots, A_n \vee \sim A_n \vdash F.$$

*Proof.* Let us take  $n = 2$ . By hypothesis,  $\sim A_1, \sim A_2 \vdash F$ ;  $\sim A_1, A_2 \vdash F$ ;  $A_1, \sim A_2 \vdash F$  and  $A_1, A_2 \vdash F$ . So, by a triple application of  $v$ -elimination (using Lemma 2), we obtain:

- |   |                |
|---|----------------|
| 1. $\sim A_1, \sim A_2 \vdash F$                      | hyp.           |
| 2. $\sim A_1, A_2 \vdash F$                           | hyp.           |
| 3. $\sim A_1, \sim F \vdash \sim \sim A_1 \vdash A_1$ | Lemma 2, 1     |
| 4. $\sim A_1, \sim F \vdash \sim A_2$                 | Lemma 2, 2     |
| 5. $\sim A_1, A_2 \vee \sim A_2 \vdash F$             | $v$ -el. 3, 4  |
| 6. $A_1, \sim A_2 \vdash F$                           | hyp.           |
| 7. $A_1, A_2 \vdash F$                                | hyp.           |
| 8. $A_1, \sim F \vdash A_2$                           | Lemma 2, 6     |
| 9. $A_1, \sim F \vdash \sim A_2$                      | Lemma 2, 7     |
| 10. $A_1, A_2 \vee \sim A_2 \vdash F$                 | $v$ -el. 8, 9  |
| 11. $A_1 \vee \sim A_1, A_2 \vee \sim A_2 \vdash F$   | $v$ -el. 5, 10 |

*Proof of Theorem 5.* Let  $F$  be a contradiction and  $A_1, \dots, A_n$  all of its letters. By the last lemma:

$$A_1 \vee \sim A_1, \dots, A_n \vee \sim A_n \vdash F$$

and, since we have  $\vdash A \vee \sim A$  by Lemma 1, we should finally have  $\vdash F$ .

This proof follows the method exposed in [1].

#### REFERENCES

- [1] S. C. Kleene, *Introduction to Metamathematics*, North-Holland, Amsterdam, 1952.

Matematički Institut  
Univerzitet u Skoplju  
91000 Skoplje  
Jugoslavija

(Received 09 03 1990)  
(Revised 24 06 1991)