

**MORE EXAMPLES AND COUNTEREXAMPLES  
FOR A CONJECTURE  
OF MERRIFIELD AND SIMMONS**

**Yong Wang, Xuelian Li, and Ivan Gutman**

*Communicated by Slobodan Simić*

ABSTRACT. Let  $\sigma(G)$  be the number of independent-vertex sets of a graph  $G$ . Merrifield and Simmons conjectured that for any connected graph  $G$  and any pair of its non-adjacent vertices  $u$  and  $v$ ,  $\Delta_{uv}(G) := \sigma(G - u)\sigma(G - v) - \sigma(G)\sigma(G - u - v)$  is positive if the distance between  $u$  and  $v$  is odd, and negative otherwise. In earlier works by the authors the conjecture was shown to be true for trees, cycles and several other types of graphs, but a few counterexamples were discovered among dense graphs. We now prove that the conjecture is true for all bipartite and some non-bipartite connected unicyclic graphs, but not for all connected unicyclic graphs. Moreover, we find a graph  $G$  for which  $\Delta_{uv}(G) = 0$ .

**1. Introduction**

Two vertices of a graph  $G$  are said to be independent if they are not adjacent. The  $k$ -th independence number of  $G$  is denoted by  $\sigma_k(G)$ . By definition, for  $k \geq 2$ ,  $\sigma_k(G)$  is equal to the number of ways in which  $k$  pairwise independent vertices can be selected in the graph  $G$ . In addition to this,  $\sigma_0(G) = 1$  and  $\sigma_1(G) =$  number of vertices of  $G$ . Thus  $\sigma(G) = \sum_{k \geq 0} \sigma_k(G)$  is the number of all independent-vertex sets of  $G$ .

On page 144 of the book [5], Merrifield and Simmons claimed without proof a property of  $\sigma(G)$ , which for non-adjacent vertices  $u$  and  $v$  of a connected graph  $G$  may be formulated as follows. Abbreviate  $\sigma(G - u)\sigma(G - v) - \sigma(G)\sigma(G - u - v)$  by  $\Delta_{uv}(G)$ . Then

$$(1) \quad \Delta_{uv}(G) \begin{cases} < 0 & \text{if } d(u, v) \text{ is even} \\ > 0 & \text{if } d(u, v) \text{ is odd} \end{cases}$$

where  $d(u, v)$  denotes the distance of  $u$  and  $v$  in  $G$ .

Gutman [1] proved that the conjecture is true for all trees. Li [4] proved that the conjecture is true for all cycles and for many other kinds of graphs. Although Li [4] found two families of counterexamples for this conjecture, the respective graphs were very dense, i.e., had relatively many edges. On the other hand, the considerations in the book [5] were limited to sparse graphs, i.e., graphs having relatively few edges. Therefore, it may be understood that the property (1) was believed to hold only for sparse graphs. If so, then the counterexamples reported in [4] cannot be considered as fully satisfactory. What one would need to convincingly disprove the validity of the Merrifield–Simmons conjecture are counterexamples based on sparse graphs. In order to accomplish this task, we examined (connected) unicyclic graphs, which, apart from trees, have the least number of edges. We show that although the majority of connected unicyclic graphs satisfies the relation (1), counterexamples for the Merrifield–Simmons conjecture exist already within this class.

## 2. Preliminary considerations

We first state without proof some properties and lemmas, which are either given in [1–4] or are immediate consequences of results found in [1–4].

PROPERTY 1. If  $v$  is a vertex of  $G$ , then

$$\sigma(G) = \sigma(G - v) + \sigma(G - [v])$$

where  $G - v$  is the subgraph obtained by deleting  $v$  from  $G$ , and  $G - [v]$  is the subgraph obtained by deleting from  $G$  both  $v$  and the vertices adjacent to  $v$ .

PROPERTY 2. Let  $G_1 \cup G_2$  denote a graph composed of disjoint components  $G_1$  and  $G_2$ . Then

$$\sigma(G_1 \cup G_2) = \sigma(G_1) \sigma(G_2)$$

An immediate consequence of Property 2 is that if the graph  $G$  is disconnected and the vertices  $u$  and  $v$  belong to its different components, then  $\Delta_{uv}(G) = 0$ . If, in turn,  $G = G_1 \cup G_2$  and both  $u$  and  $v$  belong to  $G_1$ , then  $\Delta_{uv}(G) = \sigma^2(G_2) \Delta_{uv}(G_1)$ . Therefore, without loss of generality the examination of the quantity  $\Delta_{uv}(G)$  may be restricted to connected graphs.

In what follows, if it is not explicitly stated otherwise, it is assumed that the graph considered is connected.

PROPERTY 3. Let  $P_n$  be the path on  $n$  vertices. From Property 1 follows that

$$\sigma(P_n) = \sigma(P_{n-1}) + \sigma(P_{n-2})$$

with  $\sigma(P_0) = 1$  and  $\sigma(P_1) = 2$ . Hence the  $\sigma$ -values of the path-graphs are just the Fibonacci numbers.

Label the vertices of  $P_n$  by  $v_1, v_2, \dots, v_n$  so that  $v_i$  and  $v_{i+1}$  are adjacent,  $i = 1, \dots, n - 1$ . Let  $R_1, R_2, \dots, R_n$  be distinct rooted graphs with mutually disjoint vertex sets. Then the compound graph  $P_n(R_1, \dots, R_n)$  is obtained by identifying the root  $r_i$  of  $R_i$  with the vertex  $v_i$  of  $P_n$  for  $i = 1, 2, \dots, n$ .

LEMMA 1. *If  $n \geq 2$ , then*

$$\Delta_{v_1, v_n}(P_n(R_1, \dots, R_n)) = (-1)^n \prod_{i=1}^n \sigma(R_i - r_i) \sigma(R_i - [r_i])$$

In order to proceed we need to fix our notation and terminology. Let  $U_n$  be a (connected) unicyclic graph, such that its (unique) cycle possess  $n$  vertices. Denote this cycle by  $C_n$  and label its vertices consecutively by  $a_1, a_2, \dots, a_n$ . We call the subtree of  $U_n$  attached to the vertex  $a_i$  the  $a_i$ -tree; the vertex  $a_i$  belongs to the  $a_i$ -tree.

Denote by  $T_{a_i}$  the forest obtained by deleting  $a_i$  from the  $a_i$ -tree, and by  $T_{i,n}$  the subgraph obtained by deleting from  $U_n$  the union of the  $a_1$ -tree,  $a_2$ -tree,  $\dots$ ,  $a_{i-1}$ -tree. Then we have

$$U_n - a_1 = T_{a_1} \cup T_{2,n}$$

Denote by  $T_{[a_i]}$  the forest obtained by deleting from the  $a_i$ -tree both the vertex  $a_i$  and the vertices adjacent to it. Then we have

$$U_n - [a_1] = T_{[a_1]} \cup T_{3,n-1} \cup T_{a_2} \cup T_{a_n}$$

Note that  $T_{3,n-1}$  is the subgraph obtained by deleting from  $U_{n-1}$  the union of the  $a_1$ - and  $a_2$ -trees. In harmony with the above specified notation, the graph  $U_{n-1}$  is unicyclic, containing a cycle on  $n-1$  vertices. The vertices of this cycle are  $a_1, \dots, a_{n-1}$ , and the subtrees attached to them are the same as in the case of  $U_n$ .

If  $u$  is a vertex of the  $a_i$ -tree and  $d(u, a_i) \geq 2$ , then we denote by  $\bar{a}_i$  the vertex adjacent to  $a_i$  in the unique path of the  $a_i$ -tree, connecting  $u$  and  $a_i$ . The connected component of  $T_{a_i}$  containing the vertex  $u$  is denoted by  $T_u$ .

LEMMA 2. *Let  $U$  be a unicyclic graph,  $u$  a vertex of the  $a_i$ -tree, and  $d(u, a_i) \geq 2$ , then*

$$\sigma(T_{a_i} - u) \sigma(T_{[a_i]}) - \sigma(T_{a_i}) \sigma(T_{[a_i]} - u) \begin{cases} > 0 & \text{if } d(u, a_i) \text{ is even} \\ < 0 & \text{if } d(u, a_i) \text{ is odd} \end{cases}$$

PROOF. By Properties 1 and 2,

$$\begin{aligned} & \sigma(T_{a_i} - u) \sigma(T_{[a_i]}) - \sigma(T_{a_i}) \sigma(T_{[a_i]} - u) \\ &= \sigma(T_u - u) \sigma(T_{a_i} - T_u) \sigma(T_u - \bar{a}_i) \sigma(T_{[a_i]} - (T_u - \bar{a}_i)) \\ & \quad - \sigma(T_u) \sigma(T_{a_i} - T_u) \sigma(T_u - \bar{a}_i - u) \sigma(T_{[a_i]} - (T_u - \bar{a}_i)) \\ &= \sigma(T_{a_i} - T_u) \sigma(T_{[a_i]} - (T_u - \bar{a}_i)) [\sigma(T_u - u) \sigma(T_u - \bar{a}_i) - \sigma(T_u) \sigma(T_u - \bar{a}_i - u)] \end{aligned}$$

Because  $T_u$  is a tree, by Lemma 1 we have

$$\sigma(T_u - u) \sigma(T_u - \bar{a}_i) - \sigma(T_u) \sigma(T_u - \bar{a}_i - u) \begin{cases} < 0 & \text{if } d(u, \bar{a}_i) \text{ is even} \\ > 0 & \text{if } d(u, \bar{a}_i) \text{ is odd} \end{cases}$$

Since  $\sigma(T_{a_i} - T_u) \sigma(T_{[a_i]} - (T_u - \bar{a}_i)) > 0$ , we have

$$\sigma(T_{a_i} - u) \sigma(T_{[a_i]}) - \sigma(T_{a_i}) \sigma(T_{[a_i]} - u) \begin{cases} > 0 & \text{if } d(u, a_i) = d(u, \bar{a}_i) + 1 \text{ is even} \\ < 0 & \text{if } d(u, a_i) = d(u, \bar{a}_i) + 1 \text{ is odd} \end{cases}$$

The proof is complete.  $\square$

LEMMA 3. Assume that  $T$  is a tree, and  $u$  and  $v$  are distinct vertices of  $T$ . Let  $P$  be the unique path connecting  $u$  and  $v$ . Then,

$$\Delta_{uv}(T) = (-1)^{d(u,v)} \sigma(T - P) \sigma(T - [P])$$

where  $T - [P]$  stands for the subgraph obtained by deleting from  $T$  the vertices of  $P$  and their first neighbors.

### 3. The main results

We start with two counterexamples.

COUNTEREXAMPLE 1. Let  $G_a$  be the graph depicted in Figure 1. This graph is connected and the distance between its vertices  $u$  and  $v$  is 4. By direct calculation we check that  $\sigma(G_a - u) = 92$ ,  $\sigma(G_a - v) = 114$ ,  $\sigma(G_a) = 152$ ,  $\sigma(G_a - u - v) = 69$ , and thus  $\Delta_{uv}(G_a) = \sigma(G_a - u) \sigma(G_a - v) - \sigma(G_a) \sigma(G_a - u - v) = 0$ . It seems that the possibility that for a connected graph  $\Delta_{uv}(G)$  may be equal to zero was not anticipated by Merrifield and Simmons [5].




Figure 1

COUNTEREXAMPLE 2. Let  $G_b$  be the graph shown in Figure 1. Here  $d(u, v) = 3$ , an odd number. It is not difficult to see that  $\sigma(G_b - u) = 35$ ,  $\sigma(G_b - v) = 38$ ,  $\sigma(G_b) = 58$ ,  $\sigma(G_b - u - v) = 23$ , and thus  $\Delta_{uv}(G_b) = -4 < 0$ , contradicting to relation (1). This example shows that the Merrifield–Simmons conjecture is not generally valid already in the case of sparse graphs.

Our first theorem gives a family of counterexamples for the conjecture, including Counterexample 2 as a special case.

Let  $U_n$  be a unicyclic graph with a cycle  $C_n$ , such that  $e_i$  pendant vertices are attached to the vertex  $a_i$  of  $C_n$ ,  $i = 1, 2, \dots, n$ . In other words, the subtree of  $U_n$  attached to  $a_i$  is an  $(e_i + 1)$ -vertex star.

Let  $u$  be a vertex of  $U_n$ , belonging to the  $a_1$ -tree,  $u \neq a_1$ , and let  $v = a_3$ . Therefore  $d(u, v) = 3$ .

THEOREM 1. *If  $n$  is odd,  $n \geq 5$ , if  $e_3 = e_4 = \dots = e_n = 0$ ;  $e_1, e_2 > 0$  and if*

$$(2) \quad 1 + \sum_{i=1}^{n-4} \sigma(P_i) \sigma(P_{i-1}) < 2^{e_2}$$

*then the above described graph  $U_n$  is a counterexample for the Merrifield–Simmons conjecture.*

The structure of the graphs for which Theorem 1 applies is shown in Figure 2.




Figure 2

PROOF. Consider first the general case, when  $e_i \geq 0$ ,  $i = 1, 2, \dots, n$ . Denote by  $P$  the shortest path of  $U_n$  connecting  $a_2$  and  $v$ . Then by Properties 1 and 2 and Lemma 3,

$$\begin{aligned} \Delta_{uv}(U_n) &= \sigma(U_n - u) \sigma(U_n - v) - \sigma(U_n) \sigma(U_n - u - v) \\ &= [\sigma(U_n - a_1 - u) + \sigma(U_n - [a_1] - u)] [\sigma(U_n - a_1 - v) + \sigma(U_n - [a_1] - v)] \\ &\quad - [\sigma(U_n - a_1) + \sigma(U_n - [a_1])] [\sigma(U_n - a_1 - u - v) + \sigma(U_n - [a_1] - u - v)] \\ &= \Delta_{uv}(U_n - a_1) + \Delta_{uv}(U_n - [a_1]) + [\sigma(U_n - a_1 - u) \sigma(U_n - [a_1] - v) \\ &\quad + \sigma(U_n - a_1 - v) \sigma(U_n - [a_1] - u) - \sigma(U_n - a_1) \sigma(U_n - [a_1] - u - v) \\ &\quad - \sigma(U_n - [a_1]) \sigma(U_n - a_1 - u - v)] \\ &= 0 + 0 + 2^{e_1 + e_n - 1} [\sigma(T_{2,n-1} - a_2) \sigma(T_{2,n} - v) - \sigma(T_{2,n}) \sigma(T_{2,n-1} - a_2 - v)] \\ &= 2^{e_1 + e_n - 1} [2^{e_n} [\sigma(T_{2,n-1} - v) \sigma(T_{2,n-1} - a_2) - \sigma(T_{2,n-1}) \sigma(T_{2,n-1} - a_2 - v)] \\ &\quad + 2^{e_n - 1} [\sigma(T_{2,n-2} - v) \sigma(T_{2,n-1} - a_2) - \sigma(T_{2,n-2}) \sigma(T_{2,n-1} - a_2 - v)]] \\ &= 2^{e_1 + e_n - 1} [2^{e_n} \sigma(T_{2,n-1} - P) \sigma(T_{2,n-1} - [P]) \\ &\quad + 2^{e_n - 1} [\sigma(T_{2,n-2} - v) \sigma(T_{2,n-1} - a_2) - \sigma(T_{2,n-2}) \sigma(T_{2,n-1} - a_2 - v)]] \\ &= 2^{e_1 + e_n - 1} [2^{e_n} \sigma(T_{2,n-1} - P) \sigma(T_{2,n-1} - [P]) \\ &\quad + 2^{2e_n - 1} \sigma(T_{2,n-2} - P) \sigma(T_{2,n-2} - [P]) \\ &\quad + 2^{e_n - 1 + 2e_n - 2} \sigma(T_{2,n-3} - P) \sigma(T_{2,n-3} - [P]) \\ &\quad + \dots + 2^{e_n - 1 + e_n - 2 + \dots + e_6 + 2e_5} \sigma(T_{2,4} - P) \sigma(T_{2,4} - [P]) \end{aligned}$$

$$\begin{aligned}
& + 2^{e_{n-1}+e_{n-2}+\dots+e_4} [\sigma(T_{2,3} - v) \sigma(T_{2,4} - a_2) - \sigma(T_{2,3}) \sigma(T_{2,4} - a_2 - v)] \\
= & 2^{e_1+e_n-1} [2^{e_n} \sigma(T_{4,n-1}) \sigma(T_{5,n-1}) 2^{e_2+e_3+e_4} \\
& + 2^{2e_{n-1}} \sigma(T_{4,n-2}) \sigma(T_{5,n-2}) 2^{e_2+e_3+e_4} \\
& + 2^{e_{n-1}+2e_{n-2}} \sigma(T_{4,n-3}) \sigma(T_{5,n-3}) 2^{e_2+e_3+e_4} \\
& + \dots + 2^{e_{n-1}+e_{n-2}+\dots+e_6+2e_5+e_4+e_3+e_2} (2^{e_4} + 1) \\
& + 2^{e_{n-1}+e_{n-2}+\dots+e_5+2e_4+e_3+e_2} - 2^{e_{n-1}+e_{n-2}+\dots+e_3+2e_2}]
\end{aligned}$$

Now, in our case  $e_n = e_{n-1} = \dots = e_3 = 0$ , and therefore

$$\begin{aligned}
\Delta_{uv}(U_n) = & 2^{e_2+e_1-1} [\sigma(P_{n-4}) \sigma(P_{n-5}) + \sigma(P_{n-5}) \sigma(P_{n-6}) + \dots + \sigma(P_2) \sigma(P_1) \\
& + \sigma(P_1) \sigma(P_0) + 1 - 2^{e_2}]
\end{aligned}$$

It is now evident that  $\Delta_{uv}(U_n)$  will be negative-valued whenever the condition (2) is obeyed; this can be achieved by choosing  $e_2$  sufficiently large. On the other hand,  $d(u, v) = 3$  is odd, which contradicts to (1).  $\square$

The next two theorems show that, nevertheless, many unicyclic graphs obey the Merrifield–Simmons conjecture.

**THEOREM 2.** *Let  $U_n$  be the previously described (connected) unicyclic graph. Let the parameter  $n$  be even, i.e.,  $U_n$  is bipartite. Let  $u$  and  $v$  be distinct vertices of  $U_n$ . Then relations (1) are satisfied.*

**PROOF.** have to distinguish between the following three cases.

**Case 1.**  $u$  and  $v$  belong to the same tree.

**Case 2.**  $u$  and  $v$  belong to different trees and at least one of them is not in  $C_n$ .

**Case 3.**  $u$  and  $v$  belong to  $C_n$ .

**Case 1.** Without loss of generality we may assume that  $u$  and  $v$  belong to the  $a_1$ -tree. From Lemma 1 we know that

$$\Delta_{uv}(U_n) = (-1)^{d(u,v)+1} \prod_{i=1}^n \sigma(R_i - r_i) \sigma(R_i - [r_i]) \begin{cases} < 0 & \text{if } d(u, v) \text{ is even} \\ > 0 & \text{if } d(u, v) \text{ is odd} \end{cases}$$

Therefore, Theorem 2 is true in the Case 1.

**Case 2** needs to be divided into two subcases:

**Subcase 2.1.**  $u$  belongs to the  $a_1$ -tree,  $u \neq a_1$  and  $v = a_i$ ;

**Subcase 2.2.**  $u$  belongs to the  $a_1$ -tree,  $u \neq a_1$  and  $v$  is in the  $a_i$ -tree,  $v \neq a_i$ .

Both subcases need to be further divided into:

**Subcase 2.1.1.**  $u$  belongs to the  $a_1$ -tree,  $d(u, a_1) \geq 2$  and  $v = a_i$  ( $3 \leq i \leq n/2+1$ );

**Subcase 2.1.2.**  $u$  belongs to the  $a_1$ -tree,  $v$  is in the unique cycle, and  $v = a_2$ ,  $d(u, a_1) \geq 2$ ;

**Subcase 2.1.3.**  $u$  belongs to the  $a_1$ -tree and  $v = a_2$ ,  $d(u, a_1) = 1$ ,  $d(u, v) = 2$ ;

**Subcase 2.1.4.**  $u$  belongs to the  $a_1$ -tree,  $v$  is in the unique cycle and  $d(u, a_1) = 1$ ,  $v = a_i$ ,  $3 \leq i \leq n/2 + 1$ ;

**Subcase 2.2.1.**  $u$  belongs to the  $a_1$ -tree,  $d(u, a_1) \geq 2$  and  $v$  belongs to the  $a_i$ -tree,

$3 \leq i \leq n/2 + 1$ ,  $v \neq a_i$ ;

**Subcase 2.2.2.**  $u$  belongs to the  $a_1$ -tree,  $d(u, a_1) \geq 2$  and  $v$  belongs to the  $a_2$ -tree,  $d(v, a_2) \geq 2$ ;

**Subcase 2.2.3.**  $u$  belongs to the  $a_1$ -tree,  $d(u, a_1) = 1$  and  $v$  belongs to the  $a_2$ -tree,  $d(v, a_2) \geq 2$ ;

**Subcase 2.2.4.**  $u$  belongs to the  $a_1$ -tree,  $d(u, a_1) = 1$  and  $v$  belongs to the  $a_2$ -tree,  $d(v, a_2) = 1$ .

*Proof of Subcase 2.1.1.* Under the conditions specified in Subcase 2.1.1 we have

$$\begin{aligned}
\Delta_{uv}(U_n) &= \sigma(U_n - u) \sigma(U_n - v) - \sigma(U_n) \sigma(U_n - u - v) \\
&= [\sigma(U_n - a_1 - u) + \sigma(U_n - [a_1] - u)][\sigma(U_n - a_1 - v) + \sigma(U_n - [a_1] - v)] \\
&\quad - [\sigma(U_n - a_1) + \sigma(U_n - [a_1])][\sigma(U_n - a_1 - u - v) + \sigma(U_n - [a_1] - u - v)] \\
&= \Delta_{uv}(U_n - a_1) + \Delta_{uv}(U_n - [a_1]) + [\sigma(U_n - a_1 - u) \sigma(U_n - [a_1] - v) \\
&\quad + \sigma(U_n - a_1 - v) \sigma(U_n - [a_1] - u) - \sigma(U_n - a_1) \sigma(U_n - [a_1] - u - v) \\
&\quad - \sigma(U_n - [a_1]) \sigma(U_n - a_1 - u - v)] \\
&= [\sigma(T_{a_1} - u) \sigma(T_{2,n}) \sigma(T_{a_1}) \sigma(T_{2,n} - v) \\
&\quad - \sigma(T_{a_1}) \sigma(T_{2,n}) \sigma(T_{a_1} - u) \sigma(T_{2,n} - v)] \\
&\quad + [\sigma(T_{[a_1]} - u) \sigma(T_{3,n-1}) \sigma(T_{a_2}) \sigma(T_{a_n}) \sigma(T_{[a_1]}) \sigma(T_{3,n-1}) \sigma(T_{a_2}) \sigma(T_{a_n}) \\
&\quad - \sigma(T_{[a_1]}) \sigma(T_{3,n-1}) \sigma(T_{a_2}) \sigma(T_{a_n}) \sigma(T_{[a_1]} - u) \sigma(T_{3,n-1} - v) \sigma(T_{a_2}) \sigma(T_{a_n})] \\
&\quad + [\sigma(T_{a_1} - u) \sigma(T_{2,n}) \sigma(T_{[a_1]}) \sigma(T_{3,n-1} - v) \sigma(T_{a_2}) \sigma(T_{a_n}) \\
&\quad + \sigma(T_{[a_1]} - u) \sigma(T_{3,n-1}) \sigma(T_{a_2}) \sigma(T_{a_n}) \sigma(T_{a_1}) \sigma(T_{2,n} - v) \\
&\quad - \sigma(T_{a_1}) \sigma(T_{2,n}) \sigma(T_{[a_1]} - u) \sigma(T_{3,n-1} - v) \sigma(T_{a_2}) \sigma(T_{a_n}) \\
&\quad - \sigma(T_{[a_1]}) \sigma(T_{3,n-1}) \sigma(T_{a_2}) \sigma(T_{a_n}) \sigma(T_{a_1} - u) \sigma(T_{2,n} - v)] \\
&= 0 + 0 + \sigma(T_{a_2}) \sigma(T_{a_n}) \sigma(T_{2,n}) \sigma(T_{3,n-1} - v) \\
&\quad \times [\sigma(T_{a_1} - u) \sigma(T_{a_1}) - \sigma(T_{a_1}) \sigma(T_{[a_1]} - u)] \\
&\quad + \sigma(T_{a_2}) \sigma(T_{a_n}) \sigma(T_{3,n-1}) \sigma(T_{2,n} - v) \\
&\quad \times [\sigma(T_{[a_1]} - u) \sigma(T_{a_1}) - \sigma(T_{[a_1]}) \sigma(T_{a_1} - u)] \\
&= \sigma(T_{a_2}) \sigma(T_{a_n}) [\sigma(T_{2,n}) \sigma(T_{3,n-1} - v) - \sigma(T_{3,n-1}) \sigma(T_{2,n} - v)] \\
&\quad \times [\sigma(T_{a_1} - u) \sigma(T_{[a_1]}) - \sigma(T_{a_1}) \sigma(T_{[a_1]} - u)] \\
&= \sigma(T_{a_2}) [\sigma(T_{2,n}) \sigma(T_{3,n} - a_n - v) - \sigma(T_{3,n} - a_n) \sigma(T_{2,n} - v)] \\
&\quad \times [\sigma(T_{a_1} - u) \sigma(T_{[a_1]}) - \sigma(T_{a_1}) \sigma(T_{[a_1]} - u)] \\
&= \sigma(T_{a_2}) X_1 X_2
\end{aligned}$$

where

$$\begin{aligned}
X_1 &= \sigma(T_{2,n}) \sigma(T_{3,n} - a_n - v) - \sigma(T_{3,n} - a_n) \sigma(T_{2,n} - v) \\
X_2 &= \sigma(T_{a_1} - u) \sigma(T_{[a_1]}) - \sigma(T_{a_1}) \sigma(T_{[a_1]} - u)
\end{aligned}$$

Since  $u$  is in the  $a_1$ -tree and  $d(u, a_1) \geq 2$ , by Lemma 2,

$$X_2 \begin{cases} > 0 & \text{if } d(u, a_1) \text{ is even} \\ < 0 & \text{if } d(u, a_1) \text{ is odd} \end{cases}$$

When  $i = 2k$ ,  $3 \leq i \leq n/2 + 1$  and  $v = a_i$ , we have

$$\begin{aligned} X_1 &= \sigma(T_{2,n}) \sigma(T_{3,n} - a_n - v) - \sigma(T_{3,n} - a_n) \sigma(T_{2,n} - v) \\ &= \sigma(T_{3,n} - a_n - v) [\sigma(T_{3,n}) \sigma(T_{a_2}) + \sigma(T_{4,n}) \sigma(T_{[a_2]}) \sigma(T_{a_3})] \\ &\quad - \sigma(T_{3,n} - a_n) [\sigma(T_{3,n} - v) \sigma(T_{a_2}) + \sigma(T_{4,n} - v) \sigma(T_{[a_2]}) \sigma(T_{a_3})] \\ &= [\sigma(T_{3,n} - a_n - v) \sigma(T_{3,n}) - \sigma(T_{3,n} - a_n) \sigma(T_{3,n} - v)] \sigma(T_{a_2}) \\ &\quad + \sigma(T_{a_3}) \sigma(T_{[a_2]}) [\sigma(T_{3,n} - a_n - v) \sigma(T_{4,n}) - \sigma(T_{3,n} - a_n) \sigma(T_{4,n} - v)] \\ &= q_1 \sigma(T_{a_2}) + \sigma(T_{a_3}) \sigma(T_{[a_2]}) [\sigma(T_{3,n} - a_n - v) \sigma(T_{4,n}) \\ &\quad - \sigma(T_{3,n} - a_n) \sigma(T_{4,n} - v)] \\ &= q_1 \sigma(T_{a_2}) + \sigma(T_{a_3}) \sigma(T_{[a_2]}) [[\sigma(T_{4,n} - a_n - v) \sigma(T_{a_3}) \\ &\quad + \sigma(T_{5,n} - a_n - v) \sigma(T_{[a_3]}) \sigma(T_{a_4})] \sigma(T_{4,n}) \\ &\quad - \sigma(T_{4,n} - v) [\sigma(T_{4,n} - a_n) \sigma(T_{a_3}) + \sigma(T_{5,n} - a_n) \sigma(T_{[a_3]}) \sigma(T_{a_4})]] \\ &= q_1 \sigma(T_{a_2}) + (\sigma(T_{a_3}))^2 \sigma(T_{[a_2]}) [\sigma(T_{4,n} - a_n - v) \sigma(T_{4,n}) \\ &\quad - \sigma(T_{4,n} - v) \sigma(T_{4,n} - a_n)] \\ &\quad + \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \sigma(T_{a_4}) [\sigma(T_{4,n}) \sigma(T_{5,n} - a_n - v) \\ &\quad - \sigma(T_{4,n} - v) \sigma(T_{5,n} - a_n)] \\ &= q_1 \sigma(T_{a_2}) + q_2 (\sigma(T_{a_3}))^2 \sigma(T_{[a_2]}) + q_3 (\sigma(T_{a_4}))^2 \sigma(T_{[a_2]}) \\ &\quad \times [\sigma(T_{a_3}) \sigma(T_{[a_3]}) + \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \sigma(T_{a_4}) \sigma(T_{[a_4]}) \sigma(T_{a_5}) \\ &\quad \times [\sigma(T_{5,n} - a_n - v) \sigma(T_{6,n}) - \sigma(T_{5,n} - a_n) \sigma(T_{6,n} - v)]] \\ &= q_1 \sigma(T_{a_2}) + q_2 (\sigma(T_{a_3}))^2 \sigma(T_{[a_2]}) + \cdots + q_{2k-3} (\sigma(T_{a_{2k-2}}))^2 \sigma(T_{[a_2]}) \\ &\quad \times \sigma(T_{a_3}) \sigma(T_{[a_3]}) \cdots \sigma(T_{[a_{2k-3}]} + \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \cdots \\ &\quad \times \sigma(T_{a_{2k-2}}) \sigma(T_{[a_{2k-2}]} \sigma(T_{a_{2k-1}}) [\sigma(T_{2k-1,n} - a_n - v) \sigma(T_{2k,n}) \\ &\quad - \sigma(T_{2k-1,n} - a_n) \sigma(T_{2k,n} - v)] \\ &= q_1 \sigma(T_{a_2}) + \cdots + q_{2k-2} (\sigma(T_{a_{2k-1}}))^2 \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \cdots \sigma(T_{[a_{2k-2}]} \\ &\quad + \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \cdots \sigma(T_{[a_{2k-1}]} \sigma(T_{a_{2k}}) \\ &\quad \times [\sigma(T_{2k+1,n} - a_n) \sigma(T_{2k,n}) - \sigma(T_{2k+1,n} - a_n) \sigma(T_{2k,n} - v)] \end{aligned}$$

where for  $j = 1, 2, \dots, 2k - 2$ ,

$$\begin{aligned} q_j &= \sigma(T_{j+2,n} - a_n - v) \sigma(T_{j+2,n}) - \sigma(T_{j+2,n} - a_n) \sigma(T_{j+2,n} - v) \\ &= -[\sigma(T_{j+2,n} - a_n) \sigma(T_{j+2,n} - v) - \sigma(T_{j+2,n} - a_n - v) \sigma(T_{j+2,n})] \end{aligned}$$

Note that the last term in the above expression for  $X_1$  is equal to

$$\begin{aligned} &\sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \cdots \sigma(T_{[a_{2k-1}]} \sigma(T_{a_{2k}}) [\sigma(T_{2k+1,n} - a_n) \sigma(T_{2k,n}) \\ &\quad - \sigma(T_{2k+1,n} - a_n) \sigma(T_{2k,n} - v)] > 0 \end{aligned}$$



Since  $T_{j+2,n}$  is a tree, by Lemma 1 we know that

$$q_j \begin{cases} > 0 & \text{if } d(v, a_n) \text{ is even} \\ < 0 & \text{if } d(v, a_n) \text{ is odd} \end{cases}$$

Because  $d(v, a_n) = d(a_{2k}, a_n)$  is even, it must be  $q_j > 0$  for  $j = 1, 2, \dots, 2k-2$ . So we have  $X_1 > 0$  when  $i = 2k$ ,  $3 \leq i \leq n/2 + 1$  and  $v = a_i$ .

The conclusions obtained so far can be summarized in the following table.

$d(u, a_1)$	$X_1$	$X_2$	$d_{U_n}(u, v) = d(u, a_1) + 2k - 1$	$\Delta_{uv}(U_n) = \sigma(T_{a_2}) X_1 X_2$
odd	$> 0$	$< 0$	even	$< 0$
even	$> 0$	$> 0$	odd	$> 0$

Hence, when  $i = 2k$ ,  $3 \leq i \leq n/2 + 1$  and  $v = a_i$ ,  $d(u, a_1) \geq 2$ , Theorem 2 is true.

If  $i = 2k + 1$ ,  $3 \leq i \leq n/2 + 1$  and  $v = a_i$ , then we have

$$\begin{aligned} X_1 &= \sigma(T_{2,n}) \sigma(T_{3,n} - a_n - v) - \sigma(T_{3,n} - a_n) \sigma(T_{2,n} - v) \\ &= q_1 \sigma(T_{a_2}) + q_2 (\sigma(T_{a_3}))^2 \sigma(T_{[a_2]}) + q_3 (\sigma(T_{a_4}))^2 \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) + \dots \\ &\quad + q_{2k-2} (\sigma(T_{a_{2k-2}}))^2 \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \dots \sigma(T_{[a_{2k-2}]} \\ &\quad + \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \dots \sigma(T_{a_{2k-1}}) \sigma(T_{a_{2k}}) \\ &\quad \times [\sigma(T_{2k+1,n} - a_n - v) \sigma(T_{2k,n}) - \sigma(T_{2k,n} - a_n) \sigma(T_{2k,n} - v)] \\ &= q_1 \sigma(T_{a_2}) + q_2 (\sigma(T_{a_3}))^2 \sigma(T_{[a_2]}) \\ &\quad + q_3 (\sigma(T_{a_4}))^2 \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \\ &\quad + \dots + q_{2k-2} (\sigma(T_{a_{2k-2}}))^2 \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \dots \sigma(T_{[a_{2k-2}]} \\ &\quad + q_{2k-1} (\sigma(T_{a_{2k}}))^2 \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \dots \sigma(T_{a_{2k-1}}) \sigma(T_{[a_{2k-1}]} + \\ &\quad + \sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \dots \sigma(T_{a_{2k}}) \sigma(T_{[a_{2k}]} \sigma(T_{a_{2k+1}}) \sigma(T_{2k+2,n}) \\ &\quad \times [\sigma(T_{2k+1,n} - a_n - v) - \sigma(T_{2k+1,n} - a_n)] \end{aligned}$$

where for  $j = 1, 2, \dots, 2k-1$ ,

$$\begin{aligned} q_j &= \sigma(T_{j+2,n} - a_n - v) \sigma(T_{j+2,n}) - \sigma(T_{j+2,n} - a_n) \sigma(T_{j+2,n} - v) \\ &= -[\sigma(T_{j+2,n} - a_n) \sigma(T_{j+2,n} - v) - \sigma(T_{j+2,n} - a_n - v) \sigma(T_{j+2,n})] \end{aligned}$$

Note that the last term in the above expression for  $X_1$  is:

$$\begin{aligned} &\sigma(T_{[a_2]}) \sigma(T_{a_3}) \sigma(T_{[a_3]}) \dots \sigma(T_{a_{2k}}) \sigma(T_{[a_{2k}]} \sigma(T_{a_{2k+1}}) \sigma(T_{2k+2,n}) \\ &\quad \times [\sigma(T_{2k+1,n} - a_n - v) - \sigma(T_{2k+1,n} - a_n)] < 0 \end{aligned}$$

Since  $T_{j+2,n}$  is a tree, by Lemma 1,

$$q_j \begin{cases} > 0 & \text{if } d(v, a_n) \text{ is even} \\ < 0 & \text{if } d(v, a_n) \text{ is odd} \end{cases}$$

Because  $d(v, a_n) = d(a_{2k+1}, a_n)$  is odd, it must be  $q_j < 0$  for  $j = 1, 2, \dots, 2k-1$ . So we have  $X_1 < 0$  when  $i = 2k + 1$ ,  $3 \leq i \leq n/2 + 1$  and  $v = a_i$ . Now we have the following table:

$d(u, a_1)$	$X_1$	$X_2$	$d_{U_n}(u, v) = d(u, a_1) + 2k$	$\Delta_{uv}(U_n) = \sigma(T_{a_2}) X_1 X_2$
odd	$< 0$	$< 0$	odd	$> 0$
even	$< 0$	$> 0$	even	$< 0$

Hence, when  $v = a_i$ ,  $i = 2k + 1$ ,  $3 \leq i \leq n/2 + 1$ ,  $d(u, a_1) \geq 2$ , Theorem 2 is also true.

This completes the analysis of Subcase 2.1.1.  $\square$

Based on a suggestion by the referee, we omit the equally lengthy and to a great extent analogous considerations needed to verify the validity of Theorem 2 in the remaining subcases. For the same reason we also skip the proof of Theorem 2 for Case 3. These omitted parts of the proof are available from the authors (X. L.) upon request.

By a reasoning similar to that used in the proof of Theorem 2, we obtain the following result.

Let  $U_n$  be as before and let  $n$  be odd. Let  $u$  and  $v$  be distinct, nonadjacent vertices of  $U_n$ .

**THEOREM 3.** *If  $e_1 = e_2 = \dots = e_n = e$ , a positive integer, then for the above described graph  $U_n$  the relations (1) are satisfied.*

### References

- [1] I. Gutman, *An identity for the independence polynomials of trees*, Publ. Inst. Math. (Beograd) **50** (1991), 19–23.
- [2] I. Gutman, F. Harary, *Generalizations of the matching polynomial*, Utilitas Math. **24** (1983), 97–106.
- [3] C. Hoede, X. Li, *Clique polynomials and independent set polynomials of graphs*, Discrete Math. **125** (1994), 219–228.
- [4] X. Li, *On a conjecture of Merrifield and Simmons*, Australasian J. Comb. **14** (1991), 15–20.
- [5] R. E. Merrifield, H. E. Simmons, *Topological Methods in Chemistry*, Wiley, New York, 1989.

Yong Wang and Xueliang Li  
 Department of Applied Mathematics  
 Northwestern Polytechnical University  
 Xi'an, Shaanxi 710072  
 P.R. China

(Received 08 12 1999)  
 (Revised 31 07 2000)  
 (Revised 27 04 2001)

`iamlx1@nwpu.edu.cn`

Prirodno-matematički fakultet  
 34000 Kragujevac, p.p. 60  
 Yugoslavia

`gutman@knez.uis.kg.ac.yu`