

SOME QUESTIONS ON METRIZABILITY

Ying Ge and Jian-Hua Shen

Communicated by Rade Živaljević

ABSTRACT. Let us say that a g -function $g(n, x)$ on a space X satisfies the condition $(*)$ provided: If $\{x_n\} \rightarrow p \in X$ and $x_n \in g(n, y_n)$ for every $n \in N$, then $y_n \rightarrow p$. We prove that a k -space X is a metrizable space (a metrizable space with property ACF) if and only if there exists a strongly decreasing g -function $g(n, x)$ on X such that $\{g(n, x) : x \in X\}$ is CF ($\{g(n, x) : x \in X\}$ is CF^*) in X for every $n \in N$ and the condition $(*)$ is satisfied. Our results give a partial answer to a question posed by Z. Yun, X. Yang and Y. Ge and a positive answer to a conjecture posed by S. Lin, respectively.

1. Introduction

How to characterize metrizable spaces in terms of g -functions is an important question of metrizability. In [6], Z. Yun, X. Yang and Y. Ge gave the following result.

THEOREM 1.1. [6, Theorem 4] *A Fréchet space X is metrizable if and only if there exists a strongly decreasing g -function $g(n, x)$ on X such that $\{g(n, x) : x \in X\}$ is CF in X for every $n \in N$ and the following condition is satisfied.*

$(*)$ *If $\{x_n\} \rightarrow p \in X$ and $x_n \in g(n, y_n)$ for every $n \in N$, then $y_n \rightarrow p$.*

The authors of [6] noted that the condition “Fréchet” in Theorem 1.1 can be relaxed to “ k' ”, but it can not be omitted. However, they still do not know whether the condition “Fréchet” in Theorem 1.1 can be relaxed to “ k ”. So they raised the following question.

QUESTION 1.2. [6, Question 1]. For a k -space X , are the following (1) and (2) equivalent?

(1) X is a metrizable space.

2000 *Mathematics Subject Classification*: 54D50, 54E35.

Key words and phrases: strongly decreasing g -function, CF -family, metrizable space, k -space.

This project was supported by NSF of the Education Committee of Jiangsu Province in China (No.02KJB110001).

(2) There exists a strongly decreasing g -function $g(n, x)$ on X such that $\{g(n, x) : x \in X\}$ is CF in X for every $n \in N$ and the condition $(*)$ is satisfied.

Notice that $CF^* \Rightarrow CF$. Taking Question 1.2 into account, Lin [3] raised a conjecture.

CONJECTURE 1.3. [3, Conjecture 1] A k -space X is metrizable (with some property) if and only if there exists a strongly decreasing g -function $g(n, x)$ on X such that $\{g(n, x) : x \in X\}$ is CF^* in X for every $n \in N$ and the condition $(*)$ is satisfied.

Here we investigate Question 1.2 and Conjecture 1.3. We prove that a k -space X is a metrizable space (a metrizable space with property ACF) if and only if there exists a strongly decreasing g -function $g(n, x)$ on X such that $\{\overline{g(n, x)} : x \in X\}$ is CF ($\{g(n, x) : x \in X\}$ is CF^*) in X for every $n \in N$ and the condition $(*)$ is satisfied. This gives a partial answer to Question 1.2 and a positive answer to Conjecture 1.3. As a corollary of the above results, a space X is a metrizable space with property ACF if and only if there exists a strongly decreasing g -function $g(n, x)$ on X such that $\{g(n, x) : x \in X\}$ is HCP in X for every $n \in N$ and the condition $(*)$ is satisfied.

Throughout this paper, all spaces are assumed to be regular. N and ω denote the set of all natural numbers and the first infinite ordinal, respectively. For a set A , $|A|$ denotes the cardinality of A . Let A be a subset of a space X and let \mathcal{F} be a family of subsets of X . \overline{A} , $\overline{\mathcal{F}}$, $\bigcup \mathcal{F}$ and $A \wedge \mathcal{F}$ denote the closure of A , the family $\{\overline{F} : F \in \mathcal{F}\}$, the union $\bigcup \{F : F \in \mathcal{F}\}$ and the family $\{A \cap F : F \in \mathcal{F}\}$, respectively. If also $x \in X$, $(\mathcal{F})_x$ denotes the subfamily $\{F \in \mathcal{F} : x \in F\}$ of \mathcal{F} and $\bigcup (\mathcal{F})_x$ is replaced by $st(x, \mathcal{F})$. One may consult [1] for undefined notation and terminology.

DEFINITION 1.4. [5] A space X is said to have property ACF if every compact subset of X is finite.

REMARK 1.5. Any space is the quotient space of a space with the property ACF [5].

DEFINITION 1.6. [6] Let X be a space and let τ be the topology on X . A function $g : N \times X \rightarrow \tau$ is called a g -function on X (we write $g(n, x)$ for short) if $x \in g(n, x)$ for every $n \in N$ and every $x \in X$. A g -function $g(n, x)$ on X is called strongly decreasing if $\overline{g(n+1, x)} \subset g(n, x)$ for every $n \in N$ and every $x \in X$.

DEFINITION 1.7. [4] Let \mathcal{F} be a family of subsets of a space X . \mathcal{F} is called CF in X if for every compact subset $K \subset X$, $K \wedge \mathcal{F} = \{F_1, F_2, \dots, F_k\}$, that is, $|K \wedge \mathcal{F}| < \omega$, and called CF^* in X if also only finitely many $F \in \mathcal{F}$ have infinite intersections with K . \mathcal{F} is called closure-preserving in X if for every subfamily \mathcal{F}' of \mathcal{F} , $\bigcup \overline{\mathcal{F}'} = \overline{\bigcup \mathcal{F}'}$. $\mathcal{F} = \{F_\alpha : \alpha \in A\}$ is called hereditarily closure-preserving (hereditarily CF , hereditarily CF^*) in X if for any choice $E_\alpha \subset F_\alpha$, the family $\{E_\alpha : \alpha \in A\}$ is closure-preserving (CF , CF^*) in X .

Throughout this paper, we use brief notations for the following terms.

CP – closure-preserving; HCP – hereditarily closure-preserving;
 HCF – hereditarily CF ; HCF^* – hereditarily CF^* .

Also we call a g -function $g(n, x)$ on a space X to be a P g -function (\overline{P} g -function) for short, if $\{g(n, x) : x \in X\}$ ($\{\overline{g(n, x)} : x \in X\}$) is P in X for every $n \in N$, where P is CP , HCP , CF , HCF , CF^* and HCF^* , respectively.

DEFINITION 1.8. [4] A space X is called a Fréchet space if for every $H \subset X$ and for every $x \in \overline{H}$, there exists a sequence $\{x_n\} \subset H$ such that $\{x_n\} \rightarrow x$; is called a k' -space if for every nonclosed subset $H \subset X$ and for every point $x \in \overline{H} - H$, there exists a compact subset $K \subset X$ such that $x \in \overline{H \cap K}$; is called a k -space if X has the weak topology with respect to the family of all compact subsets of X .

REMARK 1.9. It is well known that Fréchet $\Rightarrow k' \Rightarrow k$ and none of the implications can be reversed.

2. The main results

We start by giving some lemmas.

LEMMA 2.1. [6, Lemma 1] Let \mathcal{F} be a family of subsets of a space X . If \mathcal{F} is CF in X , then $\{\bigcup \mathcal{F}' : \mathcal{F}' \subset \mathcal{F}\}$ is also CF in X .

LEMMA 2.2. (1) For a family of subsets of a space, locally finite $\Rightarrow HCP \Rightarrow CP$, $HCP \Rightarrow CF^* \Rightarrow CF$ [4, Proposition 3.7] and $HCF \Leftrightarrow CF^* \Leftrightarrow HCF^*$ [2, Theorem 1].

(2) For a family of closed subsets of a k -space, $CF \Rightarrow CP$ [2, Lemma 2] and $CF^* \Leftrightarrow HCP$ [2, Theorem 4].

(3) For a family of subsets of a k' -space, $CF^* \Leftrightarrow HCP$ [2, Theorem 6].

LEMMA 2.3. [6, Theorem 3] A space X is metrizable if and only if there exists a CP strongly decreasing g -function $g(n, x)$ on X such that the condition (*) is satisfied.

The proof of the following lemma is trivial, and we omit it.

LEMMA 2.4. Let \mathcal{F} be a family of subsets of a space with property ACF . Then \mathcal{F} is CF^* in X .

LEMMA 2.5. Let X be a space. If there exists a CF^* g -function $g(n, x)$ on X , then X has property ACF .

PROOF. Let K be a compact subset of X and let $n \in N$. Then $\{g(n, x) : x \in X\}$ is HCF in X from Lemma 2.2(1). Note that $\{x\} \subset g(n, x)$ for every $x \in X$. $\{\{x\} : x \in X\}$ is CF in X , so $\{\{x\} : x \in K\} = K \wedge \{\{x\} : x \in X\}$ is finite, that is, K is finite. This proves that X has property ACF . \square

The following theorem gives an almost positive answer to Question 1.2.

THEOREM 2.6. A k -space X is metrizable if and only if there exists a \overline{CF} strongly decreasing g -function $g(n, x)$ on X such that the condition (*) is satisfied.

PROOF. Necessity: Assume that X is a metrizable space. We denote the diameter of subset A of X by $d(A)$. Let \mathcal{U}_1 be a locally finite cover of X such that $d(U) < 1$ for every $U \in \mathcal{U}_1$. Put $g(1, x) = st(x, \mathcal{U}_1)$ for every $x \in X$. Let \mathcal{U}_2 be a locally finite cover of X such that $d(U) < 1/2$ for every $U \in \mathcal{U}_2$, and $\{\overline{U} : U \in \mathcal{U}_2\}$ is a refinement of \mathcal{U}_1 . Put $g(2, x) = st(x, \mathcal{U}_2)$ for every $x \in X$. Generally, Let \mathcal{U}_n be a locally finite cover of X such that $d(U) < 1/n$ for every $U \in \mathcal{U}_n$, and $\{\overline{U} : U \in \mathcal{U}_n\}$ is a refinement of \mathcal{U}_{n-1} . Put $g(n, x) = st(x, \mathcal{U}_n)$ for every $x \in X$.

Thus we obtain a g -function $g(n, x)$ on X . By the proof of (a) \Rightarrow (b) in [6, Theorem 3], the g -function $g(n, x)$ is strongly decreasing and satisfies the condition (*). Now we only need to prove that $\{\overline{g(n, x)} : x \in X\}$ is CF in X for every $n \in N$.

In fact, for every $n \in N$, \mathcal{U}_n is locally-finite, so $\overline{\mathcal{U}_n}$ is locally finite. Put $F(n, x) = \bigcup\{\overline{U} \in \overline{\mathcal{U}_n} : x \in U\}$. Then $\{F(n, x) : x \in X\}$ is CF in X by Lemma 2.2(1) and Lemma 2.1. For every $x \in X$, note that $(\mathcal{U}_n)_x$ is a finite subfamily of \mathcal{U}_n . $F(n, x) = \bigcup\{\overline{U} \in \overline{\mathcal{U}_n} : x \in U\} = \overline{\bigcup\{U \in \mathcal{U}_n : x \in U\}} = \overline{st(x, \mathcal{U}_n)}$. Thus $\overline{g(n, x)} = \overline{st(x, \mathcal{U}_n)} = F(n, x)$. So $\{\overline{g(n, x)} : x \in X\}$ is CF in X .

Sufficiency: Let X be a k -space. Assume that there exists a \overline{CF} strongly decreasing g -function $g(n, x)$ on X such that the condition (*) is satisfied. Then for every $n \in N$, $\{\overline{g(n, x)} : x \in X\}$ is CF in X . Since X is a k -space, $\{\overline{g(n, x)} : x \in X\}$ is CP in X by Lemma 2.2(2). Note that a family \mathcal{F} of subsets of a space is CP in X if and only if $\overline{\mathcal{F}}$ is CP in X . $\{g(n, x) : x \in X\}$ is CP in X , so there exists a CP strongly decreasing g -function $g(n, x)$ on X such that the condition (*) is satisfied. Thus X is a metrizable space by Lemma 2.3 \square

The following theorem gives a positive answer to Conjecture 1.3.

THEOREM 2.7. *A k -space X is a metrizable space with property ACF if and only if there exists a CF^* strongly decreasing g -function $g(n, x)$ on X such that the condition (*) is satisfied.*

PROOF. Necessity: Assume that X is a metrizable space with property ACF . By Theorem 2.6, there exists a strongly decreasing g -function $g(n, x)$ on X such that the condition (*) is satisfied. Since X has property ACF , $\{g(n, x) : x \in X\}$ is CF^* in X by Lemma 2.4 for every $n \in N$. So there exists a CF^* strongly decreasing g -function $g(n, x)$ on X such that the condition (*) is satisfied.

Sufficiency: Let X be a k -space. Assume that there exists a CF^* strongly decreasing g -function $g(n, x)$ on X such that the condition (*) is satisfied. At first, it is obvious that X has the property ACF by Lemma 2.5. For every $n \in N$, since $\{g(n, x) : x \in X\}$ is CF^* in X and $\overline{g(n+1, x)} \subset g(n, x)$ for every $x \in X$, $\{\overline{g(n+1, x)} : x \in X\}$ is CF in X by Lemma 2.2(1). Thus there exists a \overline{CF} strongly decreasing g -function $g'(n, x)$ on X such that the condition (*) is satisfied, where $g'(n, x) = g(n+1, x)$. So X is metrizable by Theorem 2.6. \square

COROLLARY 2.8. *A space X is a metrizable space with property ACF if and only if there exists an HCP strongly decreasing g -function $g(n, x)$ on X such that the condition (*) is satisfied.*

PROOF. Necessity: Assume that X is a metrizable space with property ACF . By Theorem 2.7, there exists a CF^* strongly decreasing g -function $g(n, x)$ on X such that the condition $(*)$ is satisfied. Note that X is a k' -space. By Lemma 2.2(3), there exists an HCP strongly decreasing g -function $g(n, x)$ on X such that the condition $(*)$ is satisfied.

Sufficiency: Assume that there exists an HCP strongly decreasing g -function $g(n, x)$ on X such that the condition $(*)$ is satisfied. Since $HCP \Rightarrow CP$ by Lemma 2.2(1), X is metrizable by Lemma 2.3. Since $HCP \Rightarrow CF^*$ by Lemma 2.2(1), there exists a CF^* strongly decreasing g -function $g(n, x)$ on k -space X such that the condition $(*)$ is satisfied. Thus X is a metrizable space with property ACF by Theorem 2.7. \square

The authors would like to thank the referee for his valuable amendments.

References

- [1] R. Engelking, *General Topology*, (rev. ed.) Sigma Series in Pure Mathematics 6, Heldermann, Berlin, 1989.
- [2] Y. Ge, *Remarks On CF families*, Questions Answers in General Topology, 18 (2000), 103–110.
- [3] S. Lin, *g-functions and metrizable spaces*, preprint.
- [4] T. Mizokami, *On CF families and hyperspaces of compact subsets*, Top. Appl. 35 (1990), 75–92.
- [5] T. Mizokami and S. Lin, *On spaces with a σ - CF^* pseudo-base*, Math. Japonica 46 (1997), 377–384.
- [6] Z. Yun, X. Yang and Y. Ge, *Metrizable, \aleph -, Lašnev spaces and g-functions*, Math. Japonica 52 (2000), 225–229.

Department of Mathematics
Suzhou University
Suzhou, 215006
P. R. China
geying@pub.sz.jsinfo.net

(Received 15 03 2004)

Department of Mathematics
Suzhou Science and Technology College
Suzhou, Jiangsu, 215009
P. R. China.