

# From Quantum $A_N$ (Calogero) to $H_4$ (Rational) Model\*

Alexander V. TURBINER

Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México,  
Apartado Postal 70-543, 04510 México, D.F., Mexico

E-mail: [turbiner@nucleares.unam.mx](mailto:turbiner@nucleares.unam.mx)

Received February 28, 2011, in final form July 12, 2011; Published online July 18, 2011

[doi:10.3842/SIGMA.2011.071](https://doi.org/10.3842/SIGMA.2011.071)

**Abstract.** A brief and incomplete review of known integrable and (quasi)-exactly-solvable quantum models with rational (meromorphic in Cartesian coordinates) potentials is given. All of them are characterized by (i) a discrete symmetry of the Hamiltonian, (ii) a number of polynomial eigenfunctions, (iii) a factorization property for eigenfunctions, and admit (iv) the separation of the radial coordinate and, hence, the existence of the 2nd order integral, (v) an algebraic form in invariants of a discrete symmetry group (in space of orbits).

*Key words:* (quasi)-exact-solvability; rational models; algebraic forms; Coxeter (Weyl) invariants, hidden algebra

*2010 Mathematics Subject Classification:* 35P99; 47A15; 47A67; 47A75

## 1 Introduction

In this paper we will make an attempt to overview our constructive knowledge about (quasi)-exactly-solvable potentials having a form of a meromorphic function in Cartesian coordinates. All these models have a discrete group of symmetry, admit separation of variable(s), possess an (in)finite set of polynomial eigenfunctions. They have an infinite discrete spectrum which is linear in the quantum numbers. All of them are characterized by the presence of a hidden (Lie) algebraic structure. Each of them is a type of isospectral deformation of the isotropic harmonic oscillator.

Let us consider the Hamiltonian = the Schrödinger operator

$$\mathcal{H} = -\Delta + V(x), \quad x \in \mathbb{R}^d.$$

The main problem of quantum mechanics is to solve the Schrödinger equation

$$\mathcal{H}\Psi(x) = E\Psi(x), \quad \Psi(x) \in L^2(\mathbb{R}^d),$$

finding the spectrum (the energies  $E$  and eigenfunctions  $\Psi$ ). Since the Hamiltonian is an infinite-dimensional matrix, solving the Schrödinger equation is equivalent to diagonalizing the infinite-dimensional matrix. It is a transcendental problem: the characteristic polynomial is of infinite order and it has infinitely-many roots. Usually, we do not know how to make such a diagonalization exactly (explicitly) but we can ask: *Do models exist for which the roots (energies), some of the them or all, can be found explicitly (exactly)?* Such models do exist and we call them *solvable*. If all energies are known they are called *exactly-solvable* (ES), if only some number of them is known we call them *quasi-exactly-solvable* (QES). Surprisingly, all such models the

---

\*This paper is a contribution to the Special Issue "Symmetry, Separation, Super-integrability and Special Functions (S<sup>4</sup>)". The full collection is available at <http://www.emis.de/journals/SIGMA/S4.html>

present author familiar with, are provided by integrable systems. The Hamiltonians of these models are of the form

$$\mathcal{H}_{\text{ES}} = -\frac{1}{2}\Delta + \omega^2 r^2 + \frac{W(\Omega)}{r^2},$$

in the exactly-solvable case and

$$\mathcal{H}_{\text{QES}} = -\frac{1}{2}\Delta + \tilde{\omega}_k^2 r^2 + \frac{W(\Omega) + \Gamma}{r^2} + ar^6 + br^4,$$

in the quasi-exactly-solvable case, where  $\omega$ ,  $\tilde{\omega}_k$ ,  $\Gamma$  are parameters,  $W(\Omega)$  is a function on unit sphere and  $r$  is the radial coordinate. In both cases there exists the integral

$$\mathcal{F} = \frac{1}{2}\mathcal{L}^2 + W(\Omega), \quad (1)$$

where  $\mathcal{L}$  is the angular momentum operator, due to the separation of variables in spherical coordinates.

Now we consider some examples from the ones known so far.

## 2 Solvable models

### 2.1 Case $O(N)$

The Hamiltonian reads

$$\mathcal{H}_{O(N)} = \frac{1}{2} \sum_{i=1}^N \left( -\frac{\partial^2}{\partial x_i^2} + \omega^2 x_i^2 \right) + \frac{\nu(\nu-1)}{\sum_{i=1}^N x_i^2}, \quad (2)$$

or, in spherical coordinates,

$$\mathcal{H}_{O(N)} = -\frac{1}{2r^N} \frac{\partial}{\partial r} \left( r^N \frac{\partial}{\partial r} \right) + \frac{1}{2}\omega^2 r^2 + \frac{\mathcal{F} + \nu(\nu-1)}{r^2}, \quad (3)$$

$$\mathcal{F} = \frac{1}{2}\mathcal{L}^2. \quad (4)$$

The Hamiltonian (2) is  $O(N)$  symmetric. It describes a spherical-symmetric harmonic oscillator with a generalized centrifugal potential. Needless to say that the Hamiltonian  $\mathcal{H}_{O(N)}$  and  $\mathcal{F}$  commute,

$$[\mathcal{H}_{O(N)}, \mathcal{F}] = 0.$$

Thus,  $\mathcal{F}$  has common eigenfunctions with the Hamiltonian  $\mathcal{H}_{O(N)}$ . The spectrum can be immediately found explicitly, and all eigenfunctions are of the type

$$P_n(r^2) r^{\tilde{\ell}} Y_{\{\ell\}}(\Omega) e^{-\frac{\omega r^2}{2}},$$

where  $Y_{\{\ell\}}(\Omega)$  is a  $N$ -dimensional spherical harmonics,  $\mathcal{F}Y_{\{\ell\}}(\Omega) = \gamma Y_{\{\ell\}}(\Omega)$ . The Hamiltonian (2) describes an  $N$ -dimensional harmonic oscillator with generalized centrifugal term. Substituting in (3) the operator  $\mathcal{F}$  by its eigenvalue  $\gamma$  and gauging away  $\Psi_0 = r^{\tilde{\ell}} e^{-\frac{\omega r^2}{2}}$  we arrive at the Laguerre operator

$$h_{O(N)} \equiv (\Psi_0)^{-1} (\mathcal{H}_{O(N)} - E_0) \Psi_0|_{r^2=t} = -2t\partial_t^2 + \left( 2\omega - 1 - \frac{N}{2} - \tilde{\ell} \right) \partial_t, \quad (5)$$

where  $E_0$  is the lowest energy and the parameter  $\tilde{\ell}$  is chosen in such a way as to remove singular term  $\propto \frac{1}{r^2}$  in the potential in (3). (5) is the algebraic form of the Hamiltonian (3). The gauge-rotated Hamiltonian  $h_{O(N)}$  (5) is  $sl(2)$ -Lie-algebraic (see below), it has infinitely-many finite-dimensional invariant subspaces in polynomials  $\mathcal{P}_n$ ,  $n = 0, 1, \dots$  forming the infinite flag (see below), its eigenfunctions  $P_n(r^2 = t)$  are nothing but the associated Laguerre polynomials.

By adding to  $h_{O(N)}$  (5) the operator

$$\delta h^{(\text{qes})} = 4(at^2 - \gamma) \frac{\partial}{\partial t} - 4akt + 2\omega k, \quad (6)$$

we get the operator  $h_{O(N)} + \delta h^{(\text{qes})}$  which has a single finite-dimensional invariant subspace

$$\mathcal{P}_k = \langle t^p \mid 0 \leq p \leq k \rangle,$$

of the dimension  $(k + 1)$ . Hence, this operator is quasi-exactly-solvable. Making the change of variable  $t = r^2$  and gauge rotation with  $\tilde{\Psi}_0 = t^{\hat{\gamma}} e^{-\frac{\omega t}{2} - \frac{at^2}{4}}$  we arrive at the  $O(N)$ -symmetric QES Hamiltonian [2]

$$\mathcal{H}_{O(N)} = -\frac{1}{2r^N} \frac{\partial}{\partial r} \left( r^N \frac{\partial}{\partial r} \right) + a^2 r^6 + 2a\omega r^4 + \frac{1}{2} \tilde{\omega}^2 r^2 + \frac{\mathcal{F} + \Gamma}{r^2}, \quad (7)$$

where  $\hat{\gamma}$ ,  $\Gamma$ ,  $\tilde{\omega}$  are parameters and  $\gamma$  is replaced by the operator  $\mathcal{F}$ . In (7) a finite number of the eigenfunctions is of the form

$$P_k(r^2) r^{2\hat{\gamma}} Y_{\{\ell\}}(\Omega) e^{-\frac{\omega r^2}{2} - \frac{at^2}{4}},$$

they can be found algebraically. It is worth noting that at  $a = 0$  the operator  $h_{O(N)} + \delta h^{(\text{qes})}$  remains exactly-solvable, it preserves the infinite flag of polynomials  $\mathcal{P}$  and the emerging Hamiltonian has a form of (3).

## 2.2 Case $(\mathbb{Z}_2)^N$

The Hamiltonian reads

$$\mathcal{H}_{(\mathbb{Z}_2)^N} = \frac{1}{2} \sum_{i=1}^N \left( -\frac{\partial^2}{\partial x_i^2} + \omega^2 x_i^2 \right) + \frac{1}{2} \sum_{i=1}^N \frac{\nu_i(\nu_i - 1)}{x_i^2}, \quad (8)$$


or, in spherical coordinates,

$$\mathcal{H}_{(\mathbb{Z}_2)^N} = -\frac{1}{2r^N} \frac{\partial}{\partial r} \left( r^N \frac{\partial}{\partial r} \right) + \frac{1}{2} \omega^2 r^2 + \frac{\mathcal{F} + W_{(\mathbb{Z}_2)^N}(\Omega)}{r^2},$$

where

$$W_{(\mathbb{Z}_2)^N}(\Omega) = \frac{1}{2} \sum_{i=1}^N \nu_i(\nu_i - 1) \left( \frac{r}{x_i} \right)^2,$$

and  $\mathcal{F}$  is given by (4). The Hamiltonian (8) is  $(\mathbb{Z}_2)^N$  symmetric. It defines the so-called Smorodinsky–Winternitz integrable system [1] which is in reality the maximally-superintegrable (there exist  $(2N - 1)$  integrals including the Hamiltonian) and exactly-solvable. Gauging away in (8) the ground state,  $\Psi_0 = \prod_{i=1}^N (x_i^2)^{\frac{\nu_i}{2}} \exp(-\frac{\omega x_i^2}{2})$ , and changing variables to  $t_i = x_i^2$  we arrive at the algebraic form. Also it admits QES extension. The system described by the Hamiltonian (8) at  $\nu_i = \nu$  is a particular case of the  $BC_N$ -rational system (see below).



**Figure 1.**  $N$ -body Calogero model.

### 2.3 Case $A_{N-1}$

This is the celebrated Calogero model ( $A_{N-1}$  rational model) which was found in [3]. It describes  $N$  identical particles on a line (see Fig. 1) with singular pairwise interaction.

The Hamiltonian is

$$\mathcal{H}_{\text{Cal}} = \frac{1}{2} \sum_{i=1}^N \left( -\frac{\partial^2}{\partial x_i^2} + \omega^2 x_i^2 \right) + \nu(\nu - 1) \sum_{i>j}^N \frac{1}{(x_i - x_j)^2}, \quad (9)$$

where the singular part of the potential can be written as

$$\sum_{i>j}^N \frac{1}{(x_i - x_j)^2} = \frac{W_{A_{N-1}}(\Omega)}{r^2}, \quad W_{A_{N-1}}(\Omega) = \sum_{i=1}^N \left( \frac{1}{\frac{x_i}{r} - \frac{x_j}{r}} \right)^2, \quad (10)$$

Here  $r$  is the radial coordinate in the space of relative coordinates (see below for a definition) and  $W_{A_{N-1}}(\Omega)$  is a function on the unit sphere.

Symmetry:  $S_n$  (permutations  $x_i \rightarrow x_j$ ) plus  $\mathbb{Z}_2$  (all  $x_i \rightarrow -x_i$ ). The ground state of the Hamiltonian (9) reads

$$\Psi_0(x) = \prod_{i<j} |x_i - x_j|^\nu e^{-\frac{\omega}{2} \sum x_i^2}. \quad (11)$$

Let us make the gauge rotation

$$h_{\text{Cal}} = 2\Psi_0^{-1}(\mathcal{H}_{\text{Cal}} - E_0)\Psi_0,$$

and introduce center-of-mass variables

$$Y = \sum x_i, \quad y_i = x_i - \frac{1}{N}Y, \quad i = 1, \dots, N,$$

and then permutationally-symmetric, translationally-invariant variables

$$(x_1, x_2, \dots, x_N) \rightarrow (Y, t_n(x) = \sigma_n(y(x)) \mid n = 2, 3, \dots, N),$$

where

$$\sigma_k(x) = \sum_{i_1 < i_2 < \dots < i_k} x_{i_1} x_{i_2} \dots x_{i_k}, \quad \sigma_k(-x) = (-)^k \sigma_k(x),$$

are elementary symmetric polynomials, and

$$t_1 = 0, \quad t_2 \sim \sum_{i<j} (x_i - x_j)^2 = r^2,$$

hence, the variable  $t_2$ , which plays fundamental role, is defined by radius in space of relative coordinates. After the center-of-mass separation, the gauge rotated Hamiltonian takes the algebraic form [4]

$$h_{\text{Cal}} = \mathcal{A}_{ij}(t) \frac{\partial^2}{\partial t_i \partial t_j} + \mathcal{B}_i(t) \frac{\partial}{\partial t_i}, \quad (12)$$

where

$$\mathcal{A}_{ij} = \frac{(N-i+1)(1-j)}{N} t_{i-1} t_{j-1} + \sum_{l \geq \max(1, j-i)} (2l-j+i) t_{i+l-1} t_{j-l-1},$$

$$\mathcal{B}_i = \frac{1}{N} (1 + \nu N)(N-i+2)(N-i+1) t_{i-2} + 2\omega(i-1) t_i.$$

Eigenvalues of (12) are

$$\epsilon_{\{p\}} = 2\omega \sum_{i=2}^N (i-1) p_i,$$

hence, the spectrum is linear in the quantum numbers  $p_{2,3,\dots,N} = 0, 1, \dots$ , it corresponds to *anisotropic* harmonic oscillator with frequency ratios  $1 : 2 : 3 : \dots : (N-1)$ .

It is easy to check that the gauge-rotated Hamiltonian  $h_{\text{Cal}}$  has infinitely many finite-dimensional invariant subspaces

$$\mathcal{P}_n^{(N-1)} = \langle t_2^{p_2} t_3^{p_3} \dots t_N^{p_N} \mid 0 \leq \sum p_i \leq n \rangle.$$

where  $n = 0, 1, 2, \dots$ . As a function of  $n$  the spaces  $\mathcal{P}_n^{(N-1)}$  form the infinite flag (see below).

### 2.3.1 The $gl_{d+1}$ -algebra acting by 1st order differential operators in $\mathbb{R}^d$

It can be checked by the direct calculation that the  $gl_{d+1}$  algebra realized by the first order differential operators acting in  $\mathbb{R}^d$  in the representation given by the Young tableaux as a row  $(n, \underbrace{0, 0, \dots, 0}_{d-1})$  has a form

$$\mathcal{J}_i^- = \frac{\partial}{\partial t_i}, \quad i = 1, 2, \dots, d, \quad \mathcal{J}_{ij}^0 = t_i \frac{\partial}{\partial t_j}, \quad i, j = 1, 2, \dots, d, \quad \mathcal{J}^0 = \sum_{i=1}^d t_i \frac{\partial}{\partial t_i} - n,$$

$$\mathcal{J}_i^+ = t_i \mathcal{J}^0 = t_i \left( \sum_{j=1}^d t_j \frac{\partial}{\partial t_j} - n \right), \quad i = 1, 2, \dots, d. \quad (13)$$

where  $n$  is an arbitrary number. The total number of generators is  $(d+1)^2$ . If  $n$  takes the integer values,  $n = 0, 1, 2, \dots$ , the finite-dimensional irreps occur

$$\mathcal{P}_n^{(d)} = \langle t_1^{p_1} t_2^{p_2} \dots t_d^{p_d} \mid 0 \leq \sum p_i \leq n \rangle.$$

It is a common invariant subspace for (13). The spaces  $\mathcal{P}_n$  at  $n = 0, 1, 2, \dots$  can be ordered

$$\mathcal{P}_0 \subset \mathcal{P}_1 \subset \mathcal{P}_2 \subset \dots \subset \mathcal{P}_n \subset \dots \subset \mathcal{P}.$$

Such a nested construction is called *infinite flag (filtration)  $\mathcal{P}$* . It is worth noting that the flag  $\mathcal{P}^{(d)}$  is made out of finite-dimensional irreducible representation spaces  $\mathcal{P}_n^{(d)}$  of the algebra  $gl_{d+1}$  taken in realization (13). It is evident that *any operator made out of generators (13) has finite-dimensional invariant subspace which is finite-dimensional irreducible representation space*.

### 2.3.2 Algebraic properties of the Calogero model

It seems evident that the Hamiltonian (12) has to have a representation as a second order polynomial in generators (13) at  $d = N - 1$  acting in  $\mathbb{R}^{N-1}$ ,

$$h_{\text{Cal}} = \text{Pol}_2(\mathcal{J}_i^-, \mathcal{J}_{ij}^0),$$

where the raising generators  $\mathcal{J}_i^+$  are absent. Thus,  $gl(N)$  (or, strictly speaking, its maximal affine subalgebra) is the hidden algebra of the  $N$ -body Calogero model. Hence,  $h_{\text{Cal}}$  is an element of the universal enveloping algebra  $\mathcal{U}_{gl(N)}$ . The eigenfunctions of the  $N$ -body Calogero model are elements of the flag of polynomials  $\mathcal{P}^{(N-1)}$ . Each subspace  $\mathcal{P}_n^{(N-1)}$  is represented by the Newton polytope (pyramid). It contains  $C_{n+N-1}^{N-1}$  eigenfunctions, which is equal to the volume of the Newton polytope.

Making the gauge rotation of the integral (1) with  $W_{A_{N-1}}(\Omega)$  given by (10)

$$f_{\text{Cal}} = \Psi_0^{-1}(\mathcal{F}_{\text{Cal}} - F_0)\Psi_0,$$

where  $F_0$  is the lowest eigenvalue of the integral,  $\mathcal{F}_{\text{Cal}}\Psi_0 = F_0\Psi_0$ , the integral gets the algebraic form

$$f_{\text{Cal}} = f_{ij}(t) \frac{\partial^2}{\partial t_i \partial t_j} + g_i(t) \frac{\partial}{\partial t_i},$$

where  $f_{ij}$  is 2nd degree polynomial in  $t$ ,  $f_{2j} = 0$ , and  $g_i$  is 1st degree polynomial in  $t$ ,  $g_2 = 0$ . It also can be rewritten as the second degree polynomial in the  $gl(N)$  generators,

$$f_{\text{Cal}} = \text{Pol}_2(\mathcal{J}_i^-, \mathcal{J}_{ij}^0).$$

### 2.3.3 $sl(2)$ -quasi-exactly-solvable generalization of the Calogero model

By adding to  $h_{\text{Cal}}$  (12), the operator

$$\delta h^{(\text{qes})} = 4(at_2^2 - \gamma) \frac{\partial}{\partial t_2} - 4akt_2 + 2\omega k,$$

we get the operator  $h_{\text{Cal}} + \delta h^{(\text{qes})}$  having finite-dimensional invariant subspace

$$\mathcal{P}_k = \langle t_2^p \mid 0 \leq p \leq k \rangle.$$

By making a gauge rotation of  $h_{\text{Cal}} + \delta h^{(\text{qes})}$  and changing of variables to Cartesian ones we arrive at the Hamiltonian [5]

$$\begin{aligned} \mathcal{H}_{\text{Cal}}^{(\text{qes})} &= \frac{1}{2} \sum_{i=1}^N \left( -\frac{\partial^2}{\partial x_i^2} + \omega^2 x_i^2 \right) + \nu(\nu - 1) \sum_{j < i}^N \frac{1}{(x_i - x_j)^2} + \frac{2\gamma [\gamma - 2n(1 + \nu + \nu n) + 3]}{r^2} \\ &+ a^2 r^6 + 2a\omega r^4 - a [2k + 2n(1 + \nu + \nu n) - \gamma - 1] r^2. \end{aligned}$$

For the Hamiltonian,  $(k + 1)$  eigenfunctions are of the form

$$\begin{aligned} \Psi_k^{(\text{qes})}(x) &= \prod_{i < j}^n |x_i - x_j|^\nu (r^2)^\gamma P_k(r^2) \exp \left[ -\frac{\omega}{2} \sum_{i=1}^n x_i^2 - \frac{a}{4} r^4 \right] \\ &= (r^2)^\gamma P_k(r^2) \exp \left( -\frac{a}{4} r^4 \right) \Psi_0, \end{aligned}$$

where  $\Psi_0$  is given by (11),  $P_k$  is a polynomial of degree  $k$  in  $r^2 = \sum_{i < j} (x_i - x_j)^2 = t_2$ . All remaining eigenfunctions can be represented in the same form but  $P_k$ 's are not polynomials anymore being functions depending on all variables  $x_i$ . It is worth noting that at  $a = 0$  the operator  $h_{\text{Cal}} + \delta h^{(\text{res})}$  remains exactly-solvable, it preserves the flag of polynomials  $\mathcal{P}^{(N-1)}$  and the emerging Hamiltonian has a form of (9) with the extra term  $\frac{\Gamma}{r^2}$  in the potential. Its ground state eigenfunction is  $(r^2)^\gamma \Psi_0$ . It is the exactly-solvable generalization of the Calogero model (9) with the Weyl group  $W(A_{N-1})$  as the discrete symmetry group,

$$\mathcal{H}_{W(A_{N-1})} = \mathcal{H}_{\text{Cal}} + \frac{\Gamma}{r^2}.$$

## 2.4 Case: Hamiltonian reduction method

In this method<sup>1</sup> a family of integrable and exactly-solvable Hamiltonians associated with Weyl (Coxeter) symmetry was found with the Calogero model as one of its representatives. The idea of the method is beautiful and sufficient transparent,

- Take a simple group  $G$ ,
- Define the Laplace–Beltrami (invariant) operator on its symmetric space (free/harmonic oscillator motion),
- Radial part of Laplace–Beltrami operator is the Olshanetsky–Perelomov Hamiltonian relevant from physical point of view. The emerging Hamiltonian is the Weyl-symmetric, it can be associated with root system, it is integrable with integrals given by the invariant operators of higher than two orders with a property of solvability.

**Rational case.** This case appears when the coordinates of the symmetric space are introduced in such a way that the zero-curvature surface occurs. Emerging the Calogero–Moser–Sutherland–Olshanetsky–Perelomov Hamiltonian in the Cartesian coordinates has the form,

$$\mathcal{H} = \frac{1}{2} \sum_{k=1}^N \left[ -\frac{\partial^2}{\partial x_k^2} + \omega^2 x_k^2 \right] + \frac{1}{2} \sum_{\alpha \in R_+} \nu_{|\alpha|} (\nu_{|\alpha|} - 1) \frac{|\alpha|^2}{(\alpha \cdot x)^2}, \quad (14)$$

where  $R_+$  is a set of positive roots,  $x$  is a position vector and  $\nu_{|\alpha|}$  are coupling constants (parameters) which depend on the root length. If roots are of the same length, then  $\nu_{|\alpha|}$  have to be equal, if all roots are of the same length like for  $A_n$ , then all  $\nu_{|\alpha|} = \nu$ . In the Hamiltonian Reduction the parameters  $\nu_{|\alpha|}$  take a set of discrete values, however, they can be generalized to any real value without losing a property of integrability as well as of solvability with the only constraint of the existence of  $L^2$ -solutions of the corresponding Schrödinger equation. The configuration space for (14) is the Weyl chamber. The ground state wave function is written explicitly,

$$\Psi_0(x) = \prod_{\alpha \in R_+} |(\alpha \cdot x)|^{\nu_{|\alpha|}} e^{-\omega x^2/2}. \quad (15)$$

The Hamiltonian (14) is completely-integrable: there exists a commutative algebra of integrals (including the Hamiltonian) of dimension which is equal to the dimension of the configuration space (for integrals, see Oshima [7] with explicit forms of those). For each Hamiltonian (14) after separation of center-of-mass coordinate (if applicable) the radial coordinate (in the space of relative coordinates) can be also separated. It gives rise to the existence of one more integral

<sup>1</sup>For review and references see e.g. [6].

of the second order (1). Hence, the Hamiltonian (14) is super-integrable. The Hamiltonian (14) is invariant with respect to the Weyl (Coxeter) group transformation, which is the discrete symmetry group of the corresponding root space.

The Hamiltonian (14) has a hidden (Lie)-algebraic structure. In order to reveal it we need to

- Gauge away the ground state eigenfunction making *similarity transformation*  $(\Psi_0)^{-1}(\mathcal{H} - E_0)\Psi_0 = h$ ,
- Consider the Hamiltonian in the space of orbits of Weyl (Coxeter) group by taking the *Weyl (Coxeter) polynomial invariants as new coordinates*, these invariants are

$$t_a^{(\Omega)}(x) = \sum_{\alpha \in \Omega} (\alpha \cdot x)^a,$$

where  $a$ 's are the *degrees* of the Weyl (Coxeter) group,  $\Omega$  is an orbit.

The invariants  $t$  are defined ambiguously, up to invariants of lower degrees, they depend on chosen orbit.

## 2.5 Case $BC_N$

The  $BC_N$ -rational model is defined by the Hamiltonian,

$$\begin{aligned} \mathcal{H}_{BC_N} = & -\frac{1}{2} \sum_{i=1}^N \left( \frac{\partial^2}{\partial x_i^2} - \omega^2 x_i^2 \right) \\ & + \nu(\nu - 1) \sum_{i < j} \left[ \frac{1}{(x_i - x_j)^2} + \frac{1}{(x_i + x_j)^2} \right] + \frac{\nu_2(\nu_2 - 1)}{2} \sum_{i=1}^N \frac{1}{x_i^2}, \end{aligned} \quad (16)$$

where  $\omega$ ,  $\nu$ ,  $\nu_2$  are parameters. If  $\nu = 0$ , the Hamiltonian (16) is reduced to (8). The symmetry of the system is  $S_N \oplus (\mathbb{Z}_2)^N$  (permutations  $x_i \rightarrow x_j$  and  $x_i \rightarrow -x_i$ ).

The ground state function for (16) reads

$$\Psi_0 = \left[ \prod_{i < j} |x_i - x_j|^\nu |x_i + x_j|^\nu \prod_{i=1}^N |x_i|^{\nu_2} \right] e^{-\frac{\omega}{2} \sum_{i=1}^N x_i^2},$$

(cf. (15)). Making the gauge rotation

$$h_{BC_N} = (\Psi_0)^{-1} (\mathcal{H}_{BC_N} - E_0) \Psi_0,$$

and changing variables

$$(x_1, x_2, \dots, x_N) \rightarrow (\sigma_k(x^2) \mid k = 1, 2, \dots, N),$$

where

$$\sigma_k(x^2) = \sum_{i_1 < i_2 < \dots < i_k} x_{i_1}^2 x_{i_2}^2 \dots x_{i_k}^2, \quad \sigma_1(x^2) = x_1^2 + x_2^2 + \dots + x_N^2 = r^2,$$

where  $r$  is radius, we arrive at [8]

$$h_{BC_N} = \mathcal{A}_{ij}(\sigma) \frac{\partial^2}{\partial \sigma_i \partial \sigma_j} + \mathcal{B}_i(\sigma) \frac{\partial}{\partial \sigma_i},$$



with coefficients

$$\begin{aligned} \mathcal{A}_{ij} &= -2 \sum_{l \geq 0} (2l + 1 + j - i) \sigma_{i-l-1} \sigma_{j+l}, \\ \mathcal{B}_i &= [1 + \nu_2 + 2\nu(N - i)] (N - i + 1) \sigma_{i-1} + 2\omega i \sigma_i. \end{aligned}$$

This is the algebraic form of the  $BC_N$  Hamiltonian. Assuming polynomiality of the eigenfunctions we find the eigenvalues:

$$\epsilon_n = 2\omega \sum_{i=1}^N i n_i,$$

hence, the spectrum is equidistant, linear in the quantum numbers and corresponds to *anisotropic* harmonic oscillator with frequency ratios  $1 : 2 : 3 : \dots : N$ . The Hamiltonian  $h_{BC_N}$  has infinitely many finite-dimensional invariant subspaces of the form

$$\mathcal{P}_n^{(N)} = \langle \sigma_1^{p_1} \sigma_2^{p_2} \dots \sigma_N^{p_N} \mid 0 \leq \sum p_i \leq n \rangle,$$

where  $n = 0, 1, 2, \dots$ . They naturally form the flag  $\mathcal{P}^{(N)}$ . The Hamiltonian can be immediately rewritten in terms of generators (13) as a polynomial of the second degree,

$$h_{BC_N} = \text{Pol}_2(\mathcal{J}_i^-, \mathcal{J}_{ij}^0),$$

where the raising generators  $\mathcal{J}_i^+$  are absent. Hence,  $gl(N + 1)$  is the hidden algebra of the  $BC_N$  rational model, the same algebra as for the  $A_N$ -rational model. The eigenfunctions of the  $BC_N$ -rational model are elements of the flag of polynomials  $\mathcal{P}^{(N)}$ . Each subspace  $\mathcal{P}_n^{(N)}$  contains  $C_{n+N}^N$  eigenfunctions (volume of the Newton polytope (pyramid)  $\mathcal{P}_n^{(N)}$ ).

The  $BC_N$  Hamiltonian admits 2nd order integral as result of separation of radial variable

$$\mathcal{H}_{BC_N} = -\frac{1}{2r^{N-1}} \frac{\partial}{\partial r} \left( r^{N-1} \frac{\partial}{\partial r} \right) + \omega^2 r^2 + \frac{1}{2r^2} \underbrace{(-\Delta_\Omega^{(N-1)} + \mathcal{W}(\Omega))}_{\mathcal{F}_{BC_N}}.$$

Evidently, the commutator

$$[\mathcal{H}_{BC_N}, \mathcal{F}_{BC_N}] = 0.$$

Gauge-rotated integral

$$f_{BC_N} = \Psi_0^{-1} (\mathcal{F}_{BC_N} - F_0) \Psi_0,$$

where  $\mathcal{F}_{BC_N} \Psi_0 = F_0 \Psi_0$ , takes the algebraic form in  $t$ -coordinates,

$$f_{BC_N} = f_{ij}(t) \frac{\partial^2}{\partial t_i \partial t_j} + g_i(t) \frac{\partial}{\partial t_i},$$

where  $f_{ij}$  is 2nd degree polynomial,  $f_{1j} = 0$ , and  $g_i$  is 1st degree polynomial,  $g_1 = 0$ ,

$$f_{BC_N} = \text{Pol}_2(\mathcal{J}_i^-, \mathcal{J}_{ij}^0),$$

in terms of the  $gl(N + 1)$  generators. It is worth mentioning that the commutator of  $[h, f]$  vanishes only in the realization (13), otherwise,

$$[h_{BC_N}(\mathcal{J}), f_{BC_N}(\mathcal{J})] \neq 0.$$

### 2.5.1 $sl(2)$ -quasi-exactly-solvable generalization of the $BC_N$ rational model

By adding to  $h_{BC_N}$  the operator

$$\delta h^{(\text{qes})} = 4(a\sigma_1^2 - \gamma) \frac{\partial}{\partial \sigma_1} - 4ak\sigma_1 + 2\omega k,$$

which is similar to one for the Calogero model, we get the operator  $h_{BC_N} + \delta h^{(\text{qes})}$  which has the finite-dimensional invariant subspace

$$\mathcal{P}_k = \langle \sigma_1^p \mid 0 \leq p \leq k \rangle.$$

Making a gauge rotation of  $h_{BC_N} + \delta h^{(\text{qes})}$  and changing the variables  $\sigma$ 's back to the Cartesian ones the Hamiltonian becomes

$$\begin{aligned} \mathcal{H}_{BC_N}^{(\text{qes})} = & -\frac{1}{2} \sum_{i=1}^N \left( \frac{\partial^2}{\partial x_i^2} - \omega^2 x_i^2 \right) + \nu(\nu-1) \sum_{i<j} \left[ \frac{1}{(x_i - x_j)^2} + \frac{1}{(x_i + x_j)^2} \right] \\ & + \frac{\nu_2(\nu_2-1)}{2} \sum_{i=1}^N \frac{1}{x_i^2} + \frac{2\gamma[\gamma - 2N(1 + 2\nu(N-1) + \nu_2) + 3]}{r^2} \\ & + a^2 r^6 + 2a\omega r^4 - a[2k + 2N(1 + 2\nu(N-1) + \nu_2) - \gamma - 1] r^2, \end{aligned}$$

for which  $(k+1)$  eigenfunctions are of the form

$$\Psi_k^{(\text{qes})}(x) = \prod_{i<j}^n |x_i^2 - x_j^2|^\nu \prod_{i=1}^n |x_i|^{\nu_2} (r^2)^\gamma P_k(r^2) \exp \left[ -\frac{\omega r^2}{2} - \frac{a}{4} r^4 \right],$$

where  $P_k$  is a polynomial of degree  $k$  in  $r^2 = \sum_{i=1}^N x_i^2$ .

It is worth noting that at  $a = 0$  the operator  $h_{BC_N} + \delta h^{(\text{qes})}$  remains exactly-solvable, it preserves the flag of polynomials  $\mathcal{P}^{(N)}$  and the emerging Hamiltonian has a form of (16) with the extra term  $\frac{\Gamma}{r^2}$  in the potential. Its ground state eigenfunction is  $(r^2)^\gamma \Psi_0$ . It is the exactly-solvable generalization of the  $BC_N$ -rational model (16) with the Weyl group  $W(BC_N)$  as the discrete symmetry group,

$$\mathcal{H}_{W(BC_N)} = \mathcal{H}_{BC_N} + \frac{\Gamma}{r^2}.$$

Now we are in a position to draw an intermediate conclusion about  $A_N$  and  $BC_N$  rational models.

- Both  $A_N$ - and  $BC_N$ -rational (and trigonometric) models possess *algebraic* forms associated with preservation of the *same* flag of polynomials  $\mathcal{P}^{(N)}$ . The flag is invariant w.r.t. linear transformations in space of orbits  $t \mapsto t+A$ . It preserves the algebraic form of Hamiltonian.
- Their Hamiltonians (as well as higher integrals) can be written in the Lie-algebraic form

$$h = \text{Pol}_2(\mathcal{J}(b \subset gl_{N+1})),$$

where  $\text{Pol}_2$  is a polynomial of 2nd degree in generators  $\mathcal{J}$  of the maximal affine subalgebra of the algebra  $b$  of the algebra  $gl_{N+1}$  in realization (13). Hence,  $gl_{N+1}$  is their *hidden algebra*. From this viewpoint all four models are different faces of a *single* model.

- *Supersymmetric  $A_N$ - and  $BC_N$ -rational (and trigonometric) models possess algebraic forms, preserve the same flag of (super)polynomials and their hidden algebra is the super-algebra  $gl(N+1|N)$  (see [8]).*

In a connection to flags of polynomials we introduce a notion ‘*characteristic vector*’. Let us consider a flag made out of “triangular” linear space of polynomials

$$\mathcal{P}_{n,\vec{f}}^{(d)} = \langle x_1^{p_1} x_2^{p_2} \cdots x_d^{p_d} \mid 0 \leq f_1 p_1 + f_2 p_2 + \cdots + f_d p_d \leq n \rangle ,$$

where the “grades”  $f$ ’s are positive integer numbers and  $n = 0, 1, 2, \dots$ . In lattice space  $\mathcal{P}_{n,\vec{f}}^{(d)}$  defines a Newton pyramid.

**Definition 1.** Characteristic vector is a vector with components  $f_i$ :

$$\vec{f} = (f_1, f_2, \dots, f_d).$$

From geometrical point of view  $\vec{f}$  is normal vector to the base of the Newton pyramid. The characteristic vector for flag  $\mathcal{P}^{(d)}$  is

$$\vec{f}_0 = \underbrace{(1, 1, \dots, 1)}_d.$$

## 2.6 Case $G_2$

Take the Hamiltonian

$$\begin{aligned} \mathcal{H}_{G_2} = & \frac{1}{2} \sum_{i=1}^3 \left( -\frac{\partial^2}{\partial x_i^2} + \omega^2 x_i^2 \right) + \nu(\nu - 1) \sum_{i<j}^3 \frac{1}{(x_i - x_j)^2} \\ & + 3\mu(\mu - 1) \sum_{k<l, k,l \neq m}^3 \frac{1}{(x_k + x_l - 2x_m)^2}, \end{aligned} \quad (17)$$

where  $\omega, \nu, \mu$  are parameters. It describes the Wolfes model of three-body interacting system [9] or, in the Hamiltonian reduction nomenclature, the  $G_2$ -rational model. The symmetry of the model is dihedral group  $D_6$ . The ground state function is

$$\Psi_0 = \prod_{i<j}^3 |x_i - x_j|^\nu \prod_{k<l, k,l \neq m}^3 |x_i + x_j - 2x_k|^\mu e^{-\frac{1}{2}\omega \sum_{i=1}^3 x_i^2}.$$

Making the gauge rotation

$$h_{G_2} = (\Psi_0)^{-1} (\mathcal{H}_{G_2} - E) \Psi_0,$$

and changing variables

$$Y = \sum x_i, \quad y_i = x_i - \frac{1}{3}Y, \quad i = 1, 2, 3, \quad (x_1, x_2, x_3) \rightarrow (Y, \lambda_1, \lambda_2),$$

where

$$\lambda_1 = -y_1^2 - y_2^2 - y_1 y_2 \sim -r^2, \quad \lambda_2 = [y_1 y_2 (y_1 + y_2)]^2,$$

and separating the center-of-mass coordinate we arrive at

$$\begin{aligned} h_{G_2} = & \lambda_1 \partial_{\lambda_1 \lambda_1}^2 + 6\lambda_2 \partial_{\lambda_1 \lambda_2}^2 - \frac{4}{3} \lambda_1^2 \lambda_2 \partial_{\lambda_2 \lambda_2}^2 \\ & + \{2\omega \lambda_1 + 2[1 + 3(\mu + \nu)]\} \partial_{\lambda_1} + \left[ 6\omega \lambda_2 - \frac{4}{3}(1 + 2\mu)\lambda_1^2 \right] \partial_{\lambda_2}, \end{aligned}$$

which is the algebraic form of the Wolfes model. The eigenvalues of  $h_{G_2}$  are

$$\epsilon_{\{p\}} = 2\omega(p_1 + 3p_2).$$

It coincides to the spectrum of *anisotropic* harmonic oscillator with frequency ratio 1 : 3.

Separating the center-of-mass in (17) and introducing the polar coordinates  $(\varrho, \varphi)$  in the space of relative coordinates we arrive at the Hamiltonian

$$\tilde{\mathcal{H}}_{G_2}(\varrho, \varphi; \nu, \mu) = -\partial_r^2 - \frac{1}{r}\partial_r - \frac{1}{r^2}\partial_\varphi^2 + \omega^2 r^2 + \frac{9\nu(\nu-1)}{r^2 \cos^2 3\varphi} + \frac{9\mu(\mu-1)}{r^2 \sin^2 3\varphi}.$$

It is evident that the integral of motion which appears due to separation of variables in polar coordinates (cf. (1)) has the form

$$\mathcal{F} = -\partial_\varphi^2 + \frac{9\nu(\nu-1)}{\cos^2 3\varphi} + \frac{9\mu(\mu-1)}{\sin^2 3\varphi}. \quad (18)$$

It is evident that after gauge rotation with  $\Psi_0$  and change of variables to  $(\lambda_1, \lambda_2)$  the integral  $\mathcal{F}$  takes algebraic form.

The Hamiltonian  $h_{G_2}$  has infinitely many finite-dimensional invariant subspaces

$$\mathcal{P}_{n,(1,2)}^{(2)} = \langle \lambda_1^{p_1} \lambda_2^{p_2} \mid 0 \leq p_1 + 2p_2 \leq n \rangle, \quad n = 0, 1, 2, \dots,$$

hence the flag  $\mathcal{P}_{(1,2)}^{(2)}$  with the characteristic vector  $\vec{f} = (1, 2)$  is preserved by  $h_{G_2}$ . The eigenfunctions of  $h_{G_2}$  are elements of the flag of polynomials  $\mathcal{P}_{(1,2)}^{(2)}$ . Each subspace  $\mathcal{P}_{n,(1,2)}^{(2)} - \mathcal{P}_{n-1,(1,2)}^{(2)}$  contains  $\sim n$  eigenfunctions which is equal to length of the Newton line  $\mathcal{L}_n n = n \langle \lambda_1^{p_1} \lambda_2^{p_2} \mid p_1 + 2p_2 = n \rangle$ .

A natural question to ask: *What about hidden algebra?* Namely: *Does algebra exist for which  $\mathcal{P}_{n,(1,2)}^{(2)}$  is the space of (irreducible) representation?* Surprisingly, this algebra exists and it is, in fact, known.

Let us consider the Lie algebra spanned by seven generators

$$\begin{aligned} J^1 &= \partial_t, & J_n^2 &= t\partial_t - \frac{n}{3}, & J_n^3 &= 2u\partial_u - \frac{n}{3}, \\ J_n^4 &= t^2\partial_t + 2tu\partial_u - nt, & R_i &= t^i\partial_u, \quad i = 0, 1, 2, & L &\equiv (R_0, R_1, R_2). \end{aligned} \quad (19)$$

It is non-semi-simple algebra  $gl(2, \mathbb{R}) \ltimes \mathcal{R}^{(2)}$  (S. Lie [10] at  $n = 0$  and A. González-López et al. [11] at  $n \neq 0$  (case 24)). If the parameter  $n$  in (19) is a non-negative integer, it has

$$\mathcal{P}_n^{(2)} = \langle t^p u^q \mid 0 \leq (p + 2q) \leq n \rangle,$$

as common (reducible) invariant subspace. By adding

$$T_0^{(2)} = u\partial_t^2,$$

to  $gl(2, \mathbb{R}) \ltimes \mathcal{R}^{(2)}$  (see (19)), the action on  $\mathcal{P}_{n,(1,2)}^{(2)}$  gets irreducible. Multiple commutators of  $J_n^4$  with  $T_0^{(2)}$  generate new operators acting on  $\mathcal{P}_{n,(1,2)}^{(2)}$ ,

$$T_i^{(2)} \equiv \underbrace{[J^4, [J^4, [\dots J^4, T_0^{(2)}] \dots]]}_{i} = u\partial_t^{2-i} J_0 (J_0 + 1) \cdots (J_0 + i - 1), \quad i = 0, 1, 2,$$


where  $J_0 = t\partial_t + 2u\partial_u - n$ , and all of them are of degree 2. These new generators have a property of nilpotency,

$$T_i^{(2)} = 0, \quad i > 2,$$

and commutativity:

$$[T_i^{(2)}, T_j^{(2)}] = 0, \quad i, j = 0, 1, 2, \quad U \equiv (T_0^{(2)}, T_1^{(2)}, T_2^{(2)}). \quad (20)$$

(19) plus (20) span a linear space with a property of decomposition:  $g^{(2)} \doteq L \times (gl_2 \oplus J_0) \times U$  (see Fig. 2).



**Figure 2.** Triangular diagram relating the subalgebras  $L$ ,  $U$  and  $gl_2$ .  $P_2(gl_2)$  is a polynomial of the 2nd degree in  $gl_2$  generators. It is a generalization of the Gauss decomposition for semi-simple algebras.

Eventually, *infinite-dimensional, eleven-generated algebra (by (19) and  $J_0$  plus (20), so that the eight generators are the 1st order and three generators are of the 2nd order differential operators)* occurs. The Hamiltonian  $h_{G_2}$  can be rewritten in terms of the generators (19), (20) with the absence of the highest weight generator  $J_n^4$ ,

$$h_{G_2} = (J^2 + 3J^3)J^1 - \frac{2}{3}J^3R_2 + 2[3(\mu + \nu) + 1]J^1 + 2\omega J^2 + 3\omega J^3 - \frac{4}{3}(1 + 2\mu)R_2,$$

where  $J^{2,3} = J_0^{2,3}$ . Hence,  $gl(2, \mathbb{R}) \times \mathcal{R}^{(2)}$  is the hidden algebra of the Wolfes model.

- (i)  $G_2$  Hamiltonian admits two mutually-*non*-commuting integrals: of 2nd order as the result of the separation of radial variable  $r^2$  (see (18)) and of the 6th order. If  $\omega = 0$  the latter integral degenerates to the 3rd order integral (the square root can be calculated in closed form).
- (ii) Both integrals after gauge rotation with  $\Psi_0$  take in variables  $\lambda_{1,2}$  the algebraic form. Both preserve the same flag  $\mathcal{P}_{(1,2)}^{(2)}$ .
- (iii) Both integrals can be rewritten in term of generators of the algebra  $g^{(2)}$ : integral of 2nd order in terms of  $gl(2, \mathbb{R}) \times \mathcal{R}^{(2)}$  generators only and while one of the 6th order contains generators from  $\mathfrak{L}$  as well [13].

### 2.6.1 $sl(2)$ -quasi-exactly-solvable generalization

By adding to  $h_{G_2}$ , the operator (the same as for the Calogero and the  $BC_N$  models)

$$\delta h^{(\text{qes})} = 4(a\lambda_1^2 - \gamma) \frac{\partial}{\partial \lambda_1} - 4ak\lambda_1 + 2\omega k,$$

we get the operator  $h_{G_2} + \delta h^{(\text{qes})}$  having single finite-dimensional invariant subspace

$$\mathcal{P}_k = \langle \lambda_1^p \mid 0 \leq p \leq k \rangle.$$

Making a gauge rotation of  $h_{G_2} + \delta h^{(\text{qes})}$ , changing of variables  $(Y, \lambda_{1,2})$  back to the Cartesian coordinates and adding the center-of-mass the Hamiltonian becomes

$$\mathcal{H}_{G_2}^{(\text{qes})} = -\frac{1}{2} \sum_{i=1}^3 \left( \frac{\partial^2}{\partial x_i^2} - \omega^2 x_i^2 \right) + \nu(\nu - 1) \sum_{i < j}^3 \frac{1}{(x_i - x_j)^2}$$

$$\begin{aligned}
& + 3\mu(\mu - 1) \sum_{i < l, i, l \neq m}^3 \frac{1}{(x_i + x_l - 2x_m)^2} + \frac{4\gamma(\gamma + 3\mu + 3\nu)}{r^2} \\
& + a^2 r^6 + 2a\omega r^4 + 2a[2k - 3(\mu + \nu) - 2(\gamma + 1)]r^2,
\end{aligned}$$

for which  $(k + 1)$  eigenfunctions are of the form

$$\Psi_k^{(\text{qes})} = \prod_{i < j}^3 |x_i - x_j|^\nu \prod_{i < j; i, j \neq p}^3 |x_i + x_j - 2x_p|^\mu (r^2)^\gamma P_k(r^2) \exp \left[ -\frac{\omega}{2} \sum_{i=1}^3 x_i^2 - \frac{a}{4} r^4 \right],$$

where  $P_k$  is a polynomial of degree  $k$  in  $r^2$ .

It is worth noting that at  $a = 0$  the operator  $h_{G_2} + \delta h^{(\text{qes})}$  remains exactly-solvable, it preserves the flag of polynomials  $\mathcal{P}_{(1,2)}^{(2)}$  and the emerging Hamiltonian has a form of (17) with the extra term  $\frac{\Gamma}{r^2}$  in the potential. Its ground state eigenfunction is  $(r^2)^\gamma \Psi_0$ . It is the exactly-solvable generalization of the  $G_2$ -rational model (17) with the Weyl group  $W(G_2)$  as the discrete symmetry group,

$$\mathcal{H}_{W(G_2)} = \mathcal{H}_{G_2} + \frac{\Gamma}{r^2}.$$

## 2.7 Cases $F_4$ and $E_{6,7,8}$

In some details these four cases are described in [12].

## 2.8 Case $I_2(k)$

In some details this case is described in [13]. It is worth noting that although the Hamiltonian Reduction nomenclature is assigned to this case the parameter  $k$  takes any real value. Discrete symmetry group  $D_{2k}$  of the Hamiltonian appears for integer  $k$ .

## 2.9 Case $H_3$

The  $H_3$  rational Hamiltonian reads

$$\begin{aligned}
\mathcal{H}_{H_3} = & \frac{1}{2} \sum_{k=1}^3 \left[ -\frac{\partial^2}{\partial x_k^2} + \omega^2 x_k^2 + \frac{\nu(\nu - 1)}{x_k^2} \right] \\
& + 2\nu(\nu - 1) \sum_{\{i,j,k\}} \sum_{\mu_{1,2}=0,1} \frac{1}{[x_i + (-1)^{\mu_1} \varphi_+ x_j + (-1)^{\mu_2} \varphi_- x_k]^2},
\end{aligned} \tag{21}$$

where  $\{i, j, k\} = \{1, 2, 3\}$  and all even permutations,  $\omega, \nu$  are parameters and

$$\varphi_\pm = \frac{1 \pm \sqrt{5}}{2},$$

the golden ratio and its algebraic conjugate. Symmetry of the Hamiltonian (21) is the  $H_3$  Coxeter group (the full symmetry group of the icosahedron). It has the order 120. In total, the Hamiltonian (21) is symmetric with respect to the transformation

$$x_i \longleftrightarrow x_j, \quad \varphi_+ \longleftrightarrow \varphi_-.$$

The ground state is given by

$$\Psi_0 = \Delta_1^\nu \Delta_2^\nu \exp \left( -\frac{\omega}{2} \sum_{k=1}^3 x_k^2 \right), \quad E_0 = \frac{3}{2} \omega (1 + 10\nu),$$

where

$$\Delta_1 = \prod_{k=1}^3 x_k, \quad \Delta_2 = \prod_{\{i,j,k\}} \prod_{\mu_{1,2}=0,1} [x_i + (-1)^{\mu_1} \varphi_+ x_j + (-1)^{\mu_2} \varphi_- x_k].$$

Making the gauge rotation

$$h_{H_3} = -2(\Psi_0)^{-1}(\mathcal{H}_{H_3} - E_0)(\Psi_0),$$

we arrive at new spectral problem

$$h_{H_3} \phi(x) = -2\epsilon \phi(x).$$

After changing variables ( $x_{1,2,3} \rightarrow \tau_{1,2,3}$ ):

$$\begin{aligned} \tau_1 &= x_1^2 + x_2^2 + x_3^2 = r^2, \\ \tau_2 &= -\frac{3}{10}(x_1^6 + x_2^6 + x_3^6) + \frac{3}{10}(2 - 5\varphi_+)(x_1^2 x_2^4 + x_2^2 x_3^4 + x_3^2 x_1^4) \\ &\quad + \frac{3}{10}(2 - 5\varphi_-)(x_1^2 x_3^4 + x_2^2 x_1^4 + x_3^2 x_2^4) - \frac{39}{5}, \\ \tau_3 &= \frac{2}{125}(x_1^{10} + x_2^{10} + x_3^{10}) + \frac{2}{25}(1 + 5\varphi_-)(x_1^8 x_2^2 + x_2^8 x_3^2 + x_3^8 x_1^2) \\ &\quad + \frac{2}{25}(1 + 5\varphi_+)(x_1^8 x_3^2 + x_2^8 x_1^2 + x_3^8 x_2^2) + \frac{4}{25}(1 - 5\varphi_-)(x_1^6 x_2^4 + x_2^6 x_3^4 + x_3^6 x_1^4) \\ &\quad + \frac{4}{25}(1 - 5\varphi_+)(x_1^6 x_3^4 + x_2^6 x_1^4 + x_3^6 x_2^4) - \frac{112}{25}(x_1^6 x_2^2 x_3^2 + x_2^6 x_3^2 x_1^2 + x_3^6 x_1^2 x_2^2) \\ &\quad + \frac{212}{25}(x_1^2 x_2^4 x_3^4 + x_2^2 x_3^4 x_1^4 + x_3^2 x_1^4 x_2^4), \end{aligned}$$

in the gauge-rotated Hamiltonian, it emerges in the algebraic form [14]

$$h_{H_3} = \sum_{i,j=1}^3 A_{ij} \frac{\partial^2}{\partial \tau_i \partial \tau_j} + \sum_{j=1}^3 B_j \frac{\partial}{\partial \tau_j},$$

where

$$\begin{aligned} A_{11} &= 4\tau_1, & A_{12} &= 12\tau_2, & A_{13} &= 20\tau_3, \\ A_{22} &= -\frac{48}{5}\tau_1^2 \tau_2 + \frac{45}{2}\tau_3, & A_{23} &= \frac{16}{15}\tau_1 \tau_2^2 - 24\tau_1^2 \tau_3, & A_{33} &= -\frac{64}{3}\tau_1 \tau_2 \tau_3 + \frac{128}{45}\tau_2^3, \\ B_1 &= 6 + 60\nu - 4\omega\tau_1, & B_2 &= -\frac{48}{5}(1 + 5\nu)\tau_1^2 - 12\omega\tau_2, \\ B_3 &= -\frac{64}{15}(2 + 5\nu)\tau_1 \tau_2 - 20\omega\tau_3, \end{aligned}$$

which is amazingly simple comparing the quite complicated and lengthy form of the original Hamiltonian (21). The Hamiltonian  $h_{H_3}$  preserves infinitely-many spaces

$$\mathcal{P}_n^{(1,2,3)} = \langle \tau_1^{n_1} \tau_2^{n_2} \tau_3^{n_3} \mid 0 \leq n_1 + 2n_2 + 3n_3 \leq n \rangle, \quad n \in \mathbb{N},$$

with characteristic vector is  $(1, 2, 3)$ , they form an infinite flag. The spectrum of  $h_{H_3}$  is given by

$$\epsilon_{p_1, p_2, p_3} = 2\omega(p_1 + 3p_2 + 5p_3), \quad p_i = 0, 1, 2, \dots,$$

with degeneracy  $p_1 + 3p_2 + 5p_3 = \text{integer}$ . It corresponds to the anisotropic harmonic oscillator with frequency ratios  $1 : 3 : 5$ . Eigenfunctions  $\phi_{n,i}$  of  $h_{H_3}$  are elements of  $\mathcal{P}_n^{(1,2,3)}$ , The number of eigenfunctions in  $\mathcal{P}_n^{(1,2,3)}$  is maximal possible – it is equal to dimension of  $\mathcal{P}_n^{(1,2,3)}$ .

The space  $\mathcal{P}_n^{(1,2,3)}$  is finite-dimensional representation space of a Lie algebra of differential operators which we call the  $h^{(3)}$  algebra. It is infinite-dimensional but finitely generated algebra of differential operators with 30 generating elements of 1<sup>st</sup> (14), 2<sup>nd</sup> (10) and 3<sup>rd</sup> (5) orders, respectively, plus one of zeroth order. They span 5 + 5 Abelian (conjugated) subalgebras of lowering and raising generators  $L$  and  $U^2$  and one of the Cartan type algebra  $B$  (for details see [14]). The algebra  $B = g\ell_2 \oplus I_2$ , where  $I_2$  is two-dimensional ideal. The Hamiltonian  $h_{H_3}$  can be rewritten in terms of the generators of the  $h^{(3)}$ -algebra.

By adding to  $h_{H_3}$  (21) the operator (6) in the variable  $\tau_1$  (of the same type as for the Calogero,  $BC_N$  and  $G_2$  models) we get the operator  $h_{H_3} + \delta h^{(\text{qes})}$  which has the finite-dimensional invariant subspace

$$\mathcal{P}_k = \langle \tau_1^p \mid 0 \leq p \leq k \rangle,$$

of the dimension  $(k + 1)$ . Hence, this operator is quasi-exactly-solvable. Making the gauge rotation of this operator and changing variables  $\tau$  back to Cartesian ones we arrive at the quasi-exactly-solvable Hamiltonian of a similar type as for the Calogero,  $BC_N$  and  $G_2$  models [14]. By adding to  $h_{H_3}$  (21) the operator  $4\gamma \frac{\partial}{\partial \tau_1}$  we preserve the property of exact-solvability. This operator preserves the flag  $\mathcal{P}^{(1,2,3)}$  and the emerging Hamiltonian has a form of (21) with the extra term  $\frac{\Gamma}{r^2}$  in the potential. Its ground state eigenfunction is  $(r^2)^\gamma \Psi_0$ . It is the exactly-solvable generalization of the  $H_3$ -rational model (21) with the Coxeter group  $H_3$  as the discrete symmetry group,

$$\mathcal{H}_{H_3} = \mathcal{H}_{H_3} + \frac{\Gamma}{r^2}.$$

## 2.10 Case $H_4$

The  $H_4$  rational Hamiltonian reads

$$\begin{aligned} \mathcal{H}_{H_4} = & \frac{1}{2} \sum_{k=1}^4 \left[ -\frac{\partial^2}{\partial x_k^2} + \omega^2 x_k^2 + \frac{\nu(\nu-1)}{x_k^2} \right] \\ & + 2\nu(\nu-1) \sum_{\mu_{2,3,4}=0,1} \frac{1}{[x_1 + (-1)^{\mu_2} x_2 + (-1)^{\mu_3} x_3 + (-1)^{\mu_4} x_4]^2} \\ & + 2\nu(\nu-1) \sum_{\{i,j,k,l\}} \sum_{\mu_{1,2}=0,1} \frac{1}{[x_i + (-1)^{\mu_1} \varphi_+ x_j + (-1)^{\mu_2} \varphi_- x_k + 0 \cdot x_l]^2}, \end{aligned} \quad (22)$$

where  $\{i, j, k, l\} = \{1, 2, 3, 4\}$  and all even permutations,  $\omega, \nu$  are parameters and

$$\varphi_{\pm} = \frac{1 \pm \sqrt{5}}{2},$$

the golden ratio and its algebraic conjugate. Symmetry of the Hamiltonian (22) is the  $H_4$  Coxeter group (the symmetry group of the 600-cell). It has order 14400. In total, the Hamiltonian (22) is symmetric with respect to the transformation

$$x_i \longleftrightarrow x_j, \quad \varphi_+ \longleftrightarrow \varphi_-.$$

---

<sup>2</sup>These subalgebras can be divided into pairs. In every pair the elements of different subalgebras are related via a certain conjugation.



The ground state function and its eigenvalue are

$$\Psi_0 = \Delta_1^\nu \Delta_2^\nu \Delta_3^\nu \exp\left(-\frac{\omega}{2} \sum_{k=1}^4 x_k^2\right), \quad E_0 = 2\omega(1 + 30\nu),$$

where

$$\begin{aligned} \Delta_1 &= \prod_{k=1}^4 x_k, & \Delta_2 &= \prod_{\mu_{2,3,4}=0,1} [x_1 + (-1)^{\mu_2} x_2 + (-1)^{\mu_3} x_3 + (-1)^{\mu_4} x_4], \\ \Delta_3 &= \prod_{\{i,j,k,l\}} \prod_{\mu_{1,2}=0,1} [x_i + (-1)^{\mu_1} \varphi_+ x_j + (-1)^{\mu_2} \varphi_- x_k + 0 \cdot x_l]. \end{aligned}$$

Making a gauge rotation of the Hamiltonian

$$h_{H_4} = -2(\Psi_0)^{-1}(\mathcal{H}_{H_4} - E_0)(\Psi_0),$$

and introducing new variables  $\tau_{1,2,3,4}$ , which are invariant with respect to the  $H_4$  Coxeter group, in a form of polynomials in  $x$  of degrees 2, 12, 20, 30 (degrees of  $H_4$ ), we arrive at the Hamiltonian in the algebraic form [15]

$$h_{H_4} = \sum_{i,j=1}^4 A_{ij} \frac{\partial^2}{\partial \tau_i \partial \tau_j} + \sum_{j=1}^4 B_j \frac{\partial}{\partial \tau_j}, \quad (23)$$

where

$$\begin{aligned} A_{11} &= 4\tau_1, & A_{12} &= 24\tau_2, & A_{13} &= 40\tau_3, & A_{14} &= 60\tau_4, \\ A_{22} &= 88\tau_1\tau_3 + 8\tau_1^5\tau_2, & A_{23} &= -4\tau_1^3\tau_2^2 + 24\tau_1^5\tau_3 - 8\tau_4, \\ A_{24} &= 10\tau_1^2\tau_2^3 + 60\tau_1^4\tau_2\tau_3 + 40\tau_1^5\tau_4 - 600\tau_3^2, & A_{33} &= -\frac{38}{3}\tau_1\tau_2^3 + 28\tau_1^3\tau_2\tau_3 - \frac{8}{3}\tau_1^4\tau_4, \\ A_{34} &= 210\tau_1^2\tau_2^2\tau_3 + 60\tau_1^3\tau_2\tau_4 - 180\tau_1^4\tau_3^2 + 30\tau_2^4, \\ A_{44} &= -2175\tau_1\tau_2^3\tau_3 - 450\tau_1^2\tau_2^2\tau_4 - 1350\tau_1^3\tau_2\tau_3^2 - 600\tau_1^4\tau_3\tau_4, \\ B_1 &= 8(1 + 30\nu) - 4\omega\tau_1, & B_2 &= 12(1 + 10\nu)\tau_1^5 - 24\omega\tau_2, \\ B_3 &= 20(1 + 6\nu)\tau_1^3\tau_2 - 4\omega\tau_3, & B_4 &= 15(1 - 30\nu)\tau_1^2\tau_2^2 - 450(1 + 2\nu)\tau_1^4\tau_3 - 60\omega\tau_4, \end{aligned}$$

which is amazingly simple comparing the very complicated and lengthy form of the original Hamiltonian (22). It is easy to check that the algebraic operator  $h_{H_4}$  preserves infinitely-many finite-dimensional invariant subspaces


$$\mathcal{P}_n^{(1,5,8,12)} = \langle \tau_1^{n_1} \tau_2^{n_2} \tau_3^{n_3} \tau_4^{n_4} \mid 0 \leq n_1 + 5n_2 + 8n_3 + 12n_4 \leq n \rangle, \quad n \in \mathbb{N},$$

all of them with the same characteristic vector  $(1, 5, 8, 12)$ , they form the infinite flag. The spectrum of the Hamiltonian  $h_{H_4}$  (23) has a form

$$\epsilon_{k_1, k_2, k_3, k_4} = 2\omega(k_1 + 6k_2 + 10k_3 + 15k_4), \quad k_i = 0, 1, 2, \dots,$$

with degeneracy  $k_1 + 6k_2 + 10k_3 + 15k_4 = \text{integer}$ . It corresponds to the anisotropic harmonic oscillator with frequency ratios  $1 : 6 : 10 : 15$ . Eigenfunctions  $\phi_{n,i}$  of  $h_{H_4}$  are elements of  $\mathcal{P}_n^{(1,5,8,12)}$ . The number of eigenfunctions in  $\mathcal{P}_n^{(1,5,8,12)}$  is equal to dimension of  $\mathcal{P}_n^{(1,5,8,12)}$ .

The space  $\mathcal{P}_n^{(1,5,8,12)}$  is a finite-dimensional representation space of a Lie algebra of differential operators which we call the  $h^{(4)}$  algebra. It is infinite-dimensional but finitely generated algebra



**Figure 3.** Triangular diagram relating the subalgebras  $L$ ,  $U$  and  $B$ .  $P_p(B)$  is a polynomial of the  $p$ th degree in  $B$  generators. It is a generalization of the Gauss decomposition for semi-simple algebras where  $p = 1$ .

of differential operators, with 183 generating elements of 1<sup>st</sup> (54), 2<sup>nd</sup> (24), 3<sup>rd</sup> (18), 4<sup>rd</sup> (18), 5<sup>rd</sup> (28), 6<sup>rd</sup> (5), 8<sup>rd</sup> (20), 9<sup>rd</sup> (1), 12<sup>rd</sup> (14) orders, respectively, plus one of zeroth order. They span 26 + 26 Abelian (conjugated) subalgebras of lowering and raising generators  $L$  and  $U$ <sup>3</sup> and one of the Cartan type algebra  $B$  (for details see [16]). The algebra  $B = \mathfrak{gl}_2 \oplus I_3$ , where  $I_3$  is three-dimensional ideal. The Hamiltonian  $h_{H_4}$  can be rewritten in terms of the generators of the  $h^{(4)}$ -algebra.

By adding to  $h_{H_4}$  (22) the operator (6) in the variable  $\tau_1$  (of the same type as for the Calogero,  $BC_N$  and  $G_2$ ,  $H_3$  models) we get the operator  $h_{H_4} + \delta h^{(\text{ges})}$ , which has the finite-dimensional invariant subspace

$$\mathcal{P}_k = \langle \tau_1^p \mid 0 \leq p \leq k \rangle,$$

of dimension  $(k + 1)$ . Hence, this operator is quasi-exactly-solvable. Making the gauge rotation of this operator and changing variables  $\tau$  back to Cartesian ones we arrive at the quasi-exactly-solvable Hamiltonian of a similar type as for the Calogero,  $BC_N$  and  $G_2$  models [15]. By adding to  $h_{H_4}$  (22) the operator  $4\gamma \frac{\partial}{\partial \tau_1}$  we preserve the property of exact-solvability. This operator preserves the flag  $\mathcal{P}^{(1,5,8,12)}$  and the emerging Hamiltonian has a form of (22) with the extra term  $\frac{\Gamma}{r^2}$  in the potential. Its ground state eigenfunction is  $(r^2)^\gamma \Psi_0$ . It is the exactly-solvable generalization of the  $H_4$ -rational model (22) with the Coxeter group  $H_4$  as the discrete symmetry group,

$$\mathcal{H}_{H_4} = \mathcal{H}_{H_4} + \frac{\Gamma}{r^2}.$$

### 3 Conclusions

- For rational Hamiltonians for all classical  $A_N$ ,  $BC_N$  and exceptional root spaces  $G_2$ ,  $F_4$ ,  $E_{6,7,8}$  (also trigonometric) and non-crystallographic  $H_{3,4}$ ,  $I_2(k)$  there exists an algebraic form after gauging away the ground state eigenfunction, and changing variables to symmetric (invariant) variables. Their eigenfunctions are polynomials in these variables. They are orthogonal with the squared ground state eigenfunction as the weight factor.
- Their hidden algebras are  $\mathfrak{gl}(N + 1)$  for the case of classical  $A_N$ ,  $BC_N$  and *new* infinite-dimensional but finite-generated algebras of differential operators for all other cases. All these algebras have finite-dimensional invariant subspace(s) in polynomials.
- The generating elements of any such hidden algebra can be grouped into an even number of (conjugated) Abelian algebras  $L_i$ ,  $U_i$  and one Lie algebra  $B$ . They obey a (generalized) Gauss decomposition (see Fig. 3). A description of all these algebras will be given elsewhere.

There are two solvable potentials in  $1D$  in  $[0, \infty)$  which can be generalized to  $D$ .

<sup>3</sup>These subalgebras can be divided into pairs. In every pair the elements of different subalgebras are related via a certain conjugation.

### 3.1 General view (exact-solvability)

ES-case generalization:

$$\omega^2 r^2 + \frac{\gamma}{r^2} \rightarrow \omega^2 r^2 + \frac{\gamma(\Omega)}{r^2}$$

(a generalization by replacing the symmetry  $O(N)$  by its discrete subgroup of symmetry given by Weyl (Coxeter) group).

### 3.2 General view (quasi-exact-solvability)

QES-case generalization:

$$\omega^2 r^2 + \frac{\gamma}{r^2} + ar^6 + br^4 \rightarrow \tilde{\omega}^2 r^2 + \frac{\tilde{\gamma}(\Omega)}{r^2} + ar^6 + br^4$$

(a generalization by replacing the symmetry  $O(N)$  by its discrete subgroup of symmetry given by Weyl (Coxeter) group).

In conclusion I must emphasize that the algebraic nature of all above-considered systems was revealed when:

*The invariants of the discrete group of symmetry of a system  
are taken as variables (space of orbits).*

### Acknowledgements

This paper is based on the talk given to honor Professor Willard Miller, who was always an example for me of love and dedication for science.

### References

- [1] Evans N.W., Group theory of the Smorodinsky–Winternitz system, *J. Math. Phys.* **32** (1991), 3369–3375.
- [2] Turbiner A.V., Quasi-exactly-solvable problems and the  $sl(2)$  group, *Comm. Math. Phys.* **118** (1988), 467–474.
- [3] Calogero F., Solution of a three-body problem in one dimension, *J. Math. Phys.* **10** (1969), 2191–2196.  
Calogero F., Solution of the one-dimensional  $N$ -body problem with quadratic and/or inversely quadratic pair potentials, *J. Math. Phys.* **12** (1971), 419–436.
- [4] Rühl W., Turbiner A., Exact solvability of the Calogero and Sutherland models, *Modern Phys. Lett. A* **10** (1995), 2213–2221, [hep-th/9506105](#).
- [5] Minzoni A., Rosenbaum M., Turbiner A., Quasi-exactly-solvable many-body problems, *Modern Phys. Lett. A* **11** (1996), 1977–1984, [hep-th/9606092](#).
- [6] Olshanetsky M.A., Perelomov A.M., Quantum integrable systems related to Lie algebras, *Phys. Rep.* **94** (1983), 313–404.
- [7] Oshima T., Completely integrable systems associated with classical root systems, *SIGMA* **3** (2007), 061, 50 pages, [math-ph/0502028](#).
- [8] Brink L., Turbiner A., Wyllard N., Hidden algebras of the (super) Calogero and Sutherland models, *J. Math. Phys.* **39** (1998), 1285–1315, [hep-th/9705219](#).
- [9] Wolfes J., On the three-body linear problem with three-body interaction, *J. Math. Phys.* **15** (1974), 1420–1424.
- [10] Lie S., Gruppenregister, Vol. 5, B.G. Teubner, Leipzig, 1924, 767–773.

- 
- [11] González-López A., Kamran N., Olver P.J., Quasi-exactly-solvable Lie algebras of the first order differential operators in two complex variables, *J. Phys. A: Math. Gen.* **24** (1991), 3995–4008.  
González-López A., Kamran N., Olver P.J., Lie algebras of differential operators in two complex variables, *Amer. J. Math.* **114** (1992), 1163–1185.
- [12] Borekov K.G., Turbiner A.V., Lopez Vieyra J.C., Solvability of the Hamiltonians related to exceptional root spaces: rational case, *Comm. Math. Phys.* **260** (2005), 17–44, [hep-th/0407204](#).
- [13] Tremblay F., Turbiner A.V., Winternitz P., An infinite family of solvable and integrable quantum systems on a plane, *J. Phys. A: Math. Theor.* **42** (2009), 242001, 10 pages, [arXiv:0904.0738](#).
- [14] García M.A.G., Turbiner A.V., The quantum  $H_3$  integrable system, *Internat. J. Modern Phys. A* **25** (2010), 5567–5594, [arXiv:1007.0737](#).
- [15] García M.A.G., Turbiner A.V., The quantum  $H_4$  integrable system, *Modern Phys. Lett. A* **26** (2011), 433–447, [arXiv:1011.2127](#).
- [16] García M.A.G. Los Sistemas Integrables  $H_3$  y  $H_4$ , PhD Thesis, UNAM, 2011 (in Spanish).