

T -Path Formula and Atomic Bases for Cluster Algebras of Type D^*

Emily GUNAWAN and Gregg MUSIKER

School of Mathematics, University of Minnesota, Minneapolis, MN 55455, USA

E-mail: egunawan@umn.edu, musiker@umn.edu

URL: <http://umn.edu/home/egunawan>, <http://math.umn.edu/~musiker>

Received September 24, 2014, in final form July 09, 2015; Published online July 28, 2015

<http://dx.doi.org/10.3842/SIGMA.2015.060>

Abstract. We extend a T -path expansion formula for arcs on an unpunctured surface to the case of arcs on a once-punctured polygon and use this formula to give a combinatorial proof that cluster monomials form the atomic basis of a cluster algebra of type D .

Key words: cluster algebra; triangulated surface; atomic basis

2010 Mathematics Subject Classification: 13F60; 05E15; 16S99

1 Introduction

Cluster algebras, introduced by Fomin and Zelevinsky [11] in 2000, are commutative algebras equipped with a distinguished set of generators, the *cluster variables*. The cluster variables are grouped into sets of constant cardinality n , the *clusters*, and the integer n is called the *rank* of the cluster algebra. Starting with an initial *seed* (\mathbf{x}, B) , that is, an initial cluster \mathbf{x} together with a skew-symmetrizable integer $n \times n$ *exchange matrix* $B = (b_{ij})$, the set of cluster variables is obtained by repeated application of so called *mutations*. To be more precise, let $\{x_1, x_2, \dots, x_n\}$ be indeterminates over \mathbb{Z} and let $\mathcal{F} = \mathbb{Q}(x_1, x_2, \dots, x_n)$. For every $k = 1, 2, \dots, n$, the mutation $\mu_k(\mathbf{x})$ of the cluster $\mathbf{x} = \{x_1, x_2, \dots, x_n\}$ is a new cluster $\mu(\mathbf{x}) = (\mathbf{x} \setminus \{x_k\}) \cup \{x'_k\}$ obtained from \mathbf{x} by replacing the cluster variable x_k with the new cluster variable

$$x'_k = \frac{1}{x_k} \left(\prod_{b_{ik} > 0} x_i^{b_{ik}} + \prod_{b_{ik} < 0} x_i^{-b_{ik}} \right)$$

in \mathcal{F} . Mutations also change the attached matrix B , see [11].

The set of all cluster variables is the union of all clusters obtained from an initial cluster \mathbf{x} by repeated mutations. Note that this set may be infinite.

It is clear from the construction that every cluster variable is a rational function in the initial cluster variables x_1, x_2, \dots, x_n . In [11] it is shown that every cluster variable u is actually a Laurent polynomial in the x_i , that is, u can be written as a reduced fraction

$$u = \frac{f(x_1, x_2, \dots, x_n)}{\prod_{i=1}^n x_i^{d_i}}, \tag{1.1}$$

where $f \in \mathbb{Z}[x_1, x_2, \dots, x_n]$ and $d_i \geq 0$. The right hand side of equation (1.1) is called the *cluster expansion of u in \mathbf{x}* .

*This paper is a contribution to the Special Issue on New Directions in Lie Theory. The full collection is available at <http://www.emis.de/journals/SIGMA/LieTheory2014.html>

The *coefficient-free cluster algebra* $\mathcal{A}(\mathbf{x}, B)$ is the subring of $\mathcal{F} = \mathbb{Q}(x_1, x_2, \dots, x_n)$ generated by the cluster variables. When the set of cluster variables is finite, we say that $\mathcal{A}(\mathbf{x}, B)$ is of *finite type*.

We are interested in cluster algebras arising from *bordered surfaces with marked points* [8, 9, 10, 13]. In particular, we study cluster algebras of type D_n (type D for short), which are of finite type (as classified in [12]) and correspond to once-punctured n -gons, as also described in detail in [20]. Other, related work on type D cluster algebra combinatorial models include [1, 3, 12].

Our first result is a Laurent polynomial expansion formula for cluster variables arising from a once-punctured polygon in terms of certain paths (called T^o -paths) on an ideal triangulation T^o of the surface. This is an extension of the T -path formula (which we call the T^o -path) given for any unpunctured surface by Schiffler and Thomas [21, 22]. Our proof takes advantage of two facts proven by the second author, Schiffler, and Williams: (1) a Laurent polynomial expansion formula for cluster variables arising from any surface in terms of perfect matchings of a *snake graph* [17], and (2) a bijection between these perfect matchings and T^o -paths arising from any unpunctured surface [16]. An application of the T^o -path formula for type D is discussed in the next paragraph.

Our second result is a specific case of a result of [4, 6], proven by representation theoretic methods, that the basis consisting of all *cluster monomials* is in fact the *atomic basis* for any skew-symmetric cluster algebra of finite type (see Section 5.1). We give a combinatorial proof of this fact for *coefficient-free* cluster algebras of type D . Our proof relies heavily on the T^o -path formula for type D and is inspired by Dupont and Thomas' work in [7] on atomic bases for cluster algebras of type A and \tilde{A} .

In Section 2, we provide background material on ideal triangulations and tagged triangulations, focusing on the case of once-punctured polygons. Section 3 presents our first result (Theorem 3.8), an extension of the T^o -path formula of [21, 22] to once-punctured polygons. We give the proof of this T^o -path formula in Section 4. Finally, in Section 5, we give our second and main result (Theorem 5.8), which is a combinatorial proof, using the T^o -path formula, that the cluster monomials form the atomic basis for a coefficient-free type D cluster algebra.

2 Background: cluster algebras arising from once-punctured disks

For the reader's convenience, we begin by reviewing terminology arising in the theory of cluster algebras from marked surfaces from [10, Sections 2 and 7]. We restrict our attention to the case of a once-punctured polygon, which often simplifies the notation. Let \mathcal{C}_n denote a once-punctured n -gon, i.e., a disk with a marked point (called the puncture) in the interior and n marked points on the boundary.

Definition 2.1 (ordinary arcs). A *boundary edge* of \mathcal{C}_n is a segment of the boundary between two consecutive boundary marked points. An *ordinary arc* γ of \mathcal{C}_n is a curve (considered up to isotopy) in \mathcal{C}_n such that the endpoints of γ are marked points, γ does not cross itself except possibly at its endpoints, γ does not cross the boundary of \mathcal{C}_n except possibly at its endpoints, and γ is not contractible to a marked point or homotopic to a boundary edge.

A *radius* is an arc between a boundary marked point and the puncture. Following [7], a *peripheral arc* is an arc with both endpoints on the boundary. An ℓ -*loop* is a loop cutting out a monogon with a sole puncture inside it (i.e., as illustrated by loop ℓ in Fig. 1(c)). An ℓ -loop is considered a peripheral, ordinary arc.

Definition 2.2 (compatibility of ordinary arcs, ideal triangulations). Two distinct ordinary arcs are said to be *compatible* if they do not intersect except possibly at endpoints. Also, each arc




Figure 1. Possible types of ideal triangles.




Figure 2. An ideal triangulation T^o and the corresponding tagged triangulation $T = \iota(T^o)$. The ℓ -loop ℓ and the corresponding tagged radius $\iota(\ell)$ (tagged notched at the puncture) are both in gray.

is compatible with itself. A maximal (by inclusion) collection of distinct, pairwise compatible ordinary arcs is called an *ideal triangulation*. The ordinary arcs of an ideal triangulation cut the surface into *ideal triangles* (see Fig. 1). We call an ℓ -loop and the radius it encloses a *self-folded triangle* (Fig. 1(c)).

Remark 2.3. Three possible types of ideal triangles can appear in an ideal triangulation of \mathcal{C}_n : an ordinary triangle with 3 distinct vertices and 3 distinct sides (Fig. 1(a)), an ideal triangle with 2 distinct vertices and 3 distinct sides (Fig. 1(b)), and finally a self-folded triangle (Fig. 1(c)). The ideal triangulation of Fig. 2 (left) contains all 3 types of \mathcal{C}_n -ideal triangles. Note that the one-vertex ideal triangle (Fig. 1(d)) cannot appear.

Definition 2.4 (tagged arcs). A *tagged arc* of \mathcal{C}_n is obtained by marking (“tagging”) each endpoint of an ordinary arc (that is not an ℓ -loop) β either *plain* or *notched* such that the endpoints of β on the boundary must be tagged plain. A notching is indicated by a bow tie (see Fig. 2). Note that a tagged arc never cuts out a once-punctured monogon, i.e., an ℓ -loop is *not* a tagged arc (even for other punctured surfaces).

Remark 2.5. Every tagged arc β of \mathcal{C}_n belongs to one of the following three classes:

- β is a radius tagged plain at both endpoints (which we call a plain radius).
- β is a radius tagged notched at the puncture and plain at the boundary (which we call a notched radius).
- β is a peripheral arc connecting distinct endpoints tagged plain at both endpoints.

Definition 2.6 (compatibility of tagged arcs, tagged triangulations, and multi-tagged triangulations of \mathcal{C}_n). The following is a complete list of possible compatible pairs $\{\alpha, \beta\}$ of tagged arcs of \mathcal{C}_n :

- α and β are two peripheral arcs (tagged plain at all endpoints) that do not intersect in the interior of \mathcal{C}_n .
- α and β are a peripheral arc and a radius (tagged plain at boundary endpoints) that do not intersect in the interior of \mathcal{C}_n .



(a) Wheel-like triangulation around the puncture. (b) A self-folded triangle around the puncture where \aleph, \beth are peripheral arcs or boundary edges.

Figure 3. Local triangulations around the puncture. The shaded gray area consists only of peripheral arcs and boundary edges, each sector forming a polygon.

- α and β are two radii both adjacent to the same boundary marked point (tagged plain) but α is tagged plain at the puncture and β is tagged notched at the puncture.
- α and β are two radii with distinct boundary endpoints (tagged plain at boundary endpoints) and tagged the same way at the puncture.
- α and β are equal.

A maximal (by inclusion) collection of distinct, pairwise compatible tagged arcs is called a *tagged triangulation*. A collection of pairwise compatible tagged arcs (considered with multiplicity) is called a *multi-tagged triangulation*. A multi-tagged triangulation Σ is *compatible* with T if $\sigma \in T$ for every tagged arc $\sigma \in \Sigma$.

Definition 2.7 (representing ordinary arcs as tagged arcs). Any ordinary arc β can be represented by a tagged arc $\iota(\beta)$ as follows. Suppose β is an ℓ -loop (based at marked point v) which encloses a radius r , where r is the unique (ordinary) arc connecting v and the puncture P . Then $\iota(\beta)$ is obtained by tagging r plain at v and notched at P . Otherwise, $\iota(\beta)$ is simply β with both endpoints tagged plain. Fig. 2 shows an ideal triangulation T° of a once-punctured quadrilateral and its corresponding tagged triangulation $\iota(T^\circ)$.

As a convention, we will usually denote a tagged triangulation by T and an ideal triangulation by T° . Unless otherwise stated, $T = \iota(T^\circ)$.

Theorem 2.8 ([10, Theorem 7.11, Example 6.7]). *A cluster algebra is associated to \mathcal{C}_n as follows. Choose a tagged triangulation $T_{\text{init}} = \{\tau_1, \dots, \tau_n\}$ of \mathcal{C}_n . Let \mathcal{A} be the cluster algebra given by the initial seed $(\mathbf{x}_{T_{\text{init}}}, B_{T_{\text{init}}})$ where $\mathbf{x}_{T_{\text{init}}} = \{x_{\tau_1}, \dots, x_{\tau_n}\}$ is the cluster seed associated to T_{init} and $B_{T_{\text{init}}}$ is the exchange matrix corresponding to T_{init} (see Definition 4.1 in [10]). Then the tagged triangulations of \mathcal{C}_n are in bijection with the (unlabeled) seeds (\mathbf{x}_T, B_T) of \mathcal{A} , and the tagged arcs γ of \mathcal{C}_n are in bijection with the cluster variables (so we can denote each cluster variable by x_γ or $x(\gamma)$, where γ is a tagged arc). Therefore, a multi-tagged triangulation Γ corresponds to a cluster monomial (denoted x_Γ , see Definition 5.1).*

If r is a plain radius and ℓ is the ℓ -loop enclosing r , denote $x_\ell := x_r x_{r(p)}$. If β is a boundary edge, set $x_\beta := 1$. When we say the T -expansion of a cluster variable $x(\gamma)$, we mean the cluster expansion of $x(\gamma)$ in the variables of the seed \mathbf{x}_T , i.e., a Laurent polynomial in the variables of \mathbf{x}_T , see equation (1.1).

Remark 2.9. Due to the following proposition, it is enough to work with only two types of tagged triangulations T : one where T has all plain-tagged radii (so that T° has a local wheel-like triangulation as in Fig. 3(a)), and one where T has two parallel radii, one tagged plain and the other tagged notched at the puncture (so that T° has a self-folded triangle as in Fig. 3(b)).




Figure 4. A triangulation T° and an arc γ of a once-punctured square. The first, second, third, and fourth ideal triangles crossed by γ are denoted by Δ_0 , Δ_1 , Δ_2 , and Δ_3 .

Proposition 2.10 ([17, Proposition 3.15]). *Suppose $T = \{\tau_1, \dots, \tau_n\}$ is a tagged triangulation of \mathcal{C}_n . Let $\gamma^{(p)}$ denote the arc obtained from γ by changing the notching at the puncture p . Let $T^{(p)}$ denote the tagged triangulation that is obtained from T by replacing each $\tau \in T$ by $\tau^{(p)}$. Let $x(\gamma)$ be the T -expansion of the cluster variable corresponding to γ . Then*

$$x(\gamma^{(p)}) = x(\gamma)|_{x(\tau_i) \leftarrow x(\tau_i^{(p)})}$$

is the $T^{(p)}$ -expansion of the cluster variable corresponding to $\gamma^{(p)}$.

3 (T°, γ) -path expansion formula

We extend Schiffler and Thomas' work [21, 22] to once-punctured disks \mathcal{C}_n . Following [21, Section 3], we will use the following setup throughout the rest of the paper.

- Let S be an unpunctured surface or $S = \mathcal{C}_n$. Let T° be an ideal triangulation of S and let $\gamma \notin T^\circ$ be an ordinary arc of S . Recall that an ℓ -loop is considered an ordinary arc.
- Choose an orientation on γ , and let s^γ and t^γ be the starting point and the finishing point of γ . Denote by

$$s^\gamma = p_0^\gamma, p_1^\gamma, p_2^\gamma, \dots, p_{d+1}^\gamma = t^\gamma$$

the points of intersection of γ and T° in order. Since γ is considered up to homotopy, we pick a representative so that d is minimal. Let i_1, i_2, \dots, i_d be such that $\tau_{i_k}^\gamma$ is the arc of T° containing p_k^γ . See Fig. 4, where $\tau_{i_1}^\gamma = 1$, $\tau_{i_2}^\gamma = 2$, and $\tau_{i_3}^\gamma = 3$, and see Fig. 9(a), where $\tau_{i_k}^\gamma$ ($k = 1, \dots, 5$) are labeled 1, 2, ℓ , r and ℓ .

- For $k = 0, 1, \dots, d$, let γ_k denote the segment of γ from the point p_k to the point p_{k+1} , and let Δ_k^γ denote the (unique) ideal triangle of T° that γ_k crosses. When it is clear from the context which arc we mean, we simply write p_k^γ as p_k , $\tau_{i_k}^\gamma$ as τ_{i_k} , Δ_k^γ as Δ_k .
- The side/s of Δ_k that is not labeled τ_{i_k} or $\tau_{i_{k+1}}$ is labeled as in Figs. 5 and 6. In particular, for $k = 1, \dots, d - 1$, define arc $\tau_{[\gamma_k]}$ to be

$$\tau_{[\gamma_k]} = \begin{cases} \text{the 3rd arc in } \Delta_k & \text{if } \Delta_k \text{ is not self-folded,} \\ \text{the radius in } \Delta_k & \text{if } \Delta_k \text{ is self-folded.} \end{cases}$$

Define $\tau_{[\gamma_0]}$, $\tau_{[\gamma_{-1}]}$, $\tau_{[\gamma_d]}$, and $\tau_{[\gamma_{d+1}]}$ as follows:

- If the ideal triangle Δ_0 has three distinct edges, then Δ_0 is formed by arc τ_{i_1} and two distinct arcs/boundary edges $\tau_{[\gamma_0]}, \tau_{[\gamma_{-1}]}$ (not equal to τ_{i_1}) such that $\tau_{[\gamma_0]}, \tau_{[\gamma_{-1}]}, \tau_{i_1}$ are arranged in clockwise order around Δ_0 . Similarly, if the ideal triangle Δ_d has three distinct edges, then it is formed by the arcs τ_{i_d} and two distinct arcs/boundary edges $\tau_{[\gamma_d]}, \tau_{[\gamma_{d+1}]}$ (not equal to τ_{i_d}) such that $\tau_{[\gamma_d]}, \tau_{[\gamma_{d+1}]}, \tau_{i_d}$ are arranged in clockwise order around Δ_d .
- If Δ_0 (respectively, Δ_d) is self-folded, then $\tau_{[\gamma_0]} = \tau_{[\gamma_{-1}]}$ (respectively, $\tau_{[\gamma_d]} = \tau_{[\gamma_{d+1}]}$) is the radius.

In Fig. 9(a), $\tau_{[\gamma_0]} = b_1$, $\tau_{[\gamma_{-1}]} = b_4$, $\tau_{[\gamma_d]} = 2$, $\tau_{[\gamma_{d+1}]} = b_3$.

Definition 3.1 (Quasi-arc). If τ is an ordinary radius of \mathcal{C}_n between a marked point v on the boundary and the puncture p , let an associated *quasi-arc* τ' be a curve (not passing through p) which satisfies the following:

1. τ' is between v and a (non-marked) point p' in the vicinity of p . (Note that another quasi-arc associated to τ may use a different point p'' .)
2. τ' agrees with the arc τ outside of a radius- ϵ disk D_ϵ around p , where ϵ is chosen small enough so that the intersection of τ with any other arc is outside of D_ϵ .

If τ is a peripheral arc, we let the associated quasi-arc be τ itself. We label a quasi-arc with the label of the arc that it is associated to.

By abuse of notation, whenever we say we are going along an arc or a side of an ideal triangle Δ_k (as part of a T^o -path), we mean traversing an associated quasi-arc. Note that except for the case when τ is a radius, no abuse of notation is actually needed. The following is an extension of the complete (T^o, γ) -path definition as stated in [21, Definition 2], [16, Section 4.1].

Definition 3.2 (complete (T^o, γ) -path). A *path* $w = (w_1, w_2, \dots, w_{\text{length}(w)})$ on T^o is a concatenation of *steps*, i.e., oriented quasi-arcs and boundary edges of the ideal triangulation T^o of S , such that the starting point of a step w_i is the finishing point of the previous step w_{i-1} . We say that $w = (w_1, \dots, w_{2d+1})$ is a *complete (T^o, γ) -path* if the following axioms hold:

- (T1) Each even step w_{2k} ($k = 1, \dots, d$) is a quasi-arc associated to arc τ_{i_k} . Recall that $\tau_{i_1}, \dots, \tau_{i_d}$ is the sequence of arcs crossed by γ in order.
- (T2) For $k = 0, \dots, d$, each w_{2k+1} traverses a side of the ideal triangle Δ_k . In addition, w_1 traverses the edge $\tau_{[\gamma_0]}$ or $\tau_{[\gamma_{-1}]}$ (which is adjacent to s) and w_{2d+1} traverses the edge $\tau_{[\gamma_d]}$ or $\tau_{[\gamma_{d+1}]}$ (which is adjacent to t).
 - i) Moreover, for $k = 1, 2, \dots, d-1$, let $[p_k, p_{k+1}]_w$ denote the segment of w starting at the point p_k following w_{2k} , continuing along w_{2k+1} and $w_{2(k+1)}$ until the point p_{k+1} . Then the segment γ_k is homotopic to $[p_k, p_{k+1}]_w$. If $S = \mathcal{C}_n$, then we mean homotopy in the disk minus the puncture.
 - ii) The segment γ_0 is homotopic to the segment $[s, p_1]_w$ of the path starting at the point $s = p_0$ following w_1 and w_2 until the point p_1 ;
 - iii) The segment γ_d is homotopic to the segment $[p_d, t]_w$ of the path starting at the point p_d following w_{2d} and w_{2d+1} until the point $p_{d+1} = t$.
- (T3) The step w_{2k+1} starts and finishes in the interior of Δ_k or at a boundary marked point. This means that, if w_{2k+1} goes along a quasi-arc τ' associated to a radius, τ' must be chosen so that its endpoint p' near the puncture is located in the interior of Δ_k .




Figure 5. Ways for γ to cross the ideal triangle Δ_0 or Δ_d .




Figure 6. Ways for γ to cross an ideal triangle Δ_k for $k = 1, \dots, k-1$.

See Examples 3.9 and 3.10.

It is clear that this definition agrees with the complete T^o -paths of [16, 21] for unpunctured surfaces. For short, we will refer to a complete (T^o, γ) -path as simply a (T^o, γ) -path (or a T^o -path) for the rest of this paper.

Remark 3.3. Per (T2), a (T^o, γ) -path is homotopic to γ . It is possible to have $w_j = w_{j+1}$ and, if j is odd, to have $w_j = w_{j+1} = w_{j+2}$. However, since for each k we have $w_{2k} = \tau_{i_k}$ and $w_{2(k+1)} = \tau_{i_{k+1}}$ by (T1) but $\tau_{i_k} \neq \tau_{i_{k+1}}$, no more than three consecutive steps can coincide.

Definition 3.4 (backtrack cycle, non-backtrack cycle, quasi-backtrack). Let $\gamma \notin T^o$ and let $w = (w_1, \dots, w_{2d+1})$ be a (T^o, γ) -path. Let τ be an arc of T^o , and let (w_j, w_{j+1}) be a pair of consecutive quasi-arcs going along τ . We say that (w_j, w_{j+1}) is a *cycle* if the starting point of w_j coincides with the ending point of w_{j+1} .

- i) A cycle (w_j, w_{j+1}) is called a *backtrack*, denoted by $(\underline{\tau}, \underline{\tau})$, if it is contractible.
- ii) A cycle (w_j, w_{j+1}) is called a *non-backtrack*, denoted by $(\overline{\tau}, \overline{\tau})$, otherwise.

In the case that τ is a radius between the puncture P and a marked point v on the boundary, we say that (w_j, w_{j+1}) , denoted by $(\overline{\tau}, \overline{\tau})$, is a *quasi-backtrack* if it is a concatenation of two quasi-arcs $P' \rightsquigarrow v \rightsquigarrow P''$ where P' and P'' are distinct points in the vicinity of P .

Remark 3.5. Assume (w_j, w_{j+1}) is a pair of steps going along τ , as in Definition 3.4.




Figure 7. The three possibilities of a pair (w_j, w_{j+1}) of consecutive steps along a radius τ if T^o has a self-folded triangle with r as the radius.




Figure 8. A pair (w_j, w_{j+1}) of consecutive steps along a radius τ is either a quasi-backtrack or a backtrack cycle if every ideal triangle of T^o is an ordinary triangle.

1) Suppose τ is peripheral.

- i) Then (w_j, w_{j+1}) is a backtrack if and only if w_j and w_{j+1} are opposite orientations of τ . In particular, if τ has two distinct endpoints, (w_j, w_{j+1}) must be a backtrack cycle.
- ii) If τ is an ℓ -loop, the concatenation of two steps going along the same orientation of τ would form a non-contractible cycle. However, we show in Proposition 3.13(1) that it is impossible for (w_j, w_{j+1}) to go along the same orientation of an ℓ -loop twice as part of a (T^o, γ) -path.

2) Suppose τ is a radius between the puncture P and a marked point v on the boundary.

- a) Suppose w_j begins at v , so that (w_j, w_{j+1}) is a concatenation of two associated quasi-arcs $v \rightsquigarrow P' \rightsquigarrow v$.
 - i) Then (w_j, w_{j+1}) is a backtrack cycle if it is contractible to v . See Fig. 7(a) for the case where T^o has a self-folded triangle and τ is its radius and Fig. 8(a) for the case where τ is not the only radius of T^o .
 - ii) If (w_j, w_{j+1}) is not contractible, then due to (T2) it must be homotopic to a loop which goes around the puncture once. We say that (w_j, w_{j+1}) is a *counterclockwise non-backtrack* if it goes counterclockwise (Fig. 7(b)), and a *clockwise non-backtrack* if it goes clockwise (Fig. 7(c)). We show in Proposition 3.13(2a) that in this case, τ must be the only radius of T^o , i.e., T^o contains a self-folded triangle.
- b) Suppose w_j ends at v , so that (w_j, w_{j+1}) is a concatenation of two associated quasi-arcs $P' \rightsquigarrow v \rightsquigarrow P''$, where P' and P'' are in the vicinity of the puncture. We show in Proposition 3.13(2b) that P' and P'' must be distinct, so (w_j, w_{j+1}) is a quasi-backtrack. See Fig. 8(b).

Definition 3.6 (Laurent monomial from a T° -path). We identify each step with the label of the quasi-arc/boundary edge which it traverses and define the Laurent monomial $x(w)$ corresponding to a complete (T°, γ) -path w by

$$x(w) = \prod_{i \text{ odd}} x_{w_i} \prod_{i \text{ even}} x_{w_i}^{-1}.$$

Remark 3.7. Two or more (T°, γ) -paths may correspond to the same Laurent monomial, e.g., see Example 3.10. For each (T°, γ) -path w , the denominator of $x(w)$, before reducing, is equal to $x_{i_1}, x_{i_2}, \dots, x_{i_d}$ which corresponds to the arcs $\tau_{i_1}, \dots, \tau_{i_d}$ of T° which cross γ .

Theorem 3.8 (T° -path formula for \mathcal{C}_n , an extension of [21, Theorem 3.1], [22, Theorem 3.2]). *Let T° be an ideal triangulation of \mathcal{C}_n , let $\gamma \notin T^\circ$ be an ordinary arc of \mathcal{C}_n , and let x_γ denote the corresponding element in the cluster algebra which arises from \mathcal{C}_n (see Theorem 2.8). Then*

$$x_\gamma = \sum_w x(w),$$

where the sum is taken over all (T°, γ) -paths. The formula does not depend on the choice of orientation on γ .

The proof of Theorem 3.8 is given in Section 4.4. Note that, since $x_\ell := x_r x_{r(p)}$, Theorem 3.8 also provides a formula for the cluster variable associated to every tagged arc of \mathcal{C}_n .

Example 3.9. The following are the five (T°, γ) -paths for the situation of Fig. 4. Note that γ crosses 1, 2, 3 in order so $\tau_{i_1} = 1, \tau_{i_2} = 2$, and $\tau_{i_3} = 3$ in this example.

- 1) $(b_1, 1, \underline{2}, 2, 4, 3, b_3)$, 3) $(b_4, 1, b_2, 2, 2, 3, 4)$, 5) $(b_4, \underline{1}, \underline{1}, 2, \underline{3}, 3, b_3)$.
 2) $(b_1, 1, \underline{2}, 2, 2, 3, 4)$, 4) $(b_4, 1, b_2, 2, 4, 3, b_3)$,

Theorem 3.8 thus implies that

$$x_\gamma = \frac{x_{b_1} x_2 x_4 x_{b_3} + x_{b_1} x_2 x_2 x_4 + x_{b_4} x_{b_2} x_2 x_4 + x_{b_4} x_{b_2} x_4 x_{b_3} + x_{b_4} x_1 x_3 x_{b_3}}{x_1 x_2 x_3},$$

where $x_{b_j} = 1$ for each boundary edge b_j .

Example 3.10. The nine (T°, γ) -paths from Fig. 9(a) are as follows (with the backtracks underlined):

- (1) $(b_4, 1, b_2, 2, b_3, \underline{\ell}, \underline{r}, \underline{r}, \underline{\ell}, \ell, b_3)$, (6) $(b_4, 1, b_2, \underline{2}, \underline{2}, \underline{\ell}, \underline{\ell}, \mathbf{r}, \mathbf{r}, \ell, b_3)$,
 (2) $(b_4, \underline{1}, \underline{1}, 2, \underline{\ell}, \underline{\ell}, \underline{r}, \underline{r}, \underline{\ell}, \ell, b_3)$, (7) $(b_4, 1, b_2, \underline{2}, \underline{2}, \underline{\ell}, \underline{\ell}, \mathbf{r}, \mathbf{r}, \underline{\ell}, \ell, b_3)$,
 (3) $(b_1, 1, \underline{2}, \underline{2}, \underline{2}, \underline{\ell}, \underline{\ell}, \underline{r}, \underline{r}, \ell, 2)$, (8) $(b_1, 1, \underline{2}, \underline{2}, \underline{2}, \underline{\ell}, \underline{\ell}, \mathbf{r}, \mathbf{r}, \ell, b_3)$,
 (4) $(b_1, 1, \underline{2}, \underline{2}, b_3, \underline{\ell}, \underline{r}, \underline{r}, \underline{\ell}, \ell, b_3)$, (9) $(b_1, 1, \underline{2}, \underline{2}, \underline{2}, \underline{\ell}, \underline{\ell}, \mathbf{r}, \mathbf{r}, \underline{\ell}, \ell, b_3)$.
 (5) $(b_4, 1, b_2, \underline{2}, \underline{2}, \underline{\ell}, \underline{\ell}, \underline{r}, \underline{r}, \ell, 2)$,

The last four of these (T°, γ) -paths contain a (counterclockwise) non-backtrack (\mathbf{r}, \mathbf{r}) and are illustrated in Fig. 9. They are drawn so that the backtrack cycles $(2, 2)$ and (ℓ, ℓ) are ignored.

We apply Theorem 3.8 and replace each $(x_r x_\ell)/(x_\ell x_r x_\ell)$ with $1/x_\ell$ and each x_ℓ with $x_r x_{r(p)}$ to get

$$x_\gamma = (x_{b_4} x_{b_2} x_{b_3} x_{b_3} + x_{b_4} x_1 x_r x_{r(p)} x_{b_3} + x_{b_1} x_2^3 + x_{b_1} x_{b_3}^2 x_2 + x_{b_4} x_{b_2} x_2^2 + 2x_{b_4} x_{b_2} x_{b_3} x_2 + 2x_{b_1} x_{b_3} x_2^2)/(x_1 x_2 x_r x_{r(p)}),$$

where $x_{b_j} = 1$ for each boundary edge b_j .





Figure 9. The four (T°, γ) -paths (of T° and γ from Fig. 9(a)) which contain a (counterclockwise) non-backtrack cycle (\mathbf{r}, \mathbf{r}) . All backtracks $(\underline{2}, \underline{2})$ and $(\underline{\ell}, \underline{\ell})$ have been omitted, and steps are not drawn exactly along the arcs/boundary edges for illustration purposes.




Figure 10. The four (T°, γ) -paths of the ideal triangulation T° and the ℓ -loop γ of Fig. 10(a). Each path contains a quasi-backtrack $(\overline{1}, \overline{1})$, $(\overline{2}, \overline{2})$, or $(\overline{3}, \overline{3})$.

Example 3.11. Fig. 10 illustrates the four (T°, γ) -paths for the case where the ideal triangulation T° and the ℓ -loop $\gamma \notin T^\circ$ are illustrated in Fig. 10(a). In this example, no (T°, γ) -path contains any cycle (w_j, w_{j+1}) .

Fig. 11 illustrates three of the five (T°, γ) -paths for the case where the ideal triangulation T° and the arc $\gamma \notin T^\circ$ are illustrated in Fig. 11(a). The backtrack cycle $(3, 3)$ in Fig. 11(d) is not drawn. The other two (T°, γ) -paths are $(0, 1, b_2, 2, 0, 3, b_3)$ and $(b_1, 1, \overline{2}, \overline{2}, 0, 3, b_3)$.




Figure 11. Three of the five (T°, γ) -paths of the ideal triangulation T° and the arc γ of Fig. 11(a).




Figure 12. Examples of $non-(T^\circ, \gamma)$ -paths for the situations in Figs. 9(a), 10(a), and 11(a). Each path is homotopic to γ and satisfies (T1) but fails (T2) or (T3).

3.1 A non-backtrack cycle can only go along a self-folded triangle's radius

We prove the assertions from Remark 3.5.

Definition 3.12 (Crossing a self-folded triangle). Suppose that T° contains a self-folded triangle with radius r . We say that γ crosses r in the *counterclockwise direction* (respectively, *clockwise direction*) if it matches (respectively, if γ has the opposite orientation of) Fig. 6(d).

Proposition 3.13. *Let T° be an ideal triangulation of \mathcal{C}_n . Suppose γ is an ordinary arc which crosses $\tau \in T^\circ$, and let $w = (w_1, \dots, w_{2d+1})$ be a (T°, γ) -path. Suppose (w_j, w_{j+1}) is a pair of steps both going along quasi-arcs associated to τ .*

- 1) *Suppose $\tau \in T^\circ$ is an ℓ -loop ℓ . Then (w_j, w_{j+1}) are two opposite orientations of ℓ , so that (w_j, w_{j+1}) is a backtrack cycle. See Remark 3.5(1).*
- 2) *Suppose T° does not contain any self-folded triangle, and let $\tau \in T^\circ$ be a radius between the puncture p and a marked point v on the boundary.*
 - a) *Suppose that w_j begins at v , so that (w_j, w_{j+1}) is a concatenation of two quasi-arcs $v \rightsquigarrow p' \rightsquigarrow v$ where p' is in the vicinity of the puncture. Then (w_j, w_{j+1}) is a backtrack cycle. See Remark 3.5(2a).*
 - b) *Suppose that w_j ends at v , so that (w_j, w_{j+1}) is a concatenation of two quasi-arcs $p' \rightsquigarrow v \rightsquigarrow p''$ where p' and p'' are in the vicinity of the puncture. Then $p' \neq p''$, and hence (w_j, w_{j+1}) is a quasi-backtrack. See Remark 3.5(2b).*

Remark 3.14. In Examples 3.9 and 3.11, none of the (T°, γ) -paths include a non-backtrack cycle along a radius since T° contains no self-folded triangle (see Proposition 3.13(2a)). In

contrast, four of the (T^o, γ) -paths in Example 3.10 for (T^o, γ) -paths contains a non-backtrack cycle. Furthermore, no cycle $p' \rightsquigarrow v \rightsquigarrow p'$ can appear in general (see Proposition 3.13(2b)). See Fig. 12 and Example 3.17 for examples of non- (T^o, γ) -paths.

Proof of Proposition 3.13. First, we point out that the pair of steps (w_j, w_{j+1}) must be in the middle of the (T^o, γ) -path, i.e.,

$$\text{the first step of } w \text{ cannot be } w_j \text{ and the last step of } w \text{ cannot be } w_{j+1}. \quad (3.1)$$

To see this, note that, by (T2), w_1 goes along $\tau_{[\gamma_0]}$ or $\tau_{[\gamma_{-1}]}$, and w_{2d+1} goes along $\tau_{[\gamma_d]}$ or $\tau_{[\gamma_{d+1}]}$. But, as illustrated in Fig. 5, $\tau_{[\gamma_j]} \neq \tau_{i_1}$ if $j \in \{0, -1\}$, and $\tau_{[\gamma_j]} \neq \tau_{i_d}$ if $j \in \{d, d+1\}$. Since, by (T1), w_2 goes along τ_{i_1} and w_{2d} goes along τ_{i_d} , we must have $1 < j$ and $j+1 < d$.

We prove case (1): Suppose T^o has a self-folded triangle with radius r and ℓ -loop ℓ , and $\tau = \ell$. For the sake of argument, suppose that w_j and w_{j+1} are the same orientation of ℓ . Note that every arc of T^o other than ℓ has two distinct endpoints.

– First, suppose $j = 2k$ is even. Per (3.1), another even-indexed step $w_{2(k+1)}$ follows (w_{2k}, w_{2k+1}) . Since w_{2k+1} ends at v , the step w_{2k+2} (crossing γ) must go from v along a different arc λ to a different marked point. Using the labels of Figs. 13(a), λ can be either r , \aleph , or \beth . We check all possible cases against (T1) and (T2) but only show one of the arguments here. Suppose $\lambda = r$ and γ is a peripheral arc crossing ℓ , r , and ℓ again in the counterclockwise direction as in Fig. 13(a) with $j = 2k$ and $j+1 = 2k+1$ so that $w_{2k} = \ell$ and $w_{2k+2} = r$. For contradiction, suppose (w_{2k}, w_{2k+1}) go along the clockwise orientation of ℓ twice. Recall that γ_k is the segment of γ from p_k (a point in the interior of ℓ) and p_{k+1} (a point in the interior of r) as in Fig. 13(b). By (T2), the segment γ_k must be homotopic to $[p_k, p_{k+1}]_w$ (see Definition 3.2 and Fig. 13(c)). However, as illustrated in Fig. 13(d), the concatenation of $[p_k, p_{k+1}]_w$ and the opposite orientation γ_k^- of γ_k is not contractible. Hence γ_k is not homotopic to $[p_k, p_{k+1}]_w$.

– Second, suppose $j = 2k+1$ is odd. Again, per (3.1), there is another even-indexed step w_{2k} (crossing γ , and going from a different marked point to v along a peripheral arc) which precedes (w_{2k+1}, w_{2k+2}) . Using the same logic as in the previous paragraph, we show that (w_{2k+1}, w_{2k+2}) must go along two opposite orientations of ℓ in order for γ_k to be homotopic to $[p_k, p_{k+1}]_w$.

We prove case (2a): Suppose τ is a radius and a side of a regular triangle (hence T^o has no self-folded triangle), and every arc of T^o has two distinct endpoints.

– First, suppose $j = 2k$ is even. Per (3.1), another even-indexed step $w_{2(k+1)}$ follows (w_{2k}, w_{2k+1}) . Since w_{2k+1} ends at v , the step w_{2k+2} (crossing γ) must be a peripheral step λ to a different point. But (w_{2k}, w_{2k+1}) would need to be contractible in order for γ_k to be homotopic to $[p_k, p_{k+1}]_w$.

– Second, suppose $j = 2k+1$ is odd. Again, per (3.1), there is another even-indexed step w_{2k} (crossing γ , and going from a different marked point to v along a peripheral arc λ) which precedes (w_{2k+1}, w_{2k+2}) . But (w_{2k+1}, w_{2k+2}) would need to be contractible in order for γ_k to be homotopic to $[p_k, p_{k+1}]_w$.

We prove case (2b): First suppose $j = 2k+1$ is odd. Then p' is in the interior of Δ_k since w_{2k+1} must start in the interior of Δ_k by (T3). The next step w_{2k+2} then goes along τ from v to p'' . Since the last step of w is an odd-indexed step, after w_{2k+2} there must be another step w_{2k+3} which starts at p'' . Again by (T3), w_{2k+3} starts in the interior of Δ_{k+1} . But Δ_k and Δ_{k+1} are distinct triangles since T^o has no self-folded triangle. Hence $p' \neq p''$, and so (w_{2k+1}, w_{2k+2}) is a quasi-backtrack which starts in the interior of Δ_k and ends in the interior of Δ_{k+1} .

If $j = 2k$ is even, then by a similar argument (w_{2k}, w_{2k+1}) is a quasi-backtrack which starts in the interior of Δ_{k-1} and ends in the interior of Δ_k . See Fig. 8(b). ■




Figure 13. (ℓ_+, ℓ_+) is not a valid (T°, γ) -subpath.

3.2 A (T°, γ) -path w on \mathcal{C}_n is uniquely determined by its sequence of labels

Notation 3.15. Let r and ℓ be the radius and ℓ -loop of a self-folded triangle, and let v be the boundary vertex on the boundary that is adjacent to ℓ . Let ℓ_+ (respectively, ℓ_-) denote the clockwise (respectively, counterclockwise) orientation along ℓ . (We mark each ℓ curve in our illustrations with an arrow pointing clockwise to remind the reader that ℓ_+ denotes the clockwise direction.)

Consider the cycle (r, r)

$$v \rightsquigarrow r' \rightsquigarrow v \text{ along } r$$

from Remark 3.5(2a). Let $(\underline{r}, \underline{r})$ denote the backtrack cycle and let (\mathbf{r}, \mathbf{r}) denote a (counterclockwise or clockwise) non-backtrack cycle.

The following proposition is an analogue of a remark from [21, Section 3.1].

Proposition 3.16. *A (T°, γ) -path w on \mathcal{C}_n is uniquely determined by its sequence of labels (w_1, \dots, w_{2d+1}) , forgetting the orientations of the steps and whether a consecutive pair is a non-backtrack or a backtrack.*

Example 3.17. To illustrate Proposition 3.16, consider the (T°, γ) -path of Fig. 9(b)

$$w = (b_4, 1, b_2, 2, 2, \ell_+, \ell_-, \mathbf{r}, \mathbf{r}, \ell_+, b_3).$$

The pair $(w_8, w_9) = (\mathbf{r}, \mathbf{r})$ is a *counterclockwise* non-backtrack along the radius r , and $w_{10} = \ell_+$ goes clockwise around ℓ . Consider a different path w' (see Fig. 12(a)) which goes along the same sequence of arcs such that $(w'_8, w'_9) = (\mathbf{r}, \mathbf{r})$ is a *clockwise* non-backtrack along the radius, and $w'_{10} = \ell_-$ goes counterclockwise around ℓ . Even though w' is homotopic to γ , the segment γ_4 is not homotopic to $[p_4, p_5]_w$, violating Definition 3.2(T2).

The paths illustrated in Figs. 12(b) and 12(c) go along associated quasi-arcs of the same arcs/edges as the (T°, γ) -paths of Figs. 10(b) and 11(d), but they fail axiom (T3), which requires each odd-indexed step w_{2k+1} to start and finish in the interior of Δ_k or at a boundary marked point.




Figure 14. The subpaths (w_1, \dots, w_4) of Remark 3.18(A1), (A2) when γ starts at P and crosses ℓ and \aleph . See Fig. 21.

Proof of Proposition 3.16. For short, let the arcs $\tau_{i_1}, \dots, \tau_{i_d}$ be denoted by arcs $1, \dots, d$.

First, consider the subsequence (w_1, w_2, w_3) . There are four possibilities:

- i) Δ_0 is an ordinary triangle (Fig. 5(a)).
- ii) Δ_0 is an ideal triangle with two vertices where arc 1 is the loop (Fig. 5(b)).
- iii) Δ_0 is an ideal triangle with two vertices where arc 1 is not the loop (Fig. 5(c)).
- iv) Δ_0 is a self-folded triangle where arc 1 is the ℓ -loop (Fig. 5(d)).

For each of the first three cases, Δ_0 has three distinct edges k , $\tau_{[\gamma_0]}$, and $\tau_{[\gamma_{-1}]}$. There are exactly two possible subpaths for (w_1, w_2) , which are represented by two distinct sequences $(\tau_{[\gamma_0]}, 1)$ and $(\tau_{[\gamma_{-1}]}, 1)$.

In the fourth case, Δ_0 is a self-folded triangle with radius r and ℓ -loop ℓ (see Fig. 5(d), top), so that $\tau_{[\gamma_0]} = \tau_{[\gamma_{-1}]} = r$. However, there are exactly two possible subpaths in this case as well.

Remark 3.18. Suppose γ is a radius starting at the puncture, as in Fig. 5(d). Let v denote the vertex that ℓ is based at. Due to (T2), the first step w_1 must go $P' \rightsquigarrow v$ along $\tau_{[\gamma_0]} = \tau_{[\gamma_{-1}]} = r$, but there are two valid options for w_2 , to go counterclockwise or clockwise along ℓ , so that there are exactly two possibilities for (w_1, w_2) :

A1) (r, ℓ_-) ,

A2) (r, ℓ_+) .

We continue with the proof of Proposition 3.16 in this fourth case. In particular, we show that (A1) and (A2) lead to distinct ways for the (T^o, γ) -path to be finished. These two possible choices for (w_1, w_2) are represented by the same sequence $(\tau_{[\gamma_0]}, 1) = (\tau_{[\gamma_{-1}]}, 1)$, but we claim that the orientation of w_2 determines the possible choices for the next term w_3 . Suppose that \aleph (respectively, \beth) is the side of Δ_1 which lies clockwise (respectively, counterclockwise) of ℓ , as illustrated in Fig. 14(a).

Suppose γ ends at the vertex y . If $w_2 = \ell_-$ is counterclockwise, then $w_3 = \aleph$ (Fig. 14(b)), otherwise $w_3 = \beth$ (Fig. 14(c)).

If γ does not end at y , then the second arc that γ crosses is either \aleph or \beth (say, \aleph). If $w_2 = \ell_-$ is counterclockwise, then $(w_3, w_4) = (\aleph, \aleph)$ (Fig. 14(b)). If $w_2 = \ell_+$ is clockwise, then $(w_3, w_4) = (\beth, \aleph)$ (Fig. 14(c)) or (ℓ_-, \aleph) (Fig. 14(d)). Hence, the subpath (w_1, w_2, w_3) is either (r, ℓ_-, \aleph) , (r, ℓ_+, \beth) , or (r, ℓ_+, ℓ_-) . As these are represented by three distinct subsequences, each subsequence uniquely determines the first three steps of w .

By the same logic, the subsequence $(w_{2d-1}, w_{2d}, w_{d+1})$ uniquely determines the last three steps of w .

Next, consider any triple $(w_{2k}, w_{2k+1}, w_{2k+2})$. There are four possibilities:

- i) Δ_k is an ordinary triangle (Fig. 6(a)).
- ii) Δ_k is an ideal triangle with two vertices where neither arc k nor arc $k+1$ are loops (Fig. 6(b)).
- iii) Δ_k is an ideal triangle with two vertices where one of arcs k and $k+1$ (say, $k+1$) is a loop (Fig. 6(c)).
- iv) Δ_k is a self-folded triangle with radius r and ℓ -loop ℓ where one of arcs k and $k+1$ (say, $k+1$) is the radius r (Fig. 6(d)).

For each of the first three cases, Δ_k has three distinct edges k , $k+1$, and $\tau_{[\gamma_k]}$. There are exactly three legal subpaths for $(w_{2k}, w_{2k+1}, w_{2k+2})$, and they are represented by three distinct sequences $(k, k+1, k+1)$, $(k, k, k+1)$, and $(k, \tau_{[\gamma_k]}, k+1)$. In the case that one of these steps is an ℓ -loop, it is also clear that (T2) forces a specific orientation along ℓ as part of the (T^o, γ) -path.

In the fourth case, γ crosses ℓ , r , then ℓ in either the counterclockwise or clockwise (say, the former) direction.

Lemma 3.19. *Suppose T^o contains a self-folded triangle with radius r and ℓ -loop ℓ , and suppose that γ crosses r , say, in the counterclockwise direction. Let v be the boundary vertex on the boundary that is adjacent to ℓ . Suppose ℓ , r , ℓ are the k -th, $(k+1)$ -th, and $(k+2)$ -th arcs crossed by γ (see Fig. 6(d)). Then there are exactly three possible subpaths for $(w_{2k}, w_{2k+1}, w_{2(k+1)})$:*

- a1) $(\ell_-, \underline{r}, r)$: follow ℓ_- then backtrack cycle (r, r) (Fig. 15(1)),
- a2) (ℓ_+, ℓ_-, r) : follow (ℓ_+, ℓ_-) then follow r from v to P' (a point in the vicinity of the puncture) (Fig. 15(2)),
- a3) $(\ell_+, \mathbf{r}, \mathbf{r})$: follow ℓ_+ then counterclockwise non-backtrack cycle (\mathbf{r}, \mathbf{r}) (Fig. 15(3)).

There are also exactly three possible subpaths for $(w_{2(k+1)}, w_{2k+3}, w_{2(k+2)})$:

- b1) (r, ℓ_-, ℓ_+) : follow r from P' (a point in the vicinity of the puncture) to v then (ℓ_-, ℓ_+) (see Fig. 16(1)),
- b2) $(\underline{r}, r, \ell_-)$: backtrack cycle (r, r) then ℓ_- (Fig. 16(2)),
- b3) $(\mathbf{r}, \mathbf{r}, \ell_+)$: counterclockwise non-backtrack cycle (\mathbf{r}, \mathbf{r}) then ℓ_+ (Fig. 16(3)).

The (ℓ, r, r, ℓ, ℓ) -subsequence corresponds to exactly 2 valid subpaths for $(w_{2k}, \dots, w_{2k+2})$, one where (\underline{r}, r) is a backtrack and the other where (\mathbf{r}, \mathbf{r}) is a non-backtrack. We get them by combining (a1) with (b1) and combining (a3) with (b1):

- I) $(\ell_-, \underline{r}, r, \ell_-, \ell_+)$: follow ℓ_- then backtrack cycle (r, r) then (ℓ_-, ℓ_+) .
- II) $(\ell_+, \mathbf{r}, \mathbf{r}, \ell_-, \ell_+)$: follow ℓ_+ then counterclockwise non-backtrack cycle (\mathbf{r}, \mathbf{r}) then (ℓ_-, ℓ_+) .

Similarly, the (ℓ, ℓ, r, r, ℓ) -subsequence would correspond to exactly two valid subpaths, one where (\underline{r}, r) is a backtrack and the other where (\mathbf{r}, \mathbf{r}) is a non-backtrack. We get them by combining (a2) with (b2), and combining (a2) with (b3).




Figure 15. The subpaths $(w_{2k}, w_{2k+1}, w_{2k+2})$ of Lemma 3.19(a1), (a2), (a3) if γ crosses arcs $\tau_{i_k} = \ell$, $\tau_{i_{k+1}} = r$, and $\tau_{i_{k+2}} = \ell$ in the counterclockwise order. See Fig. 22.




Figure 16. The subpaths $(w_{2k+2}, w_{2k+3}, w_{2k+4})$ of Lemma 3.19(b1), (b2), (b3) if γ crosses arcs $\tau_{i_k} = \ell$, $\tau_{i_{k+1}} = r$, and $\tau_{i_{k+2}} = \ell$ in the counterclockwise order. See Fig. 23.

III) $(\ell_+, \ell_-, \underline{r}, r, \ell_-)$: follow ℓ_+ , then ℓ_- then backtrack cycle (\underline{r}, r) then ℓ_- .

IV) $(\ell_+, \ell_-, \mathbf{r}, \mathbf{r}, \ell_+)$: follow ℓ_+ , then ℓ_- then counterclockwise non-backtrack cycle (\mathbf{r}, \mathbf{r}) then ℓ_+ .

Remark 3.20. In Lemma 3.19, note that the subpaths (I) and (III) are homotopic to a counterclockwise orientation of ℓ . The subpaths (II) and (IV) are contractible to v .

Proof of Lemma 3.19. By (T1), since the k -th, $(k+1)$ -th, $(k+2)$ -th arcs crossed by γ are ℓ , r , and ℓ , the sequence


$$(w_{2k}, w_{2k+1}, w_{2(k+1)}, w_{2k+3}, w_{2(k+2)})$$

must be

$$(\ell, \quad, r, \quad, \ell).$$

For this to be a connected path, we must fill in the odd steps with ℓ then r (or r then ℓ) by (T2), so that the 5-term subsequence is either (ℓ, r, r, ℓ, ℓ) or (ℓ, ℓ, r, r, ℓ) .

The two subpaths (a1) and (a3) (both represented by (ℓ, r, r)) satisfy (T2):



(a) Concatenation of $[p_k, p_{k+1}]_w$ ($\ell_-, r, r(\text{backtrack})$) and γ_k^- is contractible, so $[p_k, p_{k+1}]_w$ is homotopic to γ_k .
 (b) Similarly, $[p_k, p_{k+1}]_w$ ($\ell_+, \mathbf{r}, \mathbf{r}$ (counterclockwise non-backtrack)) is homotopic to γ_k .
 (c) $[p_k, p_{k+1}]_w$ ($\ell_+, r, r(\text{backtrack})$) is *not* homotopic to γ_k .

Figure 17. Concatenation of $[p_k, p_{k+1}]_w$ (ℓ, r, r) and the opposite orientation of γ_k^- of γ_k .

Case (a1) If $(w_{2k}, w_{2k+1}, w_{2(k+1)}) = (\ell_-, r, r)$ (where (r, r) is a backtrack cycle), the concatenation of $[p_k, p_{k+1}]_w$ and the opposite orientation γ_k^- of γ_k is contractible, so $[p_k, p_{k+1}]_w$ and γ_k are homotopic. See Fig. 17(a).

Case (a3) If $(w_{2k}, w_{2k+1}, w_{2(k+1)}) = (\ell_+, \mathbf{r}, \mathbf{r})$ (where (\mathbf{r}, \mathbf{r}) is a counterclockwise non-backtrack cycle), the concatenation of $[p_k, p_{k+1}]_w$ and the opposite orientation γ_k^- of γ_k is contractible, so $[p_k, p_{k+1}]_w$ and γ_k are homotopic. See Fig. 17(b).

The sequence (ℓ, r, r) may also represent a subpath (ℓ_+, r, r) where (r, r) is a backtrack or $(\ell_-, \mathbf{r}, \mathbf{r})$ where (\mathbf{r}, \mathbf{r}) is a non-backtrack cycle, but we claim that these are not valid T^o -subpath: For contradiction, suppose $(w_{2k}, w_{2k+1}, w_{2k+2}) = (\ell_+, r, r)$ where (r, r) is a backtrack. By (T2), the segment γ_k (Fig. 13(b)) must be homotopic to $[p_k, p_{k+1}]_w$. However, as illustrated in Fig. 17(c), the concatenation of $[p_k, p_{k+1}]_w$ and the opposite orientation γ_k^- of γ_k is not contractible. Hence γ_k is not homotopic to $[p_k, p_{k+1}]_w$, and so this subpath does not appear. Similarly, no subpath $(\ell_-, \mathbf{r}, \mathbf{r})$ with a non-backtrack cycle (\mathbf{r}, \mathbf{r}) can appear.

By similar logic,

- (a2) is the only valid T^o -subpath which can be represented by (ℓ, ℓ, r) ,
- (b1) is the only valid T^o -subpath which can be represented by (r, ℓ, ℓ) , and
- (b2) and (b3) are the only two valid T^o -subpaths which can be represented by (r, r, ℓ) .

This concludes our proof of Lemma 3.19. ■

We continue with the proof of Proposition 3.16. Note that the subpaths (I) and (II) are represented by the same 5-term subsequence (ℓ, r, r, ℓ, ℓ) . We claim that the steps w_{2k-1} which precede (I) and (II) go along distinct arcs/edges. Note that the ideal triangle Δ_{k-1} is a two-vertex, three-edge triangle (Fig. 1(b)).

Case 1: Suppose $k = 1$, so that γ starts at the marked point y . Recall that the side of Δ_0 that lies clockwise of ℓ is $\tau_{[\gamma_0]}$, and the side of Δ_0 that lies counterclockwise of ℓ is $\tau_{[\gamma_{-1}]}$ (see Fig. 5(b)). In Fig. 15(4), $\tau_{[\gamma_0]} = \aleph$ and $\tau_{[\gamma_{-1}]} = \beth$.

- Since (I) starts with a counterclockwise ℓ_- step, the first step w_1 must go along $\tau_{[\gamma_{-1}]}$. So the first six-term subsequence is $(\tau_{[\gamma_{-1}]}, \ell, r, r, \ell, \ell)$.
- Since (II) starts with a clockwise ℓ_+ step, the first step w_1 must go along $\tau_{[\gamma_0]}$. So the first six-term subsequence is $(\tau_{[\gamma_0]}, \ell, r, r, \ell, \ell)$.

Case 2: Suppose $1 < k$, so that γ crosses arc $k - 1$ (in Fig. 15(4), this arc is either \aleph or \beth). Suppose (without loss of generality) that arc $k - 1$ lies clockwise of ℓ (i.e., arc $k - 1$ is \aleph in Fig. 15(4)). Then the third arc $\tau_{[\gamma_{k-1}]}$ of Δ_{k-1} lies counterclockwise of ℓ (i.e., side $\tau_{[\gamma_{k-1}]}$ is \beth in Fig. 15(4)). Recall that w_{2k-2} must traverse arc $k - 1$ by (T1).

- Since (I) starts with a counterclockwise ℓ_- step, the previous step w_{2k-1} must go along either ℓ or $\tau_{[\gamma_{k-1}]}$ (i.e., side \beth in Fig. 15(4)), so that $(w_{2k-1}, \dots, w_{2(k+2)})$ form a six-term subsequence $(\ell, \ell, r, r, \ell, \ell)$ or $(\tau_{[\gamma_{k-1}]}, \ell, r, r, \ell, \ell)$.
- Since (II) starts with a clockwise ℓ_+ step, the previous step w_{2k-1} must go along arc $k - 1$ (i.e., arc \aleph in Fig. 15(4)), so that $(w_{2k-1}, \dots, w_{2(k+2)}) = (k - 1, \ell, r, r, \ell, \ell)$.

Similarly, the subpaths (III) and (IV) are represented by the same 5-term subsequence (ℓ, ℓ, r, r, ℓ) . We claim that the steps $w_{2(k+2)+1}$ following (III) and (IV) go along distinct arc/edges. Note that the ideal triangle Δ_{k+2} is a two-vertex, three-edge triangle (Fig. 1(b)).

Case 1: Suppose $d = k + 2$, so that γ ends at the marked point y . Recall that the side of Δ_d that lies clockwise of ℓ is $\tau_{[\gamma_d]}$, and the side of Δ_0 that lies counterclockwise of ℓ is $\tau_{[\gamma_{d+1}]}$ (see Fig. 5(b)). In Fig. 15(4), $\tau_{[\gamma_d]} = \aleph$ and $\tau_{[\gamma_{d+1}]} = \beth$.

- Since (III) ends with a counterclockwise ℓ_- , the last step must go along $\tau_{[\gamma_d]}$, so that the last six-term subsequence is $(\ell, \ell, r, r, \ell, \tau_{[\gamma_d]})$
- Since (IV) ends with a clockwise ℓ_+ , the last step must go along $\tau_{[\gamma_{d+1}]}$, so that the last six-term subsequence is $(\ell, \ell, r, r, \ell, \tau_{[\gamma_{d+1}]})$.

Case 2: Suppose $d > k + 2$, so that γ crosses arc $k + 3$ (in Fig. 15(4), this arc is either \aleph or \beth). Suppose (without loss of generality) that arc $k + 3$ lies clockwise of ℓ (i.e., arc $k + 3$ is \aleph in Fig. 15(4)). Then the third arc $\tau_{[\gamma_{k+2}]}$ of Δ_{k+2} lies counterclockwise of ℓ (i.e., side $\tau_{[\gamma_{k+2}]}$ is \beth in Fig. 15(4)).

- Since (III) ends with a counterclockwise ℓ_- , the next step w_{2k+5} must go along arc $k + 3$ (i.e., arc \aleph in Fig. 15(4)), so that $(w_{2k}, \dots, w_{2k+5}) = (\ell, \ell, r, r, \ell, k + 3)$.
- Since (IV) ends with a clockwise ℓ_+ , the next step w_{2k+5} must go along either counterclockwise ℓ_- or $\tau_{[\gamma_{k+2}]}$ (i.e., arc/edge labeled \beth in Fig. 15(4)), so that $(w_{2k}, \dots, w_{2k+5})$ form a six-term subsequence $(\ell, \ell, r, r, \ell, \ell)$ or $(\ell, \ell, r, r, \ell, \tau_{[\gamma_{k+2}]})$. ■

4 Proof of Theorem 3.8, the (T°, γ) -path expansion formula for a once-punctured disk

We begin with an outline of our proof. Let T° be an ideal triangulation of a once-punctured n -gon \mathcal{C}_n and let $\gamma \notin T^\circ$ be an oriented ordinary arc (possibly an ℓ -loop).

Step 1: Following [17, Section 7], we construct a triangulated $(d + 3)$ -gon \widetilde{S}_γ (modeled after $\Delta_0, \dots, \Delta_d$). See Section 4.1.

Step 2: In Section 4.2, we give a bijection $\bar{\pi}: \{(\widetilde{S}_\gamma, \widetilde{\gamma})\text{-paths}\} \rightarrow \{(T^\circ, \gamma)\text{-paths}\}$.

Step 3: Section 4.3 recaps [17, Theorem 4.10], a T° -expansion of x_γ in terms of perfect matchings of a snake graph $G_{T^\circ, \gamma}$.

Step 4: [16, Lemma 4.5] gives a bijection $F: \{\text{perfect matchings of } G_{T^\circ, \gamma}\} \rightarrow \{(\widetilde{S}_\gamma, \widetilde{\gamma})\text{-paths}\}$. See Section 4.4.

Step 5: The composition $\bar{\pi} \circ F$ gives a bijection between perfect matchings of $G_{T^\circ, \gamma}$ and (T°, γ) -paths. Applying $\bar{\pi} \circ F$ to [17, Theorem 4.10] yields Theorem 3.8.

4.1 A triangulated polygon \widetilde{S}_γ and a lifted arc $\widetilde{\gamma}$

Let T° be an ideal triangulation of a once-punctured n -gon \mathcal{C}_n and let $\gamma \notin T^\circ$ be an oriented ordinary arc (possibly an ℓ -loop) from the point s to the point t . Let S_γ be the union of the ideal triangles Δ_k ($k = 0, \dots, d$) crossed by γ . First, for each $k = 0, \dots, d$, we build a triangle $\widetilde{\Delta}_k$ with three distinct labels.

Definition 4.1 (triangles $\widetilde{\Delta}_k$). If Δ_k has three distinct sides, its lift $\widetilde{\Delta}_k$ is an ordinary triangle with the edge labels $\widetilde{\tau}_{i_k} = \tau_{i_k}$, $\widetilde{\tau}_{i_{k+1}} = \tau_{i_{k+1}}$, $\widetilde{\tau}_{[\gamma_k]} = \tau_{[\gamma_k]}$, and the same orientation as Δ_k .

When Δ_k is a self-folded triangle with radius r and ℓ -loop ℓ , its lift $\widetilde{\Delta}_k$ is formed by two lifts of r and a lift of ℓ , as follows:

Case $k = 1$ or d : If $k = 0$, then $\widetilde{\Delta}_0$ has edge labels $\widetilde{\tau}_{i_1} = \ell$, $\widetilde{\tau}_{[\gamma_0]} = r$, $\widetilde{\tau}_{[\gamma_{-1}]} = r^{cc}$ (arranged in clockwise order). Similarly, if $k = d$, then $\widetilde{\Delta}_d$ has edge labels $\widetilde{\tau}_{i_d} = \ell$, $\widetilde{\tau}_{[\gamma_d]} = r$, $\widetilde{\tau}_{[\gamma_{d+1}]} = r^{cc}$ (arranged in clockwise order). See Fig. 18(b).

Case $0 < k < d$: Suppose $k = \ell$, $k + 1 = r$, and $k + 2 = \ell$. Then $\widetilde{\Delta}_k$ has three distinct edge labels $\widetilde{\tau}_{i_k}$, $\widetilde{\tau}_{i_{k+1}}$, and $\widetilde{\tau}_{[\gamma_k]}$: If γ crosses r in counterclockwise (respectively, clockwise) direction, let $\widetilde{\tau}_{i_{k+1}} = r$ be the label of the edge of $\widetilde{\Delta}_k$ which lies counterclockwise (respectively, clockwise) of $\widetilde{\tau}_{i_k} = \ell$. Let $\widetilde{\tau}_{[\gamma_k]} = \dot{r}$ be the label of the third edge of $\widetilde{\Delta}_k$.

Similarly, $\widetilde{\Delta}_{k+1}$ has three distinct edge labels $\widetilde{\tau}_{i_{k+1}}$, $\widetilde{\tau}_{i_{k+2}}$, and $\widetilde{\tau}_{[\gamma_{k+1}]}$: If γ crosses r in counterclockwise (respectively, clockwise) direction, let $\widetilde{\tau}_{[\gamma_{k+1}]} = \ddot{r}$ be the label of the edge of $\widetilde{\Delta}_{k+1}$ which lies counterclockwise (respectively, clockwise) of $\widetilde{\tau}_{i_{k+2}} = \ell$. Let r be the label of the third edge of $\widetilde{\Delta}_{k+1}$. See Figs. 18(e) and 18(f).

Definition 4.2 (triangulated $(d + 3)$ -gon). Glue $\widetilde{\Delta}_1$ to $\widetilde{\Delta}_0$ along τ_{i_1} , glue $\widetilde{\Delta}_2$ to $\widetilde{\Delta}_1$ along τ_{i_2} , and so forth to form a triangulated $(d + 3)$ -gon \widetilde{S}_γ with internal edges

$$\widetilde{\tau}_{i_1}, \dots, \widetilde{\tau}_{i_d}$$

and boundary edges

$$\widetilde{\tau}_{[\gamma_{-1}]}, \widetilde{\tau}_{[\gamma_0]}, \dots, \widetilde{\tau}_{[\gamma_{d+1}]}.$$





Let the lift $\widetilde{s} = \widetilde{p}_0$ of $s = p_0$ be the vertex of $\widetilde{\Delta}_0$ that is adjacent to the lifted edges $\widetilde{\tau}_{[\gamma_0]}$ and $\widetilde{\tau}_{[\gamma_{-1}]}$, and the lift $\widetilde{t} = \widetilde{p}_{d+1}$ of $t = p_{d+1}$ be the vertex of $\widetilde{\Delta}_d$ that is adjacent to the lifted edges $\widetilde{\tau}_{[\gamma_d]}$ and $\widetilde{\tau}_{[\gamma_{d+1}]}$. If $k = 1, \dots, d - 1$, let $\widetilde{p}_k, \widetilde{p}_{k+1}$ denote the lifts of p_k, p_{k+1} on $\widetilde{\Delta}_k$ which lies on the interior of $\widetilde{\tau}_{i_k}, \widetilde{\tau}_{i_{k+1}}$ (respectively).

Let $\widetilde{\gamma}$ be the arc in \widetilde{S}_γ from \widetilde{s} to \widetilde{t} . We call $\widetilde{\gamma}$ the *lift* of γ . Let $\pi: \widetilde{S}_\gamma \rightarrow S_\gamma$ denote the covering map of $\widetilde{S}_\gamma \rightarrow S_\gamma$. See Fig. 19(b) for the lifts $\widetilde{\gamma}$ and \widetilde{S}_γ of γ and S_γ of Fig. 19(a).

Remark 4.3. By construction, $\widetilde{\tau}_{i_k} = \tau_{i_k}$ for each k . Furthermore, $\widetilde{\tau}_{[\gamma_k]} = \tau_{[\gamma_k]}$ unless $\widetilde{\tau}_{[\gamma_k]}$ has label \dot{r}, \ddot{r} , or r^{cc} . In addition, $\widetilde{\tau}_{[\gamma_0]}, \widetilde{\tau}_{[\gamma_d]} \notin \{\dot{r}, \ddot{r}, r^{cc}\}$, so $\widetilde{\tau}_{[\gamma_0]} = \tau_{[\gamma_0]}$, $\widetilde{\tau}_{[\gamma_d]} = \tau_{[\gamma_d]}$.

4.2 A bijection $\overline{\pi}$ between $(\widetilde{S}_\gamma, \widetilde{\gamma})$ -paths and (T°, γ) -paths

Keep the same setup as in the previous section, where T° is an ideal triangulation of \mathcal{C}_n and $\gamma \notin T^\circ$ is an ordinary arc of \mathcal{C}_n . As \widetilde{S}_γ is a triangulation of a polygon, we can consider a $(\widetilde{S}_\gamma, \widetilde{\gamma})$ -path $\widetilde{w} = (\widetilde{w}_1, \dots, \widetilde{w}_{2d+1})$ as defined in Definition 3.2. For example, the $(\widetilde{S}_\gamma, \widetilde{\gamma})$ -paths of Fig. 20 correspond to the four (T°, γ) -paths of Fig. 9.

(a) γ is a radius starting or ending at the puncture.(b) The triangle $\tilde{\Delta}_0$ or $\tilde{\Delta}_d$ when γ is a radius starting or ending at the puncture.(c) γ crosses r in the counterclockwise direction.(d) γ crosses r in the clockwise direction.(e) The triangles $\tilde{\Delta}_k$, $\tilde{\Delta}_{k+1}$ when γ crosses a self-folded triangle's radius r in the counterclockwise direction.(f) The triangles $\tilde{\Delta}_k$, $\tilde{\Delta}_{k+1}$ when γ crosses a self-folded triangle's radius r in the clockwise direction.**Figure 18.** The triangles $\tilde{\Delta}_k$ when γ crosses an ℓ -loop.

Lemma 4.4. *The covering map $\pi: \tilde{S}_\gamma \rightarrow S_\gamma$ which gives*

$$\pi(\tilde{w}_j) = \begin{cases} r & \text{if } \tilde{w}_j \text{ has label } \dot{r}, \ddot{r}, \text{ or } r^{cc}, \\ \text{the same label as } \tilde{w}_j & \text{otherwise.} \end{cases}$$

induces a bijection

$$\begin{aligned} \bar{\pi}: \{(\tilde{S}_\gamma, \tilde{\gamma})\text{-paths in } \tilde{S}_\gamma\} &\rightarrow \{(T^o, \gamma)\text{-paths in } S_\gamma\}, \\ \bar{\pi}(\tilde{w}_1, \tilde{w}_2, \dots, \tilde{w}_{2d+1}) &= (\pi(\tilde{w}_1), \pi(\tilde{w}_2), \dots, \pi(\tilde{w}_{2d+1})). \end{aligned}$$

Proof. Let \tilde{w} be a $(\tilde{S}_\gamma, \tilde{\gamma})$ -path. Note that, since \tilde{S}_γ is a polygon, there is only one way to concatenate a pair $(\tilde{w}_j, \tilde{w}_{j+1})$ of steps. By (T1), every \tilde{w}_{2k} has label $\tilde{\tau}_{i_k}$, and every w_{2k} has




Figure 19. The construction of the triangulated polygon \widetilde{S}_γ and lifted arc $\widetilde{\gamma}$ for T^o and γ from Fig. 19(a).




Figure 20. The $(\widetilde{T}^o, \widetilde{\gamma})$ -paths corresponding to the four (T^o, γ) -paths from Fig. 9. All backtracks have been omitted.

label τ_{i_k} . Per Remark 4.3, $\widetilde{\tau}_{i_k}$ has the same label as τ_{i_k} , and $\widetilde{\tau}_{[\gamma_j]}$ has the same label as $\tau_{[\gamma_j]}$ if $j \in \{0, d\}$.

By construction, $\widetilde{\Delta}_0$ has three distinct labels, τ_{i_1} , $\tau_{[\gamma_0]}$, and $\widetilde{\tau}_{[\gamma_{-1}]}$, and $\widetilde{\Delta}_d$ has three distinct labels, τ_{i_d} , $\tau_{[\gamma_d]}$, and $\widetilde{\tau}_{[\gamma_{d+1}]}$. Hence, as discussed in Section 3.2, the subpath $(\widetilde{w}_1, \widetilde{w}_2)$ is either $(\tau_{[\gamma_0]}, \tau_{i_1})$ or $(\widetilde{\tau}_{[\gamma_{-1}]}, \tau_{i_1})$, and the subpath $(\widetilde{w}_{2d}, \widetilde{w}_{2d+1})$ is either $(\tau_{i_d}, \tau_{[\gamma_d]})$ or $(\tau_{i_d}, \widetilde{\tau}_{i_{d+1}})$.

- If Δ_0 has three distinct sides, τ_{i_1} , $\tau_{[\gamma_0]}$, and $\tau_{[\gamma_{-1}]}$, then $\widetilde{\tau}_{[\gamma_{-1}]} = \tau_{[\gamma_{-1}]}$, and $\widetilde{\pi}$ maps $(\widetilde{w}_1, \widetilde{w}_2)$ to the (T^o, γ) -subpath (w_1, w_2) with the same labels.

If Δ_0 is a self-folded triangle with radius r and ℓ -loop ℓ , then recall that $\widetilde{\Delta}_0$ is an ordinary triangle with edge labels ℓ, r^{cc}, r (in counterclockwise order) as in Fig. 21(a). By (T1) and (T2), \widetilde{w}_1 goes along $\widetilde{\tau}_{[\gamma_0]}$ or $\widetilde{\tau}_{[\gamma_{-1}]}$ and \widetilde{w}_2 goes along ℓ . Hence, either $(\widetilde{w}_1, \widetilde{w}_2) = (r^{cc}, \ell)$




Figure 21. The lifts $(\tilde{w}_1, \dots, \tilde{w}_4)$ of Remark 3.18(A1), (A2) when γ starts at the puncture, see Fig. 14.




Figure 22. The lifts $(\tilde{w}_{2k}, \tilde{w}_{2k+1}, \tilde{w}_{2k+2})$ of the subpaths of Lemma 3.19(a1), (a2), (a3), see Fig. 15.




Figure 23. The lifts $(\tilde{w}_{2k+2}, \tilde{w}_{2k+3}, \tilde{w}_{2k+4})$ of the subpaths of Lemma 3.19(b1), (b2), (b3), see Fig. 16.

or $(\tilde{w}_1, \tilde{w}_2) = (r, \ell)$ (see Fig. 21). By Remark 3.18, there are two possible (T^o, γ) -subpaths (w_1, w_2) , either (r, ℓ_-) or (r, ℓ_+) . We see that $\bar{\pi}$ maps

$$\begin{aligned} (r, \ell) &\longleftrightarrow \text{Remark 3.18(A1): } (r, \ell_-), \\ (r^{cc}, \ell) &\longleftrightarrow \text{Remark 3.18(A2): } (r, \ell_+). \end{aligned}$$

- Similarly, if Δ_d has three distinct sides, then $(\tilde{w}_{2d}, \tilde{w}_{2d+1})$ is mapped to the subpath with the same labels. Otherwise, if Δ_d is a self-folded triangle with radius r and ℓ -loop ℓ , then recall that $\tilde{\Delta}_d$ is an ordinary triangle with edges ℓ, r^{cc}, r (in counterclockwise order), see Fig. 18(b). Then either $(\tilde{w}_{2d}, \tilde{w}_{2d+1}) = (\ell, r^{cc})$ or $(\tilde{w}_{2d}, \tilde{w}_{2d+1}) = (\ell, r)$, and $\bar{\pi}$ maps

$$\begin{aligned} (\ell, r^{cc}) &\longleftrightarrow (\ell_-, r), \\ (\ell, r) &\longleftrightarrow (\ell_+, r). \end{aligned}$$

Similarly, each $\tilde{\Delta}_k$ has three distinct labels $\tau_{i_k}, \tau_{i_{k+1}}$, and $\tilde{\tau}_{[\gamma_k]}$. As discussed in Section 3.2, the subpath $(\tilde{w}_{2k}, \tilde{w}_{2k+1}, \tilde{w}_{2k+2})$ is one of $(\tau_{i_k}, \tau_{i_{k+1}}, \tau_{i_{k+1}})$, $(\tau_{i_k}, \tau_{i_k}, \tau_{i_{k+1}})$, or $(\tau_{i_k}, \tilde{\tau}_{[\gamma_k]}, \tau_{i_{k+1}})$.

- If Δ_k ($k = 1, \dots, d$) has three distinct sides, $\tau_{i_k}, \tau_{i_{k+1}}$, and $\tau_{[\gamma_k]}$, then $\tilde{\tau}_{[\gamma_k]} = \tau_{[\gamma_k]}$. Hence $(\tilde{w}_{2k}, \tilde{w}_{2k+1}, \tilde{w}_{2k+2})$ is mapped to the (T^o, γ) -subpath $(w_{2k}, w_{2k+1}, w_{2k+2})$ with the same labels.

Otherwise, suppose Δ_k is a self-folded triangle with radius r and ℓ -loop ℓ . Assume that ℓ, r, ℓ are the k -th, $(k + 1)$ -th, and $(k + 2)$ -th arcs crossed by γ and that γ crosses them in the counterclockwise direction (see Fig. 18(c)). By construction (see Definition 4.1 and Fig. 18(e)), $\tilde{\Delta}_k$ has sides labeled $\tilde{\tau}_{i_k} = \ell$, $\tilde{\tau}_{i_{k+1}} = r$, and $\tilde{\tau}_{[\gamma_k]} = \dot{r}$ (in counterclockwise order). The subpath $(\tilde{w}_{2k}, \tilde{w}_{2k+1}, \tilde{w}_{2k+2})$ is one of $(\tilde{\tau}_{i_k}, \tilde{\tau}_{i_{k+1}}, \tilde{\tau}_{i_{k+1}}) = (\ell, r, r)$, $(\tilde{\tau}_{i_k}, \tilde{\tau}_{i_k}, \tilde{\tau}_{i_{k+1}}) = (\ell, \ell, r)$, or $(\tilde{\tau}_{i_k}, \tau_{[\gamma_k]}, \tilde{\tau}_{i_{k+1}}) = (\ell, \dot{r}, r)$. See Fig. 22. Per Lemma 3.19, there are three possible (T^o, γ) -subpaths for $(w_{2k}, w_{2k+1}, w_{2k+2})$. We see that $\bar{\pi}$ maps

$$\begin{aligned} (\ell, r, r) &\longleftrightarrow \text{Lemma 3.19(a1): } (\ell_-, \underline{r}, r), \\ (\ell, \ell, r) &\longleftrightarrow \text{Lemma 3.19(a2): } (\ell_+, \ell_-, r), \\ (\ell, \dot{r}, r) &\longleftrightarrow \text{Lemma 3.19(a3): } (\ell_+, \mathbf{r}, \mathbf{r} \text{ (counterclockwise non-backtrack)}). \end{aligned}$$

By construction (see Definition 4.1 and Fig. 18(e)), $\tilde{\Delta}_{k+1}$ has sides labeled $\tilde{\tau}_{i_{k+1}} = r$, $\tilde{\tau}_{i_{k+2}} = \ell$, and $\tilde{\tau}_{[\gamma_{k+1}]} = \dot{r}$ (in counterclockwise order). The subpath $(\tilde{w}_{2k+2}, \tilde{w}_{2k+3}, \tilde{w}_{2k+4})$ is one of $(\tilde{\tau}_{i_{k+1}}, \tilde{\tau}_{i_{k+2}}, \tilde{\tau}_{i_{k+2}}) = (r, \ell, \ell)$, $(\tilde{\tau}_{i_{k+1}}, \tilde{\tau}_{i_{k+1}}, \tilde{\tau}_{i_{k+2}}) = (r, r, \ell)$, or $(\tilde{\tau}_{i_{k+1}}, \tilde{\tau}_{[\gamma_{k+1}]}, \tilde{\tau}_{i_{k+2}}) = (r, \dot{r}, \ell)$. See Fig. 23. Per Lemma 3.19, there are three possible (T^o, γ) -subpaths for $(w_{2k+2}, w_{2k+3}, w_{2k+4})$. We see that $\bar{\pi}$ maps

$$\begin{aligned} (r, \ell, \ell) &\longleftrightarrow \text{Lemma 3.19(b1): } (r, \ell_-, \ell_+), \\ (r, r, \ell) &\longleftrightarrow \text{Lemma 3.19(b2): } (r, \underline{r}, \ell_-) \text{ then } \ell_-, \\ (r, \dot{r}, \ell) &\longleftrightarrow \text{Lemma 3.19(b3): } (\mathbf{r}, \mathbf{r} \text{ (counterclockwise non-backtrack)}, \ell_+). \quad \blacksquare \end{aligned}$$

We expect that Lemma 4.4 can be generalized to other punctured surfaces. However, other punctured surfaces can have more complicated ideal triangulations (like those containing an ideal triangle of Fig. 1(d)) that have not been considered in our argument.

4.3 Perfect matching expansion formula


We recap the snake graph expansion formula of [17, Theorem 4.10] for ordinary arcs of any bordered surface (including once-punctured polygons). We continue to restrict our attention to the case of the once-punctured n -gon \mathcal{C}_n .

Definition 4.5 (snake graph $G_{T^o, \gamma}$). We *unfold* \tilde{S}_γ into a graph $\overline{G}_{T^o, \gamma}$, called a *snake graph*, by inserting negative-oriented copies $\tilde{\Delta}_k^{(-)}$ of $\tilde{\Delta}_k$ (for $k = 1, \dots, d - 1$) into \tilde{S}_γ . Fig. 24(a) is the graph $\overline{G}_{T^o, \gamma}$ that is built from \tilde{S}_γ of Fig. 19(b). See [17, Section 4] for details. A *tile* consists of $\tilde{\Delta}_k$ and $\tilde{\Delta}_{k+1}$ (or $\tilde{\Delta}_k^{(-)}$ and $\tilde{\Delta}_{k+1}^{(-)}$) glued together along a *diagonal* labeled $\tilde{\tau}_{i_k}$. Let $G_{T^o, \gamma}$ denote the graph obtained from $\overline{G}_{T^o, \gamma}$ by removing the diagonal from each tile.

Definition 4.6 (crossing monomial). Define the *crossing monomial* of γ with respect to T^o to be

$$\text{cross}(T^o, \gamma) = \prod_{k=1}^d x(\tau_{i_k}).$$

Definition 4.7 (perfect matchings and weights). A *perfect matching* of a graph G is a subset E of the edges of G such that each vertex of G is incident to exactly one edge of E . See Fig. 25. If the edges of a perfect matching E are labeled $\beta_{j_1}, \dots, \beta_{j_r}$, then the *weight* $x(E)$ of E is the product $x_{\beta_{j_1}} \cdots x_{\beta_{j_r}}$.



(a) The snake graph $\overline{G}_{T^o, \gamma}$ constructed from \widetilde{S}_γ in Fig. 19(b). (b) A path corresponding to the $(\widetilde{S}_\gamma, \widetilde{\gamma})$ -path of Fig. 20(c), with odd steps $(b_1, 2, 2, \ell, \check{r}, b_3)$, see Fig. 25(8), and even steps the diagonals 1, 2, ℓ , r , and ℓ .

Figure 24. The snake graph $\overline{G}_{T^o, \gamma}$ of Definition 4.5 and an example of a path on $\overline{G}_{T^o, \gamma}$, see Remark 4.9.




Figure 25. The nine perfect matchings of the snake graph $G_{T^o, \gamma}$ of Fig. 24(a). Matchings (6)–(9) correspond to the (T^o, γ) -paths of Fig. 9. Note that these four are the only matchings where their restrictions to the gray-shaded tile are perfect matchings.

Theorem 4.8 ([17, Theorem 4.10], perfect matching expansion formula). *Let T^o be an ideal triangulation of any surface S and let $\gamma \notin T^o$ be an ordinary arc (note: γ may be an ℓ -loop). Let x_γ denote the element corresponding to γ in the cluster algebra which arises from S (see Theorem 2.8). Then*

$$x_\gamma = \sum_E \frac{x(E)}{\text{cross}(T^o, \gamma)},$$

where the sum is over all perfect matchings E of $G_{T^o, \gamma}$.




Figure 26. The subpaths on $\overline{G}_{T^o, \gamma}$ corresponding to the subpaths $(\tilde{w}_{2k}, \tilde{w}_{2k+1}, \tilde{w}_{2k+2}) = (\ell, \ , r)$ from Fig. 22.




Figure 27. The subpaths on $\overline{G}_{T^o, \gamma}$ corresponding to the subpaths $(\tilde{w}_{2k+2}, \tilde{w}_{2k+3}, \tilde{w}_{2k+4}) = (r, \ , \ell)$ from Fig. 23.

4.4 Proof of the T^o -path formula for \mathcal{C}_n (Theorem 3.8)

Remark 4.9 ([16, Lemma 4.5]). There is a bijection

$$F: \{\text{perfect matchings of the graph } G_{T^o, \gamma}\} \rightarrow \{(\widetilde{S}_\gamma, \widetilde{\gamma})\text{-paths on } \widetilde{S}_\gamma\},$$

$$F: E \mapsto \tilde{w}_E,$$

where the diagonals of $\overline{G}_{T^o, \gamma}$ correspond to the even-indexed steps of each $(\widetilde{S}_\gamma, \widetilde{\gamma})$ -path (see Fig. 24(b)) and each matching E correspond to the odd-indexed steps of \tilde{w}_E . See Figs. 26 and 27.

The bijections $\bar{\pi}$ (of Lemma 4.4) and F compose to form a bijection between the perfect matchings of the snake graph $G_{T^o, \gamma}$ and the (T^o, γ) -paths on \mathcal{C}_n .

Theorem 4.10 (snake graph matchings to (T^o, γ) -paths). *Consider an arc γ of \mathcal{C}_n and the (T^o, γ) -paths and snake graph $G_{T^o, \gamma}$ corresponding to γ . The map*

$$\bar{\pi} \circ F: \{\text{perfect matchings in } G_{T^o, \gamma}\} \rightarrow \{(T^o, \gamma)\text{-paths on } \mathcal{C}_n\},$$

$$E \mapsto w_E := \bar{\pi}(\tilde{w}_E)$$

is a bijection where the edges of each matching E correspond to the odd-indexed steps of the (T^o, γ) -path w_E , see Fig. 24(b). In particular, the weight $x(E)$ of the perfect matching E is equal to the numerator of the Laurent monomial $x(w_E)$ (see Definitions 3.6 and 4.7).

Remark 4.11 (matching restrictions and (T^o, γ) -subpaths). The four (T^o, γ) -subpaths of Lemma 3.19(I)–(IV) correspond to the four possible matchings of the triple-tiles of $G_{T^o, \gamma}$. Compare Figs. 15–16, Figs. 22–23, and Figs. 26–27.

Proof of Theorem 3.8. Applying the bijection $\bar{\pi} \circ F$ to the formula of [17, Theorem 4.10] yields

$$x_\gamma = \frac{1}{\text{cross}(T^o, \gamma)} \sum_{\substack{E \text{ perfect matchings} \\ \text{of } G_{T^o, \gamma}}} x(E) = \frac{1}{\text{cross}(T^o, \gamma)} \sum_{(T^o, \gamma)\text{-paths } w} \left(\prod_{k=0}^d x(w_{2k+1}) \right).$$

Since $\prod_{k=1}^d x(w_{2k}) = \text{cross}(T^o, \gamma)$, this concludes the proof. \blacksquare

Example 4.12. The nine perfect matchings of the graph $G_{T^o, \gamma}$ (from Fig. 24(a)) are listed in Fig. 25. Per Theorem 4.10, they correspond to the nine (T^o, γ) -paths of Example 3.10.

5 Combinatorial proof of atomic bases for type D

5.1 Atomic bases for the once-punctured n -gons (type D_n)

Definition 5.1 (cluster monomials). Let \mathcal{A} be a coefficient-free cluster algebra. A *cluster monomial* is a monomial in cluster variables all belonging to a single cluster.

Recall that, in the case that \mathcal{A} arises from a surface, a cluster corresponds to a tagged triangulation and a cluster monomial corresponds to a multi-tagged triangulation (see Theorem 2.8). See Definition 2.6 for a list of compatible pairs of tagged arcs of \mathcal{C}_n .

The concepts of positive elements and atomic bases were first introduced in [23] for the case of an annulus with one marked point on each boundary.

Definition 5.2 (positive elements and atomic bases). An element $y \in \mathcal{A}$ is called *positive* if the Laurent expansion of y in the variables of every cluster of \mathcal{A} has non-negative coefficients.

A \mathbb{Z} -linear basis \mathcal{B} of \mathcal{A} is called an *atomic basis* of \mathcal{A} if any positive element of \mathcal{A} is a non-negative \mathbb{Z} -linear combination of \mathcal{B} . Note that, if such a \mathcal{B} exists, \mathcal{B} is the collection of all indecomposable positive elements (i.e., elements which cannot be written as a sum of positive elements) of \mathcal{A} , hence it is unique.

Remark 5.3. To prove that a collection $\mathcal{B} \subset \mathcal{A}$ is an atomic basis of \mathcal{A} , it suffices to verify the following:

$$\text{every element of } \mathcal{B} \text{ is a positive element of } \mathcal{A}, \quad (5.1)$$

$$\mathcal{B} \text{ is a } \mathbb{Z}\text{-linear basis of } \mathcal{A}, \quad (5.2)$$

$$\text{every positive element } y \in \mathcal{A} \text{ can be written as a } \mathbb{Z}_{\geq 0}\text{-linear combination of } \mathcal{B}. \quad (5.3)$$

Theorem 5.4 ([4]). *If \mathcal{A} is a cluster algebra of type A , D , or E , the cluster monomials of \mathcal{A} form the atomic basis of \mathcal{A} .*

Proof. [14, 19], [2], and [4, 6] give representation theoretic proofs for (5.1), (5.2), and (5.3), respectively. [17] and [18] provide combinatorial arguments for (5.1) and (5.2), respectively, and [7] gives a combinatorial proof for type A for (5.3). We present in the remainder of this paper a combinatorial proof for type D for (5.3) which relies on the T^o -path formula (Theorem 3.8) and is inspired by [7]. \blacksquare

Recall that a multi-tagged triangulation Σ is not compatible with a tagged triangulation T if Σ contains an arc σ which is not in T . Recall also that a *proper Laurent monomial in variables* u_i is a product of the form $u_1^{c_1} \cdots u_r^{c_r}$ where at least one of the c_i is negative.

Lemma 5.5. *Consider a once-punctured polygon \mathcal{C}_n . For every tagged triangulation T and every multi-tagged triangulation Σ which is not compatible with T , the T -expansion of x_Σ is a sum of proper Laurent monomials.*

Remark 5.6. A general version of Lemma 5.5 is known as the *proper Laurent monomial property* (following [6]) which was proven for cluster algebras from surfaces and any choice of coefficients in [6] and then for any skew-symmetric cluster algebras in [5], using representation theoretic arguments in both cases.

Definition 5.7. Let N be a subset of a tagged triangulation T . We write the *degree with respect to N* to mean the degree with respect to the cluster variables corresponding to N .

Outline of proof of Lemma 5.5. Suppose x_Σ is a cluster monomial not compatible with a tagged triangulation T , i.e., Σ contains an arc σ which is not in T . For brevity, suppose that all tagged arcs of $\Sigma \setminus T$ are peripheral and the corresponding ideal triangulation T° has no self-folded triangle. For full details and the rest of the cases, see the remainder of this paper.

- Step 1: Choose a tagged arc $\sigma \in \Sigma \setminus T$ such that σ is as close as possible to the puncture and, if possible, σ only crosses every arc of T at most once. Let (T, σ) -cross (respectively, (T, σ) -doublecross) be the set of arcs of T which σ crosses (respectively, the set of arcs of T which σ crosses twice).
- Step 2: For each (T, σ) -path w , we compare the number of odd-indexed steps (contributing to the numerator of $x(w)$) and the number of even-indexed steps (contributing to the denominator of $x(w)$) that belong to (T, σ) -cross and (T, σ) -doublecross. See Definition 3.6. This allows us to show that each term in the T -expansion of x_σ is of *negative* degree either with respect to the cluster variables corresponding to (T, σ) -cross or with respect to the cluster variables corresponding to (T, σ) -doublecross.
- Step 3: Similarly, for each factor x_β in the product of x_Σ , we consider a (T, β) -path and compare the number of odd steps versus the number of even steps to show that every term in the T -expansion of x_β has *non-positive* degree with respect to both the cluster variables corresponding to (T, σ) -cross and the cluster variables corresponding to (T, σ) -doublecross.
- Step 4: It follows, since $\sigma \in \Sigma$ and every term in the T -expansion of x_Σ has non-positive degree with respect to both (T, σ) -cross and (T, σ) -doublecross that every term in the T -expansion of x_Σ has *negative* degree with respect to either (T, σ) -cross or (T, σ) -doublecross.

Since (T, γ) -cross and (T, γ) -doublecross are subsets of T , the T -expansion of x_Σ is a sum of proper Laurent monomials. ■

Theorem 5.8. *If \mathcal{A} is a coefficient-free cluster algebra of type D , every positive element $y \in \mathcal{A}$ is equal to a linear combination $y = \sum_{\Gamma} m_{\Gamma} x_{\Gamma}$ of cluster monomials where each m_{Γ} is non-negative.*

Proof of Theorem 5.8. We assume (5.1) and (5.2) for $\mathcal{B} = \{\text{cluster monomials}\}$. The following argument appears in [6, 7, 23], and we include it here for completeness. Let y be a positive element of a cluster algebra of type D . Write $y = \sum_{\Gamma} m_{\Gamma} x_{\Gamma}$ as a linear combination of cluster monomials. Then we prove that every m_{Γ} is non-negative as follows.

Let x_{Γ} be a cluster monomial from the sum, and let Γ denote the multi-tagged triangulation corresponding to it. Choose a tagged triangulation $T := T_{\Gamma}$ that is compatible with the multi-tagged triangulation Γ , i.e., so that $\gamma \in T$ for every arc $\gamma \in \Gamma$. Consider a different cluster monomial x_{Σ} from the sum. We argue that

$$x_{\Gamma} \text{ does not appear in the } T\text{-expansion of } x_{\Sigma}. \tag{5.4}$$

If all the arcs of Σ were in T , then x_Σ is its own T -expansion, and we are done. Suppose that Σ has an arc that is not in T . By Lemma 5.5, every term in the T -expansion of x_Σ is a proper Laurent monomial in the cluster corresponding to T . Since x_Γ is a monomial (i.e., not a proper Laurent monomial) in this cluster, we have proven (5.4).

Hence, m_Γ is equal to the coefficient of x_Γ in the T -expansion of y . Since y is a positive element, the coefficient of x_Γ in the T -expansion of y is non-negative, as required. ■

In Section 5.2, we provide notations and lemmas which are helpful toward proving the lemmas in Sections 5.3 and 5.4. Section 5.3 (respectively, Section 5.4) discusses lemmas which are needed to prove Lemma 5.5 for the case where all arcs of $\Sigma \setminus T$ are peripheral (respectively, for the case where $\Sigma \setminus T$ contains a radius). Even though a few of the lemmas can also be proven using the existing snake graph formula [17, Theorem 4.3] (recapped in Section 4.3), many arguments (such as the proofs of Lemmas 5.23, 5.24, and 5.26) can be done more easily using the T^o -path formula. Finally, we prove Lemma 5.5 in Section 5.5: if Σ is not compatible with the tagged triangulation T , we can choose a tagged arc $\sigma \in \Sigma \setminus T$ such that every term in the T -expansion of x_Σ has negative degree with respect to some subset N_σ^T of $T \setminus \Sigma$ by Lemmas 5.23, 5.24, 5.25, 5.28, 5.26, and 5.29.

5.2 Notations and technical lemmas to prove Lemma 5.5

Recall from Remark 2.5 that every tagged arc of \mathcal{C}_n belongs to one of three classes: plain radius, notched radius, or peripheral.

Remark 5.9. Per Remark 2.9, we only need to prove Lemma 5.5 and the related statements for two cases: one where T has all plain radii and one where T has 2 parallel radii $r, r^{(p)}$.

Definition 5.10. Let T be a tagged triangulation, T^o the corresponding ideal triangulation, and λ a tagged arc. Let (T^o, λ) -cross denote the arcs of T^o that cross λ , and let (T^o, λ) -doublecross denote the arcs of T^o that cross λ twice. (Note that, If T contains two parallel radii $r, r^{(p)}$, then (T^o, λ) -cross and (T^o, λ) -doublecross may contain the associated ℓ -loop.)

If T contains two parallel radii $r, r^{(p)}$ (so that T^o contains a self-folded triangle with radius r and ℓ -loop ℓ), and if λ crosses the ℓ -loop ℓ , let

$$\begin{aligned} (T, \lambda)\text{-cross} &:= \{r^{(p)} \text{ and the peripheral arcs of } T \text{ that cross } \lambda\}, \text{ and} \\ (T, \lambda)\text{-doublecross} &:= \{r^{(p)} \text{ and the (peripheral) arcs of } T \text{ that cross } \lambda \text{ twice}\}. \end{aligned}$$

Otherwise, if λ does not cross an ℓ -loop that belongs to T^o , let

$$(T, \lambda)\text{-cross} := (T^o, \lambda)\text{-cross} \quad \text{and} \quad (T, \lambda)\text{-doublecross} := (T^o, \lambda)\text{-doublecross}.$$

Lemma 5.11. *Let T be a tagged triangulation with two parallel radii $r, r^{(p)}$ (so that T^o has a self-folded triangle with radius r and ℓ -loop ℓ). Suppose β and σ (possibly $\beta = \sigma$) are compatible tagged arcs not in T , and either σ is a plain radius or β is peripheral. Let $w = (w_1, \dots, w_{2d+1})$ be a (T^o, β) -path. Recall that $x_\ell := x_r x_{r^{(p)}}$.*

Then $x(w)$ having negative (respectively, non-positive) degree with respect to (T^o, σ) -cross implies that $x(w)$ has negative (respectively, non-positive) degree with respect to (T, σ) -cross.

Similarly, if (T^o, σ) -doublecross is nonempty and if $x(w)$ has negative (respectively, non-positive) degree with respect to (T^o, σ) -doublecross, then $x(w)$ has negative (respectively, non-positive) degree with respect to (T, σ) -doublecross.

Note that this statement is not necessarily true if σ is peripheral and β is a radius.




Figure 28. λ cuts out a smaller once-punctured disk \mathbf{C}_λ and a disk \mathbf{Disk}_λ , shaded in gray.

Proof. If σ is a radius (Fig. 18(a)), then it cannot cross r , so $r \notin (T^\circ, \sigma)$ -cross. Since $r \notin (T, \sigma)$ -cross, replacing the two gradings (T°, σ) -cross and (T, σ) -cross is equivalent to simply replacing ℓ in (T°, σ) -cross with $r^{(p)}$ in (T, σ) -cross, and the degree would stay the same.

If β is a peripheral arc crossing r (Figs. 18(d), 18(c)), then (T°, σ) -cross contains r , but w goes along r exactly twice in a row, per Lemma 3.19(I)–(IV). So $x(w)$ (as a T° -monomial) has zero degree with respect to r , and we can again replace ℓ in (T°, σ) -cross with $r^{(p)}$ in (T, σ) -cross to arrive at the same conclusion.

Since r is a radius, σ can cross it at most once, so $r \notin (T^\circ, \sigma)$ -doublecross. Since $r \notin (T, \sigma)$ -doublecross, replacing the two gradings (T°, σ) -doublecross and (T, σ) -doublecross is equivalent to simply replacing ℓ in (T°, σ) -doublecross with $r^{(p)}$ in (T, σ) -doublecross, and the degree would stay the same. ■

Notation 5.12. Every ordinary peripheral arc λ snips the surface into a smaller once-punctured disk (denoted \mathbf{C}_λ) and a region not containing the puncture (denoted \mathbf{Disk}_λ). See Fig. 28.

Let σ be a peripheral arc with endpoints s and t (possibly $s = t$), and let β be another ordinary arc. We say that β lies in \mathbf{Disk}_σ (respectively, \mathbf{C}_σ) if the interior of β lies entirely in the interior of \mathbf{Disk}_σ (respectively, \mathbf{C}_σ).

Let v be an endpoint of β . We say that v lies in \mathbf{Disk}_σ if v is a marked point on the boundary of \mathbf{Disk}_σ , possibly $v = s$ or t . We say that v lies in \mathbf{C}_σ if v is a marked point in the interior or on the boundary of \mathbf{C}_σ , possibly $v = s$ or t . We say that v lies strictly in \mathbf{Disk}_σ (respectively, \mathbf{C}_σ) if v lies in \mathbf{Disk}_σ (respectively, \mathbf{C}_σ) and v is not equal to s nor t . We say v lies outside of \mathbf{Disk}_σ if v is not a marked point of \mathbf{Disk}_σ (hence v is not equal to s nor t). We use the same language when the region is a disk cut out by two radii.

We use Lemmas 5.13 and 5.14 to prove Lemma 5.15(iii), (iv).

Lemma 5.13. *Suppose γ and σ are distinct compatible ordinary arcs not in T° . If σ crosses $\tau_{[\gamma_k]}$, then σ crosses arc k^γ or $k + 1^\gamma$, the k -th and $(k + 1)$ -th arcs crossed by γ .*

Proof. To simplify the proof, assume that $0 < k < d$. There are four possibilities for ways of γ to cross the ideal triangle Δ_k^γ , see Fig. 6. If Δ_k^γ is self-folded, then $\tau_{[\gamma_k]}$ is equal to either arc k^γ or $k + 1^\gamma$, and we are done. If arc k^γ , arc $k + 1^\gamma$, and $\tau_{[\gamma_k]}$ are all distinct, then γ_k (the segment between p_k^γ and p_{k+1}^γ contained in Δ_k^γ) cuts Δ_k^γ into two regions, and one of the regions is an ideal quadrilateral $Q = \text{Quad}(\Delta_k^\gamma)$ with sides $\tau_{[\gamma_k]}$, γ_k , part of arc k^γ , and part of arc $k + 1^\gamma$ (see Figs. 6(a), 6(b), and 6(c)). By assumption, σ crosses $\tau_{[\gamma_k]}$, so σ cuts through Q . Since σ does not cross γ_k (because σ and γ are compatible by assumption), σ has to cross either arc k^γ or $k + 1^\gamma$. ■

Lemma 5.14. *Suppose γ and σ are (distinct, compatible) peripheral arcs such that γ that is contained in the interior of \mathbf{Disk}_σ (i.e., σ is closer than γ to the puncture, see Fig. 29). If σ crosses $\tau_{[\gamma_k]}$ twice, then σ crosses arc k^γ twice or σ crosses $k + 1^\gamma$ twice.*




Figure 29. Lemma 5.14, when σ crosses $\tau_{[\gamma_k]}$ twice.

Proof. To simplify the proof, assume $0 < k < d$. Let v be the marked point adjacent to both arcs k^γ and $(k+1)^\gamma$. Suppose σ crosses $\tau_{[\gamma_k]}$ twice. There are four possibilities for ways of γ to cross the ideal triangle Δ_k , see Fig. 6. If Δ_k^γ is self-folded, then $\tau_{[\gamma_k]}$ is equal to either arc k^γ or $(k+1)^\gamma$, and we are done.

Suppose arc k^γ , arc $(k+1)^\gamma$, and $\tau_{[\gamma_k]}$ are all distinct. Since γ is peripheral, γ cuts the surface \mathcal{C}_n into two regions, the smaller once-punctured disk \mathbf{C}_γ containing σ , and the disk \mathbf{Disk}_γ containing v .

We claim that $\tau_{[\gamma_k]}$, arc k^γ , and arc $(k+1)^\gamma$ are all peripheral (one of them possibly an ℓ -loop). First, $\tau_{[\gamma_k]}$ must be peripheral because a radius can be crossed at most once. Because v is in the disk \mathbf{Disk}_γ , v is not the puncture, and so the arcs k and $(k+1)$ are also peripheral.

The puncture is either closest to arc k^γ , arc $(k+1)^\gamma$, or $\tau_{[\gamma_k]}$ (see Fig. 29). Since σ cuts out a disk containing γ , if the puncture is closest to arc k^γ (respectively, to arc $(k+1)^\gamma$), then σ has to cross arc k^γ (respectively, arc $(k+1)^\gamma$) twice. See Figs. 29(g) and 29(j) (respectively, Figs. 29(h) and 29(k)). If the puncture is closest to $\tau_{[\gamma_k]}$, then σ has to cross either arc k^γ or arc $(k+1)^\gamma$ twice (Figs. 29(i) and 29(l)). \blacksquare

Lemma 5.15. Let T° be an ideal triangulation. Suppose β and σ are compatible ordinary arcs, with $\beta \notin T^\circ$, and possibly $\beta = \sigma$. Let $w = (w_1, \dots, w_{2d+1})$ be a (T°, β) -path. Suppose we have one of these four scenarios:

- i) Let $N_\sigma^{T^\circ} := \{\text{all radii of } T^\circ\}$ and suppose β is peripheral. (Note that $N_\sigma^{T^\circ}$ does not depend on σ here.)
- ii) Let σ be a radius of T° , and let $N_\sigma^{T^\circ} := \{\text{all radii of } T^\circ\} \setminus \{\sigma\}$, and suppose β is peripheral.
- iii) Let $N_\sigma^{T^\circ}$ denote (T°, σ) -cross, if $\sigma \notin T$.
- iv) Let $N_\sigma^{T^\circ}$ denote (T°, σ) -doublecross if (T°, σ) -doublecross is non-empty, and suppose that β is a peripheral arc such that either $\beta = \sigma$ or β is contained in the interior of \mathbf{Disk}_σ .

In all four of these scenarios, if $w_{2k+1} \in N_\sigma^{T^\circ}$, then there is a step w_{2k} or w_{2k+2} which goes along an arc of $N_\sigma^{T^\circ}$.

Proof of Lemma 5.15. Suppose w_{2k+1} belongs to $N_\sigma^{T^\circ}$. To simplify the proof, we only discuss cases where $k \neq 0, d$. Recall that (T1) requires β to cross each even step w_{2i} of w .

First, we prove the assertion in scenarios (i) and (ii). If w_{2k+1} goes along a radius from the boundary to the puncture, we see that w_{2k+2} must be a radius, as needed for scenario (i). Furthermore, since β crosses w_{2k+2} but not σ , we see that w_{2k+2} is a radius that is not equal to σ , proving scenario (ii). If w_{2k+1} goes from the puncture to the boundary, we can repeat a similar argument to show that $w_{2k} \in N_\sigma^{T^\circ}$.

In scenarios (iii) and (iv), recall that the steps w_{2k} , w_{2k+1} , and w_{2k+2} must go along edges in the $(k+1)$ -th ideal triangle Δ_k^β crossed by β , and β crosses w_{2k} and w_{2k+2} . If w_{2k+1} goes along the same arc as w_{2k} or w_{2k+2} , then we are done. Hence assume that the ideal triangle Δ_k^β has three distinct edges $w_{2k} = \tau_{i_k}^\beta$, $w_{2k+1} = \tau_{[\beta_k]}$, $w_{2k+2} = \tau_{i_{k+1}}^\beta$.

We consider scenario (iii): If $\beta = \sigma$, then w_{2k} and w_{2k+2} cross $\beta = \sigma$, and we are done. Assume that $\beta \neq \sigma$. By Lemma 5.13, since σ crosses $\tau_{[\beta_k]}$, then σ crosses $\tau_{i_k}^\beta$ or $\tau_{i_{k+1}}^\beta$.

Finally, we consider scenario (iv): By assumption, $\beta = \sigma$ or β is peripheral arc contained in the interior of \mathbf{Disk}_σ .

First, suppose $\beta = \sigma$. Having distinct w_{2k} , w_{2k+1} , and w_{2k+2} means that the segment σ_k cuts Δ_k into two regions, and one of the regions contains w_{2k+1} . Since σ cuts w_{2k+1} twice, σ cuts Δ_k^σ three times. Since the surface \mathcal{C}_n only contains one puncture, this is impossible, hence either $w_{2k} = w_{2k+1}$ or $w_{2k+2} = w_{2k+1}$.

Next, suppose β is a peripheral arc ($\beta \neq \sigma$) contained in the interior of \mathbf{Disk}_σ . By Lemma 5.14 (see Fig. 29), since σ crosses $w_{2k+1} = \tau_{[\beta_k]}$ twice, σ crosses $\tau_{i_k}^\beta$ or $\tau_{i_{k+1}}^\beta$ twice. Hence σ crosses w_{2k} or $w_{2(k+1)}$, as needed. ■

In particular, Lemma 5.15 implies Corollary 5.17.

Definition 5.16. Let w be a T° -path. Let N be a subset of T° . A consecutive subpath $w' = (w_a, \dots, w_b)$ of w is called an N -subpath if each step w_j of w' comes from N . We say that w' is a *longest N -subpath* of w if the step w_{a-1} (if any) and the step w_{b+1} (if any) do not come from N .

Corollary 5.17. *Suppose we have the same setup as Lemma 5.15.*

- a) If β does not cross any arc of $N_\sigma^{T^\circ}$, then w does not contain any arc of $N_\sigma^{T^\circ}$, and the degree of $x(w)$ with respect to $N_\sigma^{T^\circ}$ is zero.
- b) If the corresponding term $x(w)$ has positive degree with respect to $N_\sigma^{T^\circ}$, there must be an odd-length (of length three or greater) longest $N_\sigma^{T^\circ}$ -subpath $\bar{w} = w_{2i-1}, \dots, w_{2j+1}$ of w .

Proof. a) If β does not cross any arc of $N_\sigma^{T^\circ}$, then by (T1) there is no even-indexed step w_{2k} from $N_\sigma^{T^\circ}$. By Lemma 5.15, there is no odd-indexed step from $N_\sigma^{T^\circ}$.

b) Suppose $x(w)$ has a positive degree with respect to $N_\sigma^{T^\circ}$. Then $|\text{odd-indexed steps from } N_\sigma^{T^\circ}| > |\text{even-indexed steps from } N_\sigma^{T^\circ}|$ by the T° -path formula, and so there is at least one




Figure 30. Lemma 5.18, when σ crosses two peripheral arcs α, τ_{i_1} of Δ_0 .

odd-indexed step from $N_\sigma^{T^\circ}$. By Lemma 5.15, each odd-indexed step $w_{2k+1} \in N_\sigma^{T^\circ}$ must either follow $w_{2k} \in N_\sigma^{T^\circ}$ or precede $w_{2k+2} \in N_\sigma^{T^\circ}$, and so the assertion follows. \blacksquare

Lemma 5.18. *Suppose T° is an ideal triangulation and $\sigma \notin T^\circ$ is a peripheral arc from v_1 to v'_1 (with $v_1 \neq v'_1$). Let $\tau := \tau_{i_1}$, the edge opposite v_1 in the first ideal triangle Δ_0 through which σ passes. Let α and λ denote the two sides (the arcs $\tau_{[\sigma_0]}, \tau_{[\sigma_{-1}]}$, not necessarily in this order) of Δ_0 adjacent to σ at v_1 . Suppose σ crosses α (see Fig. 30). Then:*

- α is a peripheral arc with distinct endpoints.
- λ is a peripheral arc (or boundary edge) with distinct endpoints.
- τ_{i_1} is peripheral (possibly an ℓ -loop) and $\tau_{i_1} \in (T^\circ, \sigma)$ -doublecross.
- There is no arc from (T°, σ) -cross that is adjacent to σ at v'_1 , and α is the only arc of (T°, σ) -cross that is adjacent to σ . Consequently, if $w = (w_1, \dots, w_{2d+1})$ is a (T°, σ) -path, then either $w_1 \notin (T^\circ, \sigma)$ -cross or $w_{2d+1} \notin (T^\circ, \sigma)$ -cross.
- σ crosses all radii of T° .

Proof. Observe that, since v_1 lies on the boundary, the first ideal triangle σ crosses, Δ_0 , is not self-folded, and so all its edges $\tau_{i_1}, \lambda, \alpha$ are pairwise distinct.

- First, we prove part (a): We know that α cannot be a radius, because a radius from v_1 cannot cut an arc adjacent to v_1 . Furthermore, since an ℓ -loop based at v_1 would have to lie entirely in \mathbf{C}_σ , no ℓ -loop based on v_1 can cross σ , hence α is a non- ℓ -loop peripheral arc.
- Second, we prove part (b) of this lemma.
 - Suppose for contradiction that λ is an ℓ -loop. Again, since no ℓ -loop based on v_1 can cross σ , we see that λ and σ do not cross. Hence λ would cut out a region \mathbf{Disk}_λ not containing the puncture but containing σ as well as α . It is impossible for α and σ to both cross and be adjacent in this region, hence λ is not an ℓ -loop.
 - Suppose for contradiction that λ is a radius. Then Δ_0 is an ordinary triangle with the puncture as one of the vertices. Hence the region outside of Δ_0 contains no puncture, so it is impossible for σ to cross Δ_0 a second time. Therefore σ cannot cross α , contradicting our assumption.

Hence λ is a peripheral arc with distinct endpoints or a boundary edge.

- We now prove part (c) of this lemma. By (a) and (b), $\tau = \tau_{i_1}$ is also a peripheral arc (possibly an ℓ -loop). Note that the puncture must be closer to τ_{i_1} than to α . Hence, the second arc crossed by σ is contained outside of \mathbf{Disk}_τ , while α lies in \mathbf{Disk}_τ . Since σ crosses α , σ must cross τ the second time, as needed for part (c).

- d) Next, we prove part (d) of this lemma. Suppose that $\alpha' \in T^o$ is adjacent to the other endpoint v'_1 of σ . Note that α cuts out a disk \mathbf{Disk}_α containing v'_1 , and no arc of T^o in the interior of \mathbf{Disk}_α can both be adjacent to σ and belong to (T^o, σ) -cross. But both α and α' belong to T^o so they do not cross, hence α' is contained in \mathbf{Disk}_α , and so α' cannot both be adjacent to σ and belong to (T^o, σ) -cross.

Furthermore, by definition the only arcs that are adjacent to σ at v_1 are $\tau_{[\sigma_0]}$ and $\tau_{[\sigma_{-1}]}$ (i.e., α and λ). But λ is contained in \mathbf{Disk}_σ , so λ cannot cross σ .

- e) Finally, we prove part (e) of this lemma. If τ_{i_1} is an ℓ -loop, then σ must cross the radius (which is the only radius of T^o) containing it. Otherwise, since $\tau_{i_1} \in T^o$ is peripheral, all radii of T^o are contained in \mathbf{C}_τ . Since the segment of σ that is contained in \mathbf{C}_τ is homotopic to the segment of the original surface's boundary along \mathbf{C}_τ and since the two endpoints of σ lie *strictly* in \mathbf{Disk}_τ , it follows that σ must cross all the radii. ■

The following lemma is also used many times throughout the rest of this paper.

Lemma 5.19. *Suppose T^o is an ideal triangulation, $\beta \notin T^o$ is an ordinary arc, and $w = (w_1, \dots, w_{2d+1})$ is a (T^o, β) -path. Let N be a subset of T^o .*

- a) *If $|\text{even steps from } N| \geq |\text{odd steps from } N|$, then $x(w)$ has non-positive degree with respect to N . Furthermore, inequality implies negative degree.*
- b) *$x(w)$ has non-positive degree with respect to (T^o, β) -cross.*
- c) *If $w_1, w_{2d+1} \notin (T^o, \beta)$ -cross, then $x(w)$ has negative degree with respect to (T^o, β) -cross. Hence, if β is not adjacent to any arc that crosses it, $x(w)$ has negative degree with respect to (T^o, β) -cross.*

Proof. Consider a (T^o, β) -path $w = (w_1, \dots, w_{2d+1})$. Part (a) follows from the T^o -path formula.

The corresponding term $x(w)$ in the T^o -expansion of x_β has $d + 1$ factors in the numerator and d factors in the denominator. All the factors in the denominator correspond to the arcs of T^o that cross β . By Lemma 5.18(d), at least one of the endpoints of β is not adjacent to any arc from (T^o, β) -cross, so either the first step w_1 or the last step w_{2d+1} of w does not come from (T^o, β) -cross by (T2). Hence, there are at most d odd steps contributing to the degree. Since there are exactly d even steps from (T^o, β) -cross, $x(w)$ has non-positive degree with respect to the arcs of T^o which cross σ , satisfying part (b).

If the first and last steps w_1, w_{2d+1} of w do not come from (T^o, σ) -cross, there are at most $d - 1$ odd steps which contribute to the degree, and (c) follows. Furthermore, if β is not adjacent to any arc from (T^o, σ) -cross, then $w_1, w_{2d+1} \notin (T^o, \sigma)$ -cross by (T2). ■


5.3 Technical lemmas to prove Lemma 5.5 for cases where all arcs of $\Sigma \setminus T$ are peripheral

5.3.1 Proving that x_Σ has non-positive degree with respect to (T, σ) -cross

Lemma 5.20. *Let T be a tagged triangulation and suppose σ is a tagged peripheral arc not in T .*


- a) *Then each term in the T -expansion of x_σ has non-positive degree with respect to (T, σ) -cross.*
- b) *If (T^o, σ) -doublecross is empty, then each term in the T -expansion of x_σ has negative degree with respect to (T, σ) -cross.*

Proof. By Lemma 5.19(b), each term in the T^o -expansion of x_σ has non-positive degree with respect to (T^o, σ) -cross.



(a) The peripheral arcs $\lambda_1, \dots, \lambda_r$ as close as possible to the puncture. (b) If (T°, σ) -doublecross is non-empty, then $r = 1$.

Figure 31. Proof of Lemma 5.5: the case where $\Sigma \setminus T$ are all peripheral.



(a) Correct setup for Lemma 5.23 but wrong setup for Lemma 5.24: σ is not the only central arc. (b) Correct setup for both Lemmas 5.23 and 5.24: σ is the only central arc.

Figure 32. Setups for Lemmas 5.23 and 5.24.

By Lemma 5.18(c), if (T°, σ) -doublecross is empty, then σ is not adjacent to any arc from (T°, σ) -cross. Therefore, by Lemma 5.19(c), $x(w)$ has negative degree with respect to (T°, σ) -cross.

By Lemma 5.11, both assertions follow. \blacksquare

Definition 5.21. Let Λ be a multi-tagged triangulation containing only peripheral arcs. We say that $\lambda \in \Lambda$ is *central in Λ* if λ is as close as possibly to the puncture, i.e.,

if $\beta \in \Lambda \setminus \{\lambda\}$, then the disk \mathbf{Disk}_β does not contain λ .

Remark 5.22. Let $\{\lambda_1, \dots, \lambda_r\}$ be the set of central arcs of Λ . Then, by definition, every $\beta \in \Lambda$ is contained in $\mathbf{Disk}_{\lambda_j}$ for some j . See Fig. 31(a).

Lemma 5.23. Suppose Σ is a multi-tagged triangulation and T is a tagged triangulation. Suppose $\sigma \in \Sigma \setminus T$ is a tagged peripheral arc that is central in $\Sigma \setminus T$, i.e.,

if $\beta \in \Sigma \setminus T$ is not σ , then the disk \mathbf{Disk}_β does not contain σ (see Fig. 32). (5.5)

Let β be any tagged arc in Σ such that, if $\beta \in \Sigma \setminus T$, then β is peripheral. See Figs. 32(a) and 32(b). Then each term in the T -expansion of x_β has non-positive degree with respect to (T, σ) -cross.

Proof. Let s^σ be the starting point and let t^σ be the finishing point of σ . Since σ is not an ℓ -loop by assumption, $s^\sigma \neq t^\sigma$.

If $\beta \in T$, then its degree with respect to any of the above gradings is zero since it cannot cross σ . If $\beta = \sigma$, we are done by Lemma 5.20.

Otherwise, suppose $\beta \in \Sigma \setminus T$ and $\beta \neq \sigma$, and so β is peripheral by assumption. Let $w = (w_1, \dots, w_{2d+1})$ be a (T°, β) -path. First, we prove that

the T° -expansion of $x(w)$ has non-positive degree with respect to (T°, σ) -cross. (5.6)

For contradiction, suppose otherwise. By Corollary 5.17(b), there must be a longest (T^σ, σ) -cross-subsequence $\bar{w} = (w_{2i-1}, \dots, w_{2j+1})$ of w (of length three or greater). Since σ and β are compatible, there are two possibilities for β : either β lies in the once-punctured disk \mathbf{C}_σ or β lies in the disk \mathbf{Disk}_σ .

Case 1: *First, suppose β lies in \mathbf{C}_σ .* See Fig. 32(a). Per (5.5), β has the property that it cuts out the disk \mathbf{Disk}_β not containing σ , so no radius can cross both β and σ . Hence each of the even-indexed arcs w_{2i}, \dots, w_{2j} is a peripheral arc because each of them crosses both β and σ . It then follows that each odd-indexed arc $w_{2i+1}, \dots, w_{2j-1}$ is also a peripheral arc. We claim that \bar{w} begins and finishes in \mathbf{C}_σ (recall that this means possibly at s^σ or t^σ):

- If $i = 1$, then w_1 starts in \mathbf{C}_σ since β lies in \mathbf{C}_σ . Otherwise, there is a previous step w_{2i-2} which crosses β . Since w_{2i-2} does not belong to (T^σ, σ) -cross by assumption, w_{2i-2} ends in \mathbf{C}_σ .
- If $j = d$, then w_{2d+1} finishes in \mathbf{C}_σ since β lies in \mathbf{C}_σ . Otherwise, there is a next step w_{2j+2} which crosses β . Since w_{2j+2} does not belong to (T^σ, σ) -cross by assumption, w_{2j+2} ends in \mathbf{C}_σ .

In fact, by induction starting from w_{2i-1} , every odd-indexed arc of \bar{w} must start from \mathbf{C}_σ . But this means that w_{2j+1} is a peripheral arc which begins and ends in \mathbf{C}_σ but also crosses σ , which is impossible.

Case 2: *Next, suppose that β lies in \mathbf{Disk}_σ .* See Figs. 32(b) and 33. To simplify our argument, consider

- a 2-fold cover \widetilde{T}^σ of T^σ , a triangulated once-punctured disk containing two lifts of every arc $\tau \in T^\sigma$ and of every boundary marked point of \mathcal{C}_n (see Fig. 34),
- every ideal triangle in \widetilde{T}^σ is an ordinary triangle, (5.7)
- a lift of γ is a radius if and only if γ is a radius of \mathcal{C}_n .

Since every pair of arcs of \mathcal{C}_n cross each other at most twice,


$$\text{every pair of lifted arcs cross each other at most once on the two-fold cover.} \quad (5.8)$$

Hence, by Lemma 5.18(c),

$$\text{a pair of adjacent lifted arcs do not cross on the two-fold cover.} \quad (5.9)$$


Choose a lift $\tilde{\sigma}$ of σ going from \tilde{s}^σ to \tilde{t}^σ . Here $\tilde{\sigma}$ cuts out a disk $\mathbf{Disk}_{\tilde{\sigma}}$ from \widetilde{T}^σ . Let $\tilde{\beta}$ denote the lift of β which lies in $\mathbf{Disk}_{\tilde{\sigma}}$, and let \tilde{w} be the lift of w which is a $(\widetilde{T}^\sigma, \tilde{\beta})$ -path. We abuse notation by writing \bar{w} to denote the subpath of \tilde{w} corresponding to \bar{w} .

- i) First, assume that \bar{w} has an earlier step w_{2i-2} and a later step w_{2j+2} . We see that \bar{w} starts from inside $\mathbf{Disk}_{\tilde{\sigma}}$ (otherwise, \tilde{w}_{2i-2} would have to cross $\tilde{\sigma}$) and \bar{w} ends inside $\mathbf{Disk}_{\tilde{\sigma}}$ (otherwise, \tilde{w}_{2j+2} would have to cross $\tilde{\sigma}$). Recall that by inside $\mathbf{Disk}_{\tilde{\sigma}}$ we mean possibly at \tilde{s}^σ or \tilde{t}^σ . In fact, by induction starting from w_{2i-1} , every odd-indexed arc of \bar{w} must start from inside $\mathbf{Disk}_{\tilde{\sigma}}$. But this means that \tilde{w}_{2j+1} begins and ends inside $\mathbf{Disk}_{\tilde{\sigma}}$ and also crosses $\tilde{\sigma}$. This requires $\tilde{\sigma}$ and \tilde{w}_{2j+1} to cross twice, contradicting (5.8).
- ii) Second, assume that $i = 1$ or $j = d$. Without loss of generality, assume $i = 1$. There are two cases: either β and σ are adjacent at σ or not.



(a) T^o contains a self-folded triangle with radius r and l -loop ℓ . (b) T^o has radii ρ_1, \dots, ρ_f ($f \geq 2$).

Figure 33. T^o when (T^o, σ) -doublecross = $\{\theta_1, \dots, \theta_h\}$ is nonempty.



(a) T^o contains a self-folded triangle with radius r and l -loop ℓ . (b) T^o has radii ρ_1, \dots, ρ_f ($f \geq 2$).

Figure 34. Two-fold cover $\mathbf{Disk}_{\tilde{\sigma}}$ of T^o when (T^o, σ) -doublecross = $\{\theta_1, \dots, \theta_h\}$ is nonempty.

- If β and σ are adjacent at s , then \tilde{w}_1 and $\tilde{\sigma}$ are adjacent and so \tilde{w}_1 cannot cross $\tilde{\sigma}$ by (5.9). But w_1 crosses σ by assumption, and so \tilde{w}_1 must cross the other lift $\tilde{\sigma}^2$ of σ which is outside of $\mathbf{Disk}_{\tilde{\sigma}}$. Therefore \tilde{w}_1 ends outside of $\mathbf{Disk}_{\tilde{\sigma}}$.
- If β and σ are not adjacent at σ , then w_1 starts inside of $\mathbf{Disk}_{\tilde{\sigma}}$ because $\tilde{\beta}$ lies in $\mathbf{Disk}_{\tilde{\sigma}}$. Since w_1 crosses σ by assumption, \tilde{w}_1 has to cross $\tilde{\sigma}$ and end outside of $\mathbf{Disk}_{\tilde{\sigma}}$.

So, either way, \tilde{w}_1 ends outside of $\mathbf{Disk}_{\tilde{\sigma}}$. Hence $\tilde{w}_2, \dots, \tilde{w}_{2j+1}$ forms a zig-zag pattern crossing $\tilde{\sigma}$ an even number ($2j$) of times, ending outside of $\mathbf{Disk}_{\tilde{\sigma}}$. Since $\tilde{\beta}$ lies in $\mathbf{Disk}_{\tilde{\sigma}}$, there must be a step \tilde{w}_{2j+2} right after \tilde{w} which crosses \tilde{w} , contradicting the assumption that \tilde{w} is a longest $(\tilde{T}^o, \tilde{\sigma})$ -cross-subpath.

This ends the proof for (5.6). The conclusion follows from Lemma 5.11. ■

5.3.2 Proving that x_Σ has non-positive degree with respect to (T, σ) -doublecross, and x_σ has negative degree with respect to (T, σ) -cross and (T, σ) -doublecross

Lemma 5.24. *Suppose Σ is a multi-tagged triangulation and T is a tagged triangulation. Let $\sigma \in \Sigma \setminus T$ be a tagged peripheral arc.*

- 1) *Assume (T, σ) -doublecross is non-empty, and σ is central in $\Sigma \setminus T$ and no other arc is central in $\Sigma \setminus T$. Let $\beta \in \Sigma$ (possibly $\beta = \sigma$). Then each term in the T -expansion of x_β has non-positive degree with respect to (T, σ) -doublecross.*
- 2) *Each term in the T -expansion of x_σ has negative degree with respect to (T, σ) -cross or (T, σ) -doublecross.*

Proof. If $\beta \in T$, then its degree with respect to (T°, σ) -doublecross is zero since it cannot cross σ . Suppose $\beta \in \Sigma \setminus T$. By Remark 5.22, $\beta = \sigma$ or β lies in \mathbf{Disk}_σ , i.e., if $\beta \neq \sigma$, then σ is closer than β to the puncture. See Notation 5.12.

Let $w^\beta = (w_1, \dots, w_{2e+1})$ be a (T°, β) -path. If β does not cross any arc from (T°, σ) -doublecross, then by Corollary 5.17(a) the degree of w with respect to (T°, σ) -doublecross is zero. Suppose β crosses at least an arc from (T°, σ) -doublecross. See Figs. 33.

Suppose $x(w^\beta)$ has positive degree with respect to the θ_k 's. Then by Corollary 5.17(b) there is a longest $\{\theta_k$'s}-subpath $\bar{w}^\beta = (w_{2i-1}, \dots, w_{2j+1})$ of w .

Let arc $\tau_{i_1}, \dots, \tau_{i_d}$ be the arcs crossed by σ , in order.

Let $\theta_1, \dots, \theta_h$ ($h \geq 1$) denote the arcs in that are crossed by σ twice (ordered so that θ_1 is the first and last arc from (T°, σ) -doublecross crossed by σ). Note that $\theta_1, \dots, \theta_{h-1}$ are contained in the disk \mathbf{Disk}_{θ_h} cut by θ_h , see Fig. 33. Let ρ_1, \dots, ρ_f ($f \geq 1$) be the radii of T° , in order of their intersections with σ . Recall that, by Lemma 5.18(e), σ crosses all radii of T° . Let $\tau_{i_1}, \dots, \tau_{i_m}$ (if any) be the peripheral arcs which σ exactly once, right before σ crosses θ_1 for the first time. Let $\tau_{i_{m+h+f+h+1}}, \dots, \tau_{i_d}$ (if any) be the peripheral arcs which σ exactly once, right after σ crosses θ_1 for the second time.

Consider a 2-fold cover \widetilde{T}° of T° , a triangulated once-punctured disk containing two lifts of every arc $\tau \in T^\circ$ and of every boundary marked point of \mathcal{C}_n , see (5.7) and Fig. 34.

Recall that (5.9) gives that

a pair of adjacent lifted arcs on \widetilde{T}° do not cross.

Let s^σ be the starting point of σ and let t^σ be the finishing point of σ . By assumption, σ is not an ℓ -loop, so $s \neq t$. Choose a lift $\tilde{\sigma}$ of σ , running from a lift \tilde{s} of s to a lift \tilde{t} of t . Let $\mathbf{Disk}_{\tilde{\sigma}}$ denote the disk that is cut out by $\tilde{\sigma}$. Let $\tilde{\sigma}^2$ be the other lift of σ , running from \tilde{s}^2 to \tilde{t}^2 . Note that $\tilde{\sigma}$ and $\tilde{\sigma}^2$ never coincide, and, in particular,

$$\tilde{\sigma} \text{ and } \tilde{\sigma}^2 \text{ share no common endpoint.} \tag{5.10}$$

Let $\mathbf{Disk}_{\tilde{\sigma}}^2$ be the disk that is cut out by $\tilde{\sigma}$.

Let $\rho_1^1, \dots, \rho_f^1$ denote the lifts of ρ_1, \dots, ρ_f which are crossed by $\tilde{\sigma}$. Let $\rho_1^2, \dots, \rho_f^2$ denote the other lifts of ρ_1, \dots, ρ_f , lying entirely outside of $\mathbf{Disk}_{\tilde{\sigma}}$. Let $\theta_1^1, \dots, \theta_h^1$ (respectively, $\theta_1^2, \dots, \theta_h^2$) denote the lifts of $\theta_1, \dots, \theta_h$ which $\tilde{\sigma}$ crosses first (respectively, second). In general, let τ^1 denote the lift of $\tau \in T^\circ$ in the region cut out by ρ_f^1 and ρ_f^2 which contains \tilde{s} , and let τ^2 denote the lift of $\tau \in T^\circ$ in the other region.

Since σ cuts $\tau_{i_1}, \dots, \tau_{i_m}, \theta_1, \dots, \theta_h, \rho_1, \dots, \rho_f, \theta_h, \dots, \theta_1, \tau_{i_{m+h+f+h+1}}, \dots, \tau_{i_d}$, in this order, we see that $\tilde{\sigma}$ cuts

$$\tau_{i_1}^1, \dots, \tau_{i_m}^1, \theta_1^1, \dots, \theta_h^1, \rho_1^1, \dots, \rho_f^1, \theta_1^2, \dots, \theta_h^2, \tau_{i_{m+h+f+h+1}}^2, \dots, \tau_{i_d}^2,$$

in this order.

If $\tau \neq \theta_k$, then by assumption $\tau \in T^\circ$ cuts σ at most once, so

$$\begin{aligned} &\text{if } \tau \neq \theta_k, \text{ then } \tilde{\sigma} \text{ cuts at most one of the two lifts } \tau^1, \tau^2 \text{ of } \tau, \\ &\text{and a lift of } \tau \text{ cuts at most one of the two lifts } \tilde{\sigma}, \tilde{\sigma}^2 \text{ of } \sigma. \end{aligned} \tag{5.11}$$

Since σ cuts each arc θ_k twice, $\tilde{\sigma}$ cuts both lifts θ_k^1 and θ_k^2 of θ_k . Due to (5.9),

$$\tilde{\sigma} \text{ cannot be adjacent to } \theta_k^1 \text{ or } \theta_k^2. \tag{5.12}$$

Let $\tilde{\beta}$ be the lift of β that is contained in $\mathbf{Disk}_{\tilde{\sigma}}$. Let \tilde{w}^β denote the $(\widetilde{T}^\circ, \tilde{\beta})$ -path corresponding to w^β , and we abuse notation by writing \bar{w}^β to refer to the subpath of \tilde{w}^β corresponding to \bar{w}^β .

If \bar{w}^β contains both θ_a^1 and θ_b^2 for some $1 \leq a, b \leq h$, then \bar{w}^β must contain the radius/radii $\rho_1^1, \dots, \rho_f^1$, contradicting that the fact that \bar{w}^β is a $\{\theta_k$'s $\}$ -subpath. Without loss of generality, assume each step of \bar{w}^β goes along θ_k^1 for some k .

Note that, because θ_k crosses σ twice,

$$\text{every } \theta_k^1 \text{ crosses both } \tilde{\sigma} \text{ and } \tilde{\sigma}^2.$$

Hence, since every step of \bar{w}^β is θ_k^1 for some k ,

$$\text{every step of } \bar{w}^\beta \text{ begins strictly inside } \mathbf{Disk}_{\tilde{\sigma}} \text{ or } \mathbf{Disk}_{\tilde{\sigma}^2}^2. \quad (5.13)$$

Recall that starting *strictly* inside $\mathbf{Disk}_{\tilde{\sigma}}$ (respectively, $\mathbf{Disk}_{\tilde{\sigma}^2}^2$) means at a marked point of $\mathbf{Disk}_{\tilde{\sigma}}$ (respectively, $\mathbf{Disk}_{\tilde{\sigma}^2}^2$) that is not \tilde{s} or \tilde{t} (respectively, \tilde{s}^2 or \tilde{t}^2).

If $\beta = \sigma$, then \bar{w}_1 starts at \tilde{s} and \bar{w}_{2d+1} finishes at \tilde{t} , so by (5.13) there must be an earlier step \tilde{w}_{2i-2} and a later step w_{2j+2} of \tilde{w} . If $\beta \neq \sigma$, then it is possible to have $i = 1$ (if β is not adjacent to σ at its starting point) or $j = e$ (if β is not adjacent to σ at its finishing point).

We claim that \bar{w}^β starts strictly inside $\mathbf{Disk}_{\tilde{\sigma}}$. For the sake of argument, suppose that $\bar{w}^\beta = (\tilde{w}_{2i-1}, \dots, \tilde{w}_{2j+1})$ starts strictly inside of $\mathbf{Disk}_{\tilde{\sigma}^2}^2$.

Case 1: First, suppose that \tilde{w}^β has an earlier step \tilde{w}_{2i-2} . Since \tilde{w}_{2i-2} must cross $\tilde{\beta}$, which lies in $\mathbf{Disk}_{\tilde{\sigma}}$, we see that \tilde{w}_{2i-2} must cross both $\tilde{\sigma}$ and $\tilde{\sigma}^2$ to get from strictly inside $\mathbf{Disk}_{\tilde{\sigma}}$ to strictly inside $\mathbf{Disk}_{\tilde{\sigma}^2}^2$. Since \tilde{w}_{2i-2} is not a lift of one of the θ_k 's, this contradicts (5.11).

Case 2: Second, assume that $\beta \neq \sigma$, and we have $i = 1$. Since $\tilde{\beta}$ lies in $\mathbf{Disk}_{\tilde{\sigma}}$ by assumption, either \tilde{w}_1 starts at \tilde{s}^σ or \tilde{t}^σ , or \tilde{w}_1 starts *strictly* inside of $\mathbf{Disk}_{\tilde{\sigma}}$ (see Notation 5.12). By (5.13), \tilde{w}_1 starts strictly in $\mathbf{Disk}_{\tilde{\sigma}}$.

Either way, \bar{w}^β starts strictly in $\mathbf{Disk}_{\tilde{\sigma}}$. Per (5.13), we see that $(\tilde{w}_{2i-1}, \dots, \tilde{w}_{2j+1})$ forms a zip-zag pattern bouncing back and forth between (strictly) $\mathbf{Disk}_{\tilde{\sigma}}$ and (strictly) $\mathbf{Disk}_{\tilde{\sigma}^2}^2$. So, by induction starting from \tilde{w}_{2i-1} , every odd-indexed arc of \bar{w} must start strictly in $\mathbf{Disk}_{\tilde{\sigma}}$ and ends strictly in $\mathbf{Disk}_{\tilde{\sigma}^2}^2$. Hence the last step \tilde{w}_{2j+1} ends strictly in $\mathbf{Disk}_{\tilde{\sigma}^2}^2$.


The step \tilde{w}_{2j+1} cannot be the last step of \tilde{w}^β because the finishing point of $\tilde{\beta}$ lies (strictly) outside of $\mathbf{Disk}_{\tilde{\sigma}^2}^2$ per (5.10). Hence \tilde{w}^β has a next step \tilde{w}_{2j+2} . Since \tilde{w}_{2j+2} must cross $\tilde{\beta}$, which lies in $\mathbf{Disk}_{\tilde{\sigma}}$, we see that \tilde{w}_{2j+2} must cross both $\tilde{\sigma}$ and $\tilde{\sigma}^2$ to go strictly from $\mathbf{Disk}_{\tilde{\sigma}^2}^2$ to strictly in $\mathbf{Disk}_{\tilde{\sigma}}$. Since \tilde{w}_{2j+2} is not a lift of one of the θ_k 's, this contradicts (5.11).

We prove part (2):

Suppose $w = (w_1, \dots, w_{2d+1})$ is a (T^o, σ) -path. First, we claim that the T^o -expansion of $x(w)$ has negative degree with respect to either (T^o, σ) -cross or (T^o, σ) -doublecross: By Lemma 5.18(d), either $w_1 \notin (T^o, \sigma)$ -cross or $w_{2d+1} \notin (T^o, \sigma)$ -cross. If necessary, reverse the orientation of w so that $w_{2d+1} \notin (T^o, \sigma)$ -cross. If not all the odd steps $w_1, w_3, \dots, w_{2d-1}$ cross σ , then $x(w)$ has negative degree with respect to (T^o, σ) -cross by Lemma 5.19(c). Hence suppose that all steps of w except w_{2d+1} come from (T^o, σ) -cross.

We claim that $w_{2h+1} \notin (T^o, \sigma)$ -doublecross:

Since $w_1 \in (T^o, \sigma)$ -cross, Lemma 5.18(c) gives us $w_2 = \theta_1$. Since w_1 crosses σ , its lift \tilde{w}_1 (which lies in $\mathbf{Disk}_{\theta_1}^1$) must cross a lift of σ . Because \tilde{w}_1 is adjacent to $\tilde{\sigma}$, they cannot cross by (5.9), so \tilde{w}_1 must cross the other lift $\tilde{\sigma}^2$ of σ which is located (strictly) outside of $\mathbf{Disk}_{\tilde{\sigma}}$. Hence \tilde{w}_1 goes from \tilde{s} to outside of $\mathbf{Disk}_{\tilde{\sigma}}$, and \tilde{w}_2 goes along θ_1^1 from outside to strictly inside of $\mathbf{Disk}_{\tilde{\sigma}}$. By (T1) and (T2), the steps $\tilde{w}_2, \tilde{w}_3, \dots, \tilde{w}_{2h-1}, \tilde{w}_{2h}$ make a zig-zag pattern along the arcs $\theta_1^1, \dots, \theta_h^1$, so that \tilde{w}_{2h} goes along θ_h^1 and ends strictly in $\mathbf{Disk}_{\tilde{\sigma}}$. Hence, by (T2), \tilde{w}_{2h+1} either goes along θ_h^1 or ρ_1^1 . If \tilde{w}_{2h+1} goes along θ_h^1 , it goes from strictly inside to outside of $\mathbf{Disk}_{\tilde{\sigma}}$. Since the point of ρ_1^1 that is adjacent to θ_1^1 is strictly in $\mathbf{Disk}_{\tilde{\sigma}}$, it is impossible for $\tilde{w}_{2(h+1)}$ to go



(a) 4-fold cover of T , which has at least two radii 1, 2, from Fig. 3(a), and σ is a radius.

(b) 4-fold cover of T° from Fig. 3(b), where T° has a self-folded triangle r, ℓ . Here σ is a radius.

Figure 35. 4-fold cover \widetilde{T}° of T° where β is a peripheral arc, and σ is a peripheral arc (respectively, a radius), and $\mathbf{Disk}_{\widetilde{\sigma}}$ is a disk cut out by the peripheral $\widetilde{\sigma}$ (respectively, the fundamental domain area bounded by the two lifts $\widetilde{\sigma}$ and $\widetilde{\sigma}'$ of σ .)

along ρ_1^1 . This contradicts the fact that $\widetilde{w}_{2(h+1)}$ must go along ρ_1^1 . Hence \widetilde{w}_{2h+1} goes along ρ_1^1 , as needed to show that $w_{2h+1} \notin (T^\circ, \sigma)$ -doublecross.

By part (1), the odd-indexed steps $w_{2(h+f+1)-1}, \dots, w_{2(h+f+h)+1}$ cannot all belong to (T°, σ) -doublecross, so there are at most h odd-indexed steps coming from $w_{2(h+f+1)-1}, \dots, w_{2(h+f+h)+1}$. As there are at most $(h-1) + h$ odd-indexed steps from $w_3, w_5, \dots, w_{2h-1}$, there are at most $(h-1) + h$ odd-indexed steps of w coming from (T°, σ) -doublecross total. There are exactly $2h$ even steps of w from (T°, σ) -doublecross by (T1). Hence, by Lemma 5.19(a), the term $x(w)$ has negative degree with respect to (T°, σ) -doublecross, as required.

Per Lemma 5.11, the conclusion follows. ■

5.4 Technical lemmas to prove Lemma 5.5 for cases where Σ has a radius not in T

5.4.1 $\Sigma \setminus T$ contains a plain radius σ

Lemma 5.25 (Corollary of Lemma 5.19(c)). *Let T be a tagged triangulation and let $\sigma \notin T$ be a plain radius, see Figs. 36(a) and 36(b). Then every term in the T -expansion of x_σ has negative degree with respect to*

$$N_\sigma^T := (T, \sigma)\text{-cross}.$$

As defined in Definition 5.10,

$$(T, \sigma)\text{-cross} = \begin{cases} (T^\circ, \sigma)\text{-cross} & \text{if } T \text{ has no parallel radii} \\ \{r^{(p)}, \text{ and the (peripheral) arcs of } T \text{ that cross } \sigma\} & \text{if } T \text{ has parallel radii } r, r^{(p)}. \end{cases}$$

Proof. Since σ is a radius, it is not adjacent to any arc from (T°, σ) -cross. Hence, by Lemma 5.19(c),

$$\begin{aligned} &\text{each term in the } T^\circ\text{-expansion of } x_\sigma \text{ has negative degree} \\ &\text{with respect to } (T^\circ, \sigma)\text{-cross,} \end{aligned} \tag{5.14}$$

and therefore, by Lemma 5.11, with respect to $N_\sigma^T := (T, \sigma)$ -cross. ■




Figure 36. Lemmas 5.25 and 5.26 assume that $\Sigma \setminus T$ contains a plain radius σ .

Lemma 5.26. *Assume the same setup of T , σ , and N_σ^T as in Lemma 5.25 above, and let β be a tagged arc that is compatible with σ . Then each term in the T -expansion of x_β has non-positive degree with respect to $N_\sigma^T = (T, \sigma)$ -cross.*

Proof of Lemma 5.26. Let ρ_1, \dots, ρ_f denote the radii of T^o and let σ run from the puncture to the boundary such that the radii ρ_1, ρ_f and peripheral arc τ_{i_1} (the first arc crossed by σ) form the first triangle that σ crosses. If T^o has a self-folded triangle r, ℓ , then $f = 1$ and $\rho_1 = r = \rho_f$ and $\tau_{i_1} = \ell$. See Figs. 36(a) and 36(b).

If $\beta = \sigma$, we are done by the previous Lemma 5.25. Otherwise, since β must be compatible with σ , there are only four possibilities: β is an arc of T , another plain radius, the notched radius $\sigma^{(p)}$, or a peripheral arc.

Case 1: *First, suppose $\beta \in T$ is compatible with σ . Then β cannot cross σ and β cannot be a notched radius, so $\beta \notin (T, \sigma)$ -cross.*

Case 2: *Second, suppose β is another plain radius (not in T^o). Let $w = (w_1, \dots, w_{2d+1})$ be a (T^o, β) -path which runs from the puncture to the boundary. Then w_1 is a radius, and hence w_1 does not cross σ . Note that ρ_1 and ρ_f (possibly $\rho_1 = \rho_f$) bound a region (say, R) containing σ and all the (peripheral) arcs in (T^o, σ) -cross. If β is outside of R , then w does not contain any arc of (T^o, σ) -cross. If β is contained in R , then the even-indexed arcs of w that are crossed by σ form a consecutive string*

$$w_2 = \tau_{i_1}, \dots, w_{2j} = \tau_{i_j}.$$

By Lemma 5.15(iii), the only odd-indexed step/s that may belong to (T^o, σ) -cross are w_3, \dots, w_{2j+1} since w_1 is a radius (and hence does not cross σ). Hence w has at j even-indexed steps and at most j odd-indexed steps from (T^o, σ) -cross, so, by Lemma 5.19(a), $x(w)$ has non-positive degree with respect to (T^o, σ) -cross, and hence, by Lemma 5.11, with respect to (T, σ) -cross.

Case 3: *Third, suppose $\beta = \sigma^{(p)}$. First, suppose T has parallel radii $r, r^{(p)}$. By (5.14), each term of the T^o -expansion of x_σ has negative degree with respect to (T^o, σ) -cross = $\{\ell, \text{(peripheral) arcs of } T \text{ which cross } \sigma\}$. Each term $x(w)$ (as a T^o -monomial) corresponding to a (T^o, σ) -path w has degree +1 with respect to r , so $x(w)|_{r \leftrightarrow r^{(p)}}$ (as a $T^{(p)O}$ -monomial) has degree +1 with respect to $r^{(p)}$ and negative degree with respect to (T^o, σ) -cross = $\{\ell, \text{(peripheral) arcs of } T \text{ which cross } \sigma\}$, and, therefore, non-positive degree (as a T -monomial) with respect to (T, σ) -cross. Since $x_{\sigma^{(p)}} = x_\sigma|_{r \leftrightarrow r^{(p)}}$ by [17, Proposition 3.15], every term in the T -expansion of $x_{\sigma^{(p)}}$ has non-positive degree with respect to (T, σ) -cross, as needed.*

Next, suppose $f \geq 2$, and we have, by [17, Proposition 3.15],

$$x_{\sigma^{(p)}} = x_{\sigma}|_{\rho_i \leftrightarrow \rho_i^{(p)}} = x_{\sigma}|_{\rho_1 \leftrightarrow \rho_1^{(p)}, \rho_f \leftrightarrow \rho_f^{(p)}}. \quad (5.15)$$

The second equality is due to the fact that no (T, σ) -path would contain the radii $\rho_2, \dots, \rho_{f-1}$, as these radii are not adjacent to σ . Let ℓ_1 (respectively, ℓ_f) be the loop surrounding the radius ρ_1 (respectively, ρ_f), see Fig. 36(c).

Since only the starting endpoint (and not the finish endpoint) of σ is adjacent to ρ_1 and ρ_f , we see that every term in the T -expansion of x_{σ} has degree +1 with respect to $\{\rho_1, \rho_f\}$, so

$$\begin{aligned} &\text{each term in the } T\text{-expansion of } \sigma^{(p)} \text{ has degree } +1 \\ &\text{with respect to } \{x_{\rho_1^{(p)}} = x_{\ell_1}/x_{\rho_1}, x_{\rho_f^{(p)}} = x_{\ell_f}/x_{\rho_f}\}. \end{aligned} \quad (5.16)$$

Furthermore, we observe that each term in the T -expansions of x_{ℓ_1} and x_{ℓ_f} has at most degree +1 with respect to (T, σ) -cross. Since $x_{\rho_i^{(p)}} = x_{\ell_i}/x_{\rho_i}$ for $i = 1, f$,

$$\begin{aligned} &\text{each term in the } T\text{-expansions of } x_{\rho_1^{(p)}} \text{ and } x_{\rho_f^{(p)}} \text{ has at most degree } +1 \\ &\text{with respect to } (T, \sigma)\text{-cross}. \end{aligned} \quad (5.17)$$

By Lemma 5.25, every term in the T -expansion of x_{σ} has negative degree with respect to (T, σ) -cross. Combining this fact with (5.15), (5.16), and (5.17), we see that each term in the T -expansion of $x_{\sigma^{(p)}}$ has non-positive degree with respect to (T, σ) -cross.

Case 4: *Fourth, suppose β is a peripheral arc (not in T) that does not cross σ .* Consider a 4-fold cover \widetilde{T}^o of T^o , a disk with one puncture (\widetilde{P}) and 4 lifts for every arc and marked point of T^o . Let $\widetilde{\beta}$ be a lift of β . Since β and σ do not cross, there are two lifts of σ , say, $\widetilde{\sigma}, \widetilde{\sigma}'$, which bound exactly one fundamental domain containing $\widetilde{\beta}$. We denote this fundamental domain by $\mathbf{Disk}_{\widetilde{\sigma}}$. Let \widetilde{t} and \widetilde{t}' be the boundary endpoints of $\widetilde{\sigma}$ and $\widetilde{\sigma}'$, and let \widetilde{P} be the puncture. See Fig. 35(a) (if T has no parallel radii) or 35(b) (if T^o contains a self-folded triangle r, ℓ).

Let $w = (w_1, \dots, w_{2d+1})$ denote a (T^o, β) -path and also (by abuse of notation) its corresponding lifted $(\widetilde{T}^o, \widetilde{\beta})$ -path. Per Corollary 5.17(b), in order for $x(w)$ to have a positive degree with respect (T^o, σ) -cross, there must be a subsequence \bar{w} of consecutive edges that cross σ , say, $\bar{w} = w_{2i-1}, \dots, w_{2j+1}$, where the edges directly prior to \bar{w} and directly after \bar{w} do not cross σ . Since no radius of T^o can cross σ , all edges in \bar{w} are peripheral arcs.

We claim \bar{w} starts and finishes strictly inside $\mathbf{Disk}_{\widetilde{\sigma}}$, i.e., at a marked point of \mathbf{Disk}_{σ} that is none of $\widetilde{P}, \widetilde{t}$, and \widetilde{t}' .

If $i = 1$, then w_1 starts strictly in \mathbf{Disk}_{σ} because w_1 is not adjacent to $\widetilde{\sigma}$ nor $\widetilde{\sigma}'$. Similarly, if $j = d$, then w_{2d+1} ends strictly \mathbf{Disk}_{σ} because w_{2d+1} is not adjacent to $\widetilde{\sigma}$ nor $\widetilde{\sigma}'$.

Otherwise, \bar{w} must start strictly in $\mathbf{Disk}_{\widetilde{\sigma}}$ because w_{2i-2} must cross $\widetilde{\beta}$ but not $\widetilde{\sigma}$ nor $\widetilde{\sigma}'$, and \bar{w} must end strictly in $\mathbf{Disk}_{\widetilde{\sigma}}$ because w_{2j+2} must cross $\widetilde{\beta}$ but not $\widetilde{\sigma}$ nor $\widetilde{\sigma}'$.

By induction, every odd edge of \bar{w} starts strictly in $\mathbf{Disk}_{\widetilde{\sigma}}$. But this means that w_{2j+1} starts and ends strictly $\mathbf{Disk}_{\widetilde{\sigma}}$ even though w_{2j+1} cuts one of $\widetilde{\sigma}$ and $\widetilde{\sigma}'$, which is impossible.

Hence, $x(w)$ has non-positive degree with respect to (T^o, σ) -cross, and, by Lemma 5.11, also with respect to (T, σ) -cross, as needed. \blacksquare




Figure 37. Three setups for Lemmas 5.28 and 5.29.

5.4.2 $\Sigma \setminus T$ contains no plain radius but there is a notched radius $\sigma^{(p)} \in \Sigma \setminus T$

The following lemma is helpful toward proving Lemmas 5.28 and 5.29.

Lemma 5.27. *Suppose T is a tagged triangulation with no parallel radii, so that $T = T^o$, and ρ is a radius of T . Then every term in the T -expansion of $x_{\rho^{(p)}}$ has negative degree with respect to $\{\text{all the radii of } T\}$.*

Proof of Lemma 5.27. Let ρ_1, \dots, ρ_f (with $f \geq 2$) denote all the (plain) radii (in consecutive order) of T such that $\rho = \rho_1$. Consider the loop ℓ_1 surrounding the radius ρ_1 which crosses ρ_2 first and crosses ρ_f last (see Fig. 36(c)). Consider a (T, ℓ_1) -path $w = (w_1, \dots, w_{2d+1})$. Here $d = f - 1$.

We first show that $x(w)$ has non-negative degree with respect to $\{\rho_1, \dots, \rho_f\}$. Since the arcs that are crossed by ℓ_1 are precisely ρ_2, \dots, ρ_f , the even steps of w are $\rho_2, \rho_3, \dots, \rho_f$ by (T1). By Lemma 5.19(a), The only way for $x(w)$ to have positive degree with respect to $\{\rho_1, \dots, \rho_f\}$ is if all the odd steps of w come from $\{\rho_1, \dots, \rho_f\}$. By induction, $w_1 = \rho_1, w_3 = \rho_2, \dots, w_{2d-1} = \rho_f$ such that each one goes from the boundary to the puncture. But this requires $w_{2d} = \rho_f$ to go from the puncture to the boundary. Hence w_{2d+1} goes from boundary to boundary, so $w_{2d+1} \notin \{\rho_1, \dots, \rho_f\}$. Hence each term in the T -expansion of x_{ℓ_1} has non-negative degree with respect to $\{\rho_1, \dots, \rho_f\}$.

Since $x_{\rho_1^{(p)}} = x_{\ell_1} / x_{\rho_1}$, every term in the T -expansion of $x_{\rho_1^{(p)}}$ has negative degree with respect to $\{\rho_1, \dots, \rho_f\}$. ■

Lemma 5.28. *Suppose Σ is a multi-tagged triangulation and T is a tagged triangulation such that $\Sigma \setminus T$ does not contain any plain radius but it contains a notched radius $\sigma^{(p)}$. There are three cases to consider, Figs. 37(a), 37(b), and 37(c).*

If T has parallel radii $r, r^{(p)}$, we denote

$$N_\sigma^T := \{r, \text{ and the (peripheral) arcs of } T \text{ that cross } \sigma\},$$

where $\sigma^{(p)} \in \Sigma \setminus T$ is a notched radius such that $\sigma \notin T$ (see Fig. 37(a)).

If T has no parallel radii, we denote

$$N_\sigma^T := \begin{cases} \{\text{all radii of } T\} & \text{if } \sigma^{(p)} \in \Sigma \setminus T \text{ is a notched radius such that } \sigma \notin T \\ & \text{(see Fig. 37(b)),} \\ \{\text{all radii of } T\} \setminus \{\sigma\} & \text{if } \Sigma \setminus T \text{ contains no notched radius that crosses } T, \text{ but} \\ & \sigma^{(p)} \in \Sigma \setminus T \text{ is a notched radius where } \sigma \in T \text{ (see Fig. 37(c)).} \end{cases}$$

Then each term in the T -expansion of $x_{\sigma^{(p)}}$ has negative degree with respect to N_σ^T .

Proof of Lemma 5.28. There are three cases to prove, Figs. 37(a), 37(b), and 37(c).

Fig. 37(a): In this case T has parallel radii $r, r^{(p)}$. The result follows from the proof of Lemma 5.25 by switching the roles of r, σ with $r^{(p)}, \sigma^{(p)}$.

For the remainder of this proof, assume T has no parallel radii. Let ρ_1, \dots, ρ_f denote the radii of T with $f \geq 2$.

Fig. 37(b): $\Sigma \setminus T$ contains a notched radius $\sigma^{(p)}$ such that $\sigma \notin T$. Let $N_\sigma^T := \{\text{all radii of } T\}$, per above assumption. Assume σ is adjacent to ρ_1 and ρ_f . First, recall that (5.15) gives us

$$x_{\sigma^{(p)}} = x_\sigma|_{\rho_i \leftrightarrow \rho_i^{(p)}} = x_\sigma|_{\rho_1 \leftrightarrow \rho_1^{(p)}, \rho_f \leftrightarrow \rho_f^{(p)}}.$$

Second, observe that every term in the T -expansion of x_σ has degree $+1$ with respect to $\{\rho_1, \rho_f\}$ and does not include any step from $\{\rho_2, \dots, \rho_{f-1}\}$. Third, by Lemma 5.27, every term in the T -expansions of $x_{\rho_1^{(p)}}$ and $x_{\rho_f^{(p)}}$ has negative degree with respect to $\{\rho_1, \dots, \rho_f\}$. Combing these three facts, every term in the T -expansion of $x_{\sigma^{(p)}} = x_\sigma|_{\rho_1 \leftrightarrow \rho_1^{(p)}, \rho_f \leftrightarrow \rho_f^{(p)}}$ has negative degree with respect to $\{\rho_1, \dots, \rho_f\}$.

Fig. 37(c): $\Sigma \setminus T$ contains no radius that crosses T , but $\sigma^{(p)} \in \Sigma \setminus T$ is a notched radius where $\sigma \in T$. Let ρ_1 denote the radius σ of T . Let $N_\sigma^T := \{\rho_2, \dots, \rho_f\}$, per above assumption. Consider the loop ℓ_1 around ρ_1 (see Fig. 36(c)) and a (T, ℓ_1) -path w . Observe that w has even steps $\rho_2, \rho_3, \dots, \rho_f$, and the first and last steps of w are not in $\{\rho_2, \dots, \rho_f\}$, so $x(w)$ has negative degree with respect to $\{\rho_2, \dots, \rho_f\}$ by Lemma 5.19(a). Since

$$x_{\rho_1^{(p)}} = \frac{x_{\ell_1}}{x_{\rho_1}},$$

each term in the T -expansion of $x_{\rho_1^{(p)}}$ also has negative degree with respect to $\{\rho_2, \dots, \rho_f\}$. ■

Lemma 5.29. *Assume the same setup and three different cases as Lemma 5.28 above. If $\beta \in \Sigma$, then each term in the T -expansion of x_β is of non-positive degree with respect to N_σ^T .*

Proof. As in the proof of Lemma 5.28, there are three cases to prove, Figs. 37(a), 37(b), and 37(c).

We prove the case of Fig. 37(a) where T has parallel radii $r, r^{(p)}$ and $N_\sigma^T = \{r, \text{ and the arcs of } T \text{ that cross } \sigma\}$ by switching the roles of r, σ with $r^{(p)}, \sigma^{(p)}$ in the proof of Lemma 5.26.

For the remainder of this proof, we prove the two setups of Figs. 37(b) and 37(c) for when T has no parallel radii. Let ρ_1, \dots, ρ_f denote the radii of T with $f \geq 2$.

If $\beta = \sigma^{(p)}$, we are done by the previous Lemma 5.28. Otherwise, since β is compatible with $\sigma^{(p)}$, there are four possibilities: β is an arc of T , a notched radius that crosses T , a notched radius $\rho_k^{(p)}$ parallel to a radius of T , or a peripheral arc.

Case 1: *First, suppose $\beta \in T$.* Since β is compatible with the notched radius $\sigma^{(p)}$, we see that $\beta \notin N_\sigma^T$ since either $N_\sigma^T = \{\text{all radii of } T\}$ (if $\sigma \notin T$) or $N_\sigma^T = \{\text{all radii of } T\} \setminus \{\sigma\}$ (if σ is a radius of T).

Case 2: *Second, suppose β is a notched radius that crosses T .* The case of Fig. 37(b) where $N_\sigma^T = \{\text{all radii of } T\}$ is done by Lemma 5.28 (for the same case). We do not need to consider the case of Fig. 37(c) because it is assumed that $\Sigma \setminus T$ does not contain any notched radius that crosses T .

Case 3: *Third, suppose that β is a notched radius $\rho_k^{(p)}$ not equal to $\sigma^{(p)}$.* The case of Fig. 37(b) where $N_\sigma^T = \{\text{all radii of } T\}$ is done by Lemma 5.27.

To prove the case of Fig. 37(c), let ρ_1 denote σ . Suppose β is a notched arc $\rho_k^{(p)}$ with $k \neq 1$, and consider the loop ℓ_k around ρ_k and a (T, ℓ_k) -path w . We claim that $x(w)$ has at most degree +1 with respect to $\{\rho_2, \dots, \rho_f\}$. To prove this, note that the $f-1$ even arcs of w are $\rho_{k+1}, \dots, \rho_f, \rho_1, \dots, \rho_{k-1}$, i.e., all the radii of T except for ρ_k . For $x(w)$ to have degree +2 or more with respect to $\{\rho_2, \dots, \rho_f\}$, we need all f odd arcs of w to be in $\{\rho_2, \dots, \rho_f\}$. But either the step right before or right after the even-step along ρ_1 of w needs to be either ρ_1 or another (peripheral) arc. Hence $x(w)$ has at most degree +1 with respect to $\{\rho_2, \dots, \rho_f\}$. Since

$$x_{\rho_k^{(p)}} = \frac{x_{\ell_k}}{x_{\rho_k}},$$

each term of x_{ρ_k} has at most degree 0 with respect to $\{\rho_2, \dots, \rho_f\}$, as needed.

Case 4: *Fourth, suppose β is a peripheral arc not crossing σ .* Let $w = (w_1, \dots, w_{2d+1})$ be a (T, β) -path. Suppose for contradiction that $x(w)$ is of positive degree with respect to N_σ^T . By Corollary 5.17(b),

$$\begin{aligned} &\text{we must have a subsequence } \bar{w} = w_{2i-1}, \dots, w_{2j+1} \text{ of } w \\ &\text{(of length three or greater) where all the steps of } \bar{w} \text{ belong to } N_\sigma^T \\ &\text{while the step before and the one after do not belong to } N_\sigma^T. \end{aligned} \quad (5.18)$$

We claim that \bar{w} must start at the boundary (as opposed to the puncture). Otherwise, w would contain an earlier step w_{2i-2} which must be a radius of T going from the boundary to the puncture. For the case where σ is not a radius of T and $N_\sigma^T = \{\text{all radii of } T\}$ (Fig. 37(b)), this shows that $w_{2i-2} \in N_\sigma^T$. For the case where $\sigma = \rho_1$ and $N_\sigma^T = \{\rho_2, \dots, \rho_f\}$ (Fig. 37(c)), since β cannot cross ρ_1 , we see that w_{2i-2} has to come from $\{\rho_2, \dots, \rho_f\} = N_\sigma^T$. For both cases, this contradicts the assumption (5.18) that no step before \bar{w} comes from N_σ^T , hence \bar{w} must start at the boundary. Similarly, \bar{w} must end at the boundary (as opposed to the puncture).

Since every step of \bar{w} is a radius, it follows by induction that every odd step of \bar{w} starts at the boundary. But this means that w_{2j+1} is a radius that begins and ends at the boundary, which is impossible. Hence $x(w)$ has non-positive degree with respect to N_σ^T . ■

5.5 Proof of Lemma 5.5

Proof of Lemma 5.5. Suppose x_Σ is a cluster monomial not compatible with a tagged triangulation T .

Suppose that all tagged arcs of $\Sigma \setminus T$ are peripheral. We shall choose a tagged arc $\sigma \in \Sigma \setminus T$ that is central and, if possible, crosses every arc of T at most once, as follows. Let $\{\lambda_1, \dots, \lambda_r\}$ be the set of central arcs in $\Sigma \setminus T$ (see Fig. 31(a)). If $r = 1$, then choose $\sigma := \lambda_r$.

If $r > 1$, we choose $\sigma \in \{\lambda_k\text{'s}\}$ such that (T°, σ) -doublecross = \emptyset . To see that we can do this, suppose that one of these λ_k (say, λ_r) cuts an arc $\tau \in T^\circ$ twice, see Fig. 31(b). Since τ must cut out a simply-connected region containing all of $\lambda_1, \dots, \lambda_{r-1}$, each of $\lambda_1, \dots, \lambda_{r-1}$ crosses every arc of T° at most once.

Hence, we satisfy the setup for

Lemma 5.23: σ is central in $\Sigma \setminus T$, and for

Lemma 5.24: If (T°, σ) -doublecross is nonempty, then σ is the only central arc in $\Sigma \setminus T$.

Lemma 5.24(2) tells us that each term in the T -expansion of x_σ is of negative degree with respect to either (T, σ) -cross or (T, σ) -doublecross. At the same time, Lemmas 5.23 and 5.24(1) tell us that every term in the T -expansion of x_Σ has non-positive degree with respect to both (T, σ) -cross and (T, σ) -doublecross. It follows, since $\sigma \in \Sigma$, that every term in the T -expansion of x_Σ has negative degree with respect to either (T, σ) -cross or (T, σ) -doublecross. Since (T, σ) -cross and (T, σ) -doublecross are subsets of T , the T -expansion of x_Σ is a sum of proper Laurent monomials.

Suppose now that $\Sigma \setminus T$ contains a radius σ . A very similar argument appears in the proof of [7, Proposition 2.3] but we repeat it here for completeness. Lemma 5.25 (if σ is a plain radius) or Lemma 5.28 (if all radii of $\Sigma \setminus T$ are notched) tells us that each term in the T -expansion of x_σ is of negative degree with respect to a subset N_σ^T . At the same time, Lemma 5.26 (if σ is a plain radius) or Lemma 5.29 (if all radii of $\Sigma \setminus T$ are notched) tells us that each term in the T -expansion of the other factors of x_Σ are of non-positive degree with respect to the same grading N_σ^T . It follows that each term in the T -expansion of x_Σ has negative degree with respect to this grading. Since N_σ^T is a subset of T , the T -expansion of x_Σ is a sum of proper Laurent monomials. ■

5.6 Further directions

It is not known for a general cluster algebra whether its set of indecomposable positive elements forms an atomic basis. Examples of cluster algebras where this fails to happen are those of rank 2 where the exchange matrix looks like $\begin{pmatrix} 0 & b \\ -c & 0 \end{pmatrix}$ with $bc \geq 5$ [15].

Per [18, Conjecture 1.5], we are exploring the existence of atomic bases for type \tilde{D}_{n-1} cluster algebras (which arise from $(n - 3)$ -gons with 2 punctures). Toward this goal, we would like to extend the T° -path expansion formula to general bordered surfaces with more than 1 puncture. Additionally, the T° -path formula presented in this paper was only done for ordinary arcs in the setting of ideal triangulations T° . We are working on proving a similar formula for tagged arcs and for tagged triangulations T .

Acknowledgements

The authors would like to thank Pasha Pylyavskyy, Vic Reiner, and Peter Webb for looking over an early version of this paper, Hugh Thomas for helpful discussions, and the referees for many useful comments. The authors were supported by NSF Grants DMS-1067183 and DMS-1148634.

References

- [1] Baur K., Marsh R.J., Frieze patterns for punctured discs, *J. Algebraic Combin.* **30** (2009), 349–379, [arXiv:0711.1443](#).
- [2] Caldero P., Keller B., From triangulated categories to cluster algebras, *Invent. Math.* **172** (2008), 169–211, [math.RT/0506018](#).
- [3] Ceballos C., Pilaud V., Cluster algebras of type D : Pseudotriangulations approach, [arXiv:1504.06377](#).
- [4] Cerulli Irelli G., Positivity in skew-symmetric cluster algebras of finite type, [arXiv:1102.3050](#).
- [5] Cerulli Irelli G., Keller B., Labardini-Fragoso D., Plamondon P.G., Linear independence of cluster monomials for skew-symmetric cluster algebras, *Compos. Math.* **149** (2013), 1753–1764, [arXiv:1203.1307](#).
- [6] Cerulli Irelli G., Labardini-Fragoso D., Quivers with potentials associated to triangulated surfaces, Part III: tagged triangulations and cluster monomials, *Compos. Math.* **148** (2012), 1833–1866, [arXiv:1108.1774](#).
- [7] Dupont G., Thomas H., Atomic bases of cluster algebras of types A and \tilde{A} , *Proc. Lond. Math. Soc.* **107** (2013), 825–850, [arXiv:1106.3758](#).
- [8] Fock V., Goncharov A., Moduli spaces of local systems and higher Teichmüller theory, *Publ. Math. Inst. Hautes Études Sci.* (2006), 1–211, [math.AG/0311149](#).

- [9] Fock V.V., Goncharov A.B., Cluster ensembles, quantization and the dilogarithm, *Ann. Sci. Éc. Norm. Supér. (4)* **42** (2009), 865–930, [math.AG/0311245](#).
- [10] Fomin S., Shapiro M., Thurston D., Cluster algebras and triangulated surfaces. I. Cluster complexes, *Acta Math.* **201** (2008), 83–146, [math.RA/0608367](#).
- [11] Fomin S., Zelevinsky A., Cluster algebras. I. Foundations, *J. Amer. Math. Soc.* **15** (2002), 497–529, [math.RT/0104151](#).
- [12] Fomin S., Zelevinsky A., Cluster algebras. II. Finite type classification, *Invent. Math.* **154** (2003), 63–121, [math.RA/0208229](#).
- [13] Gekhtman M., Shapiro M., Vainshtein A., Cluster algebras and Weil–Petersson forms, *Duke Math. J.* **127** (2005), 291–311, [math.QA/0309138](#).
- [14] Hernandez D., Leclerc B., Cluster algebras and quantum affine algebras, *Duke Math. J.* **154** (2010), 265–341, [arXiv:0903.1452](#).
- [15] Lee K., Li L., Zelevinsky A., Positivity and tameness in rank 2 cluster algebras, *J. Algebraic Combin.* **40** (2014), 823–840, [arXiv:1303.5806](#).
- [16] Musiker G., Schiffler R., Cluster expansion formulas and perfect matchings, *J. Algebraic Combin.* **32** (2010), 187–209, [arXiv:0810.3638](#).
- [17] Musiker G., Schiffler R., Williams L., Positivity for cluster algebras from surfaces, *Adv. Math.* **227** (2011), 2241–2308, [arXiv:0906.0748](#).
- [18] Musiker G., Schiffler R., Williams L., Bases for cluster algebras from surfaces, *Compos. Math.* **149** (2013), 217–263, [arXiv:1110.4364](#).
- [19] Nakajima H., Quiver varieties and cluster algebras, *Kyoto J. Math.* **51** (2011), 71–126, [arXiv:0905.0002](#).
- [20] Schiffler R., A geometric model for cluster categories of type D_n , *J. Algebraic Combin.* **27** (2008), 1–21, [math.RT/0608264](#).
- [21] Schiffler R., On cluster algebras arising from unpunctured surfaces. II, *Adv. Math.* **223** (2010), 1885–1923, [arXiv:0809.2593](#).
- [22] Schiffler R., Thomas H., On cluster algebras arising from unpunctured surfaces, *Int. Math. Res. Not.* **2009** (2009), 3160–3189, [arXiv:0712.4131](#).
- [23] Sherman P., Zelevinsky A., Positivity and canonical bases in rank 2 cluster algebras of finite and affine types, *Mosc. Math. J.* **4** (2004), 947–974, [math.RT/0307082](#).