

ON SECOND HANKEL DETERMINANT FOR TWO NEW SUBCLASSES OF ANALYTIC FUNCTIONS

T. V. Sudharsan and R. Vijaya

Abstract. In this paper, we obtain sharp upper bounds for the functional $|a_2a_4 - a_3^2|$ for functions belonging to $S^*(\alpha, \beta)$ and $C(\alpha, \beta)$. Our results extend corresponding previously known results.

[Full text](#)

References

- [1] T.O. Babalola and J.O. Opoola, *On the coefficients of certain analysis and univalent functions*, Advances in inequalities for series, (Edited by S.S. Dragomir and A. Sofo), Nova Science Publishers (2008), 5–17.
- [2] P.L. Duren, *Univalent functions*, Springer Verlag, New York Inc, 1983,. [MR0708494\(85j:30034\)](#). [Zbl 514.30001](#).
- [3] T. Hayami and S. Owa, *Generalized Hankel determinant for certain classes*, Int. Journal of Math. Analysis, **4**(52) (2010), 2473–2585. [MR2770050\(2011m:30032\)](#). [Zbl 1226.30015](#).
- [4] A. Janteng, S.A. Halim and M. Darus, *Hankel determinant for starlike and convex functions*, Int. Journal of Math. Analysis, **I**(13) (2007), 619–625. [MR2370200](#). [Zbl 1137.30308](#).
- [5] A. Janteng, S.A. Halim and M. Darus, *Estimate on the second Hankel functional for functions whose derivative has a positive real part*, Journal of Quality Measurement and Analysis, **4**(1) (2008), 189–195.

2010 Mathematics Subject Classification: Primary 30C80; Secondary 30C45.

Keywords: Coefficient bounds; Fekete-Szego functional; Hankel determinant.

This work was supported by UGC, under the grant F.MRP-4117/12 (MRP/UGC-SERO) of the second author.

<http://www.utgjiu.ro/math/sma>

- [6] N. Kharudin, A. Akbarally, D. Mohamad and S.C. Soh, *The second Hankel determinant for the class of close to convex functions*, European Journal of Scientific Research, **66**(3) (2011), 421–427.
- [7] R.J. Libera and E.J. Zlotkiewicz, *Early coefficients of the inverse of a regular convex function*, Proc. Amer. Math. Soc., **85**(2) (1982), 225–230. [MR0652447\(83h:30017\)](#). [Zbl 0464.30019](#).
- [8] R.J. Libera and E.J. Zlotkiewicz, *Coefficient bounds for the inverse of a function with derivative in P* , Proc. Amer. Math. Soc., **87**(2) (1983), 251–257. [MR0681830\(84a:30024\)](#). [Zbl 0488.30010](#).
- [9] J.W. Noonan and D.K. Thomas, *On the second Hankel determinant of areally mean p -valent functions*, Trans. Amer. Math. Soc., **223**(2) (1976), 337–346. [MR0422607\(54 #10593\)](#). [Zbl 0346.30012](#).
- [10] K.I. Noor, *Hankel determinant problem for the class of functions with bounded boundary rotation*, Rev. Roum. Math. Pures Et Appl., **28**(8) (1983), 731–739. [MR0725316\(85f:30017\)](#). [Zbl 0524.30008](#).
- [11] M.S. Robertson, *On the theory of univalent functions*, Annals of Math., **37** (1936), 374–408. [MR1503286](#). [Zbl 0014.16505](#).
- [12] C. Selvaraj and N. Vasanthi, *Coefficient bounds for certain subclasses of close-to-convex functions*, Int. Journal of Math. Analysis, **4**(37) (2010), 1807–1814. [MR2728501](#). [Zbl 1218.30052](#).
- [13] G. Shanmugam, B. Adolf Stephen and K.G. Subramanian, *Second Hankel determinant for certain classes of analytic functions*, Bonfring International Journal of Data Mining, **2**(2) (2012).

T. V. Sudharsan

Department of Mathematics,

SIVET College,

Chennai - 600 073, India.

E-mail: tvsudharsan@rediffmail.com

R. Vijaya

Department of Mathematics,

S.D.N.B. Vaishnav College,

Chennai - 600 044, India.

E-mail: viji_dorai67@yahoo.co.in

Surveys in Mathematics and its Applications **9** (2014), 131 – 138

<http://www.utgjiu.ro/math/sma>