

# CATEGORICAL STRUCTURES ENRICHED IN A QUANTALOID: CATEGORIES, DISTRIBUTORS AND FUNCTORS

ISAR STUBBE

ABSTRACT. We thoroughly treat several familiar and less familiar definitions and results concerning categories, functors and distributors enriched in a base quantaloid  $\mathcal{Q}$ . In analogy with  $\mathcal{V}$ -category theory we discuss such things as adjoint functors, (pointwise) left Kan extensions, weighted (co)limits, presheaves and free (co)completion, Cauchy completion and Morita equivalence. With an appendix on the universality of the quantaloid  $\text{Dist}(\mathcal{Q})$  of  $\mathcal{Q}$ -enriched categories and distributors.

## 1. Introduction

The theory of categories enriched in a symmetric monoidal closed category  $\mathcal{V}$  is, by now, well known [Bénabou, 1963, 1965; Eilenberg and Kelly, 1966; Lawvere, 1973; Kelly, 1982]. For such a  $\mathcal{V}$  with “enough” (co)limits the theory of  $\mathcal{V}$ -categories, distributors and functors can be pushed as far as needed: it includes such things as (weighted) (co)limits in a  $\mathcal{V}$ -category,  $\mathcal{V}$ -presheaves on a  $\mathcal{V}$ -category, Kan extensions of enriched functors, Morita theory for  $\mathcal{V}$ -categories, and so on.

Monoidal categories are precisely one-object bicategories [Bénabou, 1967]. It is thus natural to ask how far  $\mathcal{V}$ -category theory can be generalized to  $\mathcal{W}$ -category theory, for  $\mathcal{W}$  a general bicategory. But, whereas in  $\mathcal{V}$ -category theory one usually assumes the symmetry of the tensor in  $\mathcal{V}$  (which is essential for showing that  $\mathcal{V}$  is itself a  $\mathcal{V}$ -category with hom-objects given by the right adjoint to tensoring), in working over a general bicategory  $\mathcal{W}$  we will have to sacrifice this symmetry: tensoring objects in  $\mathcal{V}$  corresponds to composing morphisms in  $\mathcal{W}$  and in general it simply does not make sense for the composition  $g \circ f$  of two arrows  $f, g$  to be “symmetric”.

On the other hand, we can successfully translate the notion of closedness of a monoidal category  $\mathcal{V}$  to the more general setting of a bicategory  $\mathcal{W}$ : ask that, for any object  $X$  of  $\mathcal{W}$  and any arrow  $f : A \rightarrow B$  in  $\mathcal{W}$ , both functors

$$- \circ f : \mathcal{W}(B, X) \rightarrow \mathcal{W}(A, X) : x \mapsto x \circ f, \quad (1)$$

$$f \circ - : \mathcal{W}(X, A) \rightarrow \mathcal{W}(X, B) : x \mapsto f \circ x \quad (2)$$

have respective right adjoints

$$\{f, -\} : \mathcal{W}(A, X) \rightarrow \mathcal{W}(B, X) : y \mapsto \{f, y\}, \quad (3)$$

---

Received by the editors 2004-05-13 and, in revised form, 2004-01-14.

Transmitted by Jiri Rosicky. Published on 2005-01-18.

2000 Mathematics Subject Classification: 06F07, 18B35, 18D20.

Key words and phrases: Quantales and quantaloids, enriched categories.

© Isar Stubbe, 2004. Permission to copy for private use granted.

$$[f, -] : \mathcal{W}(X, B) \longrightarrow \mathcal{W}(X, A) : y \mapsto [f, y]. \quad (4)$$

Such a bicategory  $\mathcal{W}$  is said to be closed. Some call an arrow such as  $\{f, y\}$  a (right) extension and  $[f, y]$  a (right) lifting (of  $y$  through  $f$ ).

Finally, by saying that  $\mathcal{V}$  has “enough limits and colimits” is in practice often meant that  $\mathcal{V}$  has small limits and small colimits. In a bicategory  $\mathcal{W}$  the analogue is straightforward: now ask for  $\mathcal{W}$  to have in its hom-categories small limits and small colimits (i.e.  $\mathcal{W}$  is locally complete and cocomplete).

So, to summarize, when trying to develop category theory over a base bicategory  $\mathcal{W}$  rather than a base monoidal category  $\mathcal{V}$ , it seems reasonable to work with a base bicategory which is closed, locally complete and locally cocomplete. Note that in such a bicategory  $\mathcal{W}$ , due to its closedness, composition always distributes on both sides over colimits of morphisms:

$$f \circ (\operatorname{colim}_{i \in I} g_i) \cong \operatorname{colim}_{i \in I} (f \circ g_i), \quad (5)$$

$$(\operatorname{colim}_{j \in J} f_j) \circ g \cong \operatorname{colim}_{j \in J} (f_j \circ g). \quad (6)$$

That is to say, the local colimits are stable under composition. (But this does not hold in general for local limits!)

We will focus on a special case of these closed, locally complete and locally cocomplete bicategories: namely, we study such bicategories whose hom-categories are moreover small and skeletal. Thus the hom-categories are simply complete lattices. We will write the local structure as an order, and local limits and colimits of morphisms as their infimum, resp. supremum—so for arrows with same domain and codomain we have things like  $f \leq f'$ ,  $\bigvee_{i \in I} f_i$ ,  $\bigwedge_{j \in J} g_j$ , etc. In particular (5) and (6) become

$$f \circ (\bigvee_i g_i) = \bigvee_i (f \circ g_i), \quad (7)$$

$$(\bigvee_j f_j) \circ g = \bigvee_j (f_j \circ g). \quad (8)$$

The adjoint functor theorem says that the existence of the adjoints (3) and (4) to the composition functors (1) and (2) (not only implies but also) is implied by their distributing over suprema of morphisms as in (7) and (8). Such bicategories – whose hom-categories are complete lattices and whose composition distributes on both sides over arbitrary suprema – are called quantaloids. A one-object quantaloid is a quantale<sup>1</sup>. So a quantaloid  $\mathcal{Q}$  is a **Sup**-enriched category (and a quantale is monoid in **Sup**).

We will argue below that “ $\mathcal{V}$ -category theory” can be generalized to “ $\mathcal{Q}$ -category theory”, where now  $\mathcal{Q}$  is a quantaloid. This is a particular case of the theory of “ $\mathcal{W}$ -category theory” as pioneered by [Bénabou, 1967; Walters, 1981; Street, 1983a]. But we feel that this particular case is also of particular interest: many examples of bicategory-enriched categories are really quantaloid-enriched. Also, without becoming trivial, quantaloid-enrichment often behaves remarkably better than general bicategory-enrichment: essentially because all diagrams of 2-cells in a quantaloid commute. So, for example, when

<sup>1</sup>It was C. Mulvey [1986] who introduced the word ‘quantale’ in his work on (non-commutative)  $C^*$ -algebras as a contraction of ‘quantum’ and ‘locale’.

calculating extensions and liftings in a quantaloid, we need not keep track of the universal 2-cells; or when defining a  $\mathcal{Q}$ -category, we need not specify its composition- and identity-2-cells nor impose coherence axioms; also the composition of distributors simplifies a great deal, for we need not compute complicated colimits of 1-cells in a general bicategory but simply suprema of arrows in a quantaloid; and so on. In our further study of categorical structures enriched in a quantaloid we rely heavily on the basic theory of  $\mathcal{Q}$ -categories, distributors and functors; however, often we could not find an appropriate reference for one or another basic fact. With this text we wish to provide such a reference.

A first further study, in [Stubbe, 2004a], is concerned with “variation and enrichment”, in the sense of [Betti *et al.*, 1983; Gordon and Power, 1997, 1999]. This is a development of the notion of weighted colimit in a  $\mathcal{Q}$ -enriched category; in particular, a tensored and cotensored  $\mathcal{Q}$ -category has all weighted colimits (as in section 5 below) if and only if it has all conical colimits, if and only if its underlying order has suprema of objects of the same type. This allows for a detailed analysis of the biequivalence between  $\mathcal{Q}$ -modules and cocomplete  $\mathcal{Q}$ -categories.

The subject of [Stubbe, 2004b] is that of presheaves on  $\mathcal{Q}$ -semicategories (“categories without units”), along the lines of [Moens *et al.*, 2002]; it generalizes the theory of regular modules on an  $R$ -algebra without unit, for  $R$  a commutative ring (with unit). The point is that certain “good properties” of the Yoneda embedding for  $\mathcal{Q}$ -categories (see section 6 below) are no longer valid for  $\mathcal{Q}$ -semicategories—for example, a presheaf on a  $\mathcal{Q}$ -semicategory is not canonically the weighted colimit of representables. Enforcing precisely this latter condition defines what is called a “regular presheaf” on a  $\mathcal{Q}$ -semicategory; there is an interesting theory of “regular  $\mathcal{Q}$ -semicategories”.

[Borceux and Cruciani, 1998] gives an elementary description of ordered objects in the topos of sheaves on a locale  $\Omega$ ; this turns out to be all about enriched  $\Omega$ -semicategories that admit an appropriate Cauchy completion. In [Stubbe, 2004c] we describe more generally the theory of  $\mathcal{Q}$ -semicategories that admit a well-behaved Cauchy completion (based on the material in section 7): these Cauchy complete semicategories are what we believe to be the correct notion of “ordered sheaves on a quantaloid  $\mathcal{Q}$ ”, and we simply call them ‘ $\mathcal{Q}$ -orders’. Such  $\mathcal{Q}$ -orders can equivalently be described as categories enriched in the split-idempotent completion of  $\mathcal{Q}$ , and so provide a missing link between [Walters, 1981] and [Borceux and Cruciani, 1998]. We hope that the theory of  $\mathcal{Q}$ -orders, as a theory of “(ordered) sheaves on a non-commutative topology”, will provide new grounds for defining notions of spectra for non-commutative rings and  $C^*$ -algebras—an issue which was at the origin of the theory of quantales and quantaloids!

Acknowledgment. I thank Francis Borceux and Ross Street for their generosity, not only in mathematics, but simply in general.

## 2. Quantaloids

A sup-lattice is an antisymmetrically ordered set<sup>2</sup>  $(X, \leq)$  for which every subset  $X' \subseteq X$  has a supremum  $\bigvee X'$  in  $X$ . A morphism of sup-lattices  $f: (X, \leq) \rightarrow (Y, \leq)$  is a map  $f: X \rightarrow Y$  that preserves arbitrary suprema. It is well-known that sup-lattices and sup-morphisms constitute a symmetric monoidal closed category **Sup**.

**2.1. DEFINITION.** A quantaloid  $\mathcal{Q}$  is a **Sup**-enriched category. A homomorphism  $F: \mathcal{Q} \rightarrow \mathcal{Q}'$  of quantaloids is a **Sup**-enriched functor.

In principle we do not mind a quantaloid having a proper class of objects. Thus quantaloids and homomorphisms form an illegitimate category **QUANT**; small quantaloids define a (true) subcategory **Quant**. A quantaloid with one object is often thought of as a monoid in **Sup**, and is called a *quantale*.

In elementary terms, a quantaloid  $\mathcal{Q}$  is a category whose hom-sets are actually sup-lattices, in which composition distributes on both sides over arbitrary suprema of morphisms. In the same vein, a homomorphism  $F: \mathcal{Q} \rightarrow \mathcal{Q}'$  is a functor of (the underlying) categories that preserves arbitrary suprema of morphisms.

For arrows  $f: A \rightarrow B$ ,  $g: B \rightarrow C$ ,  $(h_i: X \rightarrow Y)_{i \in I}$  in a quantaloid  $\mathcal{Q}$  we use notations like  $g \circ f: A \rightarrow C$  for composition, and  $\bigvee_i h_i: X \rightarrow Y$  and  $h_i \leq h_j: X \rightrightarrows Y$  for its local structure;  $\mathcal{Q}(A, B)$  is the hom-lattice of arrows from  $A$  to  $B$ . The identity arrow on an object  $A \in \mathcal{Q}$  is written  $1_A: A \rightarrow A$ . The bottom element of a sup-lattice  $\mathcal{Q}(A, B)$  will typically be denoted by  $0_{A,B}$ . With these notations for identity and bottom, we can write a “Kronecker delta”

$$\delta_{A,B}: A \rightarrow B = \begin{cases} 1_A: A \rightarrow A & \text{if } A = B, \\ 0_{A,B}: A \rightarrow B & \text{otherwise.} \end{cases}$$

As we may consider **Sup** to be a “simplified version of **Cat**”, we may regard quantaloids as “simplified bicategories”. Notably, a quantaloid  $\mathcal{Q}$  has small hom-categories with stable local colimits, and therefore it is closed. Considering morphisms  $f: A \rightarrow B$  and  $g: B \rightarrow C$  we note the respective adjoints to composition in  $\mathcal{Q}$  as  $- \circ f \dashv \{f, -\}$  and  $g \circ - \dashv [g, -]$ ; that is to say, for any  $h: A \rightarrow C$  we have that  $g \circ f \leq h$  iff  $f \leq [g, h]$  iff  $g \leq \{f, h\}$ . Note furthermore that every diagram of 2-cells in a quantaloid trivially commutes.

There are a couple of lemmas involving  $[-, -]$  and  $\{-, -\}$ , that hold in any quantaloid, upon which we rely quite often. Let us give a short overview.

**2.2. LEMMA.** We work in a quantaloid  $\mathcal{Q}$ .

1. For  $f: A \rightarrow B$ ,  $g: B \rightarrow C$  and  $h: A \rightarrow C$ ,  $[g, h] = \bigvee \{x: A \rightarrow B \mid g \circ x \leq h\}$  and  $\{f, h\} = \bigvee \{y: B \rightarrow C \mid y \circ f \leq h\}$ .

2. The following are equivalent:

---

<sup>2</sup>By an “ordered set” we mean a set endowed with a transitive, reflexive relation; it is what is often called a “preordered set”. We will be explicit when we mean one such relation that is moreover antisymmetric, i.e. a “partial order”.


- (a)  $f \dashv g: B \xleftrightarrow{\quad} A$  in  $\mathcal{Q}$  (i.e.  $1_A \leq g \circ f$  and  $f \circ g \leq 1_B$ );
- (b)  $(f \circ -) \dashv (g \circ -): \mathcal{Q}(-, B) \xleftrightarrow{\quad} \mathcal{Q}(-, A)$  in  $\mathbf{Sup}$ ;
- (c)  $(- \circ g) \dashv (- \circ f): \mathcal{Q}(B, -) \xleftrightarrow{\quad} \mathcal{Q}(A, -)$  in  $\mathbf{Sup}$ ;
- (d)  $(g \circ -) = [f, -]: \mathcal{Q}(-, B) \rightarrow \mathcal{Q}(-, A)$  in  $\mathbf{Sup}$ ;
- (e)  $(- \circ f) = \{g, -\}: \mathcal{Q}(B, -) \rightarrow \mathcal{Q}(A, -)$  in  $\mathbf{Sup}$ .

3. An arrow  $f: A \rightarrow B$  has a right adjoint if and only if  $[f, 1_B] \circ f = [f, f]$ ; in this case the right adjoint to  $f$  is  $[f, 1_B]$ . Dually,  $g: B \rightarrow A$  in  $\mathcal{Q}$  has a left adjoint if and only if  $g \circ \{g, 1_B\} = \{g, g\}$ ; in this case the left adjoint to  $g$  is  $\{g, 1_B\}$ .
4. For  $f, f': A \rightrightarrows B$  and  $g, g': B \rightrightarrows A$  such that  $f \dashv g$  and  $f' \dashv g'$ ,  $f \leq f'$  if and only if  $g' \leq g$ .
5. Any  $f: A \rightarrow B$  induces, for every  $X \in \mathcal{Q}_0$ , an adjunction

$$[-, f] \dashv \{-, f\}: \mathcal{Q}(X, B) \xleftrightarrow{\quad} \mathcal{Q}(A, X)^{\text{op}}.$$

By  $\mathcal{Q}(A, X)^{\text{op}}$  is meant the sup-lattice  $\mathcal{Q}(A, X)$  with opposite order.

6. For arrows as in the diagrams below, we have the identity  $[f, [g, h]] = [g \circ f, h]$ , its dual  $\{k, \{l, m\}\} = \{k \circ l, m\}$  and the self-dual  $[x, \{y, z\}] = \{y, [x, z]\}$ .



7. For arrows as in the diagrams below, we have the identity  $[h, g] \circ [g, f] \leq [h, f]$  and its dual  $\{l, m\} \circ \{k, l\} \leq \{k, m\}$ . Also  $1_{\text{dom}(f)} \leq [f, f]$ , and dually  $1_{\text{cod}(f)} \leq \{f, f\}$ , hold.



$\mathbf{Sup}$  may also be thought of as an “infinitary version of  $\mathbf{Ab}$ ”; a quantaloid is then like a “ring(oid) with infinitary (and idempotent) sum”. This point of view helps explain some of the terminology below—especially when we talk about “matrices with elements in a quantaloid  $\mathcal{Q}$ ” in the appendix.

**2.3. EXAMPLE.** Any locale  $\Omega$  is a (very particular) quantale. Further on we shall denote by  $\mathbf{2}$  the two-element boolean algebra. Given a locale  $\Omega$ , [Walters, 1981] uses the quantaloid of relations in  $\Omega$ : the objects of  $\text{Rel}(\Omega)$  are the elements of  $\Omega$ , the hom-lattices are given by  $\text{Rel}(\Omega)(u, v) = \{w \in \Omega \mid w \leq u \wedge v\}$  with order inherited from  $\Omega$ , composition is given by infimum in  $\Omega$ , and the identity on an object  $u$  is  $u$  itself. This quantaloid is very particular: it equals its opposite.

**2.4. EXAMPLE.** The extended non-negative reals  $[0, +\infty]$  form a (commutative) quantale for the opposite order and addition as binary operation [Lawvere, 1973].

**2.5. EXAMPLE.** The ideals in a commutative ring  $R$  form a (commutative) quantale. When  $R$  is not commutative, the two-sided ideals still form a (commutative) quantale. But, as G. Van den Bossche [1995] points out (but she credits B. Lawvere for this idea), there is also quite naturally a “quantaloid of ideals” containing a lot more information than just the two-sided ideals. Denoting  $\mathcal{Q}_R$  for this structure, define that it has two objects, 0 and 1; the hom-sup-lattices are

$$\begin{aligned} \mathcal{Q}_R(0, 0) &= \text{additive subgroups of } R \text{ which are } \mathcal{Z}(R)\text{-modules,} \\ \mathcal{Q}_R(0, 1) &= \text{left-sided ideals of } R, \\ \mathcal{Q}_R(1, 0) &= \text{right-sided ideals of } R, \\ \mathcal{Q}_R(1, 1) &= \text{two-sided ideals of } R, \end{aligned}$$

with sum of additive subgroups as supremum:

$$\sum_{k \in K} I_k = \{\text{finite sums of elements in } \bigcup_{k \in K} I_k\}.$$

Composition in  $\mathcal{Q}_R$  is the multiplication of additive subgroups, as in

$$I \circ J = \{\text{finite sums } i_1 j_1 + \dots + i_n j_n \text{ with all } i_k \in I \text{ and all } j_k \in J\};$$

and the identity arrow on 1 is  $R$  and that on 0 is  $\mathcal{Z}(R)$ . (By  $\mathcal{Z}(R)$  we denote the center of  $R$ ). For a commutative  $R$ , this quantaloid  $\mathcal{Q}_R$  is equivalent as **Sup**-category to the quantale of ideals in  $R$ .

More examples can be found in the literature, e.g. [Rosenthal, 1996].

### 3. Three basic definitions

In the following  $\mathcal{Q}$  always stands for a quantaloid, and  $\mathcal{Q}_0$  for its (possibly large) class of objects. A  *$\mathcal{Q}$ -typed set*  $X$  is a (small) set  $X$  together with a mapping  $t: X \rightarrow \mathcal{Q}_0: x \mapsto tx$  sending each element  $x \in X$  to its *type*  $tx \in \mathcal{Q}_0$ . The notation with a “ $t$ ” for the types of elements in a  $\mathcal{Q}$ -typed set is generic; i.e. even for two different  $\mathcal{Q}$ -typed sets  $X$  and  $Y$ , the type of  $x \in X$  is written  $tx$ , and that of  $y \in Y$  is  $ty$ . A  $\mathcal{Q}$ -typed set  $X$  is just a way of writing a set-indexed family  $(tx)_{x \in X}$  of objects (i.e. a small discrete diagram) in  $\mathcal{Q}$ . If  $\mathcal{Q}$  is a small quantaloid, then a  $\mathcal{Q}$ -typed set is an object of the slice category  $\mathbf{Set}/\mathcal{Q}_0$ .

3.1. DEFINITION. A  $\mathcal{Q}$ -enriched category (or  $\mathcal{Q}$ -category for short)  $\mathbb{A}$  consists of

- objects: a  $\mathcal{Q}$ -typed set  $\mathbb{A}_0$ ,
- hom-arrows: for all  $a, a' \in \mathbb{A}_0$ , an arrow  $\mathbb{A}(a', a): ta \rightarrow ta'$  in  $\mathcal{Q}$ ,

satisfying

- composition-inequalities: for all  $a, a', a'' \in \mathbb{A}_0$ ,  $\mathbb{A}(a'', a') \circ \mathbb{A}(a', a) \leq \mathbb{A}(a'', a)$  in  $\mathcal{Q}$ ,
- identity-inequalities: for all  $a \in \mathbb{A}_0$ ,  $1_{ta} \leq \mathbb{A}(a, a)$  in  $\mathcal{Q}$ .

3.2. DEFINITION. A distributor  $\Phi: \mathbb{A} \rightleftarrows \mathbb{B}$  between two  $\mathcal{Q}$ -categories is given by

- distributor-arrows: for all  $a \in \mathbb{A}_0$ ,  $b \in \mathbb{B}_0$ , an arrow  $\Phi(b, a): ta \rightarrow tb$  in  $\mathcal{Q}$

satisfying

- action-inequalities: for all  $a, a' \in \mathbb{A}_0$ ,  $b, b' \in \mathbb{B}_0$ ,  $\mathbb{B}(b', b) \circ \Phi(b, a) \leq \Phi(b', a)$  and  $\Phi(b, a) \circ \mathbb{A}(a, a') \leq \Phi(b, a')$  in  $\mathcal{Q}$ .

3.3. DEFINITION. A functor  $F: \mathbb{A} \rightarrow \mathbb{B}$  between  $\mathcal{Q}$ -categories is

- object-mapping: a map  $F: \mathbb{A}_0 \rightarrow \mathbb{B}_0: a \mapsto Fa$

satisfying

- type-equalities: for all  $a \in \mathbb{A}_0$ ,  $ta = t(Fa)$  in  $\mathcal{Q}$ ,
- action-inequalities: for all  $a, a' \in \mathbb{A}_0$ ,  $\mathbb{A}(a', a) \leq \mathbb{B}(Fa', Fa)$  in  $\mathcal{Q}$ .

None of these definitions requires any of the usual diagrammatic axioms (associativity and identity axioms for the composition in a category, coherence of the action on a distributor with the composition in its (co)domain category, the functoriality of a functor) simply because those conditions, which require the commutativity of certain diagrams of 2-cells in the base quantaloid  $\mathcal{Q}$ , hold trivially! It is therefore a property of, rather than an extra structure on, a given set  $\mathbb{A}_0$  of elements with types in  $\mathcal{Q}$  together with hom-arrows  $\mathbb{A}(a', a): ta' \rightarrow ta$  (for  $a, a' \in \mathbb{A}_0$ ) to be a  $\mathcal{Q}$ -category; in other words, if these data determine a  $\mathcal{Q}$ -category, then they do so in only one way. Similar for distributors and functors: it is a property of a given collection of arrows  $\Phi(b, a): ta \rightarrow tb$  whether or not it determines a distributor between  $\mathcal{Q}$ -categories  $\mathbb{A}$  and  $\mathbb{B}$ , as it is a property of an object mapping  $F: \mathbb{A}_0 \rightarrow \mathbb{B}_0$  whether or not it determines a functor.

Our  $\mathcal{Q}$ -categories are by definition small: they have a set of objects. As a consequence, the collection of distributors between two  $\mathcal{Q}$ -categories is always a small set too, and so is the collection of functors between two  $\mathcal{Q}$ -categories. (However, we will soon run into size-related trouble: our base quantaloid  $\mathcal{Q}$  having a proper class of objects will conflict with the  $\mathcal{Q}$ -categories being small, in particular in matters related to cocompleteness of

$\mathcal{Q}$ -categories. When doing categorical algebra over  $\mathcal{Q}$  we will therefore suppose that  $\mathcal{Q}$  is small too. But this hypothesis is not necessary at this point.)

Note that for a distributor  $\Phi: \mathbb{A} \multimap \mathbb{B}$ , for all  $a \in \mathbb{A}$  and  $b \in \mathbb{B}$ ,

$$\Phi(b, a) = 1_{tb} \circ \Phi(b, a) \leq \mathbb{B}(b, b) \circ \Phi(b, a) \leq \bigvee_{b' \in \mathbb{B}} \mathbb{B}(b, b') \circ \Phi(b', a) \leq \Phi(b, a),$$

that is to say,  $\bigvee_{b' \in \mathbb{B}} \mathbb{B}(b, b') \circ \Phi(b', a) = \Phi(b, a)$ ;  $\bigvee_{a' \in \mathbb{A}} \Phi(b, a') \circ \mathbb{A}(a', a) = \Phi(b, a)$  is analogous. A  $\mathcal{Q}$ -category  $\mathbb{A}$  is itself a distributor from  $\mathbb{A}$  to  $\mathbb{A}$ ; and the identities above become, for all  $a, a'' \in \mathbb{A}$ ,  $\bigvee_{a' \in \mathbb{A}} \mathbb{A}(a'', a') \circ \mathbb{A}(a', a) = \mathbb{A}(a'', a)$ . These identities allow to say, in 3.4, with a suitable definition for the composition of distributors, that “ $\mathbb{A}$  is the identity distributor on  $\mathbb{A}$ ”.

There is a notational issue that we should comment on. We have chosen to write the composition of arrows in a base quantaloid  $\mathcal{Q}$  “from right to left”: the composite of  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  is  $g \circ f: X \rightarrow Z$ . Therefore we have chosen to write the hom-arrows in a  $\mathcal{Q}$ -enriched category  $\mathbb{A}$  also “from right to left”: for two objects  $a, a' \in \mathbb{A}$ , the hom-arrow  $\mathbb{A}(a', a)$  goes from  $ta$  to  $ta'$ . Doing so it is clear that, for example, the composition-inequality in  $\mathbb{A}$  is written  $\mathbb{A}(a'', a') \circ \mathbb{A}(a', a) \leq \mathbb{A}(a'', a)$ , with the *pivot*  $a'$  nicely in the middle, which we find very natural. Our notational conventions are thus basically those of R. Street’s seminal paper [1983a]; other authors have chosen other notations.

The next proposition displays the calculus of  $\mathcal{Q}$ -categories and distributors.

**3.4. PROPOSITION.**  *$\mathcal{Q}$ -categories are the objects, and distributors the arrows, of a quantaloid  $\text{Dist}(\mathcal{Q})$  in which*

- *the composition  $\Psi \otimes_{\mathbb{B}} \Phi: \mathbb{A} \multimap \mathbb{C}$  of two distributors  $\Phi: \mathbb{A} \multimap \mathbb{B}$  and  $\Psi: \mathbb{B} \multimap \mathbb{C}$  has as distributor-arrows, for  $a \in \mathbb{A}_0$  and  $c \in \mathbb{C}_0$ ,*

$$(\Psi \otimes_{\mathbb{B}} \Phi)(c, a) = \bigvee_{b \in \mathbb{B}_0} \Psi(c, b) \circ \Phi(b, a);$$

- *the identity distributor on a  $\mathcal{Q}$ -category  $\mathbb{A}$  has as distributor-arrows precisely the hom-arrows of the category  $\mathbb{A}$  itself, so we simply write it as  $\mathbb{A}: \mathbb{A} \multimap \mathbb{A}$ ;*
- *the supremum  $\bigvee_{i \in I} \Phi_i: \mathbb{A} \multimap \mathbb{B}$  of given distributors  $(\Phi_i: \mathbb{A} \multimap \mathbb{B})_{i \in I}$  is calculated elementwise, thus its distributor-arrows are, for  $a \in \mathbb{A}_0$  and  $b \in \mathbb{B}_0$ ,*

$$\left( \bigvee_{i \in I} \Phi_i \right)(b, a) = \bigvee_{i \in I} \Phi_i(b, a).$$

The proof of the fact that the data above define a quantaloid is straightforward. Actually,  $\text{Dist}(\mathcal{Q})$  is a universal construction on  $\mathcal{Q}$  in **QUANT**: there is a fully faithful homomorphism of quantaloids

$$\mathcal{Q} \rightarrow \text{Dist}(\mathcal{Q}): \left( f: A \rightarrow B \right) \mapsto \left( (f): *_{A} \multimap *_{B} \right) \quad (9)$$

sending an object  $A \in \mathcal{Q}$  to the  $\mathcal{Q}$ -category with only one object, say  $*$ , of type  $t* = A$ , and hom-arrow  $1_A$ , and a  $\mathcal{Q}$ -arrow  $f: A \rightarrow B$  to the distributor  $(f): *_{A} \multimap *_{B}$  whose single element is  $f$ . This turns out to be the universal direct-sum-and-split-monad completion of  $\mathcal{Q}$  in **QUANT**. The appendix gives details. (Even for a small base quantaloid  $\mathcal{Q}$ ,  $\text{Dist}(\mathcal{Q})$  has a proper class of objects; so large quantaloids arise as universal constructions on small ones. Therefore we do not wish to exclude large quantaloids *a priori*.)

Since  $\text{Dist}(\mathcal{Q})$  is a quantaloid, it is in particular closed; the importance of this fact cannot be overestimated. Let for example  $\Theta: \mathbb{A} \multimap \mathbb{C}$  and  $\Psi: \mathbb{B} \multimap \mathbb{C}$  be distributors between  $\mathcal{Q}$ -categories, then  $[\Psi, \Theta]: \mathbb{A} \multimap \mathbb{B}$  is the distributor with distributor-arrows, for  $a \in \mathbb{A}_0$  and  $b \in \mathbb{B}_0$ ,

$$[\Psi, \Theta](b, a) = \bigwedge_{c \in \mathbb{C}_0} [\Psi(c, b), \Theta(c, a)], \quad (10)$$

where the liftings on the right are calculated in  $\mathcal{Q}$ . A similar formula holds for  $\{-, -\}$ .

The category of  $\mathcal{Q}$ -categories and functors is the obvious one.

**3.5. PROPOSITION.**  *$\mathcal{Q}$ -categories are the objects, and functors the arrows, of a category  $\text{Cat}(\mathcal{Q})$  in which*

- *the composition  $G \circ F: \mathbb{A} \rightarrow \mathbb{C}$  of two functors  $F: \mathbb{A} \rightarrow \mathbb{B}$  and  $G: \mathbb{B} \rightarrow \mathbb{C}$  is determined by the composition of object maps  $G \circ F: \mathbb{A}_0 \rightarrow \mathbb{C}_0: a \mapsto G(F(a))$ ;*
- *the identity functor  $1_{\mathbb{A}}: \mathbb{A} \rightarrow \mathbb{A}$  on a  $\mathcal{Q}$ -category  $\mathbb{A}$  is determined by the identity object map  $1_{\mathbb{A}}: \mathbb{A}_0 \rightarrow \mathbb{A}_0: a \mapsto a$ .*

Every functor between  $\mathcal{Q}$ -categories induces an adjoint pair of distributors, and the resulting inclusion of the functor category in the distributor category – although straightforward – is a key element for the development of the theory of  $\mathcal{Q}$ -enriched categories.

**3.6. PROPOSITION.** *For  $\mathcal{Q}$ -categories and functors  $F: \mathbb{A} \rightarrow \mathbb{B}$  and  $G: \mathbb{C} \rightarrow \mathbb{B}$ , the  $\mathcal{Q}$ -arrows  $\mathbb{B}(Gc, Fa): ta \rightarrow tc$ , one for each  $(a, c) \in \mathbb{A}_0 \times \mathbb{C}_0$ , determine a distributor<sup>3</sup>  $\mathbb{B}(G-, F-): \mathbb{A} \multimap \mathbb{C}$ . For any functor  $F: \mathbb{A} \rightarrow \mathbb{B}$  the distributors  $\mathbb{B}(1_{\mathbb{B}}-, F-): \mathbb{A} \multimap \mathbb{B}$  and  $\mathbb{B}(F-, 1_{\mathbb{B}}-): \mathbb{B} \multimap \mathbb{A}$  are adjoint in the quantaloid  $\text{Dist}(\mathcal{Q}): \mathbb{B}(1_{\mathbb{B}}-, F-) \dashv \mathbb{B}(F-, 1_{\mathbb{B}}-)$ .*

**PROOF.** For any  $a, a' \in \mathbb{A}_0$  and  $c, c' \in \mathbb{C}_0$ ,  $\mathbb{C}(c', c) \circ \mathbb{B}(Gc, Fa) \circ \mathbb{A}(a, a') \leq \mathbb{B}(Gc', Gc) \circ \mathbb{B}(Gc, Fa) \circ \mathbb{B}(Fa, Fa') \leq \mathbb{B}(Gc', Fa)$  by functoriality of  $F$  and  $G$  and composition in  $\mathbb{B}$ . So  $\mathbb{B}(G-, F-)$  is a distributor from  $\mathbb{A}$  to  $\mathbb{B}$ .

To see that  $\mathbb{B}(1_{\mathbb{B}}-, F-) \dashv \mathbb{B}(F-, 1_{\mathbb{B}}-)$  in  $\text{Dist}(\mathcal{Q})$ , we must check two inequalities: the unit of the adjunction is due to the composition in  $\mathbb{B}$  and functoriality of  $F$ ,  $\mathbb{B}(F-, 1_{\mathbb{B}}-) \otimes_{\mathbb{B}} \mathbb{B}(1_{\mathbb{B}}-, F-) = \mathbb{B}(F-, F-) \geq \mathbb{A}(-, -)$ ; the counit follows from the fact that  $\{Fa \mid a \in \mathbb{A}_0\} \subseteq \mathbb{B}_0$  and – again – composition in  $\mathbb{B}$ ,  $\mathbb{B}(1_{\mathbb{B}}-, F-) \otimes_{\mathbb{A}} \mathbb{B}(F-, 1_{\mathbb{B}}-) \leq \mathbb{B}(1_{\mathbb{B}}-, -) \otimes_{\mathbb{B}} \mathbb{B}(-, 1_{\mathbb{B}}-) = \mathbb{B}(-, -)$ . ■

<sup>3</sup>There is a converse: for any distributor  $\Phi: \mathbb{A} \multimap \mathbb{C}$  there is a – universal – way in which  $\Phi = \mathbb{B}(G-, F-)$  for certain functors  $F: \mathbb{A} \rightarrow \mathbb{B}$  and  $G: \mathbb{C} \rightarrow \mathbb{B}$ .

In the following we use the abbreviated notations  $\mathbb{B}(-, F-) = \mathbb{B}(1_{\mathbb{B}}-, F-)$  and  $\mathbb{B}(F-, -) = \mathbb{B}(F-, 1_{\mathbb{B}}-)$ .

**3.7. PROPOSITION.** *Sending a functor to the left adjoint distributor that it induces, as in*

$$\text{Cat}(\mathcal{Q}) \longrightarrow \text{Dist}(\mathcal{Q}): (F: \mathbb{A} \longrightarrow \mathbb{B}) \mapsto (\mathbb{B}(-, F-): \mathbb{A} \dashv \!\!\! \dashv \mathbb{B}),$$

*is functorial. Sending a functor to the right adjoint determines a similar contravariant functor.*

**PROOF.** Trivially an identity functor  $1_{\mathbb{A}}: \mathbb{A} \longrightarrow \mathbb{A}$  is mapped onto the identity distributor  $\mathbb{A}(-, 1_{\mathbb{A}}-) = \mathbb{A}(-, -): \mathbb{A} \dashv \!\!\! \dashv \mathbb{A}$ . And if  $F: \mathbb{A} \longrightarrow \mathbb{B}$  and  $G: \mathbb{B} \longrightarrow \mathbb{C}$ , then we want  $\mathbb{C}(-, G \circ F-) = \mathbb{C}(-, G-) \otimes_{\mathbb{B}} \mathbb{B}(-, F-)$  but also this holds trivially: read the right hand side of the equation as the action of  $\mathbb{B}$  on  $\mathbb{C}(-, G-)$ . ■

The category  $\text{Cat}(\mathcal{Q})$  inherits the local structure from the quantaloid  $\text{Dist}(\mathcal{Q})$  via the functor  $\text{Cat}(\mathcal{Q}) \longrightarrow \text{Dist}(\mathcal{Q})$ : we put, for two functors  $F, G: \mathbb{A} \rightrightarrows \mathbb{B}$ ,

$$F \leq G \iff \mathbb{B}(-, F-) \leq \mathbb{B}(-, G-) \quad \left( \iff \mathbb{B}(G-, -) \leq \mathbb{B}(F-, -) \right).$$

Thus every hom-set  $\text{Cat}(\mathcal{Q})(\mathbb{A}, \mathbb{B})$  is (neither antisymmetrically nor cocompletely) ordered, and composition in  $\text{Cat}(\mathcal{Q})$  distributes on both sides over the local order:  $\text{Cat}(\mathcal{Q})$  is a 2-category, and the functor in 3.7 is a 2-functor which is the identity on objects, “essentially faithful” (but not full), and locally fully faithful. (The contravariant version of this functor reverses the local order!)

In general the “opposite” of a  $\mathcal{Q}$ -category  $\mathbb{A}$  is not again a  $\mathcal{Q}$ -category, but rather a  $\mathcal{Q}^{\text{op}}$ -category: of course  $\mathbb{A}^{\text{op}}$  is defined to have the same  $\mathcal{Q}$ -typed set of objects as  $\mathbb{A}$ , but the hom-arrows are reversed: for objects  $a, a'$  put  $\mathbb{A}^{\text{op}}(a', a) = \mathbb{A}(a, a')$ . Similarly, for a distributor  $\Phi: \mathbb{A} \dashv \!\!\! \dashv \mathbb{B}$  between  $\mathcal{Q}$ -categories we may define an opposite distributor between the opposite categories over the opposite base—but this distributor will go in the opposite direction:  $\Phi^{\text{op}}: \mathbb{B}^{\text{op}} \dashv \!\!\! \dashv \mathbb{A}^{\text{op}}$  is defined by  $\Phi^{\text{op}}(a, b) = \Phi(b, a)$ . For  $\Phi \leq \Psi: \mathbb{A} \dashv \!\!\! \dashv \mathbb{B}$  in  $\text{Dist}(\mathcal{Q})$  it is quite obvious that  $\Phi^{\text{op}} \leq \Psi^{\text{op}}: \mathbb{B}^{\text{op}} \dashv \!\!\! \dashv \mathbb{A}^{\text{op}}$  in  $\text{Dist}(\mathcal{Q}^{\text{op}})$ . Finally, for a functor  $F: \mathbb{A} \longrightarrow \mathbb{B}$  between  $\mathcal{Q}$ -categories, the *same* object mapping  $a \mapsto Fa$  determines an arrow  $F^{\text{op}}: \mathbb{A}^{\text{op}} \longrightarrow \mathbb{B}^{\text{op}}$  of  $\text{Cat}(\mathcal{Q}^{\text{op}})$ . But if  $F \leq G: \mathbb{A} \rightrightarrows \mathbb{B}$  in  $\text{Cat}(\mathcal{Q})$  then  $G^{\text{op}} \leq F^{\text{op}}: \mathbb{A}^{\text{op}} \rightrightarrows \mathbb{B}^{\text{op}}$  in  $\text{Cat}(\mathcal{Q}^{\text{op}})$ . It is obvious that applying the “op” twice, always gives back the original structure.

**3.8. PROPOSITION.** *“Taking opposites” determines isomorphisms*

$$\text{Dist}(\mathcal{Q}) \cong \text{Dist}(\mathcal{Q}^{\text{op}})^{\text{op}} \quad \text{and} \quad \text{Cat}(\mathcal{Q}) \cong \text{Cat}(\mathcal{Q}^{\text{op}})^{\text{co}}$$

*of 2-categories (where the “co” means: reversing order in the homs).*

These isomorphisms allow us to “dualize” all notions and results concerning  $\mathcal{Q}$ -enriched categories. For example, the dual of ‘left Kan extension’ in  $\text{Cat}(\mathcal{Q})$  is ‘right Kan extension’, and whereas the former help to characterize left adjoints in  $\text{Cat}(\mathcal{Q})$ , the latter do the same

for right adjoints. As another example, the dual of ‘weighted colimit’ in a  $\mathcal{Q}$ -category, is ‘weighted limit’; the former are ‘preserved’ by all left adjoint functors, the latter by right adjoint ones. Some notions are self-dual, like ‘equivalence’ of categories, or ‘Cauchy complete’ category.

**3.9. EXAMPLE.** In a quantaloid  $\mathcal{Q}$ , the arrows whose *codomain* is some object  $Y$  are the objects of a  $\mathcal{Q}$ -category that we will denote – anticipating 6.1 – by  $\mathcal{P}Y$ : put  $t(f: X \rightarrow Y) = X$  and  $\mathcal{P}Y(f', f) = [f', f]$ . Similarly – and again anticipating further results – we denote by  $\mathcal{P}^\dagger X$  the  $\mathcal{Q}$ -category whose objects are  $\mathcal{Q}$ -arrows with *domain*  $X$ , with types  $t(f: X \rightarrow Y) = Y$ , and  $\mathcal{P}^\dagger X(f', f) = \{f, f'\}$ .

**3.10. EXAMPLE.** Recall that  $\mathbf{2}$  is the 2-element Boolean algebra;  $\mathbf{2}$ -categories are orders, distributors are ideal relations, and functors are order-preserving maps.

**3.11. EXAMPLE.** Consider a sheaf  $F$  on a locale  $\Omega$ ; it determines a  $\text{Rel}(\Omega)$ -category  $\mathbb{F}$  whose objects are the partial sections of  $F$ , the type of a section  $s$  being the largest  $u \in \Omega$  on which it is defined, and whose hom-arrows  $\mathbb{F}(s', s)$  are, for sections  $s, s'$  of types  $u, u'$ , the largest  $v \leq u \wedge u'$  on which (restrictions of)  $s$  and  $s'$  agree [Walters, 1981].

**3.12. EXAMPLE.** A category enriched in the quantale  $[0, +\infty]$  (cf. 2.4) is a “generalized metric space” [Lawvere, 1973]: the enrichment itself is a binary distance function taking values in the positive reals. In particular is the composition-inequality in such an enriched category the triangular inequality. A functor between such generalized metric spaces is a distance decreasing application. (These metric spaces are “generalized” in that the distance function is not symmetric, that the distance between two points being zero does not imply their being identical, and that the distance between two points may be infinite.)

**3.13. EXAMPLE.** A (not necessarily commutative) ring  $R$  determines a  $\mathcal{Q}_R$ -enriched category (with  $\mathcal{Q}_R$  as in 2.5): denoting it as  $\text{Comm}_R$ , its objects of type 0 are the elements of  $R$ , its objects of type 1 are the elements of  $\mathcal{Z}(R)$ , and hom-arrows are given by commutators:  $\text{Comm}(r, s) = \{x \in R \mid rx = xs\}$ .

## 4. Some direct consequences

### UNDERLYING ORDERS

For an object  $A$  of a quantaloid  $\mathcal{Q}$ , denote by  $*_A$  the one-object  $\mathcal{Q}$ -category whose hom-arrow is the identity  $1_A$ . Given a  $\mathcal{Q}$ -category  $\mathbb{A}$ , the set  $\{a \in \mathbb{A}_0 \mid ta = A\}$  is in bijection with  $\text{Cat}(\mathcal{Q})(*_A, \mathbb{A})$ : any such object  $a$  determines a “constant” functor  $\Delta a: *_A \rightarrow \mathbb{A}$ ; and any such functor  $F: *_A \rightarrow \mathbb{A}$  “picks out” an object  $a \in \mathbb{A}$ . We may thus order the objects of  $\mathbb{A}$  by ordering the corresponding constant functors; we will speak of the *underlying order*  $(\mathbb{A}_0, \leq)$  of the  $\mathcal{Q}$ -category  $\mathbb{A}$ . Explicitly, for two objects  $a, a' \in \mathbb{A}_0$  we have that  $a' \leq a$  if and only if  $A := ta = ta'$  and for all  $x \in \mathbb{A}_0$ ,  $\mathbb{A}(x, a') \leq \mathbb{A}(x, a)$  in  $\mathcal{Q}$ , or equivalently  $\mathbb{A}(a', x) \geq \mathbb{A}(a, x)$ , or equivalently  $1_A \leq \mathbb{A}(a', a)$ . Whenever two objects of

$\mathbb{A}$  are equivalent in  $\mathbb{A}$ 's underlying order ( $a \leq a'$  and  $a' \leq a$ ) then we say that they are *isomorphic objects* (and write  $a \cong a'$ ).

4.1. EXAMPLE. In the  $\mathcal{Q}$ -category  $\mathcal{P}Y$ , whose objects are  $\mathcal{Q}$ -arrows with codomain  $Y$  and whose hom-arrows are given by  $[-, -]$ , the underlying order coincides with the local order in  $\mathcal{Q}$ . More precisely, for  $f, g: X \rightrightarrows Y$  it is the same to say that  $f \leq g$  in  $(\mathcal{P}Y)_0$  as to say that  $f \leq g$  in  $\mathcal{Q}(X, Y)$ .

It is immediate that  $F \leq G: \mathbb{A} \rightrightarrows \mathbb{B}$  in  $\text{Cat}(\mathcal{Q})$  if and only if, for all  $a \in \mathbb{A}_0$ ,  $Fa \leq Ga$  in the underlying order of  $\mathbb{B}$ . This says that the local structure in the 2-category  $\text{Cat}(\mathcal{Q})$  is “pointwise order”. Equivalently we could have written that  $F \leq G$  if and only if  $1_{ta} \leq \mathbb{B}(Fa, Ga)$  for all  $a \in \mathbb{A}_0$ , which exhibits the resemblance with the usual notion of “enriched natural transformation”.

#### ADJOINTS AND EQUIVALENCES

An arrow  $F: \mathbb{A} \rightarrow \mathbb{B}$  is *left adjoint* to an arrow  $G: \mathbb{B} \rightarrow \mathbb{A}$  in  $\text{Cat}(\mathcal{Q})$  (and  $G$  is then *right adjoint* to  $F$ ), written  $F \dashv G$ , if  $1_{\mathbb{A}} \leq G \circ F$  and  $F \circ G \leq 1_{\mathbb{B}}$ . Due to the 2-category  $\text{Cat}(\mathcal{Q})$  being locally ordered, we need not ask any of the usual triangular coherence diagrams. The unicity of adjoints in the quantaloid  $\text{Dist}(\mathcal{Q})$  and the locally fully faithful  $\text{Cat}(\mathcal{Q}) \rightarrow \text{Dist}(\mathcal{Q})$  allow for the following equivalent expression.

4.2. PROPOSITION.  $F: \mathbb{A} \rightarrow \mathbb{B}$  is left adjoint to  $G: \mathbb{B} \rightarrow \mathbb{A}$  in  $\text{Cat}(\mathcal{Q})$  if and only if  $\mathbb{B}(F-, -) = \mathbb{A}(-, G-): \mathbb{B} \rightarrow \mathbb{A}$  in  $\text{Dist}(\mathcal{Q})$ .

Further,  $F: \mathbb{A} \rightarrow \mathbb{B}$  is an *equivalence* in  $\text{Cat}(\mathcal{Q})$  if there exists a  $G: \mathbb{B} \rightarrow \mathbb{A}$  such that  $G \circ F \cong 1_{\mathbb{A}}$  and  $F \circ G \cong 1_{\mathbb{B}}$  (in which case also  $G$  is an equivalence). Again because  $\text{Cat}(\mathcal{Q})$  is locally ordered, this is the same as saying that  $F$  is both left and right adjoint to some  $G$ . Again the functor  $\text{Cat}(\mathcal{Q}) \rightarrow \text{Dist}(\mathcal{Q})$  gives equivalent expressions.

4.3. PROPOSITION. Given functors  $F: \mathbb{A} \rightarrow \mathbb{B}$  and  $G: \mathbb{B} \rightarrow \mathbb{A}$  constitute an equivalence in  $\text{Cat}(\mathcal{Q})$  if and only if  $\mathbb{B}(-, F-): \mathbb{A} \rightarrow \mathbb{B}$  and  $\mathbb{A}(-, G-): \mathbb{B} \rightarrow \mathbb{A}$  constitute an isomorphism in  $\text{Dist}(\mathcal{Q})$ , if and only if  $\mathbb{B}(F-, -): \mathbb{B} \rightarrow \mathbb{A}$  and  $\mathbb{A}(G-, -): \mathbb{A} \rightarrow \mathbb{B}$  constitute an isomorphism in  $\text{Dist}(\mathcal{Q})$ .

For ordinary categories and functors it is well known that the equivalences are precisely the fully faithful functors which are essentially surjective on objects. This holds for  $\mathcal{Q}$ -enriched categories too: say that a functor  $F: \mathbb{A} \rightarrow \mathbb{B}$  is *fully faithful* if

$$\forall a, a' \in \mathbb{A}_0 : \mathbb{A}(a', a) = \mathbb{B}(Fa', Fa) \text{ in } \mathcal{Q},$$

and that is *essentially surjective on objects* whenever

$$\forall b \in \mathbb{B}_0, \exists a \in \mathbb{A}_0 : Fa \cong b \text{ in } \mathbb{B}.$$

In fact,  $F$  is fully faithful if and only if the unit of the adjunction  $\mathbb{B}(-, F-) \dashv \mathbb{B}(F-, -)$  in  $\text{Dist}(\mathcal{Q})$  saturates to an equality; and if  $F$  is essentially surjective on objects then necessarily the co-unit of this adjunction saturates to an equality (but a functor  $F: \mathbb{A} \rightarrow \mathbb{B}$  for which the co-unit of the induced adjunction is an equality – which is sometimes said to have a “dense image” – is not necessarily essentially surjective on objects).

4.4. PROPOSITION. *An arrow  $F: \mathbb{A} \rightarrow \mathbb{B}$  in  $\text{Cat}(\mathcal{Q})$  is an equivalence if and only if it is fully faithful and essentially surjective on objects.*

PROOF. Suppose that  $F$  is an equivalence, with inverse equivalence  $G$ . Then, by functoriality of  $F$  and  $G$ , and since  $G \circ F \cong 1_{\mathbb{A}}$ ,  $\mathbb{A}(a', a) \leq \mathbb{B}(Fa', Fa) \leq \mathbb{A}(GFa', GFa) = \mathbb{A}(a', a)$  for all  $a, a' \in \mathbb{A}_0$ . Further, using  $F \circ G \cong 1_{\mathbb{B}}$ , it follows that  $b \cong FGb$ , and  $Gb \in \mathbb{A}_0$  as required. Conversely, supposing that  $F$  is fully faithful and essentially surjective on objects, choose – using the essential surjectivity of  $F$  – for any  $b \in \mathbb{B}_0$  one particular object  $Gb \in \mathbb{A}_0$  for which  $FGb \cong b$ . Then the object mapping  $\mathbb{B}_0 \rightarrow \mathbb{A}_0: b \mapsto Gb$  is fully faithful itself (and therefore also functorial) due to the fully faithfulness of  $F$ :  $\mathbb{B}(b', b) = \mathbb{B}(FGb', FGb) = \mathbb{A}(Gb', Gb)$  for all  $b, b' \in \mathbb{B}_0$ . It is the required inverse equivalence. ■

The following is well known for ordinary categories, and holds for  $\mathcal{Q}$ -categories too; it will be useful further on.

4.5. PROPOSITION. *Suppose that  $F \dashv G: \mathbb{B} \rightleftarrows \mathbb{A}$  in  $\text{Cat}(\mathcal{Q})$ . Then  $F$  is fully faithful if and only if  $G \circ F \cong 1_{\mathbb{A}}$ , and  $G$  is fully faithful if and only if  $F \circ G \cong 1_{\mathbb{B}}$ . If moreover  $G \dashv H: \mathbb{A} \rightleftarrows \mathbb{B}$  in  $\text{Cat}(\mathcal{Q})$  then  $F$  is fully faithful if and only if  $H$  is fully faithful.*

PROOF.  $F \dashv G$  in  $\text{Cat}(\mathcal{Q})$  implies  $\mathbb{B}(F-, F-) = \mathbb{A}(-, G \circ F-)$  in  $\text{Dist}(\mathcal{Q})$ ; and  $F$  is fully faithful if and only if  $\mathbb{B}(F-, F-) = \mathbb{A}(-, -)$  in  $\text{Dist}(\mathcal{Q})$ . So, obviously,  $F$  is fully faithful if and only if  $\mathbb{A}(-, G \circ F-) = \mathbb{A}(-, -)$  in  $\text{Dist}(\mathcal{Q})$ , or equivalently,  $G \circ F \cong 1_{\mathbb{A}}$  in  $\text{Cat}(\mathcal{Q})$ . Likewise for the fully faithfulness of  $G$ . Suppose now that  $F \dashv G \dashv H$ , then  $\mathbb{A}(G \circ F-, -) = \mathbb{B}(F-, H-) = \mathbb{B}(-, G \circ H-)$ , which implies that  $\mathbb{A}(G \circ F-, -) = \mathbb{A}(-, -)$  if and only if  $\mathbb{B}(-, G \circ H-) = \mathbb{A}(-, -)$  so the result follows. ■

## LEFT KAN EXTENSIONS

Given  $\mathcal{Q}$ -categories and functors  $F: \mathbb{A} \rightarrow \mathbb{B}$  and  $G: \mathbb{A} \rightarrow \mathbb{C}$ , the *left Kan extension of  $F$  along  $G$*  is – in so far it exists – a functor  $K: \mathbb{C} \rightarrow \mathbb{B}$  such that  $K \circ G \geq F$  in a universal way: whenever  $K': \mathbb{C} \rightarrow \mathbb{B}$  satisfies  $K' \circ G \geq F$ , then  $K' \geq K$ . If the left Kan extension of  $F$  along  $G$  exists, then it is essentially unique; we denote it by  $\langle F, G \rangle$ . So  $\langle F, G \rangle: \mathbb{C} \rightarrow \mathbb{B}$  is the reflection of  $F \in \text{Cat}(\mathcal{Q})(\mathbb{A}, \mathbb{B})$  along

$$- \circ G: \text{Cat}(\mathcal{Q})(\mathbb{C}, \mathbb{B}) \rightarrow \text{Cat}(\mathcal{Q})(\mathbb{A}, \mathbb{B}): H \mapsto H \circ G.$$

Left adjoint functors may be characterized in terms of left Kan extensions; this uses the idea of functors that preserve Kan extensions. Suppose that the left Kan extension  $\langle F, G \rangle: \mathbb{C} \rightarrow \mathbb{B}$  exists; then a functor  $F': \mathbb{B} \rightarrow \mathbb{B}'$  is said to *preserve  $\langle F, G \rangle$*  if  $\langle F' \circ F, G \rangle$  exists and is isomorphic to  $F' \circ \langle F, G \rangle$ . And  $\langle F, G \rangle$  is *absolute* if it is preserved by any  $F': \mathbb{B} \rightarrow \mathbb{B}'$ .

4.6. PROPOSITION. *For a functor  $F: \mathbb{A} \rightarrow \mathbb{B}$  between  $\mathcal{Q}$ -categories the following are equivalent:*

1.  $F$  has a right adjoint;

2.  $\langle 1_{\mathbb{A}}, F \rangle$  exists and is absolute;
3.  $\langle 1_{\mathbb{A}}, F \rangle$  exists and is preserved by  $F$  itself.

In this case,  $\langle 1_{\mathbb{A}}, F \rangle$  is the right adjoint of  $F$ .

PROOF. For (1 $\Rightarrow$ 2), suppose that  $F \dashv G$ ; we shall prove that  $G$  is the absolute left Kan extension of  $1_{\mathbb{A}}$  along  $F$ . The unit of the adjunction already says that  $G \circ F \geq 1_{\mathbb{A}}$ ; and if  $K: \mathbb{B} \rightarrow \mathbb{A}$  is another functor such that  $K \circ F \geq 1_{\mathbb{A}}$ , then necessarily  $K = K \circ 1_{\mathbb{B}} \geq K \circ (F \circ G) = (K \circ F) \circ G \geq 1_{\mathbb{A}} \circ G = G$ , using now the unit of the adjunction. So indeed  $G \cong \langle 1_{\mathbb{A}}, F \rangle$ . Let now  $F': \mathbb{A} \rightarrow \mathbb{B}'$  be any functor; we claim that  $F' \circ G$  is the left Kan extension of  $F' \circ 1_{\mathbb{A}}$  along  $F$ . Already  $(F' \circ G) \circ F = F' \circ (G \circ F) \geq F' \circ 1_{\mathbb{A}}$  is obvious (using the unit of the adjunction); and if  $K': \mathbb{B} \rightarrow \mathbb{B}'$  is another functor such that  $K' \circ F \geq F' \circ 1_{\mathbb{A}}$ , then  $K' = K' \circ 1_{\mathbb{B}} \geq K' \circ (F \circ G) = (K' \circ F) \circ G \geq (F' \circ 1_{\mathbb{A}}) \circ G = F' \circ G$ . Thus,  $G$  is the absolute left Kan extension of  $1_{\mathbb{A}}$  along  $F$ .

(2 $\Rightarrow$ 3) being trivial, we now prove (3 $\Rightarrow$ 1); it suffices to prove that  $F \dashv \langle 1_{\mathbb{A}}, F \rangle$ . But the wanted unit  $\langle 1_{\mathbb{A}}, F \rangle \circ F \geq 1_{\mathbb{A}}$  is part of the universal property of the left Kan extension; and using the hypothesis that  $\langle 1_{\mathbb{A}}, F \rangle$  is preserved by  $F$  itself, we have  $F \circ \langle 1_{\mathbb{A}}, F \rangle = \langle F, F \rangle$  which is smaller than  $1_{\mathbb{B}}$  by the universal property of  $\langle F, F \rangle$  (since  $1_{\mathbb{B}} \circ F \geq F$ ). ■

The dual of this result says that a functor  $F: \mathbb{A} \rightarrow \mathbb{B}$  has a left adjoint in  $\text{Cat}(\mathcal{Q})$  if and only if the *right Kan extension* of  $1_{\mathbb{A}}$  along  $F$  exists and is absolute, if and only if this right Kan extension exists and is preserved by  $F$ . Of course, the definition of *right Kan extension* is the dual of that of left Kan extension: given  $F: \mathbb{A} \rightarrow \mathbb{B}$  and  $G: \mathbb{A} \rightarrow \mathbb{C}$  in  $\text{Cat}(\mathcal{Q})$ , the right Kan extension of  $F$  along  $G$  is a functor  $(F, G): \mathbb{C} \rightarrow \mathbb{B}$  such that  $(F, G)^{\text{op}}: \mathbb{C}^{\text{op}} \rightarrow \mathbb{B}^{\text{op}}$  is the left Kan extension of  $F^{\text{op}}$  along  $G^{\text{op}}$  in  $\text{Cat}(\mathcal{Q}^{\text{op}})$ . In elementary terms: there is a universal inequality  $(F, G) \circ G \leq F$  in  $\text{Cat}(\mathcal{Q})$ .

After having introduced weighted colimits in a  $\mathcal{Q}$ -category, we will discuss *pointwise* left Kan extensions: particular colimits that enjoy the universal property given above.

## SKELETAL CATEGORIES

To any order corresponds an antisymmetric order by passing to equivalence classes of elements of the order. For  $\mathcal{Q}$ -enriched categories this can be imitated: say that a  $\mathcal{Q}$ -category  $\mathbb{A}$  is *skeletal* if no two different objects in  $\mathbb{A}$  are isomorphic; equivalently this says that, for any  $\mathcal{Q}$ -category  $\mathbb{C}$ ,  $\text{Cat}(\mathcal{Q})(\mathbb{C}, \mathbb{A})$  is an antisymmetric order, i.e. that

$$\text{Cat}(\mathcal{Q})(\mathbb{C}, \mathbb{A}) \rightarrow \text{Dist}(\mathcal{Q})(\mathbb{C}, \mathbb{A}): F \mapsto \mathbb{A}(-, F-)$$

is injective. Any  $\mathcal{Q}$ -category is then equivalent to its “skeletal quotient”.

**4.7. PROPOSITION.** *For a  $\mathcal{Q}$ -category  $\mathbb{A}$ , the following data define a skeletal  $\mathcal{Q}$ -category  $\mathbb{A}_{\text{skel}}$ :*

- *objects:*  $(\mathbb{A}_{\text{skel}})_0 = \left\{ [a] = \{x \in \mathbb{A}_0 \mid x \cong a\} \mid a \in \mathbb{A}_0 \right\}$ , with type function  $t[a] = ta$ ;
- *hom-arrows:* for any  $[a], [a'] \in (\mathbb{A}_{\text{skel}})_0$ ,  $\mathbb{A}_{\text{skel}}([a'], [a]) = \mathbb{A}(a', a)$ .

The object mapping  $[-]: \mathbb{A} \rightarrow \mathbb{A}_{\text{skel}}: x \mapsto [x]$  determines an equivalence in  $\text{Cat}(\mathcal{Q})$ .

PROOF. The construction of  $\mathbb{A}_{\text{skel}}$  is well-defined, because:

- as a set the quotient of  $\mathbb{A}_0$  by  $\cong$  is well-defined ( $\cdot \cong \cdot$  is an equivalence);
- $t[a]$  is well-defined (all elements of the equivalence class  $[a]$  necessarily have the same type since they are isomorphic);
- for any  $a' \cong b'$  and  $a \cong b$  in  $\mathbb{A}$ ,  $\mathbb{A}(b', b) = \mathbb{A}(a', a)$ , so  $\mathbb{A}_{\text{skel}}([a'], [a])$  is well-defined.

$\mathbb{A}_{\text{skel}}$  inherits “composition” and “identities” from  $\mathbb{A}$ , so it is a  $\mathcal{Q}$ -category. And  $\mathbb{A}_{\text{skel}}$  is skeletal: if  $[a] \cong [a']$  in  $(\mathbb{A}_{\text{skel}})_0$  then necessarily  $a \cong a'$  in  $\mathbb{A}_0$ , which by definition implies that  $[a] = [a']$ .

The mapping  $\mathbb{A}_0 \rightarrow (\mathbb{A}_{\text{skel}})_0: x \mapsto [x]$  determines a functor which is obviously fully faithful and essentially surjective on objects, so by 4.4 it is an equivalence in  $\text{Cat}(\mathcal{Q})$ . ■

Skeletal  $\mathcal{Q}$ -categories can be taken as objects of a full sub-2-category of  $\text{Cat}(\mathcal{Q})$ , respectively a full sub-quantaloid of  $\text{Dist}(\mathcal{Q})$ ; the proposition above then says that the embedding  $\text{Cat}_{\text{skel}}(\mathcal{Q}) \rightarrow \text{Cat}(\mathcal{Q})$  is a biequivalence of 2-categories, and  $\text{Dist}_{\text{skel}}(\mathcal{Q}) \rightarrow \text{Dist}(\mathcal{Q})$  is then an equivalence of quantaloids.  $\text{Cat}_{\text{skel}}(\mathcal{Q})$  is locally antisymmetrically ordered, so the obvious 2-functor

$$\text{Cat}_{\text{skel}}(\mathcal{Q}) \rightarrow \text{Dist}_{\text{skel}}(\mathcal{Q}): (F: \mathbb{A} \rightarrow \mathbb{B}) \mapsto (\mathbb{B}(-, F-): \mathbb{A} \rightarrow \mathbb{B})$$

is the identity on objects, faithful (but not full), and locally fully faithful.

4.8. EXAMPLE. The  $\mathcal{Q}$ -categories  $\mathcal{P}Y$  and  $\mathcal{P}^\dagger X$  as defined in 3.9 are skeletal.

4.9. EXAMPLE. Taking  $\mathcal{Q} = \mathbf{2}$ , the skeletal  $\mathbf{2}$ -categories are precisely the antisymmetric orders, i.e. the partial orders (cf. 3.10).

4.10. EXAMPLE. The  $\text{Rel}(\Omega)$ -category associated to a sheaf on  $\Omega$ , as in 3.11, is skeletal.

Later on we will encounter important  $\mathcal{Q}$ -categories which are always skeletal: the categories  $\mathcal{P}\mathbb{A}$  and  $\mathcal{P}^\dagger\mathbb{A}$  of (contravariant and covariant) presheaves on a given  $\mathcal{Q}$ -category  $\mathbb{A}$ , and also  $\mathbb{A}$ 's Cauchy completion  $\mathbb{A}_{\text{cc}}$ . (The reason, ultimately, that these  $\mathcal{Q}$ -categories are skeletal, is that the hom-objects of the quantaloid  $\mathcal{Q}$  are antisymmetrically ordered.)

## 5. Weighted (co)limits

### COLIMITS

We consider a functor  $F: \mathbb{A} \rightarrow \mathbb{B}$  and a distributor  $\Theta: \mathbb{C} \rightarrow \mathbb{A}$  between  $\mathcal{Q}$ -categories. Since  $F$  determines a distributor  $\mathbb{B}(F-, -): \mathbb{B} \rightarrow \mathbb{A}$ , and by closedness of the quantaloid  $\text{Dist}(\mathcal{Q})$ , we can calculate the universal lifting  $[\Theta, \mathbb{B}(F-, -)]: \mathbb{B} \rightarrow \mathbb{C}$ . A functor  $G: \mathbb{C} \rightarrow \mathbb{B}$  is the  $\Theta$ -weighted colimit of  $F$  if it represents the universal lifting:  $\mathbb{B}(G-, -) = [\Theta, \mathbb{B}(F-, -)]$ . If the  $\Theta$ -weighted colimit of  $F$  exists, then it is necessarily essentially unique. It therefore

makes sense to speak of “the” colimit and to denote it by  $\text{colim}(\Theta, F)$ ; its universal property is thus that

$$\mathbb{B}\left(\text{colim}(\Theta, F)-, -\right) = \left[\Theta, \mathbb{B}(F-, -)\right] \text{ in } \text{Dist}(\mathcal{Q}).$$

The following diagrams picture the situation:

We wish to speak of “those  $\mathcal{Q}$ -categories that admit all weighted colimits”, i.e. *cocomplete*  $\mathcal{Q}$ -categories. But there is a small problem (the word is well-chosen).

5.1. LEMMA. *A cocomplete  $\mathcal{Q}$ -category  $\mathbb{B}$  has at least as many objects as the base quantaloid  $\mathcal{Q}$ .*

PROOF. For each  $X \in \mathcal{Q}$ , consider the empty diagram in  $\mathbb{B}$  weighted by the empty distributor with domain  $*_X$ . Then  $\text{colim}(\emptyset, \emptyset): *_X \rightarrow \mathbb{B}$  must exist by cocompleteness of  $\mathbb{B}$ ; it “picks out” an object of type  $X$  in  $\mathbb{B}$ —thus  $\mathbb{B}$  must have such an object in the first place. ■

So, by the above, would the base quantaloid have a proper class of objects, then so would all cocomplete  $\mathcal{Q}$ -enriched categories. This is a problem in the framework of this text, because we did not even bother defining “large”  $\mathcal{Q}$ -categories, let alone develop a theory of  $\mathcal{Q}$ -categories that is sensitive to these size-related issues. *Therefore, from now on, we are happy to work with a small base quantaloid  $\mathcal{Q}$ .* The problem of “small” versus “large” then disappears: also “small”  $\mathcal{Q}$ -categories can be cocomplete.

Next up is a collection of lemmas that will help us calculate colimits. (There will be some abuse of notation: for a distributor  $\Phi: *_A \rightarrow \mathbb{A}$  and a functor  $F: \mathbb{A} \rightarrow \mathbb{B}$ ,  $\text{colim}(\Phi, F)$  is in principle a functor from  $*_A$  to  $\mathbb{B}$ . But such a functor simply “picks out” an object of type  $A$  in  $\mathbb{B}$ . Therefore we will often think of  $\text{colim}(\Phi, F)$  just as being that object. Of course, when the domain of the weight has more than one object, then any colimit with that weight is *really* a functor!)

5.2. LEMMA.

1. For  $(\Phi_i: \mathbb{C} \rightarrow \mathbb{A})_{i \in I}$  and  $F: \mathbb{A} \rightarrow \mathbb{B}$ , if all colimits involved exist, then  $\text{colim}(\bigvee_i \Phi_i, F)$  is the supremum of the  $\text{colim}(\Phi_i, F)$  in the order  $\text{Cat}(\mathcal{Q})(\mathbb{C}, \mathbb{B})$ :

$$\text{colim}\left(\bigvee_i \Phi_i, F\right) \cong \bigvee_i \text{colim}(\Phi_i, F).$$

2. For any  $F: \mathbb{A} \rightarrow \mathbb{B}$ ,  $\text{colim}(\mathbb{A}, F)$  (exists and) is isomorphic to  $F$ .

3. For  $\Phi: \mathbb{D} \multimap \mathbb{C}$ ,  $\Theta: \mathbb{C} \multimap \mathbb{A}$  and  $F: \mathbb{A} \rightarrow \mathbb{B}$ , suppose that the colimit  $\text{colim}(\Theta, F)$  exists; then  $\text{colim}(\Phi, \text{colim}(\Theta, F))$  exists if and only if  $\text{colim}(\Theta \otimes_{\mathbb{C}} \Phi, F)$  exists, in which case they are isomorphic.
4. For  $\Theta: \mathbb{C} \multimap \mathbb{A}$  and  $F: \mathbb{A} \rightarrow \mathbb{B}$ ,  $\text{colim}(\Theta, F)$  exists if and only if, for all objects  $c \in \mathbb{C}_0$ ,  $\text{colim}(\Theta(-, c), F)$  exists; then  $\text{colim}(\Theta, F)(c) \cong \text{colim}(\Theta(-, c), F)$ .

PROOF. (1) By assumption,

$$\begin{aligned} \mathbb{C}\left(\text{colim}\left(\bigvee_i \Phi_i, F\right)-, -\right) &= \left[\bigvee_i \Phi_i, \mathbb{B}(F-, -)\right] \\ &= \bigwedge_i \left[\Phi_i, \mathbb{B}(F-, -)\right] \\ &= \bigwedge_i \mathbb{B}\left(\text{colim}(\Phi_i, F)-, -\right) \end{aligned}$$

from which it follows that  $\text{colim}(\bigvee_i \Phi_i, F)$  is the supremum of the  $\text{colim}(\Phi_i, F)$  in the order  $\text{Cat}(\mathcal{Q})(\mathbb{C}, \mathbb{B})$ .

(2) Trivially,  $[\mathbb{A}(-, -), \mathbb{B}(F-, -)] = \mathbb{B}(F-, -)$ .

(3) By a simple calculation

$$\left[\Phi, \mathbb{B}\left(\text{colim}(\Theta, F)-, -\right)\right] = \left[\Phi, \left[\Theta, \mathbb{B}(F-, -)\right]\right] = \left[\Theta \otimes_{\mathbb{C}} \Phi, \mathbb{B}(F-, -)\right]$$

so the  $\Phi$ -weighted colimit of  $\text{colim}(\Theta, F)$  and the  $\Theta \otimes_{\mathbb{C}} \Phi$ -weighted colimit of  $F$  are the same thing.

(4) Necessity is easy:  $\text{colim}(\Theta, F): \mathbb{C} \rightarrow \mathbb{B}$  is a functor satisfying, for all  $c \in \mathbb{C}_0$  and  $b \in \mathbb{B}_0$ ,

$$\mathbb{B}\left(\text{colim}(\Theta, F)(c), b\right) = \left[\Theta(-, c), \mathbb{B}(F-, b)\right];$$

this literally says that  $\text{colim}(\Theta, F)(c)$  is the  $\Theta(-, c)$ -weighted colimit of  $F$  (which thus exists). As for sufficiency, we prove that the mapping

$$K: \mathbb{C}_0 \rightarrow \mathbb{B}_0: c \mapsto \text{colim}(\Theta(-, c), F)$$

is a functor:  $\mathbb{C}(c', c) \leq \mathbb{B}(Kc', Kc)$  because

$$\begin{aligned} \mathbb{B}(Kc', Kc) &= \mathbb{B}(Kc', -) \otimes_{\mathbb{B}} \mathbb{B}(-, Kc) \\ &= \left\{ \mathbb{B}(Kc, -), \mathbb{B}(Kc', -) \right\} \\ &= \left\{ \left[ \Phi(-, c), \mathbb{B}(F-, -) \right], \left[ \Phi(-, c'), \mathbb{B}(F-, -) \right] \right\} \end{aligned}$$

and, with slight abuse of notation<sup>4</sup>,

$$\begin{aligned} \mathbb{C}(c', c) &\leq \left\{ \left[ \Phi(-, c), \mathbb{B}(F-, -) \right], \left[ \Phi(-, c'), \mathbb{B}(F-, -) \right] \right\} \\ &\iff \Phi(-, c') \otimes_{*_{tc'}} \mathbb{C}(c', c) \otimes_{*_{tc}} \left[ \Phi(-, c), \mathbb{B}(F-, -) \right] \leq \mathbb{B}(F-, -) \end{aligned}$$

<sup>4</sup>The  $\mathcal{Q}$ -arrow  $\mathbb{C}(c', c): tc \rightarrow tc'$  is thought of as a one-element distributor  $*_{tc} \multimap *_{tc'}$ .

which holds because  $\Phi(-, c') \otimes_{*_{tc'}} \mathbb{C}(c', c) \leq \Phi(-, c)$  and by the universal property of the “ $[-, -]$ ”. This functor  $K$  has, by construction, the required universal property for it to be the  $\Phi$ -weighted colimit of  $F$ . ■

**5.3. LEMMA.** *For  $F \leq G: \mathbb{A} \rightrightarrows \mathbb{B}$  and  $\Theta: \mathbb{C} \dashrightarrow \mathbb{A}$ , suppose that  $\text{colim}(\Theta, F)$  and  $\text{colim}(\Theta, G)$  exist; then  $\text{colim}(\Theta, F) \leq \text{colim}(\Theta, G)$ . In particular, suppose that  $F \cong G$ ; then  $\text{colim}(\Theta, F)$  exists if and only if  $\text{colim}(\Theta, G)$  exists, in which case they are isomorphic.*

**PROOF.**  $[-, -]$  is order-preserving in its second variable, so

$$\mathbb{B}(\text{colim}(\Theta, F)-, -) = \left[ \Theta, \mathbb{B}(F-, -) \right] \geq \left[ \Theta, \mathbb{B}(G-, -) \right] = \mathbb{B}(\text{colim}(\Theta, G)-, -),$$

hence  $\text{colim}(\Theta, F) \leq \text{colim}(\Theta, G)$  as claimed. The second part of the statement is similar. ■

Let us point out two important consequences.

**5.4. COROLLARY.** *A  $\mathcal{Q}$ -category  $\mathbb{B}$  is cocomplete if and only if it admits colimits of the identity functor  $1_{\mathbb{B}}: \mathbb{B} \rightarrow \mathbb{B}$  weighted by distributors into  $\mathbb{B}$  whose domains are one-object  $\mathcal{Q}$ -categories<sup>5</sup>: for every  $\phi: *_B \dashrightarrow \mathbb{B}$  in  $\text{Dist}(\mathcal{Q})$ , there exists an object  $b \in \mathbb{B}$ , necessarily of type  $tb = B$ , such that*

$$\mathbb{B}(b, -) = \left[ \phi, \mathbb{B} \right] \text{ in } \text{Dist}(\mathcal{Q}).$$

**PROOF.** One direction is trivial. For the other, consider a weight  $\Theta: \mathbb{C} \rightarrow \mathbb{A}$  and a functor  $F: \mathbb{A} \rightarrow \mathbb{B}$ . Then

$$\left[ \Theta, \mathbb{B}(F-, -) \right] = \left[ \Theta, \left[ \mathbb{B}(-, F-), \mathbb{B} \right] \right] = \left[ \mathbb{B}(-, F-) \otimes_{\mathbb{A}} \Theta, \mathbb{B}(1_{\mathbb{B}}-, -) \right]$$

so that  $\text{colim}(\Theta, F): \mathbb{C} \rightarrow \mathbb{B}$  exists if and only if  $\text{colim}(\mathbb{B}(-, F-) \otimes_{\mathbb{A}} \Theta, 1_{\mathbb{B}}): \mathbb{C} \rightarrow \mathbb{B}$  exists. But the latter exists if and only if, for all objects  $c \in \mathbb{C}$ ,

$$\text{colim} \left( \mathbb{B}(-, F-) \otimes_{\mathbb{A}} \Theta(-, c), 1_{\mathbb{B}} \right): *_c \rightarrow \mathbb{B}$$

exists—which they do by hypothesis. ■

**5.5. EXAMPLE.** A  $\mathbf{2}$ -enriched category  $\mathbb{A}$  is an ordered set, and a distributor  $\phi: * \dashrightarrow \mathbb{A}$  is a down-closed subset of  $\mathbb{A}$ . Since  $[\phi, \mathbb{A}]$  is the up-closed set of lower bounds of  $\phi$ , it is representable if and only if the supremum of  $\phi$  exists in  $\mathbb{A}$ . Cocompleteness of  $\mathbb{A}$  as  $\mathbf{2}$ -category is thus the same thing as cocompleteness of  $\mathbb{A}$  as order (since it is equivalent to ask for the suprema of all subsets or only the suprema of down-closed sets).

<sup>5</sup>Later we will call those distributors *contravariant presheaves*, so  $\mathbb{B}$  is cocomplete if and only if it admits colimits of  $1_{\mathbb{B}}$  weighted by contravariant presheaves on  $\mathbb{B}$ .

5.6. COROLLARY. *Given  $(F_i: \mathbb{A} \rightarrow \mathbb{B})_{i \in I}$  in  $\text{Cat}(\mathcal{Q})$ , if the  $\bigvee_i \mathbb{B}(-, F_i-)$ -weighted colimit of  $1_{\mathbb{B}}$  exists, then it is their supremum in the order  $\text{Cat}(\mathcal{Q})(\mathbb{A}, \mathbb{B})$ :*

$$\bigvee_i F_i \cong \text{colim} \left( \bigvee_i \mathbb{B}(-, F_i-), 1_{\mathbb{B}} \right).$$


PROOF. The colimit, supposed to exist, satisfies

$$\begin{aligned} \mathbb{B} \left( \text{colim} \left( \bigvee_i \mathbb{B}(-, F_i-), 1_{\mathbb{B}} \right) -, - \right) &= \left[ \bigvee_i \mathbb{B}(-, F_i-), \mathbb{B} \right] \\ &= \bigwedge_i \left[ \mathbb{B}(-, F_i-), \mathbb{B} \right] \\ &= \bigwedge_i \mathbb{B}(F_i-, -). \end{aligned}$$

It follows straightforwardly that  $\text{colim}(\bigvee_i \mathbb{B}(-, F_i-), 1_{\mathbb{B}})$  is the supremum of the  $F_i$ . ■

This corollary thus says that, if there is a functor  $F: \mathbb{A} \rightarrow \mathbb{B}$  such that  $\mathbb{B}(F-, -) = \bigwedge_i \mathbb{B}(F_i-, -)$  for given  $(F_i: \mathbb{A} \rightarrow \mathbb{B})_{i \in I}$ , then  $F$  is the supremum of the  $F_i$ . In particular, for an object  $a \in \mathbb{A}$  it follows that  $Fa$  is the supremum of the  $F_i a$  in the underlying order  $(\mathbb{B}, \leq)$ .

5.7. EXAMPLE. Concerning the result in 5.6, it is *not* true that each supremum in the order  $\text{Cat}(\mathcal{Q})(\mathbb{A}, \mathbb{B})$  is necessarily a weighted colimit in  $\mathbb{B}$ . Consider the orders



then the order morphisms  $f(0) = a$ ,  $f(1) = c$  and  $g(0) = b$ ,  $g(1) = d$  admit the supremum  $h(0) = h(1) = c$ , but clearly  $h(0)$  is not the supremum of  $f(0)$  and  $g(0)$ . This implies that  $h$  is not the weighted colimit determined by  $f$  and  $g$  as in 5.6.

## LIMITS

A *limit* in a  $\mathcal{Q}$ -category  $\mathbb{A}$  is a colimit in the opposite category of  $\mathbb{A}$ . This simple definition hides the difficulty that  $\mathbb{A}$  is a  $\mathcal{Q}$ -category but  $\mathbb{A}^{\text{op}}$  is a  $\mathcal{Q}^{\text{op}}$ -category. Let us therefore make the definition of limit in  $\mathbb{A}$  elementary in terms of  $\mathcal{Q}$ -enriched structures: for a distributor  $\Phi: \mathbb{B} \rightarrow \mathbb{C}$  in  $\text{Dist}(\mathcal{Q})$  and a functor  $F: \mathbb{B} \rightarrow \mathbb{A}$  in  $\text{Cat}(\mathcal{Q})$ , the  $\Phi$ -weighted limit of  $F$  is – whenever it exists – a functor  $\lim(\Phi, F): \mathbb{C} \rightarrow \mathbb{A}$  in  $\text{Cat}(\mathcal{Q})$  such that

$$\mathbb{A} \left( -, \lim(\Phi, F) - \right) = \left\{ \Phi, \mathbb{A}(-, F-) \right\} \text{ in } \text{Dist}(\mathcal{Q}).$$

These diagrams picture the situation:

$$\begin{array}{ccc}
 \begin{array}{ccc} \mathbb{C} & & \\ \uparrow \Phi \circ & & \\ \mathbb{B} & \xrightarrow{F} & \mathbb{A} \end{array} & 
 \begin{array}{ccc} \mathbb{C} & & \\ \uparrow \Phi \circ & \searrow \{ \Phi, \mathbb{A}(-, F-) \} & \\ \mathbb{C} & \xrightarrow{\mathbb{A}(-, F-)} & \mathbb{A} \\ \wedge \downarrow & & \end{array} & 
 \begin{array}{ccc} \mathbb{C} & & \\ \uparrow \Phi \circ & \searrow \text{lim}(\Phi, F) & \\ \mathbb{B} & \xrightarrow{F} & \mathbb{A} \end{array}
 \end{array}$$

A  $\mathcal{Q}$ -category  $\mathbb{A}$  is *complete* when the  $\mathcal{Q}^{\text{op}}$ -category  $\mathbb{A}^{\text{op}}$  is cocomplete; by 5.4 we know that this is so precisely when for all  $\phi: *_A \dashv\vdash \mathbb{A}^{\text{op}}$  in  $\text{Dist}(\mathcal{Q}^{\text{op}})$ , the  $\phi$ -weighted colimit of  $1_{\mathbb{A}^{\text{op}}}$  exists in  $\text{Cat}(\mathcal{Q}^{\text{op}})$ ; in terms of  $\mathcal{Q}$ -enriched structures, this occurs precisely when for every<sup>6</sup>  $\phi: \mathbb{A} \dashv\vdash *_A$  in  $\text{Dist}(\mathcal{Q})$ , there is an object  $a \in \mathbb{A}$ , necessarily of type  $ta = A$ , such that

$$\mathbb{A}(-, a) = \{ \phi, \mathbb{A} \} \text{ in } \text{Dist}(\mathcal{Q}).$$

**5.8. PROPOSITION.** *For any  $\mathcal{Q}$ -category  $\mathbb{A}$  and any  $\phi: \mathbb{A} \dashv\vdash *_A$  in  $\text{Dist}(\mathcal{Q})$ ,  $\text{lim}(\phi, 1_{\mathbb{A}})$  exists if and only if  $\text{colim}(\{ \phi, \mathbb{A} \}, 1_{\mathbb{A}})$  exists, in which case they are isomorphic.*

**PROOF.** First suppose that  $a = \text{lim}(\phi, 1_{\mathbb{A}})$  exists:  $\mathbb{A}(-, a) = \{ \phi, \mathbb{A} \}$ . But then, using that  $\mathbb{A}(-, a) \dashv \mathbb{A}(a, -)$  in  $\text{Dist}(\mathcal{Q})$ ,  $\mathbb{A}(a, -) = [\mathbb{A}(-, a), \mathbb{A}] = [\{ \phi, \mathbb{A} \}, \mathbb{A}]$ , which says that  $a$  is the  $\{ \phi, \mathbb{A} \}$ -weighted colimit of  $1_{\mathbb{A}}$ .

Conversely, to say that  $a = \text{colim}(\{ \phi, \mathbb{A} \}, 1_{\mathbb{A}})$  exists, means that  $\mathbb{A}(a, -) = [\{ \phi, \mathbb{A} \}, \mathbb{A}]$ . Again using the adjunction  $\mathbb{A}(-, a) \dashv \mathbb{A}(a, -)$ , and by calculation with  $[-, -]$  and  $\{-, -\}$  in  $\text{Dist}(\mathcal{Q})$  it follows that  $\mathbb{A}(-, a) = \{ [\{ \phi, \mathbb{A} \}, \mathbb{A}], \mathbb{A} \} = \{ \phi, \mathbb{A} \}$ , so that indeed  $a$  is the  $\phi$ -weighted colimit of  $1_{\mathbb{A}}$ .  $\blacksquare$

This proposition is valid for categories enriched in any quantaloid  $\mathcal{Q}$ , hence also for  $\mathcal{Q}^{\text{op}}$ -categories; translating the result in 5.8 for  $\mathcal{Q}^{\text{op}}$ -categories in terms of  $\mathcal{Q}$ -categories, we obtain its dual: for any  $\psi: *_A \dashv\vdash \mathbb{A}$  in  $\text{Dist}(\mathcal{Q})$ ,  $\text{colim}(\psi, 1_{\mathbb{A}})$  exists if and only if  $\text{lim}([\psi, \mathbb{A}], 1_{\mathbb{A}})$  exists, in which case they are isomorphic.

**5.9. EXAMPLE.** Applied to  $\mathbf{2}$ -enriched categories, i.e. ordered sets, 5.8 says that the infimum of an up-closed subset equals the supremum of the down-set of its lower bounds (and dually, the supremum of a down-closed subset equals the infimum of the up-closed set of its upper bounds).

The following is then an immediate consequence of 5.4 and 5.8.

**5.10. PROPOSITION.** *A  $\mathcal{Q}$ -category is complete if and only if it is cocomplete.*

This does not mean that “the theory of complete  $\mathcal{Q}$ -categories” is *the same* as “the theory of cocomplete  $\mathcal{Q}$ -categories” (although it is its dual): even though the objects are the same, the appropriate morphisms are not. For example, the free cocompletion of a  $\mathcal{Q}$ -category is different from its free completion (see further).

<sup>6</sup>Later we will call such a distributor a *covariant presheaf* on  $\mathbb{A}$ .

## ABSOLUTE (CO)LIMITS AND (CO)CONTINUOUS FUNCTORS

Given a distributor  $\Theta: \mathbb{A} \multimap \mathbb{B}$  and a functor  $F: \mathbb{B} \rightarrow \mathbb{C}$  between  $\mathcal{Q}$ -categories for which  $\text{colim}(\Theta, F): \mathbb{C} \rightarrow \mathbb{B}$  exists, a functor  $F': \mathbb{B} \rightarrow \mathbb{B}'$  preserves this colimit if  $\text{colim}(\Theta, F' \circ F)$  exists and is isomorphic to  $F' \circ \text{colim}(\Theta, F)$ . The functor  $F'$  is *cocontinuous* if it preserves all colimits that exist in  $\mathbb{B}$ . And  $\text{colim}(\Theta, F)$  is *absolute* if it is preserved by any functor  $F': \mathbb{B} \rightarrow \mathbb{B}'$ .

*Continuous functors* and *absolute limits* are defined dually. For all of the following results we do not bother explicitly writing the dual statements, even though we may use them further on.

5.11. PROPOSITION. *If  $\Theta: \mathbb{C} \multimap \mathbb{A}$  is a left adjoint in  $\text{Dist}(\mathcal{Q})$ , then any colimit with weight  $\Theta$  is absolute<sup>7</sup>.*

PROOF. If  $\Theta \dashv \Psi$  in  $\text{Dist}(\mathcal{Q})$  and  $\text{colim}(\Theta, F)$  exists, then necessarily

$$\mathbb{B}\left(\text{colim}(\Theta, F)-, -\right) = \left[\Theta, \mathbb{B}(F-, -)\right] = \Psi \otimes_{\mathbb{A}} \mathbb{B}(F-, -).$$

From this it is obvious that, for any  $F': \mathbb{B} \rightarrow \mathbb{B}'$ ,

$$\begin{aligned} \mathbb{B}\left(F' \circ \text{colim}(\Theta, F)-, -\right) &= \mathbb{B}\left(\text{colim}(\Theta, F)-, -\right) \otimes_{\mathbb{B}} \mathbb{B}'(F'-, -) \\ &= \Psi \otimes_{\mathbb{A}} \mathbb{B}(F-, -) \otimes_{\mathbb{B}} \mathbb{B}'(F'-, -) \\ &= \Psi \otimes_{\mathbb{A}} \mathbb{B}(F' \circ F-, -) \\ &= \left[\Theta, \mathbb{B}(F' \circ F-, -)\right] \end{aligned}$$

so  $\text{colim}(\Theta, F' \circ F)$  exists and is isomorphic to  $F' \circ \text{colim}(\Theta, F)$ . ■

5.12. PROPOSITION. *If  $F': \mathbb{B} \rightarrow \mathbb{B}'$  is a left adjoint in  $\text{Cat}(\mathcal{Q})$ , then it is cocontinuous<sup>8</sup>.*

PROOF. Suppose that  $F' \dashv G$  in  $\text{Cat}(\mathcal{Q})$ —i.e.  $\mathbb{B}'(F'-, -) \dashv \mathbb{B}(G-, -)$  in  $\text{Dist}(\mathcal{Q})$ . For any colimit

$$\begin{array}{ccccc} \mathbb{A} & \xrightarrow{F} & \mathbb{B} & \xrightarrow{F'} & \mathbb{B}' \\ \uparrow \Theta & & \nearrow \text{colim}(\Theta, F) & & \\ \mathbb{C} & & & & \end{array}$$

we can then calculate

$$\mathbb{B}'\left(F' \circ \text{colim}(\Theta, F)-, -\right) = \mathbb{B}\left(\text{colim}(\Theta, F)-, -\right) \otimes_{\mathbb{B}} \mathbb{B}'(F'-, -)$$

<sup>7</sup>A converse can be proved too: if any  $\Theta$ -weighted colimit is absolute, then  $\Theta$  is left adjoint [Street, 1983b].

<sup>8</sup>If  $\mathbb{B}$  is cocomplete, then also the converse holds—see 6.8.

$$\begin{aligned}
&= [\Theta, \mathbb{B}(F-, -)] \otimes_{\mathbb{B}} \mathbb{B}'(F'-, -) \\
&= \left\{ \mathbb{B}(G-, -), [\Theta, \mathbb{B}(F-, -)] \right\} \\
&= [\Theta, \left\{ \mathbb{B}(G-, -), \mathbb{B}(F-, -) \right\}] \\
&= [\Theta, \mathbb{B}(F-, -) \otimes_{\mathbb{B}} \mathbb{B}'(F'-, -)] \\
&= [\Theta, \mathbb{B}'(F' \circ F-, -)]
\end{aligned}$$

which shows that  $\text{colim}(\Theta, F' \circ F)$  exists and equals  $F' \circ \text{colim}(\Theta, F)$ .  $\blacksquare$

5.13. COROLLARY. *Consider a fully faithful right adjoint  $G: \mathbb{A} \rightarrow \mathbb{B}$  in  $\text{Cat}(\mathcal{Q})$ ; if  $\mathbb{B}$  is cocomplete, then so is  $\mathbb{A}$ .*

PROOF. Let us write  $F: \mathbb{B} \rightarrow \mathbb{A}$  for  $G$ 's left adjoint; now consider any colimit diagram

$$\begin{array}{ccccc}
& & & & F \\
& & & & \downarrow \\
& & & & \perp \\
& & & & \downarrow \\
& & & & G \\
& & & & \downarrow \\
& & & & \mathbb{B} \\
& & & & \uparrow \\
& & & & \perp \\
& & & & \downarrow \\
& & & & F \\
\mathbb{D} & \xrightarrow{H} & \mathbb{A} & \xrightarrow{G} & \mathbb{B} \\
& & \uparrow & & \uparrow \\
& & \Phi & & \text{colim}(\Phi, G \circ H) \\
& & \uparrow & & \uparrow \\
& & \mathbb{C} & & \uparrow \\
& & & & \text{dotted arrow}
\end{array}$$

Then by cocontinuity of  $F$  we have  $F \circ \text{colim}(\Phi, G \circ H) \cong \text{colim}(\Phi, F \circ G \circ H)$  (thus in particular the existence of the right hand side, the dotted arrow in the diagram), and by fully faithfulness of  $G$  (see 4.5),  $\text{colim}(\Phi, F \circ G \circ H) \cong \text{colim}(\Phi, H)$ .  $\blacksquare$

The following two corollaries are now immediate.

5.14. COROLLARY. *Consider a string of three adjoints in  $\text{Cat}(\mathcal{Q})$ ,*

$$\begin{array}{ccc}
& & F \\
& & \downarrow \\
& & \perp \\
& & \downarrow \\
& & G \\
& & \downarrow \\
& & \perp \\
& & \downarrow \\
& & H \\
\mathbb{A} & \xleftrightarrow{G} & \mathbb{B}
\end{array}$$

*with  $F$  fully faithful. If  $\mathbb{B}$  is cocomplete, then so is  $\mathbb{A}$ .*

5.15. COROLLARY. *If  $F: \mathbb{A} \rightarrow \mathbb{B}$  is an equivalence in  $\text{Cat}(\mathcal{Q})$ , then both  $F$  and its inverse equivalence are cocontinuous, and  $\mathbb{A}$  is cocomplete if and only if  $\mathbb{B}$  is.*

## 6. Presheaves

### CONTRAVARIANT PRESHEAVES

Principally the ‘‘contravariant presheaves on a  $\mathcal{Q}$ -category  $\mathbb{A}$ ’’ should be the objects of a

$\mathcal{Q}$ -category  $\mathcal{P}\mathbb{A}$  which “classifies” distributors into  $\mathbb{A}$ :

$$\text{for every } \mathcal{Q}\text{-category } \mathbb{C}, \text{Dist}(\mathcal{Q})(\mathbb{C}, \mathbb{A}) \simeq \text{Cat}(\mathcal{Q})(\mathbb{C}, \mathcal{P}\mathbb{A}). \quad (11)$$

(Here we ask *a priori* for an equivalence of orders, but it will prove to be an isomorphism of sup-lattices.) Putting  $\mathbb{C} = *_C$  in the above equivalence dictates that a *contravariant presheaf of type  $C \in \mathcal{Q}$  on  $\mathbb{A}$*  is a distributor  $*_C \multimap \mathbb{A}$ .

In what follows we will systematically use small Greek letters – instead of capital ones – for such contravariant presheaves; the elements of such a  $\phi: *_C \multimap \mathbb{A}$  will be written as  $\phi(a): C \rightarrow ta$  (instead of  $\phi(a, *)$ , thus stressing the fact that the action of  $*_C$  on  $\phi$  is trivial).

The precise sense in which a distributor  $\phi: *_C \multimap \mathbb{A}$  is *contravariant*, is the following: the action of  $\mathbb{A}$  on  $\phi$  can be written as  $\mathbb{A}(a', a) \leq \{\phi(a), \phi(a')\}$  (for all  $a, a' \in \mathbb{A}_0$ ). A contravariant presheaf on  $\mathbb{A}$  may thus be thought of as a “functor on  $\mathbb{A}^{\text{op}}$  with values in  $\mathcal{Q}$ ” sending  $a$  to  $\phi(a)$ . To make this precise: indeed,  $a \mapsto \phi(a)$  is a functor between  $\mathbb{A}^{\text{op}}$  and  $(\mathcal{P}^\dagger C)^{\text{op}}$  (see 3.9) in  $\text{Cat}(\mathcal{Q}^{\text{op}})$ , or equivalently a functor between  $\mathbb{A}$  and  $\mathcal{P}^\dagger C$  in  $\text{Cat}(\mathcal{Q})^{\text{co}}$  (by the duality in 3.8).

Since we agreed to work over a *small* base quantaloid  $\mathcal{Q}$ , the collection of contravariant presheaves on a  $\mathcal{Q}$ -category  $\mathbb{A}$  forms a  $\mathcal{Q}$ -typed set<sup>9</sup>. We now need to enrich this set in  $\mathcal{Q}$  in such a way that the classifying property in (11) holds.

**6.1. PROPOSITION.** *For any  $\mathcal{Q}$ -category  $\mathbb{A}$ , the following defines the  $\mathcal{Q}$ -category  $\mathcal{P}\mathbb{A}$  of contravariant presheaves on  $\mathbb{A}$ :*

- *objects:*  $(\mathcal{P}\mathbb{A})_0 = \{\phi: *_C \multimap \mathbb{A} \text{ in } \text{Dist}(\mathcal{Q}) \mid C \in \mathcal{Q}\}$  with types  $t(\phi: *_C \multimap \mathbb{A}) = C$ ;
- *hom-arrows:* for  $\phi: *_C \multimap \mathbb{A}$  and  $\psi: *_C' \multimap \mathbb{A}$  put  $\mathcal{P}\mathbb{A}(\psi, \phi): C \rightarrow C'$  to be the single element of the distributor  $[\psi, \phi]: *_C \multimap *_C'$ .

*This  $\mathcal{Q}$ -category is skeletal, and it satisfies the equivalence in (11) as follows:*

- *for a distributor  $\Phi: \mathbb{C} \multimap \mathbb{A}$  the corresponding functor  $F_\Phi: \mathbb{C} \rightarrow \mathcal{P}\mathbb{A}$  maps  $c \in \mathbb{C}_0$  onto  $\Phi(-, c): tc \multimap \mathbb{A}$ ;*
- *and for a functor  $F: \mathbb{C} \rightarrow \mathcal{P}\mathbb{A}$  the corresponding distributor  $\Phi_F: \mathbb{C} \multimap \mathbb{A}$  has elements, for  $(c, a) \in \mathbb{C}_0 \times \mathbb{A}_0$ ,  $\Phi_F(a, c) = F(c)(a)$ .*

**PROOF.** With the properties of “[−, −]” in  $\text{Dist}(\mathcal{Q})$  it is easily verified that  $\mathcal{P}\mathbb{A}$  is a  $\mathcal{Q}$ -category. For  $\phi, \psi \in \mathcal{P}\mathbb{A}$ , observe that  $\phi \leq \psi$  in  $(\mathcal{P}\mathbb{A})_0$  if and only if  $t\phi = t\psi =: C$  and  $(1_C) \leq \mathcal{P}\mathbb{A}(\phi, \psi)$ , that is to say,  $\phi, \psi: *_C \multimap \mathbb{A}$  and  $1_C \leq [\phi, \psi]$  in  $\text{Dist}(\mathcal{Q})$ , thus precisely when  $\phi \leq \psi: *_C \multimap \mathbb{A}$ . This shows that the underlying order on the objects of  $\mathcal{P}\mathbb{A}$  (which are certain distributors) coincides with their order as arrows in  $\text{Dist}(\mathcal{Q})$ —so it is antisymmetric, i.e.  $\mathcal{P}\mathbb{A}$  is a skeletal  $\mathcal{Q}$ -category. As for the equivalence of orders in (11), it is easily seen that the mappings  $F \mapsto \Phi_F$  and  $\Phi \mapsto F_\Phi$  as in the statement of the proposition define a bijection:

<sup>9</sup>Would the objects of the base category  $\mathcal{Q}$  form a class, then the collection of contravariant presheaves on  $\mathbb{A}$  would be large! The reason – ultimately – is that  $\mathcal{P}\mathbb{A}$  is a cocomplete category—see 5.1 and 6.4.

- for any  $c \in \mathbb{C}_0$ ,  $F_{\Phi_F}(c) = \Phi_F(-, c) = F(c)$ , so  $F_{\Phi_F} = F$ ,
- for any  $(c, a) \in \mathbb{C}_0 \times \mathbb{A}_0$ ,  $\Phi_{F_\Phi}(a, c) = F_\Phi(c)(a) = \Phi(a, c)$ , so  $\Phi_{F_\Phi} = \Phi$ .

This bijection preserves order in both directions:

$$\begin{aligned}
& \Phi \leq \Psi \text{ in } \text{Dist}(\mathbb{C}, \mathbb{A}) \\
& \iff \forall (c, a) \in \mathbb{C}_0 \times \mathbb{A}_0 : \Phi(a, c) \leq \Psi(a, c) \text{ in } \mathcal{Q}(tc, ta) \\
& \iff \forall c \in \mathbb{C}_0 : F_\Phi(c) \leq F_\Psi(c) \text{ in } ((\mathcal{P}\mathbb{A})_0, \leq) \\
& \iff F_\Phi \leq F_\Psi \text{ in } \text{Cat}(\mathcal{Q})(\mathbb{C}, \mathcal{P}\mathbb{A})
\end{aligned}$$

and likewise

$$\begin{aligned}
& F \leq G \text{ in } \text{Cat}(\mathcal{Q})(\mathbb{C}, \mathcal{P}\mathbb{A}) \\
& \iff \forall c \in \mathbb{C}_0 : F(c) \leq G(c) \text{ in } ((\mathcal{P}\mathbb{A})_0, \leq) \\
& \iff \forall (c, a) \in \mathbb{C}_0 \times \mathbb{A}_0 : \Phi_F(a, c) \leq \Phi_G(a, c) \text{ in } \mathcal{Q}(tc, ta) \\
& \iff \Phi_F \leq \Phi_G \text{ in } \text{Dist}(\mathbb{C}, \mathbb{A}).
\end{aligned}$$

■

**6.2. EXAMPLE.** For a  $\mathcal{Q}$ -object  $Y$ , the  $\mathcal{Q}$ -category of contravariant presheaves on  $*_Y$  is denoted by  $\mathcal{P}Y$  instead of  $\mathcal{P}(*_Y)$ —see also 3.9.

Very often there will be an abuse of notation: for  $\phi: *_C \rightarrow \mathbb{A}$  and  $\psi: *_{C'} \rightarrow \mathbb{A}$  there is a *distributor*  $[\psi, \phi]: *_C \rightarrow *_{C'}$  with precisely one element (because it goes between one-object categories); therefore we defined  $\mathcal{P}\mathbb{A}(\psi, \phi)$  to be *the single element* of this distributor, for we want  $\mathcal{P}\mathbb{A}(\psi, \phi)$  to be a  $\mathcal{Q}$ -arrow. But more often than not we will simply write  $\mathcal{P}\mathbb{A}(\psi, \phi) = [\psi, \phi]$ , thus identifying a quantaloid  $\mathcal{Q}$  with the full subcategory of  $\text{Dist}(\mathcal{Q})$  determined by the one-object  $\mathcal{Q}$ -categories whose hom-arrows are identities (see also (9) and the appendix).

The equivalence of orders in (11) has a nice side-effect: because  $\mathcal{P}\mathbb{A}$  is skeletal,  $\text{Cat}(\mathcal{Q})(\mathbb{C}, \mathcal{P}\mathbb{A})$  is an antisymmetric order which is order-isomorphic to the sup-lattice  $\text{Dist}(\mathcal{Q})(\mathbb{C}, \mathbb{A})$ , thus it is a sup-lattice too. In particular are  $F \mapsto \Phi_F$  and  $\Phi \mapsto F_\Phi$  sup-morphisms.

To the identity distributor  $\mathbb{A}: \mathbb{A} \rightarrow \mathbb{A}$  corresponds, under the equivalence in (11), the functor

$$Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A} : a \mapsto \left( \mathbb{A}(-, a): *_a \rightarrow \mathbb{A} \right)$$

which is, of course, the *Yoneda embedding* for  $\mathbb{A}$ . The presheaf  $\mathbb{A}(-, a): *_a \rightarrow \mathbb{A}$  is said to be *represented* by  $a \in \mathbb{A}_0$ . We can now prove the “Yoneda lemma” for presheaves on a  $\mathcal{Q}$ -enriched category.

**6.3. PROPOSITION.** *For any  $\mathcal{Q}$ -category  $\mathbb{A}$ , any object  $a \in \mathbb{A}_0$  and any  $\phi: *_C \rightarrow \mathbb{A}$  in  $\text{Dist}(\mathcal{Q})$ ,  $\mathcal{P}\mathbb{A}(Y_{\mathbb{A}}a, \phi) = \phi(a)$ . As a consequence, the Yoneda embedding  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  is fully faithful.*

**PROOF.** Using that  $\mathbb{A}(-, a) \dashv \mathbb{A}(a, -)$  in  $\text{Dist}(\mathcal{Q})$ , we have by definition of  $\mathcal{P}\mathbb{A}(-, -)$  that  $\mathcal{P}\mathbb{A}(Y_{\mathbb{A}}a, \phi) = [\mathbb{A}(-, a), \phi] = \mathbb{A}(a, -) \otimes_{\mathbb{A}} \phi = \phi(a)$ . ■

An – almost – immediate but non the less important consequence of 6.1 is the following.

**6.4. PROPOSITION.** *The category  $\mathcal{P}\mathbb{B}$  of contravariant presheaves on a  $\mathcal{Q}$ -category  $\mathbb{B}$  is cocomplete.*

**PROOF.** Given a distributor  $\Theta: \mathbb{C} \multimap \mathbb{A}$  and a functor  $F: \mathbb{A} \rightarrow \mathcal{P}\mathbb{B}$ , to  $F$  corresponds a unique distributor  $\Phi_F: \mathbb{A} \multimap \mathbb{B}$  and, in turn, to  $\Phi_F \otimes_{\mathbb{A}} \Theta: \mathbb{C} \multimap \mathbb{B}$  corresponds a unique functor  $F_{\Phi_F \otimes_{\mathbb{A}} \Theta}: \mathbb{C} \rightarrow \mathcal{P}\mathbb{B}$ . We claim that this latter functor is the colimit of  $F$  weighted by  $\Theta$ : for any  $c \in \mathbb{C}$  and  $\beta \in \mathcal{P}\mathbb{B}$  we should verify that

$$\mathcal{P}\mathbb{B}\left(F_{\Phi_F \otimes_{\mathbb{A}} \Theta}(c), \beta\right) = \left[\Theta(-, c), \mathcal{P}\mathbb{B}(F-, \beta)\right].$$

The right hand side equals  $[\Theta(-, c), [F-, \beta]]$  and the left hand side equals

$$\left[(\Phi_F \otimes_{\mathbb{A}} \Theta)(-, c), \beta\right] = \left[\Phi_F \otimes_{\mathbb{A}} \Theta(-, c), \beta\right] = \left[\Theta(-, c), [\Phi_F, \beta]\right].$$

Since  $[F-, \beta] = [\Phi_F, \beta]$ , the required universality holds.  $\blacksquare$

**6.5. COROLLARY.** *For any  $\phi: *_C \multimap \mathbb{A}$ , the  $\phi$ -weighted colimit of  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  (exists and) is  $\phi$  itself.*

In other words, every contravariant presheaf on a  $\mathcal{Q}$ -category  $\mathbb{A}$  is canonically the colimit of representable presheaves:  $\text{colim}(\phi, Y_{\mathbb{A}}) = \phi$  (and we may write an equality instead of merely an isomorphism, for  $\mathcal{P}\mathbb{A}$  is skeletal).

Here is another result that should sound familiar.

**6.6. PROPOSITION.** *The Yoneda embedding  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  is continuous.*

**PROOF.** Consider a weighted limit (which happens to exist) in  $\mathbb{A}$ , like so:

$$\begin{array}{ccccc} & \mathbb{C} & & & \\ & \uparrow & \text{lim}(\Phi, F) & & \\ \Phi \circ & & \swarrow & & \\ \mathbb{B} & \xrightarrow{F} & \mathbb{A} & \xrightarrow{Y_{\mathbb{A}}} & \mathcal{P}\mathbb{A} \end{array}$$

Then, for any  $c \in \mathbb{C}_0$ ,

$$\left(Y_{\mathbb{A}} \circ \text{lim}(\Phi, F)\right)(c) = \mathbb{A}\left(-, \text{lim}(\Phi, F)(c)\right) = \left\{\Phi(c, -), \mathbb{A}(-, F-)\right\}$$

and therefore

$$\begin{aligned} \mathcal{P}\mathbb{A}(-, Y_{\mathbb{A}} \circ \text{lim}(\Phi, F)-) &= \left[-, \left\{\Phi, \mathbb{A}(-, F-)\right\}\right] \\ &= \left\{\Phi, \left[-, \mathbb{A}(-, F-)\right]\right\} \\ &= \left\{\Phi, \mathcal{P}\mathbb{A}(-, Y_{\mathbb{A}} \circ F-)\right\} \end{aligned}$$

so  $Y_{\mathbb{A}} \circ \text{lim}(\Phi, F)$  is  $\text{lim}(\Phi, Y_{\mathbb{A}} \circ F)$ .  $\blacksquare$

When  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  is a right adjoint, then  $\mathbb{A}$  is cocomplete (combine 5.13 with 6.3 and 6.4). Also the converse can be shown; actually,  $\mathcal{P}\mathbb{A}$  is the free cocompletion of  $\mathbb{A}$  in  $\text{Cat}(\mathcal{Q})$ . To make this precise in an elegant way, it is useful to treat pointwise left Kan extensions first.

### POINTWISE LEFT KAN EXTENSIONS

If for  $F: \mathbb{B} \rightarrow \mathbb{A}$  and  $G: \mathbb{B} \rightarrow \mathbb{C}$  in  $\text{Cat}(\mathcal{Q})$  the  $\mathbb{C}(G-, -)$ -weighted colimit of  $F$  exist, then it is the left Kan extension of  $F$  along  $G$ ; so  $\langle F, G \rangle \cong \text{colim}(\mathbb{C}(G-, -), F)$  whenever the right hand side exists. By lemma 5.2 we know that in this case the action of  $\langle F, G \rangle: \mathbb{C} \rightarrow \mathbb{A}$  on objects is given by the colimit formula  $\langle F, G \rangle(c) \cong \text{colim}(\mathbb{C}(G-, c), F)$ . Such a colimit weighted by the right adjoint distributor determined by some functor is therefore said to be a *pointwise left Kan extension*. (Note that, for a pointwise left Kan extension, the two notions of absoluteness, one relevant for Kan extensions and the other relevant for colimits, coincide.)

A  $\mathcal{Q}$ -category is said to *admit all pointwise left Kan extensions* when it admits all colimits of this special kind. A cocomplete  $\mathcal{Q}$ -category then certainly admits all pointwise left Kan extensions. (For a cocomplete  $\mathbb{B}$ , any functor  $G: \mathbb{A} \rightarrow \mathbb{C}$  thus determines an adjunction

$$\langle -, G \rangle \dashv (- \circ G): \text{Cat}(\mathcal{Q})(\mathbb{C}, \mathbb{B}) \rightleftarrows \text{Cat}(\mathcal{Q})(\mathbb{A}, \mathbb{B}).$$

This has been called the “meta-adjointness” associated to the concept of Kan extension.) But it is also true that a  $\mathcal{Q}$ -category admitting all pointwise Kan extensions, is cocomplete! This follows from 5.4 in combination with 6.9 below; note that the Yoneda embedding plays a crucial rôle.

**6.7. PROPOSITION.** *Suppose that for  $F: \mathbb{B} \rightarrow \mathbb{A}$  and  $G: \mathbb{B} \rightarrow \mathbb{C}$  in  $\text{Cat}(\mathcal{Q})$  the pointwise left Kan extension of  $F$  along  $G$  exists. If  $G$  is fully faithful, then  $\langle F, G \rangle \circ G \cong F$ .*

**PROOF.** It is easily seen that,  $\langle F, G \rangle$  being a pointwise left Kan extension,

$$\begin{aligned} \langle F, G \rangle \circ G &\cong F \\ \iff \mathbb{B}(\langle F, G \rangle \circ G-, -) &= \mathbb{B}(F-, -) \\ \iff \mathbb{C}(G-, -) \otimes_{\mathbb{C}} \mathbb{B}(\langle F, G \rangle-, -) &= \mathbb{B}(F-, -) \\ \iff \mathbb{C}(G-, -) \otimes_{\mathbb{C}} \left[ \mathbb{C}(G-, -), \mathbb{B}(F-, -) \right] &= \mathbb{B}(F-, -) \\ \iff \mathbb{C}(G-, -) \otimes_{\mathbb{C}} \mathbb{C}(-, G-) \otimes_{\mathbb{A}} \mathbb{B}(F-, -) &= \mathbb{B}(F-, -) \\ \iff \mathbb{C}(G-, G-) \otimes_{\mathbb{A}} \mathbb{B}(F-, -) &= \mathbb{B}(F-, -) \end{aligned}$$

so if  $G$  is fully faithful, i.e.  $\mathbb{C}(G-, G-) = \mathbb{A}$ , then the above holds. ■

**6.8. PROPOSITION.** *Let  $\mathbb{A}$  be a cocomplete  $\mathcal{Q}$ -category; then the cocontinuous functors with domain  $\mathbb{A}$  coincide with the left adjoints.*

PROOF. One direction is a consequence of 5.12. To prove the other, let  $F: \mathbb{A} \rightarrow \mathbb{B}$  be a cocontinuous functor. By cocompleteness of  $\mathbb{A}$ ,  $\langle 1_{\mathbb{A}}, F \rangle$  not only exists but is even pointwise, i.e. it is a colimit, so it must be preserved by  $F$ . By 4.6 we conclude that  $F$  is a left adjoint. ■

6.9. PROPOSITION. *A functor  $F: \mathbb{A} \rightarrow \mathbb{B}$  has a pointwise left Kan extension along  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  if and only if, for every  $\phi \in \mathcal{P}\mathbb{A}$ , the  $\phi$ -weighted colimit of  $F$  exists. In this case,  $\langle F, Y_{\mathbb{A}} \rangle: \mathcal{P}\mathbb{A} \rightarrow \mathbb{B}$  is left adjoint to  $\langle Y_{\mathbb{A}}, F \rangle: \mathbb{B} \rightarrow \mathcal{P}\mathbb{A}$ , and  $\langle F, Y_{\mathbb{A}} \rangle \circ Y_{\mathbb{A}} \cong F$ . Actually,  $\langle F, Y_{\mathbb{A}} \rangle$  is then the essentially unique cocontinuous factorization of  $F$  through  $Y_{\mathbb{A}}$ .*

PROOF. If by hypothesis,  $\langle F, Y_{\mathbb{A}} \rangle = \text{colim}(\mathcal{P}\mathbb{A}(Y_{\mathbb{A}}-, -), F)$  exists, then  $\langle F, Y_{\mathbb{A}} \rangle(\phi) = \text{colim}(\mathcal{P}\mathbb{A}(Y_{\mathbb{A}}-, \phi), F) = \text{colim}(\phi, F)$  because  $\mathcal{P}\mathbb{A}(Y_{\mathbb{A}}-, \phi) = \phi$ . Conversely, supposing that  $\mathbb{B}$  allows colimits of  $F$  weighted by presheaves on  $\mathbb{A}$ ,  $\text{colim}(\mathcal{P}\mathbb{A}(Y_{\mathbb{A}}-, -), F)$  exists because, for all  $\phi = \mathcal{P}\mathbb{A}(Y_{\mathbb{A}}-, \phi)$ ,  $\text{colim}(\mathcal{P}\mathbb{A}(Y_{\mathbb{A}}-, \phi), F)$  exists (see 5.2).

Since  $\mathcal{P}\mathbb{A}$  is cocomplete,  $\langle Y_{\mathbb{A}}, F \rangle$  exists, is pointwise, and can thus be computed as  $\langle Y_{\mathbb{A}}, F \rangle(b) \cong \text{colim}(\mathbb{B}(F-, b), Y_{\mathbb{A}}) \cong \mathbb{B}(F-, b)$ . It is then a matter of applying 4.2 to prove the adjunction:

$$\begin{aligned} & \langle F, Y_{\mathbb{A}} \rangle \dashv \langle Y_{\mathbb{A}}, F \rangle \\ \iff & \forall \phi \in \mathcal{P}\mathbb{A}, \forall b \in \mathbb{B} : \mathbb{B} \left( \langle F, Y_{\mathbb{A}} \rangle(\phi), b \right) = \mathcal{P}\mathbb{A} \left( \phi, \langle Y_{\mathbb{A}}, F \rangle(b) \right) \\ \iff & \forall \phi \in \mathcal{P}\mathbb{A}, \forall b \in \mathbb{B} : \mathbb{B} \left( \text{colim}(\phi, F), b \right) = \left[ \phi, \mathbb{B}(F-, b) \right] \end{aligned}$$

which holds by the definition of colimit.

Finally, by fully faithfulness of  $Y_{\mathbb{A}}$  and 6.7 it follows that  $F$  factors through  $Y_{\mathbb{A}}$  as  $\langle F, Y_{\mathbb{A}} \rangle$ . Would now  $G: \mathcal{P}\mathbb{A} \rightarrow \mathbb{B}$  be another cocontinuous functor such that  $F \cong G \circ Y_{\mathbb{A}}$  then – by cocontinuity and using the fact that any presheaf  $\phi \in \mathcal{P}\mathbb{A}$  is canonically the colimit of representables –  $G$  and  $\langle F, Y_{\mathbb{A}} \rangle$  agree on all objects of  $\mathcal{P}\mathbb{A}$ :

$$\begin{aligned} G(\phi) &= G(\text{colim}(\phi, Y_{\mathbb{A}})) \\ &\cong \text{colim}(\phi, G \circ Y_{\mathbb{A}}) \\ &\cong \text{colim}(\phi, F) \\ &= \langle F, Y_{\mathbb{A}} \rangle(\phi). \end{aligned}$$

■

6.10. COROLLARY. *A  $\mathcal{Q}$ -category  $\mathbb{A}$  is cocomplete if and only if  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  is a right adjoint in  $\text{Cat}(\mathcal{Q})$ .*

PROOF. If  $Y_{\mathbb{A}}$  is right adjoint, apply 5.13 to  $Y_{\mathbb{A}}$ , using 6.3 and 6.4. If  $\mathbb{A}$  is cocomplete, apply 6.9 to the identity functor  $1_{\mathbb{A}}: \mathbb{A} \rightarrow \mathbb{A}$ :  $\langle 1_{\mathbb{A}}, Y_{\mathbb{A}} \rangle \dashv \langle Y_{\mathbb{A}}, 1_{\mathbb{A}} \rangle = Y_{\mathbb{A}}$ . ■

## FREE COCOMPLETION

Applying 6.9 we easily obtain that, for two  $\mathcal{Q}$ -categories  $\mathbb{A}$  and  $\mathbb{B}$ , with  $\mathbb{B}$  cocomplete, any functor  $F: \mathbb{A} \rightarrow \mathbb{B}$  factors through  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  by the essentially unique cocontinuous – even left adjoint – functor  $\langle F, Y_{\mathbb{A}} \rangle: \mathcal{P}\mathbb{A} \rightarrow \mathbb{B}$ . Write now  $\mathbf{Cocont}(\mathcal{Q})$  for the sub-2-category of  $\mathbf{Cat}(\mathcal{Q})$  whose objects are the cocomplete  $\mathcal{Q}$ -categories, and whose morphisms are the cocontinuous functors, or equivalently (by 6.8) the left adjoint ones; there is a forgetful 2-functor  $\mathcal{U}: \mathbf{Cocont}(\mathcal{Q}) \rightarrow \mathbf{Cat}(\mathcal{Q})$ . The above then says that, for  $\mathcal{Q}$ -categories  $\mathbb{A}$  and  $\mathbb{B}$ , with  $\mathbb{B}$  cocomplete, the Yoneda embedding  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  induces a natural equivalence of orders

$$- \circ Y_{\mathbb{A}}: \mathbf{Cocont}(\mathcal{Q})(\mathcal{P}\mathbb{A}, \mathbb{B}) \xrightarrow{\sim} \mathbf{Cat}(\mathcal{Q})(\mathbb{A}, \mathbb{B})$$

with inverse  $F \mapsto \langle F, Y_{\mathbb{A}} \rangle$ .

**6.11. PROPOSITION.** *The presheaf construction provides a left (bi)adjoint to the inclusion of  $\mathbf{Cocont}(\mathcal{Q})$  in  $\mathbf{Cat}(\mathcal{Q})$ :  $\mathcal{P} \dashv \mathcal{U}: \mathbf{Cocont}(\mathcal{Q}) \rightleftarrows \mathbf{Cat}(\mathcal{Q})$ .*

For any two  $\mathcal{Q}$ -categories  $\mathbb{A}$  and  $\mathbb{B}$ , we have – by 6.1 and the above – equivalences

$$\mathbf{Dist}(\mathcal{Q})(\mathbb{A}, \mathbb{B}) \xrightarrow{\sim} \mathbf{Cat}(\mathcal{Q})(\mathbb{A}, \mathcal{P}\mathbb{B}) \xrightarrow{\sim} \mathbf{Cocont}(\mathcal{Q})(\mathcal{P}\mathbb{A}, \mathcal{P}\mathbb{B}) \quad (12)$$

sending a  $\Phi: \mathbb{A} \rightarrow \mathbb{B}$  to  $\langle F_{\Phi}, Y_{\mathbb{A}} \rangle: \mathcal{P}\mathbb{B} \rightarrow \mathcal{P}\mathbb{A}$  (with  $F_{\Phi}: \mathbb{B} \rightarrow \mathcal{P}\mathbb{A}: b \mapsto \Phi(-, b)$  as in 6.1). We can prove more.

**6.12. PROPOSITION.** *The locally ordered<sup>10</sup> category  $\mathbf{Cocont}(\mathcal{Q})$  has stable local colimits, and the action*

$$\mathbf{Dist}(\mathcal{Q}) \rightarrow \mathbf{Cocont}(\mathcal{Q}): \left( \Phi: \mathbb{A} \rightarrow \mathbb{B} \right) \mapsto \left( \langle F_{\Phi}, Y_{\mathbb{A}} \rangle: \mathcal{P}\mathbb{A} \rightarrow \mathcal{P}\mathbb{B} \right)$$

*is a 2-functor which is locally an equivalence.*

**PROOF.** We already know that  $\mathbf{Cocont}(\mathcal{Q})$  is a 2-category; we must show that it has stable local colimits. By 5.6 we know how to compute suprema in  $\mathbf{Cat}(\mathcal{Q})(\mathbb{A}, \mathbb{B})$  whenever  $\mathbb{B}$  is cocomplete; we may thus apply this to given arrows  $(F_i: \mathbb{A} \rightarrow \mathbb{B})_{i \in I}$  in  $\mathbf{Cocont}(\mathcal{Q})$ :

$$\bigvee_{i \in I} F_i := \text{colim} \left( \bigvee_{i \in I} \mathbb{B}(-, F_i -), 1_{\mathbb{B}} \right).$$

But we must show that  $\bigvee_i F_i$  is still a cocontinuous functor (for *a priori* it is merely a functor); equivalently (by cocompleteness of  $\mathbb{A}$ ), we may assume that each  $F_i$  is a left adjoint in  $\mathbf{Cat}(\mathcal{Q})$ , and show that  $\bigvee_i F_i$  itself is a left adjoint in  $\mathbf{Cat}(\mathcal{Q})$ . Let us write  $F_i \dashv G_i$  for such adjunctions; then, by 5.6 and its dual<sup>11</sup>,

$$\mathbb{B} \left( \bigvee_i F_i -, - \right) = \bigwedge_i \mathbb{B}(F_i -, -) = \bigwedge_i \mathbb{A}(-, G_i -) = \mathbb{A}(-, \bigwedge_i G_i -).$$

<sup>10</sup>... but not antisymmetrically so!

<sup>11</sup>The dual of 5.6 says the following: if  $\mathbb{A}$  is a complete category, then the infimum of functors  $(G_i: \mathbb{B} \rightarrow \mathbb{A})_{i \in I}$  exists in  $\mathbf{Cat}(\mathcal{Q})(\mathbb{B}, \mathbb{A})$ , and is given by the  $\bigvee_i \mathbb{A}(G_i -, -)$ -weighted limit of  $1_{\mathbb{A}}$ . This means exactly that this infimum  $\bigwedge_i G_i$  satisfies  $\mathbb{A}(-, \bigwedge_i G_i -) = \bigwedge_i \mathbb{A}(-, G_i -)$ . By 5.10 any complete category is also cocomplete, so this applies in this case!

That is to say,  $\bigvee_i F_i \dashv \bigwedge_i G_i$  in  $\text{Cat}(\mathcal{Q})$ . As for the stability of these local colimits, take for example  $(F_i: \mathbb{A} \rightarrow \mathbb{B})_{i \in I}$  and  $G: \mathbb{B} \rightarrow \mathbb{C}$  in  $\text{Cocont}(\mathcal{Q})$ ; then

$$\begin{aligned}
 G \circ \left( \bigvee_i F_i \right) &= G \circ \text{colim} \left( \bigvee_i \mathbb{B}(-, F_i -), 1_{\mathbb{B}} \right) \\
 &\cong \text{colim} \left( \bigvee_i \mathbb{B}(-, F_i -), G \right) \\
 &\cong \text{colim} \left( \mathbb{C}(-, G-) \otimes_{\mathbb{B}} \bigvee_i (-, F_i -), 1_{\mathbb{C}} \right) \\
 &\cong \text{colim} \left( \bigvee_i (-, G \circ F_i -), 1_{\mathbb{C}} \right) \\
 &= \bigvee_i (G \circ F_i)
 \end{aligned}$$

because  $G$  is cocontinuous. Now consider  $F: \mathbb{A} \rightarrow \mathbb{B}$  and  $(G_i: \mathbb{B} \rightarrow \mathbb{C})_{i \in I}$ , then for any  $a \in \mathbb{A}_0$

$$\begin{aligned}
 \left( \left( \bigvee_i G_i \right) \circ F \right) (a) &= \text{colim} \left( \bigvee_i \mathbb{C}(-, G_i -), 1_{\mathbb{C}} \right) (Fa) \\
 &\cong \text{colim} \left( \bigvee_i \mathbb{C}(-, G_i -) \otimes_{\mathbb{B}} \mathbb{B}(-, Fa), 1_{\mathbb{C}} \right) \\
 &\cong \text{colim} \left( \bigvee_i \mathbb{C}(-, G_i(Fa)), 1_{\mathbb{C}} \right) \\
 &\cong \text{colim} \left( \bigvee_i \mathbb{C}(-, G_i \circ F -), 1_{\mathbb{C}} \right) (a) \\
 &= \left( \bigvee_i (G_i \circ F) \right) (a)
 \end{aligned}$$

by properties of the colimit – see 5.2 – so  $(\bigvee_i G_i) \circ F = \bigvee_i (G_i \circ F)$ .

To prove the functoriality of the action described in the statement of the proposition, observe that for  $\Phi: \mathbb{A} \dashv\vdash \mathbb{B}$  and  $\theta \in \mathcal{P}\mathbb{A}$ ,

$$\begin{aligned}
 \langle F_{\Phi}, Y_{\mathbb{A}} \rangle (\theta) &= \text{colim} \left( \mathcal{P}\mathbb{A}(\theta, Y_{\mathbb{A}} -), F_{\Phi} \right) \\
 &= \text{colim}(\theta, F_{\Phi}) \\
 &= \Phi \otimes_{\mathbb{A}} \theta.
 \end{aligned}$$

In the first line we use the cocompleteness of  $\mathcal{P}\mathbb{B}$  to compute the (pointwise) left Kan extension as a colimit; to pass from the first line to the second we use the Yoneda lemma 6.3; the third line follows from the way in which weighted colimits are calculated in a presheaf category, cf. the proof of 6.4. Using this formula it is clear that  $\Phi \mapsto \langle F_{\Phi}, Y_{\mathbb{A}} \rangle$  is functorial. That this functor is really a 2-functor, and that it is locally an equivalence, follows from the preceding remarks, in particular (12).  $\blacksquare$

Restricting to skeletal cocomplete  $\mathcal{Q}$ -categories,  $\mathbf{Cocont}_{\text{skel}}(\mathcal{Q})$  is a quantaloid and the 2-functor above corestricts to a quantaloid homomorphism  $\mathbf{Dist}(\mathcal{Q}) \rightarrow \mathbf{Cocont}_{\text{skel}}(\mathcal{Q})$  that induces sup-isomorphisms between the hom-sup-lattices.

6.13. **EXAMPLE.** For a  $\mathbf{2}$ -category  $\mathbb{A}$ , i.e. an ordered set,  $\mathcal{P}\mathbb{A}$  is set of down-sets of  $\mathbb{A}$ , ordered by inclusion. The Yoneda embedding  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  sends an element to the principal down-set that it determines. An eventual left adjoint to  $Y_{\mathbb{A}}$  is “taking the supremum”. The isomorphism  $\mathbf{Dist}(\mathbf{2})(\mathbb{A}, \mathbb{B}) \cong \mathbf{Cocont}_{\text{skel}}(\mathbf{2})(\mathcal{P}\mathbb{A}, \mathcal{P}\mathbb{B})$  says that an ideal relation between given orders is the same thing as a sup-morphism between the sup-lattices of down-sets of those orders. In fact,  $\mathbf{Cocont}_{\text{skel}}(\mathbf{2}) = \mathbf{Sup}$ .

#### COVARIANT PRESHEAVES, FREE COMPLETION

The results in the previous sections may be dualized; one obtains that “the  $\mathcal{Q}$ -category of covariant presheaves on  $\mathbb{A}$  is  $\mathbb{A}$ ’s free completion in  $\mathbf{Cat}(\mathcal{Q})$ ”. Let us quickly do this exercise.

A *covariant presheaf of type*  $C \in \mathcal{Q}$  on a  $\mathcal{Q}$ -category  $\mathbb{A}$  is a distributor  $\phi: \mathbb{A} \rightarrow *C$ . Covariant presheaves form the object-set of a  $\mathcal{Q}$ -category  $\mathcal{P}^{\dagger}\mathbb{A}$ , whose hom-arrows are, for  $\phi: \mathbb{A} \rightarrow *C$  and  $\psi: \mathbb{A} \rightarrow *C'$ ,

$$\mathcal{P}^{\dagger}\mathbb{A}(\psi, \phi) = \text{single element of } \{\phi, \psi\} \text{ in } \mathbf{Dist}(\mathcal{Q}).$$

This  $\mathcal{Q}$ -category satisfies

$$\text{for every } \mathcal{Q}\text{-category } \mathbb{C}, \mathbf{Dist}(\mathcal{Q})(\mathbb{A}, \mathbb{C}) \simeq \mathbf{Cat}(\mathcal{Q})^{\text{co}}(\mathbb{C}, \mathcal{P}^{\dagger}\mathbb{A}), \quad (13)$$

and it is complete. There is a Yoneda embedding  $Y_{\mathbb{A}}^{\dagger}: \mathbb{A} \rightarrow \mathcal{P}^{\dagger}\mathbb{A}$ , sending an object  $a \in \mathbb{A}$  to the representable distributor  $\mathbb{A}(a, -): \mathbb{A} \rightarrow *ta$ , which is fully faithful.  $\mathbb{A}$  is complete if and only if  $Y_{\mathbb{A}}^{\dagger}$  is a left adjoint in  $\mathbf{Cat}(\mathcal{Q})$ ; but in any case is  $Y_{\mathbb{A}}^{\dagger}$  cocontinuous.

Continuous functors whose domain is a complete  $\mathcal{Q}$ -category, coincide with the right adjoints. Denoting  $\mathbf{Cont}(\mathcal{Q})$  for the sub-2-category of  $\mathbf{Cat}(\mathcal{Q})$  whose objects are complete  $\mathcal{Q}$ -categories and arrows are continuous functors, we have

$$\mathcal{P}^{\dagger} \dashv \mathcal{U}: \mathbf{Cont}(\mathcal{Q}) \xleftrightarrow{\quad} \mathbf{Cat}(\mathcal{Q});$$

that is,  $\mathcal{P}^{\dagger}\mathbb{A}$  is the free completion of  $\mathbb{A}$  in  $\mathbf{Cat}(\mathcal{Q})$ . Furthermore,  $\mathbf{Cont}(\mathcal{Q})$  has stable local limits, and there is a 2-functor  $\mathbf{Dist}(\mathcal{Q}) \rightarrow \mathbf{Cont}(\mathcal{Q})^{\text{coop}}$ , which is locally an equivalence, sending a distributor  $\Phi: \mathbb{A} \rightarrow \mathbb{B}$  to the right Kan extension  $(F^{\Phi}, Y_{\mathbb{A}}^{\dagger}): \mathcal{P}^{\dagger}\mathbb{B} \rightarrow \mathcal{P}^{\dagger}\mathbb{A}$ , where  $F^{\Phi}: \mathbb{B} \rightarrow \mathcal{P}^{\dagger}\mathbb{A}$  is determined by  $\Phi$  through the equivalence in (13).

Since a  $\mathcal{Q}$ -category is complete if and only if it is cocomplete, and since between such categories continuous (cocontinuous) functors coincide with right adjoints (left adjoints), there is an isomorphism of (locally cocompletely ordered) 2-categories

$$\mathbf{Cocont}(\mathcal{Q}) \xrightarrow{\sim} \mathbf{Cont}(\mathcal{Q})^{\text{coop}}: \left( F: \mathbb{A} \rightarrow \mathbb{B} \right) \longleftrightarrow \left( F^*: \mathbb{B} \rightarrow \mathbb{A} \right)$$

where  $F \dashv F^*$  in  $\text{Cat}(\mathcal{Q})$ . Considering only skeletal objects, this restricts to an isomorphism of quantaloids  $\text{Cocont}_{\text{skel}}(\mathcal{Q}) \cong \text{Cont}_{\text{skel}}(\mathcal{Q})^{\text{coop}}$ .

Assembling previous results we obtain that: for a  $\mathcal{Q}$ -category  $\mathbb{A}$ ,  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  is a right adjoint if and only if  $Y_{\mathbb{A}}^{\dagger}: \mathbb{A} \rightarrow \mathcal{P}^{\dagger}\mathbb{A}$  is a left adjoint; this is precisely the case when  $\mathbb{A}$  is (co)complete. It is interesting to take a closer look at this result. First observe that, in any case, there is an adjunction

$$\begin{array}{ccc} & [-, \mathbb{A}] & \\ & \curvearrowright & \\ \mathcal{P}\mathbb{A} & \perp & \mathcal{P}^{\dagger}\mathbb{A} \\ & \curvearrowleft & \\ & \{-, \mathbb{A}\} & \end{array}$$

in  $\text{Cat}(\mathcal{Q})$ , where the functors involved are obtained by calculating  $\{-, -\}$  and  $[-, -]$  in  $\text{Dist}(\mathcal{Q})$ :

$$\begin{aligned} [-, \mathbb{A}]: \mathcal{P}\mathbb{A} &\rightarrow \mathcal{P}^{\dagger}\mathbb{A}: \left( \phi: *_A \multimap \mathbb{A} \right) \mapsto \left( [\phi, \mathbb{A}]: \mathbb{A} \multimap *_A \right), \\ \{-, \mathbb{A}\}: \mathcal{P}^{\dagger}\mathbb{A} &\rightarrow \mathcal{P}\mathbb{A}: \left( \psi: \mathbb{A} \multimap *_A \right) \mapsto \left( \{\psi, \mathbb{A}\}: *_A \multimap \mathbb{A} \right). \end{aligned}$$

Now considering the diagram of  $\mathcal{Q}$ -categories and functors

$$\begin{array}{ccc} & \mathbb{A} & \\ & \swarrow \quad \searrow & \\ R & & L \\ & \swarrow \quad \searrow & \\ & \mathcal{P}^{\dagger}\mathbb{A} & \mathcal{P}\mathbb{A} \\ & \longleftarrow \quad \longrightarrow & \\ & [-, \mathbb{A}] & \end{array}$$

$Y_{\mathbb{A}}^{\dagger}$  (left arrow from  $\mathbb{A}$  to  $\mathcal{P}^{\dagger}\mathbb{A}$ ),  $Y_{\mathbb{A}}$  (right arrow from  $\mathbb{A}$  to  $\mathcal{P}\mathbb{A}$ ),  $\{-, \mathbb{A}\}$  (left arrow from  $\mathcal{P}^{\dagger}\mathbb{A}$  to  $\mathcal{P}\mathbb{A}$ ),  $[-, \mathbb{A}]$  (right arrow from  $\mathcal{P}\mathbb{A}$  to  $\mathcal{P}^{\dagger}\mathbb{A}$ ).

we have on the one hand that, if  $L \dashv Y_{\mathbb{A}}$  is known, then  $R := L \circ \{-, \mathbb{A}\}$  is right adjoint to  $Y_{\mathbb{A}}^{\dagger}$ ; and dually, if  $Y_{\mathbb{A}}^{\dagger} \dashv R$  is known, then  $L := R \circ [-, \mathbb{A}]$  is left adjoint to  $Y_{\mathbb{A}}$ . Actually,  $L(\phi) = \text{colim}(\phi, 1_{\mathbb{A}})$  and  $R(\psi) = \text{lim}(\psi, 1_{\mathbb{A}})$  (whenever one, thus also the other, exists), so that the *crux* of this construction is really the statement in 5.8.

**6.14. EXAMPLE.** Take  $\mathbf{2}$ -categories, then  $\text{Cont}(\mathbf{2}) \simeq \text{Inf} = \text{Cont}_{\text{skel}}(\mathbf{2})$ . For a  $\mathbf{2}$ -category  $\mathbb{A}$ ,  $\mathcal{P}^{\dagger}\mathbb{A}$  is the set of up-sets of  $\mathbb{A}$ , ordered by containment (reverse inclusion);  $Y_{\mathbb{A}}^{\dagger}$  sends an element of  $\mathbb{A}$  to the corresponding principal up-set. An eventual right adjoint to  $Y_{\mathbb{A}}^{\dagger}$  is “infimum”. It is well-known that  $\text{Sup} \cong \text{Inf}^{\text{coop}}$  in **QUANT** by passing from left-adjoints to right-adjoints in  $\text{Cat}_{\text{skel}}(\mathbf{2}) = \text{Ord}$ . The adjunction between  $\mathcal{P}\mathbb{A}$  (down-sets of  $\mathbb{A}$ , ordered by inclusion) and  $\mathcal{P}^{\dagger}\mathbb{A}$  (up-sets of  $\mathbb{A}$ , ordered by containment) is constituted by the left adjoint that sends a down-set to the up-set of its upper bounds, and the right adjoint that sends an up-set to the down-set of its lower bounds.

**6.15. EXAMPLE.** For a  $\mathcal{Q}$ -object  $X$ , the category of covariant presheaves on  $*_X$  is denoted by  $\mathcal{P}^{\dagger}X$  rather than  $\mathcal{P}^{\dagger}(*_X)$  (cf. 3.9).

## 7. Cauchy completion and Morita equivalence

### CAUCHY COMPLETE CATEGORIES

A left adjoint distributor  $\Phi: \mathbb{C} \dashv\vdash \mathbb{A}$  is often called a *Cauchy distributor* (into  $\mathbb{A}$ ); we will systematically denote its right adjoint by  $\Phi^*$ :

$$\begin{array}{ccc} & \Phi & \\ & \circlearrowleft & \\ \mathbb{A} & \xrightarrow{\quad} & \mathbb{B} \\ & \circlearrowright & \\ & \Phi^* & \end{array}$$

In particular, a *Cauchy presheaf*  $\phi: *_C \dashv\vdash \mathbb{A}$  is a contravariant presheaf with a right adjoint  $\phi^*: \mathbb{A} \dashv\vdash *_C$ , whose elements are  $\phi^*(a): ta \rightarrow C$ .

A Cauchy distributor  $\Phi: \mathbb{C} \dashv\vdash \mathbb{A}$  is said to *converge* (to a functor  $F: \mathbb{C} \rightarrow \mathbb{A}$ ) if (there exists a functor  $F: \mathbb{C} \rightarrow \mathbb{A}$  such that)  $\Phi = \mathbb{B}(-, F-)$ —or equivalently,  $\Phi^* = \mathbb{B}(F-, -)$ . A Cauchy distributor  $\Phi: \mathbb{C} \dashv\vdash \mathbb{A}$  need not converge, but if it does then the functor  $F: \mathbb{C} \rightarrow \mathbb{A}$  that it converges to, is essentially unique. We are interested in those  $\mathcal{Q}$ -categories in which “all Cauchy distributors converge”: say that  $\mathbb{A}$  is *Cauchy complete* if, for all  $\mathbb{C}$ , any Cauchy distributor  $\Phi: \mathbb{C} \dashv\vdash \mathbb{A}$  converges. In other words, letting  $\text{Map}(\text{Dist}(\mathcal{Q}))(\mathbb{C}, \mathbb{A})$  stand for the full subcategory of  $\text{Dist}(\mathcal{Q})(\mathbb{C}, \mathbb{A})$  whose objects are the Cauchy distributors (“maps”),  $\mathbb{A}$  is Cauchy complete if and only if, for every  $\mathbb{C}$ , the mapping

$$\text{Cat}(\mathcal{Q})(\mathbb{C}, \mathbb{A}) \rightarrow \text{Map}(\text{Dist}(\mathcal{Q}))(\mathbb{C}, \mathbb{A}): F \mapsto \mathbb{A}(-, F-)$$

is surjective; recalling that it is always “essentially injective”, we thus ask for it to be an equivalence.

Fortunately, to have Cauchy completeness, convergence of the Cauchy presheaves suffices!

**7.1. PROPOSITION.** *A  $\mathcal{Q}$ -category  $\mathbb{A}$  is Cauchy complete if and only if all Cauchy presheaves on  $\mathbb{A}$  converge (i.e. are representable).*

**PROOF.** We only need to prove the “if”. Let  $\Phi \dashv\vdash \Phi^*: \mathbb{A} \overset{\leftarrow}{\otimes} \mathbb{C}$ , then for each  $c \in \mathbb{C}_0$ ,  $\Phi(-, c) \dashv\vdash \Phi^*(c, -): \mathbb{A} \overset{\leftarrow}{\otimes} \mathbb{C}$ , because  $\Phi \otimes_{\mathbb{C}} \mathbb{C}(-, c) = \Phi(-, c)$  and  $\mathbb{C}(c, -) \otimes_{\mathbb{C}} \Phi^* = \Phi^*(c, -)$ , and by composition of adjoints. By assumption there exists for each  $c \in \mathbb{C}_0$  a functor  $Fc: *_c \rightarrow \mathbb{A}$  — that we will henceforth identify with the single element of type  $tc$  that it picks out — such that  $\mathbb{A}(-, Fc) = \Phi(-, c)$  or equivalently,  $\mathbb{A}(Fc, -) = \Phi^*(c, -)$ . But the object mapping  $\mathbb{C}_0 \rightarrow \mathbb{A}_0: c \mapsto Fc$  is functorial: since  $\mathbb{C} \leq \Phi^* \otimes_{\mathbb{A}} \Phi$  it follows that

$$\mathbb{C}(c', c) \leq \Phi^*(c', -) \otimes_{\mathbb{A}} \Phi(-, c) = \mathbb{A}(Fc', -) \otimes_{\mathbb{A}} \mathbb{A}(-, Fc) = \mathbb{A}(Fc', Fc).$$

Clearly  $\Phi \dashv\vdash \Phi^*$  converges to the functor  $F: \mathbb{C} \rightarrow \mathbb{A}: c \mapsto Fc$ . ■

The next proposition explains Cauchy completeness as “admitting all Cauchy-weighted colimits”.

**7.2. PROPOSITION.** *A  $\mathcal{Q}$ -category is Cauchy complete if and only if it admits all colimits weighted by a Cauchy distributor, if and only if it admits all colimits weighted by a Cauchy (contravariant) presheaf.*

**PROOF.** Let  $\mathbb{B}$  be Cauchy complete; given a Cauchy distributor  $\Phi: \mathbb{C} \multimap \mathbb{A}$  and a functor  $F: \mathbb{A} \rightarrow \mathbb{B}$ , the  $\Phi$ -weighted colimit of  $F$  is the functor  $K: \mathbb{C} \rightarrow \mathbb{B}$  to which the Cauchy distributor  $\mathbb{B}(-, F-) \otimes_{\mathbb{A}} \Phi$  converges:

$$\mathbb{B}(K-, -) = \Phi^* \otimes_{\mathbb{A}} \mathbb{B}(F-, -) = \left[ \Phi, \mathbb{B}(F-, -) \right].$$

Suppose now that  $\mathbb{B}$  admits all colimits weighted by a Cauchy presheaf; then every such Cauchy presheaf is representable because  $\phi: *_C \multimap \mathbb{B}$  (with  $\phi \dashv \phi^*$ ) converges to  $\text{colim}(\phi, 1_{\mathbb{B}})$ :

$$\mathbb{B}\left(\text{colim}(\phi, 1_{\mathbb{B}})-, -\right) = \left[ \phi, \mathbb{B}(1_{\mathbb{B}}-, -) \right] = \phi^* \otimes_{\mathbb{B}} \mathbb{B}(-, -) = \phi^*.$$

By 7.1 the category  $\mathbb{B}$  is Cauchy complete. (The remaining implication is trivial.) ■

‘Cauchy completeness’ is a self-dual notion: a  $\mathcal{Q}$ -category  $\mathbb{A}$  is Cauchy complete if and only if the  $\mathcal{Q}^{\text{op}}$ -category  $\mathbb{A}^{\text{op}}$  is. (This really follows directly from the definition.) Therefore we have – by dualizing the statement of 7.2, using that ‘Cauchy completeness’ is a self-dual notion – at once the following.

**7.3. PROPOSITION.** *A  $\mathcal{Q}$ -category  $\mathbb{A}$  is Cauchy complete if and only if it admits all limits weighted by a right adjoint distributor, if and only if it admits all limits weighted by a right adjoint (covariant) presheaf.*

In the rest of this section we work in terms of colimits.

**7.4. EXAMPLE.** Every  $\mathbf{2}$ -enriched category  $\mathbb{A}$  is Cauchy complete: a down-set of  $\mathbb{A}$  (i.e. a contravariant presheaf on  $\mathbb{A}$ ) is Cauchy if and only if it is principal (i.e. a representable presheaf).

**7.5. EXAMPLE.** The  $\text{Rel}(\Omega)$ -category associated to a sheaf on a locale  $\Omega$  (see 3.11) is (skeletal and) Cauchy complete. In fact, it is also *symmetric* in the sense that it equals its opposite (which makes sense because  $\text{Rel}(\Omega)^{\text{op}} = \text{Rel}(\Omega)$ ). [Walters, 1981] proves that sheaves on  $\Omega$  are *precisely* the symmetric, skeletal, Cauchy complete  $\text{Rel}(\Omega)$ -categories. Dropping symmetry and skeletality, Cauchy complete  $\text{Rel}(\Omega)$ -categories are ordered sheaves on  $\Omega$  [Betti *et al.*, 1983; Stubbe, 2004c].

7.6. **EXAMPLE.** Consider again Lawvere’s generalized metric spaces (see 3.12); denote by  $\mathbb{A}$  such a space. In [Lawvere, 1973] it is proved that a left adjoint distributor from a one-point space into  $\mathbb{A}$  is precisely (an equivalence class of) a Cauchy sequence in  $\mathbb{A}$ —and it is therefore that left adjoint distributors are also called Cauchy distributors. Cauchy completeness of  $\mathbb{A}$  as enriched category coincides with its Cauchy completeness as metric space (“all Cauchy sequences/distributors converge”).

#### CAUCHY COMPLETING A CATEGORY

Not every  $\mathcal{Q}$ -category  $\mathbb{A}$  is Cauchy complete; still we are interested in “classifying” the Cauchy distributors on  $\mathbb{A}$ : we want to construct a  $\mathcal{Q}$ -category  $\mathbb{A}_{\text{cc}}$  such that

$$\text{for every } \mathcal{Q}\text{-category } \mathbb{C}, \text{Map}(\text{Dist}(\mathcal{Q}))(\mathbb{C}, \mathbb{A}) \simeq \text{Cat}(\mathcal{Q})(\mathbb{C}, \mathbb{A}_{\text{cc}}). \quad (14)$$

Much like the construction of  $\mathcal{P}\mathbb{A}$  we have the following proposition (the proof of which is much like that of 6.1, so it is omitted).

7.7. **PROPOSITION.** *For any  $\mathcal{Q}$ -category  $\mathbb{A}$ , the “Cauchy completion  $\mathbb{A}_{\text{cc}}$  of  $\mathbb{A}$ ” is, by definition, the full subcategory of  $\mathcal{P}\mathbb{A}$  determined by the Cauchy presheaves on  $\mathbb{A}$ :*

$$(\mathbb{A}_{\text{cc}})_0 = \{\phi \in (\mathcal{P}\mathbb{A})_0 \mid \phi \text{ is Cauchy}\}.$$

*This  $\mathcal{Q}$ -category is skeletal and it satisfies (14) in the following way:*

- for a Cauchy distributor  $\Phi: \mathbb{C} \dashrightarrow \mathbb{A}$  the corresponding functor  $F_\Phi: \mathbb{C} \rightarrow \mathbb{A}_{\text{cc}}$  maps  $c \in \mathbb{C}_0$  onto the Cauchy presheaf  $\Phi(-, c): t\mathbb{C} \dashrightarrow \mathbb{A}$ ;
- and for a functor  $F: \mathbb{C} \rightarrow \mathbb{A}_{\text{cc}}$  the corresponding Cauchy distributor  $\Phi_F: \mathbb{C} \dashrightarrow \mathbb{A}$  has elements, for  $(c, a) \in \mathbb{C}_0 \times \mathbb{A}_0$ ,  $\Phi_F(a, c) = F(c)(a)$  (and its right adjoint has elements  $\Phi_F^*(c, a) = F(c)^*(a)$ ).

Note that the hom-arrows in  $\mathbb{A}_{\text{cc}}$  are identical to those in  $\mathcal{P}\mathbb{A}$ , but since objects  $\phi, \psi \in \mathbb{A}_{\text{cc}}$  have right adjoints in  $\text{Dist}(\mathcal{Q})$  we may write (with slight abuse of notation) that  $\mathbb{A}_{\text{cc}}(\psi, \phi) = \psi^* \otimes_{\mathbb{A}} \phi$ . This will be a helpful trick in many of the calculations that follow.

Plugging the identity distributor  $\mathbb{A}: \mathbb{A} \dashrightarrow \mathbb{A}$  into the equivalence (14) gives us a functor

$$i_{\mathbb{A}}: \mathbb{A} \rightarrow \mathbb{A}_{\text{cc}}: a \mapsto \left( \mathbb{A}(-, a): *_{ta} \dashrightarrow \mathbb{A} \right)$$

which is precisely the factorization of the Yoneda embedding  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  over the full inclusion  $j_{\mathbb{A}}: \mathbb{A}_{\text{cc}} \rightarrow \mathcal{P}\mathbb{A}$  that defined  $\mathbb{A}_{\text{cc}}$ . An alternative for the classifying property in (14) is then: for any  $\mathbb{C}$ , a distributor  $\Phi: \mathbb{C} \dashrightarrow \mathbb{A}$  is Cauchy if and only if the corresponding functor  $F_\Phi: \mathbb{C} \rightarrow \mathcal{P}\mathbb{A}$  (as in 6.1) factors through  $j_{\mathbb{A}}: \mathbb{A}_{\text{cc}} \rightarrow \mathcal{P}\mathbb{A}$ . It follows too that  $i_{\mathbb{A}}$  enjoys properties similar to those of  $Y_{\mathbb{A}}$  (especially 6.3 and 6.5).

7.8. **PROPOSITION.** *For any Cauchy presheaf  $\phi: *_{\mathbb{C}} \dashrightarrow \mathbb{A}$  and any object  $a \in \mathbb{A}_0$ ,  $\mathbb{A}_{\text{cc}}(i_{\mathbb{A}}(a), \phi) = \phi(a)$  and  $\mathbb{A}_{\text{cc}}(\phi, i_{\mathbb{A}}(a)) = \phi^*(a)$ . In particular is  $i_{\mathbb{A}}: \mathbb{A} \rightarrow \mathbb{A}_{\text{cc}}$  fully faithful.*

**7.9. PROPOSITION.** *Every Cauchy presheaf on a  $\mathcal{Q}$ -category  $\mathbb{A}$  is canonically the colimit of representables in  $\mathbb{A}_{\text{cc}}$ : for  $\phi: *_{\mathcal{C}} \dashrightarrow \mathbb{A}$  Cauchy, the  $\phi$ -weighted colimit of  $i_{\mathbb{A}}: \mathbb{A} \rightarrow \mathbb{A}_{\text{cc}}$  (exists and) is  $\phi$  itself. In other words,  $\text{colim}(\phi, i_{\mathbb{A}}) = \phi$  (where we may write an equality instead of merely an isomorphism, for  $\mathbb{A}_{\text{cc}}$  is skeletal).*

Up to now we have discussed many similarities between  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  and  $i_{\mathbb{A}}: \mathbb{A} \rightarrow \mathbb{A}_{\text{cc}}$ ; but here are two remarkable differences.

**7.10. PROPOSITION.** *A  $\mathcal{Q}$ -category  $\mathbb{A}$  is Cauchy complete if and only if  $i_{\mathbb{A}}: \mathbb{A} \rightarrow \mathbb{A}_{\text{cc}}$  is an essentially surjective functor, if and only if it is an equivalence in  $\text{Cat}(\mathcal{Q})$ <sup>12</sup>.*

**PROOF.** Since  $i_{\mathbb{A}}: \mathbb{A} \rightarrow \mathbb{A}_{\text{cc}}$  is fully faithful, it is clear that it is an equivalence if and only if it is essentially surjective. But the essential surjectivity, saying that any  $\phi \in \mathbb{A}_{\text{cc}}$  is isomorphic to  $i_{\mathbb{A}}(a)$  for some  $a \in \mathbb{A}$ , is precisely the statement in 7.1.  $\blacksquare$

**7.11. PROPOSITION.** *Any  $\mathcal{Q}$ -category  $\mathbb{A}$  is isomorphic to its Cauchy completion in the quantaloid  $\text{Dist}(\mathcal{Q})$ <sup>13</sup>.*

**PROOF.**  $i_{\mathbb{A}}: \mathbb{A} \rightarrow \mathbb{A}_{\text{cc}}$  is a fully faithful functor, so the unit of the induced adjoint pair of distributors  $\mathbb{A}_{\text{cc}}(-, i_{\mathbb{A}}-) \dashv \mathbb{A}_{\text{cc}}(i_{\mathbb{A}}-, -)$  is an equality. It suffices to prove that also its co-unit is an equality. Thereto, consider  $\phi, \psi \in (\mathbb{A}_{\text{cc}})_0$  and calculate that

$$\begin{aligned} \mathbb{A}_{\text{cc}}(\psi, i_{\mathbb{A}}-) \otimes_{\mathbb{A}} \mathbb{A}_{\text{cc}}(i_{\mathbb{A}}-, \phi) &= \bigvee_{a \in \mathbb{A}_0} \mathbb{A}_{\text{cc}}(\psi, \mathbb{A}(-, a)) \circ \mathbb{A}_{\text{cc}}(\mathbb{A}(-, a), \phi) \\ &= \psi^* \otimes_{\mathbb{A}} \bigvee_{a \in \mathbb{A}_0} (\mathbb{A}(-, a) \circ \mathbb{A}(a, -)) \otimes_{\mathbb{A}} \phi \\ &= \psi^* \otimes_{\mathbb{A}} \mathbb{A} \otimes_{\mathbb{A}} \phi \\ &= \psi^* \otimes_{\mathbb{A}} \phi \\ &= \mathbb{A}_{\text{cc}}(\psi, \phi). \end{aligned}$$

That is to say,  $\mathbb{A}_{\text{cc}}(-, i_{\mathbb{A}}-) \otimes_{\mathbb{A}} \mathbb{A}_{\text{cc}}(i_{\mathbb{A}}-, -) = \mathbb{A}_{\text{cc}}$ , as wanted.  $\blacksquare$

The following result, although it uses 7.11 in its proof, may be considered analogous to 6.4.

**7.12. PROPOSITION.** *For any  $\mathbb{A}$ , the Cauchy completion  $\mathbb{A}_{\text{cc}}$  is Cauchy complete.*

**PROOF.** It suffices to prove that any Cauchy presheaf on  $\mathbb{A}_{\text{cc}}$  is representable: for any Cauchy  $\Phi: *_{\mathcal{C}} \dashrightarrow \mathbb{A}_{\text{cc}}$  we must find a Cauchy  $\phi: *_{\mathcal{C}} \dashrightarrow \mathbb{A}$  such that  $\mathbb{A}_{\text{cc}}(-, \phi) = \Phi$ . Using  $\mathbb{A} \cong \mathbb{A}_{\text{cc}}$  in  $\text{Dist}(\mathcal{Q})$  (see 7.11) we put  $\phi := \mathbb{A}_{\text{cc}}(i_{\mathbb{A}}-, -) \otimes_{\mathbb{A}_{\text{cc}}} \Phi$ ,

$$\begin{array}{ccc} *_{\mathcal{C}} & \xrightarrow{\Phi} & \mathbb{A}_{\text{cc}} \xrightarrow{\mathbb{A}_{\text{cc}}(i_{\mathbb{A}}-, -)} \mathbb{A}, \\ & \searrow \phi & \nearrow \\ & & \phi \end{array}$$

<sup>12</sup>Cocompleteness of  $\mathbb{A}$  is in general not equivalent to  $Y_{\mathbb{A}}: \mathbb{A} \rightarrow \mathcal{P}\mathbb{A}$  being an equivalence in  $\text{Cat}(\mathcal{Q})$ !

<sup>13</sup>But  $\mathbb{A}$  need not be isomorphic to  $\mathcal{P}\mathbb{A}$  in  $\text{Dist}(\mathcal{Q})$ !

which is Cauchy because it is the composition of Cauchy distributors ( $\mathbb{A}_{\text{cc}}(i_{\mathbb{A}}-, -)$  is not only right but also left adjoint because it is invertible). Now for any  $\psi \in \mathbb{B}_{\text{cc}}$ ,

$$\begin{aligned} \mathbb{A}_{\text{cc}}(\psi, \phi) &= \psi^* \otimes_{\mathbb{A}} \phi \\ &= \psi^* \otimes_{\mathbb{A}} \mathbb{A}_{\text{cc}}(i_{\mathbb{A}}-, -) \otimes_{\mathbb{A}_{\text{cc}}} \Phi \\ &= \mathbb{A}_{\text{cc}}(\psi, i_{\mathbb{A}}-) \otimes_{\mathbb{A}} \mathbb{A}_{\text{cc}}(i_{\mathbb{A}}-, -) \otimes_{\mathbb{A}_{\text{cc}}} \Phi \\ &= \mathbb{A}_{\text{cc}}(\psi, -) \otimes_{\mathbb{A}_{\text{cc}}} \Phi \\ &= \Phi(\psi), \end{aligned}$$

so  $\phi$  indeed represents  $\Phi$ . ■

A direct consequence of 7.11 and 7.12 is that  $\text{Dist}(\mathcal{Q})$  is equivalent to its full subquanti-  
taloid  $\text{Dist}_{\text{cc}}(\mathcal{Q})$  determined by the Cauchy complete categories.

In view of the examples, in particular 7.5, it is tempting to think of the Cauchy completion  $\mathbb{A} \mapsto \mathbb{A}_{\text{cc}}$  of a  $\mathcal{Q}$ -category as a sheafification. The following will only strengthen that intuition.

#### THE UNIVERSALITY OF THE CAUCHY COMPLETION

The proof of the following is analogous to that of 6.9, and therefore it is omitted.

**7.13. PROPOSITION.** *A functor  $F: \mathbb{A} \rightarrow \mathbb{B}$  has a pointwise left Kan extension along  $i_{\mathbb{A}}: \mathbb{A} \rightarrow \mathbb{A}_{\text{cc}}$  if and only if, for every  $\phi \in \mathbb{A}_{\text{cc}}$ , the  $\phi$ -weighted colimit of  $F$  exists; in this case  $\langle F, i_{\mathbb{A}} \rangle \circ i_{\mathbb{A}} \cong F^{14}$ . Actually,  $\langle F, i_{\mathbb{A}} \rangle$  is the essentially unique factorization of  $F$  through  $i_{\mathbb{A}}$ .*

In other words, denoting  $\text{Cat}_{\text{cc}}(\mathcal{Q})$  for the full sub-2-category of  $\text{Cat}(\mathcal{Q})$  determined by the Cauchy complete categories, we have for  $\mathcal{Q}$ -categories  $\mathbb{A}$  and  $\mathbb{B}$ , with  $\mathbb{B}$  Cauchy complete, an equivalence induced by composition with  $i_{\mathbb{A}}: \mathbb{A} \rightarrow \mathbb{A}_{\text{cc}}$

$$\text{Cat}_{\text{cc}}(\mathcal{Q})(\mathbb{A}_{\text{cc}}, \mathbb{B}) \xrightarrow{\sim} \text{Cat}(\mathcal{Q})(\mathbb{A}, \mathbb{B}): G \mapsto G \circ i_{\mathbb{A}}$$

(with inverse  $F \mapsto \langle F, i_{\mathbb{A}} \rangle$ ).

**7.14. PROPOSITION.** *Cauchy completing provides a left (bi)adjoint to the embedding of  $\text{Cat}_{\text{cc}}(\mathcal{Q})$  in  $\text{Cat}(\mathcal{Q})$ :  $(-)\text{cc} \dashv \mathcal{U}: \text{Cat}_{\text{cc}}(\mathcal{Q}) \rightleftarrows \text{Cat}(\mathcal{Q})$ .*

The 2-functor

$$\text{Cat}_{\text{cc}}(\mathcal{Q}) \rightarrow \text{Map}(\text{Dist}_{\text{cc}}(\mathcal{Q})): (F: \mathbb{A} \rightarrow \mathbb{B}) \mapsto (\mathbb{B}(-, F-): \mathbb{A} \dashv \mathbb{B})$$

is the identity on objects, and induces equivalences of the hom-orders: it is a biequivalence. Since  $\text{Dist}(\mathcal{Q})$  itself is equivalent to  $\text{Dist}_{\text{cc}}(\mathcal{Q})$ , we may record that

$$\text{Cat}_{\text{cc}}(\mathcal{Q}) \simeq \text{Map}(\text{Dist}_{\text{cc}}(\mathcal{Q})) \simeq \text{Map}(\text{Dist}(\mathcal{Q})).$$

Finally we wish to discuss the ‘‘interplay’’ between  $\mathbb{A}$ ,  $\mathcal{P}\mathbb{A}$ ,  $\mathcal{P}^{\dagger}\mathbb{A}$  and  $\mathbb{A}_{\text{cc}}$ .

<sup>14</sup>But it is not true in general that  $\langle F, i_{\mathbb{A}} \rangle: \mathbb{A}_{\text{cc}} \rightarrow \mathbb{B}$  is a left adjoint!

7.15. PROPOSITION. For  $\mathcal{Q}$ -categories  $\mathbb{A}$  and  $\mathbb{B}$ , the following are equivalent:

1.  $\mathbb{A} \cong \mathbb{B}$  in  $\text{Dist}(\mathcal{Q})$ ;
2.  $\mathbb{A}_{\text{cc}} \simeq \mathbb{B}_{\text{cc}}$  in  $\text{Cat}(\mathcal{Q})$ ;
3.  $\mathcal{P}\mathbb{A} \simeq \mathcal{P}\mathbb{B}$  in  $\text{Cat}(\mathcal{Q})$ ;
4.  $\mathcal{P}^\dagger\mathbb{A} \simeq \mathcal{P}^\dagger\mathbb{B}$  in  $\text{Cat}(\mathcal{Q})$ .

PROOF.  $(1 \Leftrightarrow 2)$   $\mathbb{A} \cong \mathbb{B}$  in  $\text{Dist}(\mathcal{Q})$  implies and is implied by  $\mathbb{A}_{\text{cc}} \cong \mathbb{B}_{\text{cc}}$  in  $\text{Dist}_{\text{cc}}(\mathcal{Q})$  (using 7.7), which by the biequivalence  $\text{Cat}_{\text{cc}}(\mathcal{Q}) \simeq \text{Map}(\text{Dist}_{\text{cc}}(\mathcal{Q}))$  means that  $\mathbb{A}_{\text{cc}} \simeq \mathbb{B}_{\text{cc}}$  in  $\text{Cat}_{\text{cc}}(\mathcal{Q})$ , or in  $\text{Cat}(\mathcal{Q})$ .

$(1 \Leftrightarrow 3)$   $\mathbb{A} \cong \mathbb{B}$  in  $\text{Dist}(\mathcal{Q})$  if and only if  $\mathcal{P}\mathbb{A} \simeq \mathcal{P}\mathbb{B}$  in  $\text{Cocont}(\mathcal{Q})$  by the local equivalence  $\text{Dist}(\mathcal{Q}) \rightarrow \text{Cocont}(\mathcal{Q})$  of 6.12; but to give an equivalence between cocomplete categories in  $\text{Cocont}(\mathcal{Q})$  is the same as in  $\text{Cat}(\mathcal{Q})$  (because the functors constituting the equivalence are always left adjoint, thus cocontinuous).

$(1 \Leftrightarrow 4)$  is dual to the previous argument. ■

Whenever for two  $\mathcal{Q}$ -categories  $\mathbb{A}$  and  $\mathbb{B}$  the equivalent conditions of 7.15 are fulfilled, then  $\mathbb{A}$  is *Morita equivalent* to  $\mathbb{B}$ . The morality is then that “a  $\mathcal{Q}$ -category is but a presentation of its Cauchy completion”; in particular, if one only cares about  $\mathcal{Q}$ -category theory “up to Cauchy completion” (i.e. “up to Morita equivalence”), then one can forget about functors and Cauchy completion and Morita equivalence altogether, and treat  $\text{Map}(\text{Dist}(\mathcal{Q}))$  as “the 2-category of  $\mathcal{Q}$ -categories and functors”.

## 8. Appendix: Distributor calculus

### LAX (CO)LIMITS IN A QUANTALOID

Write  $\mathcal{Q}$  for a quantaloid, and  $\mathcal{D}$  for a small category. A *lax functor*  $F: \mathcal{D} \rightarrow \mathcal{Q}$  is a mapping of objects  $\mathcal{D}_0 \rightarrow \mathcal{Q}_0: D \mapsto FD$  together with actions  $\mathcal{D}(D, D') \rightarrow \mathcal{Q}(FD, FD'): f \mapsto Ff$  on hom-sets such that

- for all  $D \in \mathcal{D}_0$ ,  $1_{FD} \leq F1_D$ ;
- for all composable  $f, g \in \mathcal{D}$ ,  $(Fg \circ Ff) \leq F(g \circ f)$ .

Given two such lax functors  $F, G: \mathcal{D} \rightrightarrows \mathcal{Q}$ , a family  $(\theta_D: FD \rightarrow GD)_{D \in \mathcal{D}_0}$  of  $\mathcal{Q}$ -arrows constitutes a *lax-natural transformation*  $\theta: F \rightrightarrows G$  if

- for all  $f: D \rightarrow D'$  in  $\mathcal{D}$ ,  $(\theta_{D'} \circ Ff) \geq (Gf \circ \theta_D)$ ;

and the family is said to constitute an *oplax-natural transformation* if

- for all  $f: D \rightarrow D'$  in  $\mathcal{D}$ ,  $(\theta_{D'} \circ Ff) \leq (Gf \circ \theta_D)$ .

The collection of lax transfos, resp. oplax transfos, between two parallel lax functors  $F, G: \mathcal{D} \rightrightarrows \mathcal{Q}$  constitutes a sup-lattice (with “componentwise supremum”); we shall denote it by  $\mathbf{Lax}(F, G)$ , resp.  $\mathbf{OpLax}(F, G)$ . Lax transfos (oplax transfos) may also be composed: this too is done “componentwise”. The notions of ‘lax transfo’ and ‘oplax transfo’ are dual in the following sense: two lax functors  $F, G: \mathcal{D} \rightrightarrows \mathcal{Q}$  may be thought of as  $F^{\text{op}}, G^{\text{op}}: \mathcal{D}^{\text{op}} \rightrightarrows \mathcal{Q}^{\text{op}}$ , and then we have that  $\mathbf{Lax}(F, G) = \mathbf{OpLax}(G^{\text{op}}, F^{\text{op}})$ .

Any (ordinary) functor  $F: \mathcal{D} \rightarrow \mathcal{Q}$  is trivially lax, and any (ordinary) natural transformation  $\alpha: F \Rightarrow G$  between functors is trivially (op)lax. The set  $\mathbf{Nat}(F, G)$  of natural transformations between two functors  $F, G: \mathcal{D} \rightrightarrows \mathcal{Q}$  is a sub-sup-lattice of both the sup-lattices  $\mathbf{Lax}(F, G)$  and  $\mathbf{OpLax}(F, G)$  (it is their intersection).

In what follows, denote by  $\Delta_A: \mathcal{D} \rightarrow \mathcal{Q}$  the constant functor at the object  $A \in \mathcal{Q}$  (on whatever category  $\mathcal{D}$ ). For an arrow  $k: A \rightarrow B$  in  $\mathcal{Q}$  the constant transformation from  $\Delta_A$  to  $\Delta_B$  will be denoted by  $\Delta_k$ .

**8.1. DEFINITION.** *Let  $\mathcal{D}$  be a small category, and  $\mathcal{Q}$  any quantaloid. For a lax functor  $F: \mathcal{D} \rightarrow \mathcal{Q}$  a lax cone over  $F$  consists of an object  $L \in \mathcal{Q}$  and a lax transfo  $\pi \in \mathbf{Lax}(\Delta_L, F)$ . Such a lax cone  $(L, \pi)$  is the (necessarily essentially unique) lax conical limit of  $F$  if it is universal: for all  $A \in \mathcal{Q}$  there is an isomorphism<sup>15</sup> of sup-lattices induced by composition with  $\pi: \Delta_L \Rightarrow F$ ,*

$$\mathcal{Q}(A, L) \xrightarrow{\sim} \mathbf{Lax}(\Delta_A, F): k \mapsto \pi \circ \Delta_k. \quad (15)$$

Lax cocone and lax conical colimit are the dual notions: a lax cocone on  $F$  is an oplax transfo  $\sigma \in \mathbf{OpLax}(F, \Delta_C)$ , and it is the lax colimit if it is universal.

So a lax functor  $F: \mathcal{D} \rightarrow \mathcal{Q}$  admits a lax limit (lax colimit) in  $\mathcal{Q}$  if and only if  $F^{\text{op}}: \mathcal{D}^{\text{op}} \rightarrow \mathcal{Q}^{\text{op}}$  admits a lax colimit (lax limit) in  $\mathcal{Q}^{\text{op}}$ —this is the duality between limits and colimits.

In general, for an ordinary functor  $F: \mathcal{D} \rightarrow \mathcal{Q}$ , the limit on  $F$  is not necessarily its lax limit: whereas any cone  $\alpha \in \mathbf{Nat}(\Delta_L, F)$  is trivially also a lax cone, it is not true that the universal property for such an  $\alpha$  to be a *limit* implies the universal property for  $\alpha$  to be a *lax* limit. But in some cases lax limits do coincide with ordinary ones. (And by duality we can apply the result to lax colimits, of course.)

**8.2. PROPOSITION.** *Suppose that, for some lax functor  $F_1: \mathcal{D}_1 \rightarrow \mathcal{Q}$  and some (ordinary) functor  $F_2: \mathcal{D}_2 \rightarrow \mathcal{Q}$  into a quantaloid  $\mathcal{Q}$ , for all  $X \in \mathcal{Q}$  there is a sup-isomorphism*

$$\mathbf{Lax}(\Delta_X, F_1) \xrightarrow{\sim} \mathbf{Nat}(\Delta_X, F_2) \quad (16)$$

*which is natural in  $X$ . Then some  $\pi \in \mathbf{Lax}(\Delta_L, F_1)$  is the lax limit of  $F_1$  if and only if  $\pi' \in \mathbf{Nat}(\Delta_L, F_2)$ , the image of  $\pi$  under the isomorphism above (taking  $X = L$ ), is the limit of  $F_2$ .*

<sup>15</sup>Actually the definition of “lax (co)limit” in a general bicategory asks only for an *equivalence*, and not for an isomorphism. But an equivalence between antisymmetric orders is always an isomorphism, so we put that right away in this definition.

PROOF. To say that the functor  $F_2: \mathcal{D}_2 \rightarrow \mathcal{Q}$  has limit  $\pi' \in \mathbf{Nat}(\Delta_L, F_2)$  is to say that there are bijections

$$\beta'_X: \mathcal{Q}(X, L) \xrightarrow{\sim} \mathbf{Nat}(\Delta_X, F_2): f \mapsto \pi' \circ \Delta_f$$

defining an isomorphism of the **Set**-valued functors  $\mathcal{Q}(-, L)$  and  $\mathbf{Nat}(\Delta_-, F_2)$  on (the underlying category of)  $\mathcal{Q}^{\text{op}}$  in **CAT**; and  $\pi' = \beta'_L(1_L)$ . However, not only is each  $\mathbf{Nat}(\Delta_X, F_2)$  is a sup-lattice,  $\mathbf{Nat}(\Delta_-, F_2)$  also preserves suprema of  $\mathcal{Q}$ -arrows; it is thus a **Sup**-valued homomorphism on  $\mathcal{Q}^{\text{op}}$ . Moreover, the natural bijections above – since they are induced by composition in  $\mathcal{Q}$  – always preserve suprema. Therefore,  $F_2$  has limit  $\pi'$  if and only if  $\beta': \mathcal{Q}(-, L) \Rightarrow \mathbf{Nat}(\Delta_-, F_2)$  is an isomorphism of **Sup**-valued homomorphisms on  $\mathcal{Q}^{\text{op}}$  in **QUANT**.

On the other hand, to say that  $\pi \in \mathbf{Lax}(\Delta_X, F_1)$  is the lax limit of the lax functor  $F_1: \mathcal{D}_1 \rightarrow \mathcal{Q}$ , is to say that there are sup-isomorphisms

$$\beta_X: \mathcal{Q}(X, L) \xrightarrow{\sim} \mathbf{Lax}(\Delta_X, F_1): f \mapsto \pi \circ \Delta_f$$

defining an isomorphism  $\beta: \mathcal{Q}(-, L) \Rightarrow \mathbf{Lax}(\Delta_-, F_1)$  of **Sup**-valued homomorphisms on  $\mathcal{Q}^{\text{op}}$  in **QUANT**; and  $\pi = \beta_L(1_L)$ .

The result above then follows by composition of isomorphisms of **Sup**-valued homomorphisms on  $\mathcal{Q}^{\text{op}}$  in **QUANT**. ■

## TWO IMPORTANT EXAMPLES

A set-indexed family  $(A_i)_{i \in I}$  of objects in a quantaloid  $\mathcal{Q}$  can be identified with (the image of) the functor  $F: I \rightarrow \mathcal{Q} : i \mapsto A_i$  whose domain is the index-set regarded as discrete category.

**8.3. PROPOSITION.** *For a set-indexed family  $(A_i)_{i \in I}$  of objects in a quantaloid  $\mathcal{Q}$ , the following are equivalent:*

1.  $(A_i)_{i \in I}$  has a limit (i.e. product) in  $\mathcal{Q}$ ;
2.  $(A_i)_{i \in I}$  has a colimit (i.e. coproduct) in  $\mathcal{Q}$ ;
3.  $(A_i)_{i \in I}$  has a lax limit in  $\mathcal{Q}$ ;
4.  $(A_i)_{i \in I}$  has a lax colimit in  $\mathcal{Q}$ ;
5. there exist an object  $A$  and arrows

$$(A \xrightarrow{p_i} A_i \xrightarrow{s_i} A)_{i \in I}$$

in  $\mathcal{Q}$  such that, for all  $i, j \in I$ ,  $p_j \circ s_i = \delta_{A_i, A_j}$  and  $\bigvee_{i \in I} (s_i \circ p_i) = 1_A$ .

The cone  $(p_i: A \rightarrow A_i)_{i \in I}$  is then the (lax) limit of the given family, and the cocone  $(s_i: A_i \rightarrow A)_{i \in I}$  is its (lax) colimit. For each  $i \in I$  the coprojection  $s_i: A_i \rightarrow A$  is left adjoint to the projection  $p_i: A \rightarrow A_i$ .

PROOF. The equivalence  $(1 \Leftrightarrow 2 \Leftrightarrow 5)$  is well-known.  $(1 \Leftrightarrow 3)$  and  $(2 \Leftrightarrow 4)$  follow from 8.2: because the diagram  $F: I \rightarrow \mathcal{Q}: i \mapsto A_i$  is discrete, any lax cone on  $F$  is in fact an ordinary cone; and any lax cocone is an ordinary cocone. That is to say, in this particular case we have for any object  $A \in \mathcal{Q}$  that  $\mathbf{Nat}(\Delta_A, F) = \mathbf{Lax}(\Delta_A, F)$  and  $\mathbf{Nat}(F, \Delta_A) = \mathbf{OpLax}(F, \Delta_A)$ . The adjunctions  $s_i \dashv p_i$  hold since  $1_{A_i} = p_i \circ s_i$  and  $s_i \circ p_i \leq 1_A$ . ■

As usual, we will refer to limits-colimits (in a quantaloid  $\mathcal{Q}$ ) as *direct sums* and we write them as  $\oplus_i A_i$ .

A monad in a quantaloid  $\mathcal{Q}$  is an endo-arrow  $\widehat{A}^t$  such that  $t \circ t \leq t$  and  $1_A \leq t$ . It can be identified with (the image of) a lax functor  $F: \mathbf{Ob} \rightarrow \mathcal{Q}: * \mapsto A, 1_* \mapsto t$ . Here  $\mathbf{Ob}$  stands for the “generic object”: the category with one object  $*$  and one arrow  $1_*$ . On the other hand,  $\widehat{A}^t$  is a diagram in  $\mathcal{Q}$ , thus (the image of) an ordinary functor  $F': \mathbf{Ar} \rightarrow \mathcal{Q}: *_1 \mapsto A, *_2 \mapsto A, \alpha \mapsto t$ . Now  $\mathbf{Ar}$  stands for the “generic arrow”: the category with two objects,  $*_1$  and  $*_2$ , and exactly one arrow that is not an identity,  $\alpha: *_1 \rightarrow *_2$ .

8.4. PROPOSITION. *For a monad  $\widehat{A}^t$  in a quantaloid  $\mathcal{Q}$ , the following are equivalent:*

1.  $\widehat{A}^t$  has a limit (i.e.  $t, 1_A: A \rightrightarrows A$  have an equalizer) in  $\mathcal{Q}$ ;
2.  $\widehat{A}^t$  has a colimit (i.e.  $t, 1_A: A \rightrightarrows A$  have a coequalizer) in  $\mathcal{Q}$ ;
3.  $\widehat{A}^t$  has a lax limit in  $\mathcal{Q}$ ;
4.  $\widehat{A}^t$  has a lax colimit in  $\mathcal{Q}$ ;
5. there exist an object  $B$  and arrows

$$B \xrightarrow{p} A \xrightarrow{s} B$$

in  $\mathcal{Q}$  such that  $p \circ s = t$  and  $s \circ p = 1_B$ .

If the above is the case for the given monad  $t$ , then  $p: B \rightarrow A$  is actually its (lax) limit, and  $s: A \rightarrow B$  its (lax) colimit. And the coprojection  $s: A \rightarrow B$  is left adjoint to the projection  $p: B \rightarrow A$ .

PROOF.  $(1 \Leftrightarrow 2 \Leftrightarrow 5)$  is well-known for idempotents in a category, so in particular applies to monads in a quantaloid (it is immediate that a monad in  $\mathcal{Q}$  is an idempotent).  $(1 \Leftrightarrow 3)$  and  $(2 \Leftrightarrow 4)$  follow from 8.2: because  $1_A \leq t$ , it is the same to give a lax cone on  $F: \mathbf{Ob} \rightarrow \mathcal{Q}$  as to give an ordinary cone on  $F': \mathbf{Ar} \rightarrow \mathcal{Q}$ ; and lax cocones on  $F$  coincide with cocones on  $F'$ . More precisely, we have for any object  $X \in \mathcal{Q}$  identical sup-lattices  $\mathbf{Nat}(\Delta_X, F') = \mathbf{Lax}(\Delta_X, F)$  and  $\mathbf{Nat}(F', \Delta_X) = \mathbf{OpLax}(F, \Delta_X)$ . The adjunction  $s \dashv p$  follows because  $1_A \leq t = p \circ s$  and  $s \circ p = 1_B$ . ■

The lax limit of a monad  $\widehat{A}^t$  in a 2-category is called the *object of  $t$ -algebras*, or the *Eilenberg-Moore object* for  $t$ , and often written as  $A^t$ . And the lax colimit is the *object of free  $t$ -algebras*, or the *Kleisli object*, written  $A_t$ . In a quantaloid  $\mathcal{Q}$ ,  $A^t$  is isomorphic to  $A_t$  as soon as one (and thus also the other) exists: “every  $t$ -algebra is free”! When  $\mathcal{Q}$  admits (free) algebra objects for all of its monads, we will say that *all monads in  $\mathcal{Q}$  split*.

#### UNIVERSAL PROPERTY OF THE DISTRIBUTOR CALCULUS

For the following we refer also to [Betti *et al.*, 1983; Carboni *et al.*, 1987].

Given a quantaloid  $\mathcal{Q}$ , there is a quantaloid  $\mathbf{Matr}(\mathcal{Q})$  of “matrices with elements in  $\mathcal{Q}$ ”, with:

- objects:  $\mathcal{Q}_0$ -typed sets<sup>16</sup>;
- arrows: for two objects  $(X, t)$ ,  $(Y, t)$ , an arrow  $\mathbb{M}: (X, t) \rightarrow (Y, t)$  is a *matrix* of  $\mathcal{Q}$ -arrows  $\mathbb{M}(y, x): tx \rightarrow ty$ , one for each  $(x, y) \in X \times Y$ ;
- two-cells: for  $\mathbb{M}, \mathbb{N}: (X, t) \rightrightarrows (Y, t)$ ,  $\mathbb{M} \leq \mathbb{N}$  when  $\mathbb{M}(y, x) \leq \mathbb{N}(y, x)$  for all  $x, y$ ;
- composition: for  $\mathbb{M}: (X, t) \rightarrow (Y, t)$  and  $\mathbb{N}: (Y, t) \rightarrow (Z, t)$ , the elements of the composite matrix  $\mathbb{N} \circ \mathbb{M}: (X, t) \rightarrow (Z, t)$  are  $(\mathbb{N} \circ \mathbb{M})(z, x) = \bigvee_{y \in Y} \mathbb{N}(z, y) \circ \mathbb{M}(y, x)$ ;
- identities: the identity matrix  $\mathbb{I}_{(X, t)}: (X, t) \rightarrow (X, t)$  has elements  $\mathbb{I}_{(X, t)}(x', x) = \delta_{tx, tx'}$  (“Kronecker deltas”).

Clearly,  $\mathcal{Q}$  can be embedded in  $\mathbf{Matr}(\mathcal{Q})$ : every arrow  $f: A \rightarrow B$  in  $\mathcal{Q}$  may be viewed as a one-element matrix. We write this embedding as  $i_{\mathcal{Q}}: \mathcal{Q} \rightarrow \mathbf{Matr}(\mathcal{Q})$ . Obviously,  $i_{\mathcal{Q}}: \mathcal{Q} \rightarrow \mathbf{Matr}(\mathcal{Q})$  is a fully faithful homomorphism of quantaloids; it is an equivalence of quantaloids if and only if  $\mathcal{Q}$  has all direct sums. Actually,  $\mathbf{Matr}(\mathcal{Q})$  itself has all direct sums (so “taking matrices” is an essentially idempotent process), and  $i_{\mathcal{Q}}: \mathcal{Q} \rightarrow \mathbf{Matr}(\mathcal{Q})$  describes the universal direct-sum completion of  $\mathcal{Q}$  in **QUANT**.

Given a quantaloid  $\mathcal{Q}$ , the following defines a quantaloid  $\mathbf{Mnd}(\mathcal{Q})$  of “monads and bimodules in  $\mathcal{Q}$ ”:

- objects: monads in  $\mathcal{Q}$ ;
- arrows: an arrow  $b: t \rightrightarrows s$  between monads  $\widehat{A}^t$  and  $\widehat{B}^s$  is a *bimodule* from  $t$  to  $s$ , i.e. a  $\mathcal{Q}$ -arrow  $b: A \rightarrow B$  such that  $s \circ b \leq b$  and  $b \circ t \leq b$ ;
- two-cells: put  $b \leq c$  in  $\mathbf{Mnd}(\mathcal{Q})(t, s)$  whenever  $b \leq c$  in  $\mathcal{Q}(A, B)$ ;
- composition: the composition  $c \otimes_s b: t \rightrightarrows r$  of two bimodules  $b: t \rightrightarrows s$  and  $c: s \rightrightarrows r$  is  $c \otimes_s b = c \circ b$  in  $\mathcal{Q}$ ;

<sup>16</sup>Recall that such a  $\mathcal{Q}_0$ -typed set  $(X, t)$  is really just a way of writing a small discrete diagram  $(tx)_{x \in X}$  in  $\mathcal{Q}$ .

- identity: on a monad  $\widehat{\mathbb{A}}^t$  the identity bimodule is  $t: t \dashrightarrow t$  itself.

The “tensor” notation for composition of bimodules reminds us that the identity on a monad  $\widehat{\mathbb{A}}^t$  is not  $1_A$  but rather  $t$  itself.

An arrow  $f: A \rightarrow B$  may be regarded as bimodule from  $1_A$  to  $1_B$ ; denote the corresponding embedding of quantaloids by  $j_{\mathcal{Q}}: \mathcal{Q} \rightarrow \mathbf{Mnd}(\mathcal{Q})$ . Then  $j_{\mathcal{Q}}: \mathcal{Q} \rightarrow \mathbf{Mnd}(\mathcal{Q})$  is a fully faithful homomorphism of quantaloids which is an equivalence of quantaloids if and only if in  $\mathcal{Q}$  all monads split. In the quantaloid  $\mathbf{Mnd}(\mathcal{Q})$  all monads split (so “taking monads and bimodules” is an essentially idempotent process), actually,  $j_{\mathcal{Q}}: \mathcal{Q} \rightarrow \mathbf{Mnd}(\mathcal{Q})$  describes the universal split-monad completion of  $\mathcal{Q}$  in **QUANT**.

It is now a matter of fact that, for any quantaloid  $\mathcal{Q}$ ,  $\mathbf{Dist}(\mathcal{Q}) = \mathbf{Mnd}(\mathbf{Matr}(\mathcal{Q}))$ . Indeed, take for example the definition of “ $\mathcal{Q}$ -category”: such an  $\mathbb{A}$  consists of an endomatrix of hom-arrows  $\mathbb{A}(a', a): ta \rightarrow ta'$  on the  $\mathcal{Q}_0$ -typed set  $(\mathbb{A}_0, t)$  of objects; these data ought to satisfy, for all  $a, a', a'' \in \mathbb{A}_0$ ,

$$\mathbb{A}(a'', a') \circ \mathbb{A}(a', a) \leq \mathbb{A}(a'', a) \text{ and } 1_{ta} \leq \mathbb{A}(a, a),$$

or equivalently,

$$\bigvee_{a' \in \mathbb{A}_0} \mathbb{A}(a'', a') \circ \mathbb{A}(a', a) \leq \mathbb{A}(a'', a) \text{ and } \delta_{ta, ta'} \leq \mathbb{A}(a', a),$$

or still equivalently, but now in terms of matrices,  $\mathbb{A} \circ \mathbb{A} \leq \mathbb{A}$  and  $\mathbb{I}_{\mathbb{A}_0} \leq \mathbb{A}$ . This says precisely that  $\mathbb{A}$  is a monad in  $\mathbf{Matr}(\mathcal{Q})$ , thus an object of  $\mathbf{Mnd}(\mathbf{Matr}(\mathcal{Q}))$ . (Similar for distributors.)

Consequently we now know the universal property of  $\mathbf{Dist}(\mathcal{Q})$ : the composite embedding  $k_{\mathcal{Q}}: \mathcal{Q} \rightarrow \mathbf{Dist}(\mathcal{Q})$  given by  $j_{\mathbf{Matr}(\mathcal{Q})} \circ i_{\mathcal{Q}}$  is a fully faithful homomorphism of quantaloids which is an equivalence if and only if  $\mathcal{Q}$  has all direct sums and all monads split.  $\mathbf{Dist}(\mathcal{Q})$  itself has all direct sums and all monads split (“taking categories and distributors” is an essentially idempotent process); actually,  $k_{\mathcal{Q}}: \mathcal{Q} \rightarrow \mathbf{Dist}(\mathcal{Q})$  describes the universal direct-sum-and-split-monad completion of  $\mathcal{Q}$  in **QUANT**.

Having all direct sums and a splitting for all monads, is enough for a quantaloid  $\mathcal{Q}$  to admit all lax (co)limits.

**8.5. PROPOSITION.** *For any quantaloid  $\mathcal{Q}$ , the following are equivalent:*

1.  $\mathcal{Q}$  has direct sums and all monads split;
2.  $\mathcal{Q}$  has all lax limits;
3.  $\mathcal{Q}$  has all lax colimits;
4. for any lax functor  $F: \mathcal{D} \rightarrow \mathcal{Q}$  there exist an object  $L$  and arrows

$$(L \xrightarrow{p_D} F D \xrightarrow{s_D} L)_{D \in \mathcal{D}_0}$$

in  $\mathcal{Q}$  such that

$$\bigvee \{s_D \circ p_D \mid D \in \mathcal{D}_0\} = 1_L \text{ and } p_{D'} \circ s_D = \bigvee \{Fd \mid d \in \mathcal{D}(D, D')\}.$$

With notations as in the last sentence, the  $(p_D: L \rightarrow FD)_{D \in \mathcal{D}_0}$  form the lax limit of  $F$ , and the  $(s_D: FD \rightarrow L)_{D \in \mathcal{D}_0}$  its lax colimit; moreover, for every  $D \in \mathcal{D}_0$ , the coprojection  $s_D$  is left adjoint to the projection  $p_D$ .

PROOF. (2  $\Rightarrow$  1) and (3  $\Rightarrow$  1) follow from 8.3 and 8.4.

For (4  $\Rightarrow$  2) and (4  $\Rightarrow$  3), first observe that  $s_D \dashv p_D$ : on the one hand  $1_L = \bigvee \{s_D \circ p_D \mid D \in \mathcal{D}_0\} \geq s_D \circ p_D$ , and on the other  $p_D \circ s_D = \bigvee \{Fd \mid d \in \mathcal{D}(D, D)\} \geq F(1_D) \geq 1_{FD}$ . Knowing this, it follows from  $p_{D'} \circ s_D = \bigvee \{Fd \mid d \in \mathcal{D}(D, D')\}$  that for every  $d: D \rightarrow D'$ ,  $p_{D'} \circ s_D \geq Fd$ , hence also  $s_D \geq s_{D'} \circ Fd$  and  $p_{D'} \geq Fd \circ p_D$ . Hence  $\pi = (p_D)_D$  is a lax cone, and  $\sigma = (s_D)_D$  a lax cocone, on  $F$ . As for their universality, we must verify that for any  $X \in \mathcal{Q}$  the sup-morphisms

$$\mathcal{Q}(X, L) \rightarrow \text{Lax}(\Delta_X, F): x \mapsto \pi \circ \Delta_x,$$

$$\mathcal{Q}(L, X) \rightarrow \text{Lax}(F, \Delta_X): x \mapsto \Delta_x \circ \sigma$$

are bijective (in which case they are sup-isomorphisms). But a calculation shows that their respective inverses are

$$\pi' = (p'_D)_D \mapsto \bigvee \{s_D \circ p'_D \mid D \in \mathcal{D}_0\},$$

$$\sigma' = (s'_D)_D \mapsto \bigvee \{s'_D \circ p_D \mid D \in \mathcal{D}_0\}.$$

For (1  $\Rightarrow$  4), note that a lax functor  $F: \mathcal{D} \rightarrow \mathcal{Q}$  determines a  $\mathcal{Q}$ -category  $\mathbb{D}$  with objects  $\mathbb{D}_0 = \mathcal{D}_0$ , types  $tD = FD$ , and hom-arrows  $\mathbb{D}(D', D) = \bigvee \{Fd \mid d \in \mathcal{D}(D, D')\}$ : the inequalities  $\mathbb{D}(D'', D') \circ \mathbb{D}(D', D) \leq \mathbb{D}(D'', D)$  and  $1_{FD} \leq \mathbb{D}(D, D)$  hold precisely because  $F$  is lax! Since  $\mathcal{Q}$  has all direct sums and all monads split, this  $\mathcal{Q}$ -category  $\mathbb{D}$  must be isomorphic in  $\text{Dist}(\mathcal{Q})$  to a one-object  $\mathcal{Q}$ -category, call it  $*_L$ . This means that there exist distributors  $\sigma: \mathbb{D} \rightarrow *_L$  and  $\pi: *_L \rightarrow \mathbb{D}$  such that  $\sigma \otimes_{*_L} \pi = \mathbb{D}$  and  $\pi \otimes_{\mathbb{D}} \sigma = 1_L$ . Such  $\sigma$  and  $\pi$  are really collections of  $\mathcal{Q}$ -arrows

$$\left( L \xrightarrow{p_D := \pi(D)} FD \xrightarrow{s_D := \sigma(D)} L \right)_{D \in \mathcal{D}_0}$$

satisfying

$$p_{D'} \circ s_D = \mathbb{D}(D', D) = \bigvee \{Fd \mid d \in \mathcal{D}(D, D')\},$$

$$\bigvee \{s_D \circ p_D \mid D \in \mathbb{D}_0\} = \bigvee \{s_D \circ p_D \mid D \in \mathcal{D}_0\} = 1_L.$$

■

## References

- [1] Jean Bénabou (1963) Catégories avec multiplication, *C. R. Acad. Sci. Paris* **256**, pp. 1887–1890.
- [2] Jean Bénabou (1965) Catégories relatives, *C. R. Acad. Sci. Paris* **260**, pp. 3824–3827.
- [3] Jean Bénabou (1967) Introduction to bicategories, *Lecture Notes in Math.* **47**, pp. 1–77.
- [4] Renato Betti, Aurelio Carboni, Ross H. Street and Robert F. C. Walters (1983) Variation through enrichment, *J. Pure Appl. Algebra* **29**, pp. 109–127.
- [5] Francis Borceux and Rosanna Cruciani (1998) Skew  $\Omega$ -sets coincide with  $\Omega$ -posets, *Cahiers Topol. Géom. Différ. Catég.* **39**, pp. 205–220.
- [6] Aurelio Carboni, Stefano Kasangian and Robert F. C. Walters (1987) An axiomatics for bicategories of modules, *J. Pure Appl. Algebra* **45**, pp. 127–141.
- [7] Samuel Eilenberg and G. Max Kelly (1966) Closed categories, pp. 421–562 in *Proc. Conf. Categorical Algebra (La Jolla 1965)*, Springer, New York.
- [8] Robert Gordon and A. John Power (1997) Enrichment through variation, *J. Pure Appl. Algebra* **120**, pp. 167–185.
- [9] G. Max Kelly (1982) *Basic concepts of enriched category theory*. London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge.
- [10] Anna Labella and Vincent Schmitt (2002) Change of base, Cauchy completeness and reversibility, *Theory Appl. Categ.* **10**, pp. 187–219.
- [11] F. William Lawvere (1973) Metric spaces, generalized logic and closed categories, *Rend. Sem. Mat. Fis. Milano* **43**, pp. 135–166. Also in: Reprints in Theory and Applications of Categories, No. 1 (2002) pp. 1–37
- [12] Marie-Anne Moens, Ugo Berni-Canani and Francis Borceux (2002) On regular presheaves and regular semicategories, *Cahiers Topol. Géom. Différ. Catég.* **43**, pp. 163–190.
- [13] Christopher J. Mulvey (1986)  $\&$ , *Suppl. Rend. Circ. Mat. Palermo II* **12**, pp. 99–104.
- [14] Kimmo I. Rosenthal (1996) *The theory of quantaloids*. Pitman Research Notes in Mathematics Series. Longman, Harlow.
- [15] Ross H. Street (1972) The formal theory of monads, *J. Pure Appl. Algebra* **2**, pp. 149–168.
- [16] Ross H. Street (1981) Cauchy characterization of enriched categories, *Rend. Sem. Mat. Fis. Milano* **51**, pp. 217–233. Also in: Reprints in Theory and Applications of Categories, No. 4 (2004) pp. 1–16
- [17] Ross H. Street (1983a) Enriched categories and cohomology, *Questiones Math.* **6**, pp. 265–283.
- [18] Ross H. Street (1983b) Absolute colimits in enriched categories, *Cahiers Topol. Géom. Différ. Catég.* **24**, pp. 377–379.
- [19] Isar Stubbe (2004a) Categorical structures enriched in a quantaloid: tensored and cotensored categories, *preprint*, [arXiv:math.CT/0411366](https://arxiv.org/abs/math/0411366).

- [20] Isar Stubbe (2004b) Categorical structures enriched in a quantaloid: regular presheaves, regular semicategories, *Cahiers Topol. Géom. Différ. Catég.* (to appear), [arXiv:math.CT/0409475](https://arxiv.org/abs/math/0409475).
- [21] Isar Stubbe (2004c) Categorical structures enriched in a quantaloid: orders and ideals over a base quantaloid, *Appl. Categ. Structures* (to appear), [arXiv:math.CT/0409477](https://arxiv.org/abs/math/0409477).
- [22] Gilberte Van den Bossche (1995) Quantaloids and non-commutative ring representations, *Appl. Categ. Structures* **3**, pp. 305–320.
- [23] Robert F. C. Walters (1981) Sheaves and Cauchy-complete categories, *Cahiers Topol. Géom. Différ. Catég.* **22**, pp. 283–286.

*Isar Stubbe*  
*Département de Mathématique*  
*Université de Louvain*  
*Chemin du Cyclotron 2*  
*1348 Louvain-la-Neuve*  
*Belgique*

Email: [i.stubbe@math.ucl.ac.be](mailto:i.stubbe@math.ucl.ac.be)

This article may be accessed via WWW at <http://www.tac.mta.ca/tac/> or by anonymous ftp at [ftp://ftp.tac.mta.ca/pub/tac/html/volumes/14/1/14-01.{dvi,ps}](ftp://ftp.tac.mta.ca/pub/tac/html/volumes/14/1/14-01.dvi,ps)

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

The method of distribution of the journal is via the Internet tools `WWW/ftp`. The journal is archived electronically and in printed paper format.

**SUBSCRIPTION INFORMATION.** Individual subscribers receive (by e-mail) abstracts of articles as they are published. Full text of published articles is available in .dvi, Postscript and PDF. Details will be e-mailed to new subscribers. To subscribe, send e-mail to `tac@mta.ca` including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, `rrosebrugh@mta.ca`.

**INFORMATION FOR AUTHORS.** The typesetting language of the journal is  $\text{T}_{\text{E}}\text{X}$ , and  $\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}2\text{e}$  is the preferred flavour.  $\text{T}_{\text{E}}\text{X}$  source of articles for publication should be submitted by e-mail directly to an appropriate Editor. They are listed below. Please obtain detailed information on submission format and style files from the journal's WWW server at `http://www.tac.mta.ca/tac/`. You may also write to `tac@mta.ca` to receive details by e-mail.

**MANAGING EDITOR.** Robert Rosebrugh, Mount Allison University: `rrosebrugh@mta.ca`

**$\text{T}_{\text{E}}\text{X}$  NICAL EDITOR.** Michael Barr, McGill University: `barr@barrs.org`

**TRANSMITTING EDITORS.**

Lawrence Breen, Université Paris 13: `breen@math.univ-paris13.fr`

Ronald Brown, University of North Wales: `r.brown@bangor.ac.uk`

Jean-Luc Brylinski, Pennsylvania State University: `jlb@math.psu.edu`

Aurelio Carboni, Università dell'Insubria: `aurelio.carboni@uninsubria.it`

Valeria de Paiva, Xerox Palo Alto Research Center: `paiva@parc.xerox.com`

Martin Hyland, University of Cambridge: `M.Hyland@dpms.cam.ac.uk`

P. T. Johnstone, University of Cambridge: `ptj@dpms.cam.ac.uk`

G. Max Kelly, University of Sydney: `maxk@maths.usyd.edu.au`

Anders Kock, University of Aarhus: `kock@imf.au.dk`

Stephen Lack, University of Western Sydney: `s.lack@uws.edu.au`

F. William Lawvere, State University of New York at Buffalo: `wlawvere@acsu.buffalo.edu`

Jean-Louis Loday, Université de Strasbourg: `loday@math.u-strasbg.fr`

Ieke Moerdijk, University of Utrecht: `moerdijk@math.uu.nl`

Susan Niefield, Union College: `niefiels@union.edu`

Robert Paré, Dalhousie University: `pare@mathstat.dal.ca`

Jiri Rosicky, Masaryk University: `rosicky@math.muni.cz`

James Stasheff, University of North Carolina: `jds@math.unc.edu`

Ross Street, Macquarie University: `street@math.mq.edu.au`

Walter Tholen, York University: `tholen@mathstat.yorku.ca`

Myles Tierney, Rutgers University: `tierney@math.rutgers.edu`

Robert F. C. Walters, University of Insubria: `robert.walters@uninsubria.it`

R. J. Wood, Dalhousie University: `rjwood@mathstat.dal.ca`