

MAIN ARTICLES

SPACES WITH A LOCALLY COUNTABLE sn -NETWORK

Ying Ge

Department of Mathematics, Suzhou University,
Suzhou, 215006, P.R.China
geying@pub.sz.jsinfo.net

Abstract: In this paper, we discuss a class of spaces with a locally countable sn -network. We give some characterizations of this class and establish the relation among spaces with a locally countable weak-base, spaces with a locally countable sn -network and spaces with a locally countable cs -network. Also, we investigate variance and inverse invariance of spaces with a locally countable sn -network under certain mappings. As some applications of these results, we obtain some results relative to spaces with a locally countable weak-base.

Key words: sn -networks, cs -networks, weak-base, perfect-mappings, (strongly) Lindelöf mapping, finite subsequence-covering mapping

MSC 2000: 54C10, 54D20, 54D65, 54D80

1. Introduction

sn -networks is a class of important networks between weak-bases and cs -networks, which was introduced and called universal cs -networks by S. Lin in [20]. It is an interesting work to discuss spaces with certain sn -network. In recent years, sn -metrizable spaces (i.e. spaces with a σ -locally finite sn -network) and sn -second spaces (i.e. spaces with a countable sn -network) have attracted considerable attention and many interesting results have been obtained ([11, 12, 13, 20, 22, 23, 32]).

In this paper, we discuss a class of spaces with a locally countable sn -network. We give some characterizations of this class and establish the relation among spaces with a locally countable weak-base, spaces with a locally countable sn -network and spaces with a locally countable cs -network. Also, we investigate variance and inverse invariance of spaces with a locally countable sn -network under certain mappings. As some applications of these results, we obtain some results relative to spaces with a locally countable weak-base.

Throughout this paper, all spaces are assumed to be regular, T_1 and all mappings are continuous and onto. \mathbb{N} , ω and ω_1 denote the set of all natural numbers, the first infinite ordinal and the first uncountable ordinal respectively. For a set D , $|D|$ denotes the cardinal of D . $\{x_n\}$ denotes a sequence, where the n -th term is x_n . Let X be a space and $P \subset X$. A sequence $\{x_n\}$ converging to x in X is eventually in P if $\{x_n : n > k\} \cup \{x\} \subset P$ for some $k \in \mathbb{N}$; is frequently in P if $\{x_{n_k}\}$ is eventually in P for some subsequence $\{x_{n_k}\}$ of $\{x_n\}$. Let \mathcal{P} be a family of subsets of X , $x \in X$ and f be a mapping defined on X .

Then $f(\mathcal{P}) = \{f(P) : P \in \mathcal{P}\}$, $(\mathcal{P})_x$ denotes the subfamily $\{P \in \mathcal{P} : x \in P\}$ of \mathcal{P} , $\bigcup \mathcal{P}$ and $\bigcap \mathcal{P}$ denote the union $\bigcup\{P : P \in \mathcal{P}\}$ and the intersection $\bigcap\{P : P \in \mathcal{P}\}$ respectively.

2. Spaces with a Locally Countable sn -Network

Definition 2.1 ([8, 9]). Let X be a space and let $x \in X$.

(1) $P \subset X$ is called a sequential neighborhood of x if each sequence $\{x_n\}$ converging to x is eventually in P .

(2) A subset U of X is called sequentially open if U is a sequential neighborhood of each of its points; a subset F of X is called sequentially closed if $X - F$ is sequentially open.

(3) X is called a sequential space if each sequentially open subset of X is open in X , equivalently, if each sequentially closed subset of X is closed in X .

(4) X is called a k -space if for each $A \subset X$, A is closed in X iff $A \cap K$ is closed in K for each compact subset K of X .

Remark 2.2 (1) P is a sequential neighborhood of x iff each sequence $\{x_n\}$ converging to x is frequently in P .

(2) The intersection of finite sequential neighborhoods of x is a sequential neighborhood of x .

(3) sequential spaces $\implies k$ -spaces.

Definition 2.3 Let $\mathcal{P} = \bigcup\{\mathcal{P}_x : x \in X\}$ be a cover of a space X , where $\mathcal{P}_x \subset (\mathcal{P})_x$.

(1) \mathcal{P} is called a network of X , if whenever $x \in U \subset X$ with U open in X , there is $P \in \mathcal{P}_x$ such that $x \in P \subset U$, where \mathcal{P}_x is called a network at x in X .

(2) \mathcal{P} is called a k -network of X ([27]), if whenever $K \subset U$ with K compact in X and U open in X , there is a finite $\mathcal{F} \subset \mathcal{P}$ such that $K \subset \bigcup \mathcal{F} \subset U$.

(3) \mathcal{P} is called a cs^* -network of X ([10]), if each convergent sequence S converging to a point $x \in U$ with U open in X , then S is frequently in $P \subset U$ for some $P \in \mathcal{P}$.

Definition 2.4 Let $\mathcal{P} = \bigcup\{\mathcal{P}_x : x \in X\}$ be a network of a space X , where $\mathcal{P}_x \subset (\mathcal{P})_x$.

(1) \mathcal{P} is called a cs -network of X ([15]), if each convergent sequence S converging to a point $x \in U$ with U open in X , then S is eventually in $P \subset U$ for some $P \in \mathcal{P}_x$, where \mathcal{P}_x is called a cs -network at x in X .

Assume \mathcal{P}_x also satisfies the following Condition (*) for each $x \in X$ in the following (2) and (3).

Condition (*): If $P_1, P_2 \in \mathcal{P}_x$, then there is $P \in \mathcal{P}_x$ such that $P \subset P_1 \cap P_2$.

(2) \mathcal{P} is called a weak-base of X ([1]), if for $G \subset X$, G is open in X iff for each $x \in G$ there is $P \in \mathcal{P}_x$ such that $P \subset G$, where \mathcal{P}_x is called a weak neighborhood base at x in X .

(3) \mathcal{P} is called an *sn-network* of X ([11]), if each element of \mathcal{P}_x is a sequential neighborhood of x for each $x \in X$, where \mathcal{P}_x is called an *sn-network* at x in X .

Remark 2.5 ([23]). (1) *weak-bases* \implies *sn-networks* \implies *cs-networks* \implies *cs*-networks*.

(2) In a sequential space, *weak-bases* \iff *sn-networks*.

(3) *sn-networks* are called *universal cs-networks* in [20].

The following example belongs to S. Lin.

Example 2.6 In a k -space, *sn-networks* $\not\iff$ *weak-bases*.

Proof. Let X be the *Stone – Čech* compactification $\beta\mathbb{N}$ of \mathbb{N} . Then X is compact, and so it is a k -space. Since each convergent sequence in $\beta\mathbb{N}$ is trivial, $\mathcal{P} = \{\{x\} : x \in X\}$ is an *sn-network* of X . It is clear that \mathcal{P} is not a weak-base. ■

Definition 2.7 (1) A space X is called *g-metrizable* ([8]) (resp. *sn-metrizable* ([12]), \aleph ([27])) if X has a σ -locally finite weak-base (resp. *sn-network*, *k-network*).

(2) A space X is called *g-second countable* ([29]) (resp. *sn-second countable* ([13]), \aleph_0 ([26])) if X has a countable weak-base (resp. *sn-network*, *k-network*).

(3) A space X is called *g-first countable* ([1]) (resp. *sn-first countable* ([12]), *cs-first countable* ([20])), if X has a weak-base (resp. *sn-network*, *cs-network*) $\mathcal{P} = \bigcup\{\mathcal{P}_x : x \in X\}$ such that \mathcal{P}_x is countable for each $x \in X$.

Remark 2.8 (1) By Remark 2.5, a space X is *g-metrizable* (resp. *g-second countable*, *g-first countable*) iff it is sequential and *sn-metrizable* (resp. sequential and *sn-second countable*, sequential and *sn-first countable*).

(2) If X has a point countable weak-base (resp. *sn-network*, *cs-network*), then X is *g-first countable* (resp. *sn-first countable*, *cs-first countable*).

(3) It is well known that a space is a \aleph_0 -space iff it has a countable *cs-network*, iff it has a countable *cs*-network*.

(4) *sn-first countable* is called *universally csf-countable* in [20].

The following lemma is obtained by combining [19, Theorem 2.8.6] and [22, Corollary 5.1.13].

Lemma 2.9 The following are equivalent for a space X .

(1) X has a locally countable *cs-network*.

(1) X has a locally countable *cs*-network*.

(1) X has a locally countable *k-network*.

Theorem 2.10 *The following are equivalent for a space X .*

- (1) X has a locally countable sn -network.
- (2) X is an sn -first countable space with a locally countable cs -network (resp. k -network, cs^* -network).
- (3) X is a locally sn -second countable space with a σ -locally countable sn -network
- (4) X is a locally \aleph_0 -space with a σ -locally countable sn -network.
- (5) $(MA + \neg CH + TOP)$ X is a locally hereditarily separable space with a σ -locally countable sn -network.
- (6) X is a locally (hereditarily) Lindelöf space with a σ -locally countable sn -network.

Proof. (1) \implies (2). Note that a space with a locally countable sn -network is sn -first countable. So (1) \implies (2) from Remark 2.5(1) and Lemma 2.9.

(2) \implies (1). By Lemma 2.9, let \mathcal{P} be a locally countable cs -network of X . We can assume that \mathcal{P} is closed under finite intersections. For each $x \in X$, let $\{B_n(x) : n \in \mathbb{N}\}$ be an sn -network at x in X , and let $\mathcal{P}_x = \{P \in \mathcal{P} : B_n(x) \subset P \text{ for some } n \in \mathbb{N}\}$, then each element of \mathcal{P}_x is a sequential neighborhood of x . Put $\mathcal{P}' = \bigcup \{\mathcal{P}_x : x \in X\}$, then $\mathcal{P}' \subset \mathcal{P}$ is locally countable. It suffices to prove that \mathcal{P}_x is a network at x in X for each $x \in X$. If not, there is an open neighborhood U of x such that $P \not\subset U$ for each $P \in \mathcal{P}_x$. Let $\{P \in \mathcal{P} : x \in P \subset U\} = \{P_m(x) : m \in \mathbb{N}\}$. Then $B_n(x) \subset P_m(x)$ for each $n, m \in \mathbb{N}$. Choose $x_{n,m} \in B_n(x) - P_m(x)$. For $n \geq m$, let $x_{n,m} = y_k$, where $k = m + n(n-1)/2$. Then the sequence $\{y_k : k \in \mathbb{N}\}$ converges to x . Thus, there is $m, i \in \mathbb{N}$ such that $\{y_k : k \geq i\} \cup \{x\} \subset P_m(x) \subset U$. Take $j \geq i$ with $y_j = x_{n,m}$ for some $n \geq m$. Then $x_{n,m} \in P_m(x)$. This is a contradiction.

(1) \implies (3). Let \mathcal{P} be a locally countable sn -network of X . For each $x \in X$, there is an open neighborhood U of x such that $\mathcal{P}_U = \{P \cap U : P \in \mathcal{P}\}$ is countable. It is easy to prove that \mathcal{P}_U is a countable sn -network of subspace U . So U is an sn -second countable space. Hence, X is a locally sn -second countable space.

(3) \implies (4) \implies (5). It is clear that sn -second countable $\implies \aleph_0 \implies$ hereditarily separable. So (3) \implies (4) \implies (5).

(5) \implies (6). It suffices to prove that X is locally hereditarily Lindelöf. Let $x \in X$ and U be a hereditarily separable neighborhood of x . Recalled a space is an S -space if it is a hereditarily separable and not hereditarily Lindelöf. Since $(MA + \neg CH + TOP)$ there are no S -spaces ([28, Theorem 7.2.3]), U is hereditarily Lindelöf. So X is locally hereditarily Lindelöf.

(6) \implies (1). Let $\mathcal{P} = \bigcup \{\mathcal{P}_n : n \in \mathbb{N}\}$ be a σ -locally countable sn -network of a Locally Lindelöf space X , where each \mathcal{P}_n is locally countable in X . Let $x \in X$ and let U be a Lindelöf neighborhood of x . Let $n \in \mathbb{N}$. For each $y \in U$, there is an open neighborhood U_y of y such that U_y intersects at most countable many elements of \mathcal{P}_n . The open cover $\{U_y : y \in U\}$ of U has countable subcover \mathcal{V} . Put $V = \bigcup \mathcal{V}$, then $U \subset V$ and V intersects at

most countable many elements of \mathcal{P}_n . So U intersects at most countable many elements of \mathcal{P}_n . Moreover, U intersects at most countable many elements of \mathcal{P} . Thus \mathcal{P} is a locally countable sn -network of X . ■

Question 2.11 Can “ $(MA + \neg CH + TOP)$ ” in Theorem 2.10(5) be omitted?

We give some partial answers of Question 2.11 by assuming X is a k -space.

Lemma 2.12 ([14, 22]). *The following hold for a space X .*

- (1) *If X is a compact space with a point countable k -network, then X is metrizable.*
- (2) *If X is a k -space with a point countable k -network, then X is sequential.*
- (3) *If X has a point countable cs^* -network and each compact subset of X is metrizable, then X has a point countable k -network.*

Lemma 2.13 *If X is a k -space with a σ -locally countable cs^* -network, then X is sequential.*

Proof. Let \mathcal{P} be a σ -locally countable cs^* -network of X . Whenever K is a compact subset of X , put $\mathcal{P}_K = \{P \cap K : P \in \mathcal{P}\}$, then \mathcal{P}_K is a σ -locally countable cs^* -network of K . It is easy to see that \mathcal{P}_K is a countable cs^* -network of K , and so K has a countable k -network from Remark 2.8(3). By Lemma 2.12(1), K is metrizable. So X has a point-countable k -network from Remark 2.12(3), hence X is sequential from Remark 2.12(2). ■

Theorem 2.14 *The following are equivalent for a k -space X .*

- (1) *X has a locally countable sn -network.*
- (2) *X is a topological sum of sn -second countable spaces.*
- (3) *X is a sn -metrizable, locally (hereditarily) separable space.*
- (4) *X is a locally (hereditarily) separable space with a σ -locally countable sn -network.*

Proof. (1) \implies (2). X is a k -space with a locally countable cs -network, so X is a topological sum of \aleph_0 -spaces ([17, Theorem 1]). It is easy to see that sn -first countability is hereditary to subspace. Note that each sn -first countable, \aleph_0 -space is sn -second countable ([13, Theorem 2.1]). So X is a topological sum of sn -second countable spaces.

(2) \implies (3). Let $X = \oplus\{X_\alpha : \alpha \in \Lambda\}$, where each X_α is sn -second countable. Note that each X_α is a (hereditarily) separable, open subspace of X , So X is locally (hereditarily) separable. For each $\alpha \in \Lambda$, let $\{P_{\alpha,n} : n \in \mathbb{N}\}$ be a countable sn -network of X_α . Put $\mathcal{P}_n = \{P_{\alpha,n} : \alpha \in \Lambda\}$ for each $n \in \mathbb{N}$, and put $\mathcal{P} = \bigcup\{\mathcal{P}_n : n \in \mathbb{N}\}$, then \mathcal{P} is a locally finite sn -network of X . So X is an sn -metrizable space.

(3) \implies (4). It is clear.

(4) \implies (1). By Theorem 2.10, it suffices to prove that X is locally Lindelöf. Let \mathcal{P} be a σ -locally countable sn -network of X . X is a sequential

space from Lemma 2.13, so \mathcal{P} is a σ -locally countable k -network of X ([30, Corollary 1.5]). Recalled a space is meta-*Lindelöf* if each open cover of it has a point countable open refinement. Thus X is hereditarily meta-*Lindelöf* ([17, Proposition 1]). Each hereditarily meta-*Lindelöf*, locally separable space is locally *Lindelöf* ([14, Proposition 8.7]), so X is locally *Lindelöf*. ■

Corollary 2.15 *A space X is a k -space with a locally countable sn -network iff X has a locally countable weak-base.*

The following examples to shows that “ k ” in Theorem 2.14 can not be omitted.

Example 2.16 *There is a space with a locally countable sn -network, even it is not a topological sum of \aleph_0 -spaces.*

Proof. Let D is a discrete space, where $|D| = 2^\omega$. By [3, Example 4.2], there is an almost disjoint family $\{P_\alpha : \alpha < 2^\omega\}$ consisting of countable infinite subsets of D such that for each uncountable subset P of D , there is $\alpha < 2^\omega$ such that $P_\alpha \subset P$. Let $\{P_{\alpha,n} : n \in \mathbb{N}\}$ be a mutually disjoint family consisting of infinite subsets of P_α . For each $\alpha < 2^\omega$ and each $n \in \mathbb{N}$, choose $p_{\alpha,n} \in \overline{P_{\alpha,n}} - P_{\alpha,n}$, where $\overline{P_{\alpha,n}}$ is the closure of $P_{\alpha,n}$ in the Stone – Čech compactification βD of D . Put $X = D \cup \{p_{\alpha,n} : \alpha < 2^\omega, n \in \mathbb{N}\}$, and X is endowed the subspace topology of βD .

Claim 1. X has a σ -locally countable sn -network.

In fact, since each compact subset of X is finite ([22, Example 1.5.5]), and so each convergent sequence of X is finite. Then, it is easy to see that each cs -network of X is an sn -network. X has a σ -locally countable cs -network ([22, Example 5.1.18(1)]), so X has a σ -locally countable sn -network.

Claim 2. X is not a topological sum of \aleph_0 -spaces ([22, Example 5.1.18(1)]). ■

Example 2.17 *There is a space with a locally countable sn -network, even it is not an \aleph -spaces.*

Proof. Let $X = \omega_1 \cup (\omega_1 \times \{1/n : n \in \mathbb{N}\})$. Define a neighborhood base \mathcal{B}_x for each $x \in X$ for the desired topology on X as follows.

(1) If $x \in X - \omega_1$, then $\mathcal{B}_x = \{\{x\}\}$.

(2) If $x \in \omega_1$, then $\mathcal{B}_x = \{\{x\} \cup (\cup \{V(n, x) \times \{1/n\} : n \geq m\}) : m \in \mathbb{N} \text{ and } V(n, x) \text{ is a neighborhood of } x \text{ in } \omega_1 \text{ with the order topology}\}$.

By [17, Example 1], X has a locally countable k -network, which is not an \aleph -space. It suffices to prove that X is sn -first countable from Theorem 2.10.

Let $x \in X$. If $x \in X - \omega_1$, then $\{\{x\}\}$ is a countable sn -network at x in X . If $x \in \omega_1$, put $\mathcal{P}_x = \{P_{x,m} : m \in \mathbb{N}\}$, where $P_{x,m} = \{x\} \cup \{(x, 1/n) : n \geq m\}$. Then \mathcal{P}_x is a countable network at x in X . We only need to prove that each $P_{x,m}$ is a sequential neighborhood of x .

Let $\{x_n\}$ be a sequence converging to x . Put $K = \{x\} \cup \{x_n : n \in \mathbb{N}\}$, then K is a compact subset of X . By [17, Example 1], we have the following facts.

Fact 1. $K \cap \omega_1$ is finite.

Fact 2. $K - \bigcup\{\{y\} \cup \{(y, 1/n) : n \in \mathbb{N}\} : y \in K \cap \omega_1\}$ is finite.

Case 1. If there is $y \in K \cap \omega_1$ such that $y = x_n$ for infinite many $n \in \mathbb{N}$, i.e., there is a subsequence $\{x_{n_k}\}$ of $\{x_n\}$ such that $y = x_{n_k}$ for each $k \in \mathbb{N}$, then $y = x$. So $\{x_n\}$ is frequently in $P_{x,m}$.

Case 2. If Case 1 does not hold, without loss of the generalization, we may assume $K \cap \omega_1 = \{x\}$ from Fact 1. By Fact 2, $K - \{x\} \cup \{(x, 1/n) : n \in \mathbb{N}\}$ is finite. If there is $y \in K - \{x\} \cup \{(x, 1/n) : n \in \mathbb{N}\}$ such that $y = x_n$ for infinite many $n \in \mathbb{N}$, then $\{x_n\}$ is frequently in $P_{x,m}$ by a similar way in the proof of Case 1. Conversely, there is $k_0 \in \mathbb{N}$ such that $\{x\} \cup \{x_n : n \geq k_0\} \subset \{x\} \cup \{(x, 1/n) : n \in \mathbb{N}\}$. So $\{x_n\}$ is eventually in $P_{x,m}$.

By the above Case 1 and Case 2, $P_{x,m}$ is a sequential neighborhood of x from Remark 2.2(1). ■

Recalled a space X is sequentially separable ([6]) if X has a countable subset D such that for each $x \in X$, there is a sequence $\{x_n\}$ in D converging to x , where D is a sequentially dense subset of X . It is know that each sequentially separable space is separable.

Proposition 2.18 *Let X have a point countable sn -network \mathcal{P} . If X is sequentially separable, then \mathcal{P} is countable. So X is sn -second countable.*

Proof. Let D be a sequentially dense subset of X , and let $\mathcal{P} = \{\mathcal{P}_x : x \in X\}$, where \mathcal{P}_x is an sn -network at x in X for each $x \in X$. For each $x \in D$, since \mathcal{P} is point countable, $(\mathcal{P})_x$ is countable. Hence $\bigcup\{(\mathcal{P})_x : x \in D\}$ is countable. For each $x \in X$ and $P \in \mathcal{P}_x$, there is a sequence S in D converging to x . Note that P is a sequential neighborhood of x . S is eventually in P . This proves that each element of \mathcal{P} intersects with D . Thus, it is easy to see that $\mathcal{P} = \bigcup\{(\mathcal{P})_x : x \in D\}$. So \mathcal{P} is countable. ■

Corollary 2.19 *Let X have a σ -locally countable (or point countable) sn -network \mathcal{P} . If X is locally sequentially separable, then \mathcal{P} is locally countable in X . So X has a locally countable sn -network.*

Proof. Since σ -locally countable \implies point countable, we only need to prove parenthetic part.

Let X be locally sequentially separable. For each $x \in X$, there is an open neighborhood of x such that U is sequentially separable. It is clear that $\{P \cap U : P \in \mathcal{P}\}$ is a point countable sn -network of U . $\{P \cap U : P \in \mathcal{P}\}$ is countable from Proposition 2.18, So \mathcal{P} is locally countable in X . ■

The following example shows that “sequentially separable” in Proposition 2.18 can not be relaxed to “separable”, which is due to [16, Example 1].

Example 2.20 *There is a separable, sn -metrizable space. But it is not an \aleph_0 -spaces, and so it is not an sn -second countable space.*

Proof. Let $\mathbb{Q} \subset X \subset \mathbb{R}$ and $|X| > \omega$, where \mathbb{Q} and \mathbb{R} are the set of all rational numbers and the set of all real numbers respectively. Let $Y = X \cup (\cup\{\mathbb{Q} \times \{1/n\} : n \in \mathbb{N}\})$. Define a neighborhood base \mathcal{B}_y for each $y \in Y$ for the desired topology on Y as follows.

(1) If $y \in Y - X$, then $\mathcal{B}_y = \{\{y\}\}$.

(2) If $y \in X$, then $\mathcal{B}_y = \{\{y\} \cup (\cup\{([a_{y,n}, y) \cap \mathbb{Q}] \times \{1/m\} : n \geq m\}) : m \in \mathbb{N} \text{ and } y > a_{y,n} \in \mathbb{R}\}$.

Then Y is a separable, \aleph -space and not an \aleph_0 -space ([16, Example 1]). On the other hand, each compact subset of Y is finite ([16, Example 1]). By a similar way as in the proof of Example 2.16(claim 1), we can prove Y has a σ -locally finite sn -network. That is, Y is an sn -metric space. ■

3. Mappings on Spaces with a Locally Countable sn -Network

In this section, we discuss invariance and inverse invariance of spaces with a locally countable sn -network under certain mappings

Definition 3.21 Let $f : X \rightarrow Y$ be a mapping.

(1) f is called a perfect mapping ([7]) if f is closed and $f^{-1}(y)$ is a compact subset of X for each $y \in Y$;

(2) f is called a Lindelöf mapping ([31]) (resp. strongly Lindelöf mapping ([31]) if for each $y \in Y$, $f^{-1}(y)$ is a Lindelöf subset of X (resp. $f^{-1}(\bar{U})$ is a Lindelöf subset of X for some neighborhood U of y in Y).

(3) f is called a 1-sequence-covering mapping ([23]) if for each $y \in Y$ there is $x \in f^{-1}(y)$, such that whenever $\{y_n\}$ is a sequence converging to y in Y , there is a sequence $\{x_n\}$ converging to x in X with each $x_n \in f^{-1}(y_n)$.

(4) f is called a finite subsequence-covering mapping ([25]) if for each $y \in Y$ there is a finite subset F of $f^{-1}(y)$, such that for any sequence S in Y converging to y , there is a sequence L in X converging to some $x \in F$ and $f(L)$ is a subsequence of S .

(5) f is a sequentially-quotient mapping ([4]) if whenever S is a convergent sequence in Y there is a convergent sequence L in X such that $f(L)$ is a subsequence of S .

(6) f is a quotient mapping ([7]) if whenever $U \subset Y$, $f^{-1}(U)$ is open in X iff U is open in Y .

We call a space X to be point- G_δ if for each $x \in X$, there is a sequence $\{U_n\}$ of neighborhoods of x in X such that $\{x\} = \bigcap\{U_n : n \in \mathbb{N}\}$. It is clear that if a space X has a locally countable cs -network, then X is point- G_δ (see [26, (D)], for example).

Remark 3.22 ([19]). (1) 1-sequence-covering mappings or sequentially-quotient, finite-to-one mappings \implies finite subsequence-covering mappings \implies sequentially-quotient mappings.

(2) Closed mappings \implies quotient mappings.

(3) If the domain is point- G_δ , then closed mappings \implies sequentially-quotient mappings

(4) If the domain is sequential, then quotient mappings \implies sequentially-quotient mappings.

(5) Quotient mappings preserve k -spaces and perfect mappings inversely preserve k -spaces.

Definition 3.23 ([20]). Let X be a space. Put $\sigma = \{P \subset X : P \text{ is sequentially open in } X\}$. The (X, σ) , the set X with the topology σ , is called the sequential coreflection of X , which is denoted by σX .

Definition 3.24 ([2]). Let $T_0 = \{a_n : n \in \mathbb{N}\}$ be a sequence converging to $x_0 \notin T_0$, and let T_n be a sequence converging to $a_n \notin T_n$ for every $n \in \mathbb{N}$. Let T be the topological sum of $\{T_n \cup \{a_n\} : n \in \mathbb{N}\}$. S_ω is defined as a quotient space obtained from T by identifying all point $a_n \in T$ to the point x_0 .

The following lemma is obtained by combining [20, Theorem 3.6] and [20, Theorem 3.13].

Lemma 3.25 ([20]). A point- G_δ space X is sn -first countable iff X is cs -first countable and contains no closed subspace having S_ω as its sequential coreflection.

Lemma 3.26 ([21]). Let $f : X \longrightarrow Y$ be a perfect mapping and X have a G_δ -diagonal. If Y has a locally countable k -network, then X has a locally countable k -network.

Lemma 3.27 ([11]). Let $f : X \longrightarrow Y$ be a closed mapping and X be point- G_δ . If F is sequentially closed in X , then $f(F)$ is sequentially closed in Y .

Theorem 3.28 Let $f : X \longrightarrow Y$ be a perfect mapping and X have a G_δ -diagonal. If Y has a locally countable sn -network, then X has a locally countable sn -network.

Proof. If Y has a locally countable sn -network, then X has a locally countable cs -network from Remark 2.5(1), Lemma 2.9 and Lemma 3.6. It is clear that X is cs -first countable. Since X has a G_δ -diagonal, X is point- G_δ . It suffices to prove that X contains no closed subspace having S_ω as its sequential coreflection from Theorem 2.10 and Lemma 3.5.

Assume X contains closed subspace S having S_ω as its sequential coreflection. Put $g = f|_{\sigma S} : \sigma S \longrightarrow \sigma f(S)$.

Claim 1. g is closed.

Proof. Let A be a closed subset of σS , then A is sequentially closed in S . It is clear $f : S \longrightarrow f(S)$ is closed and S is point- G_δ . So $f(A)$ is sequentially closed in $f(S)$ from Lemma 3.7, thus $f(A)$ is closed in $\sigma f(S)$.

Claim 2. $g^{-1}(y)$ is compact in σS for each $y \in \sigma f(S)$.

Proof. Let $y \in \sigma f(S)$. Note that X has a G_δ -diagonal and $f^{-1}(y)$ is compact in X , so $f^{-1}(y)$ is metrizable ([5]). Therefore, the topology on the sequential coreflection of $f^{-1}(y) \cap S$ is equivalent to the induced topology of subspace S of X . Thus $g^{-1}(y) = f^{-1}(y) \cap S$ is compact in σS .

By the above two claims, g is perfect. Since S_ω , which is homeomorphic to σS , is a *Fréchet*, \aleph -space and perfect mappings preserve *Fréchet*, \aleph -spaces, $\sigma f(S)$ is a *Fréchet*, \aleph -space. On the other hand, Y is *sn*-first countable, so $f(S)$, as a subspace of Y , is *sn*-first countable. By [20, Theorem 3.13], $\sigma f(S)$ is g -first countable, so $\sigma f(S)$ is *sn*-first countable. Thus $\sigma f(S)$ is a metric space ([11, Theorem 2.4]), and so σS is a metric space ([5]). This contradicts that S_ω is not metrizable. ■

We have the following corollary from Corollary 2.15 and Remark 3.2(5) and Theorem 3.8.

Corollary 3.29 *Let $f : X \longrightarrow Y$ be a perfect mapping and X have a G_δ -diagonal. If Y has a locally countable weak-base, then X has a locally countable weak-base.*

Example 3.30 *A perfect image of a g -second countable space has not any locally countable *sn*-network.*

Proof. Let $X = \{0\} \cup \mathbb{N} \cup (\mathbb{N} \times \mathbb{N})$, $\mathcal{F} = \{F \subset \mathbb{N} : F \text{ is finite}\}$, $\mathbb{N}^\mathbb{N} = \{f : f \text{ is a correspondence from } \mathbb{N} \text{ to } \mathbb{N}\}$. For $n, m, k \in \mathbb{N}$, $F \in \mathcal{F}$ and $f \in \mathbb{N}^\mathbb{N}$, put $V(n, m) = \{n\} \cup \{k : k \geq m\}$, $H(F, f) = \bigcup \{V(n, f(n)) : n \in \mathbb{N} - F\}$. Define a neighborhood base \mathcal{B}_x for each $x \in X$ for the desired topology on X as follows.

- (1) If $x \in \mathbb{N} \times \mathbb{N}$, then $\mathcal{B}_x = \{\{x\}\}$.
- (2) If $x \in \mathbb{N}$, then $\mathcal{B}_x = \{V(x, m) : m \in \mathbb{N}\}$.
- (3) If $x = 0$, then $\mathcal{B}_x = \{\{x\} \cup H(F, f) : F \in \mathcal{F}, f \in \mathbb{N}^\mathbb{N}\}$.

Let Y be the quotient space obtained from X by shrinking the set $\{0\} \cup \mathbb{N}$ to a point, $f : X \longrightarrow Y$ be a natural mapping. Then

Claim 1. f is perfect and X is g -second countable ([18, Example 3.1]).

Claim 2. Y is not *sn*-first countable ([11, Example 3.2]), so Y has not any locally countable *sn*-network from Theorem 2.10. ■

Which mappings preserve spaces with a locally countable *sn*-network? We give some answers for this question.

Lemma 3.31 *Let $f : X \longrightarrow Y$ be a finite subsequence-covering mapping. If X is *sn*-first countable, then Y is *sn*-first countable.*

Proof. Let $y \in Y$. Then there is a finite subset F of $f^{-1}(y)$, such that for any sequence S in Y converging to y , there is a sequence L in X converging to some $x \in F$ and $f(L)$ is a subsequence of S . X is *sn*-first countable, for each

$x \in F$, let $\mathcal{P}_x = \{P_{x,n} : n \in \mathbb{N}\}$ be a decreasing sn -network at x in X . Put $\mathcal{F}_y = \{\bigcup\{f(P_{x,n}) : x \in F\} : n \in \mathbb{N}\}$. Then \mathcal{F}_y is countable decreasing.

(1) \mathcal{F}_y is a network at y in Y . In fact, let U be an open neighborhood of y , then $F \subset f^{-1}(y) \subset f^{-1}(U)$. For each $x \in F$, there is $n_x \in \mathbb{N}$ such that $x \in P_{x,n_x} \subset f^{-1}(U)$, so $y \in f(P_{x,n_x}) \subset U$. Put $n_0 = \max\{n_x : x \in F\}$, then $P_{x,n_0} \subset P_{x,n_x}$ for each $x \in F$. So $y \in \bigcup\{f(P_{x,n_0}) : x \in F\} \subset \bigcup\{f(P_{x,n_x}) : x \in F\} \subset U$.

(2) Let $\bigcup\{f(P_{x,n_1}) : x \in F\}, \bigcup\{f(P_{x,n_2}) : x \in F\} \in \mathcal{F}_y$. Put $n_0 = \max\{n_1, n_2\}$, then $\bigcup\{f(P_{x,n_0}) : x \in F\} \in \mathcal{F}_y$ and $\bigcup\{f(P_{x,n_0}) : x \in F\} \subset (\bigcup\{f(P_{x,n_1}) : x \in F\}) \cap (\bigcup\{f(P_{x,n_2}) : x \in F\})$.

(3) $\bigcup\{f(P_{x,n}) : x \in F\}$ is a sequential neighborhood of y for each $n \in \mathbb{N}$. In fact, let S be a sequence in Y converging to y . Then there is a sequence L in X converging to some $x_0 \in F$ and $f(L)$ is a subsequence of S . For each $n \in \mathbb{N}$. Since $P_{x_0,n}$ is a sequential neighborhood of x_0 , L is eventually in $P_{x_0,n}$. So $f(L)$ is eventually in $f(P_{x_0,n})$, hence S is frequently in $f(P_{x_0,n})$. Moreover, S is frequently in $\bigcup\{f(P_{x,n}) : x \in F\}$. By Remark 2.2(1), $\bigcup\{f(P_{x,n}) : x \in F\}$ is a sequential neighborhood of y . ■

Lemma 3.32 *Let $f : X \rightarrow Y$ be a closed, Lindelöf mapping. If \mathcal{P} is a locally countable family of subsets of X , then $f(\mathcal{P})$ is a locally countable family of subsets of Y .*

Proof. Let $\mathcal{P} = \{P_\alpha : \alpha \in \Lambda\}$ be a locally countable family of subsets of X and let $y \in Y$. For each $x \in f^{-1}(y)$, there is an open neighborhood U_x of x such that $\{\alpha \in \Lambda : U_x \cap P_\alpha \neq \emptyset\}$ is countable. $f^{-1}(y) \subset \bigcup\{U_x : x \in f^{-1}(y)\}$ and $f^{-1}(y)$ is Lindelöf, so there is a countable subset B of $f^{-1}(y)$ such that $f^{-1}(y) \subset \bigcup\{U_x : x \in B\}$. Put $U = \bigcup\{U_x : x \in B\}$. It is clear that $\{\alpha \in \Lambda : U \cap P_\alpha \neq \emptyset\}$ is countable. Note that f is closed. By [7, Theorem 1.4.13], there is an open neighborhood V of y such that $f^{-1}(V) \subset U$. Thus $\Lambda' = \{\alpha \in \Lambda : f^{-1}(V) \cap P_\alpha \neq \emptyset\}$ is countable. It is easy to check that $\{\alpha \in \Lambda : V \cap f(P_\alpha) \neq \emptyset\} = \Lambda'$. So $\{\alpha \in \Lambda : V \cap f(P_\alpha) \neq \emptyset\}$ is countable. This proves that $f(\mathcal{P})$ is a locally countable family of subsets of Y . ■

Theorem 3.33 *Let $f : X \rightarrow Y$ be a closed, finite-to-one mapping. If X has a locally countable sn -network, then Y has a locally countable sn -network.*

Proof. Let \mathcal{P} be a locally countable sn -network of X . Then f is sequentially quotient from Remark 3.2(3), and so Y is sn -first countable from Remark 3.2(1) and Lemma 3.11. Since sequentially quotient mappings preserve cs^* -networks ([19, Proposition 2.7.3]), $f(\mathcal{P})$ is a cs^* -network of Y . $f(\mathcal{P})$ is locally countable from Lemma 3.12, so $f(\mathcal{P})$ is a locally countable cs^* -network of Y . Thus Y has a locally countable sn -network from Theorem 2.10. ■

Question 3.34 *Do closed, countable-to-one mappings preserve spaces with a locally countable sn -network?*

A clopen mapping means an open and closed mapping.

Theorem 3.35 *Let $f : X \longrightarrow Y$ be a clopen, Lindelöf mapping. If X has a locally countable sn-network, then Y has a locally countable sn-network.*

Proof. Let $\mathcal{P} = \bigcup\{\mathcal{P}_x : x \in X\}$ be a locally countable sn-network of X . Since f is closed, Lindelöf, by a similar way as in the proof of Theorem 3.13, $f(\mathcal{P})$ is a locally countable cs^* -network of Y . It suffices to prove that Y is sn-first countable from Theorem 2.10. Let $y \in Y$. Put $\mathcal{F}_y = \{f(P) : P \in \mathcal{P}_x \text{ and } x \in f^{-1}(y)\}$, then $\mathcal{F}_y \subset f(\mathcal{P})$, so \mathcal{F}_y is locally countable. Note that $y \in \bigcap \mathcal{F}_y$, \mathcal{F}_y is countable. It is clear that \mathcal{F}_y is a network at y in Y . We only need to prove that each element of \mathcal{F}_y is a sequential neighborhood of y . Let $f(P) \in \mathcal{F}_y$ and $\{y_k\}$ be a sequence in Y converging to y . Then there is $x \in f^{-1}(y)$ such that $P \in \mathcal{P}_x$. Since X is point- G_δ , $\{x\} = \bigcap\{U_n : n \in \mathbb{N}\}$, where each U_n is open in X and $\overline{U_{n+1}} \subset U_n$. For each $n \in \mathbb{N}$, $y \in f(U_n)$ and $f(U_n)$ is open as f is open, so there is $m_n \in \mathbb{N}$ such that $y_k \in f(U_n)$ for each $k \geq m_n$. Pick $x_n \in U_n$ such that $f(x_n) = y_{m_n}$. Since f is closed, it is not difficult to prove that the sequence $\{x_n\}$ converges to $x \in P$. P is a sequential neighborhood of x , so $\{x_n\}$ is eventually in P . Consequently, $\{f(x_n)\}$ is eventually in $f(P)$, so $\{y_k\}$ is frequently in $f(P)$. By Remark 2.2(1), $f(P)$ is a sequential neighborhood of y . ■

Corollary 3.36 *Let $f : X \longrightarrow Y$ be an open, perfect mapping. If X has a locally countable sn-network, then Y has a locally countable sn-network.*

Clopen mappings preserve spaces with a locally countable weak-base ([24, Theorem 4.7]). But the following question is still open.

Question 3.37 *Do clopen mappings preserve spaces with a locally countable sn-network (resp. cs-network)?*

Lemma 3.38 *Let $f : X \longrightarrow Y$ be a strongly Lindelöf-mapping. If \mathcal{P} is a locally countable family of subsets of X , then $f(\mathcal{P})$ is a locally countable family of subsets of Y .*

Proof. Let $\mathcal{P} = \{P_\alpha : \alpha \in \Lambda\}$ be a locally countable family of subsets of X and let $y \in Y$. Then there is a neighborhood W of y in Y such that $f^{-1}(\overline{W})$ is a Lindelöf subset of X . It suffices to prove that $\{\alpha \in \Lambda : W \cap f(P_\alpha) \neq \emptyset\}$ is countable. For each $x \in f^{-1}(\overline{W})$, there is an open neighborhood U_x of x such that $\{\alpha \in \Lambda : U_x \cap P_\alpha \neq \emptyset\}$ is countable. $f^{-1}(\overline{W}) \subset \bigcup\{U_x : x \in f^{-1}(\overline{W})\}$ and $f^{-1}(\overline{W})$ is Lindelöf, so there is a countable subset B of $f^{-1}(\overline{W})$ such that $f^{-1}(\overline{W}) \subset \bigcup\{U_x : x \in B\}$. It is easy to see that $\{\alpha \in \Lambda : (\bigcup\{U_x : x \in B\}) \cap P_\alpha \neq \emptyset\}$ is countable, so $\Lambda' = \{\alpha \in \Lambda : (f^{-1}(W) \cap P_\alpha) \neq \emptyset\}$ is countable. It is easy to check that $\{\alpha \in \Lambda : W \cap f(P_\alpha) \neq \emptyset\} = \Lambda'$. This completes the proof. ■

Theorem 3.39 *Let X have a locally countable sn -network. If one of the following holds, then Y has a locally countable sn -network.*

- (1) *f is finite subsequence-covering, strongly Lindelöf.*
- (2) *f is 1-sequence-covering, strongly Lindelöf.*
- (3) *f is sequentially-quotient, finite-to-one, strongly Lindelöf.*

Proof. We only need to prove part (1) from Remark 3.2(1). Let $f : X \rightarrow Y$ be a finite subsequence-covering, strongly Lindelöf-mapping and \mathcal{P} be a locally countable sn -network of X . Then Y is sn -first countable from lemma 3.11 and $f(\mathcal{P})$ is a locally countable family of subsets of Y from Lemma 3.18. By a similar way as in the proof of Theorem 3.13, we can prove $f(\mathcal{P})$ is a cs^* -network of Y . So Y has a locally countable sn -network from Theorem 2.10. ■

The following corollary is obtained from Remark 3.2(2),(4),(5), Corollary 2.15, Theorem 3.13 and Theorem 3.19.

Corollary 3.40 *Let X have a locally countable weak-base. If one of the following holds, then Y has a locally countable weak-base.*

- (1) *f is closed, finite-to-one.*
- (2) *f is finite subsequence-covering, quotient, strongly Lindelöf.*
- (3) *f is 1-sequence-covering, quotient, strongly Lindelöf.*
- (4) *f is quotient, finite-to-one, strongly Lindelöf.*

Acknowledgement. This project was supported by NSF of the Education Committee of Jiangsu Province in China(No.02KJB110001)

References

- [1] Arhangel'skii, A.V., Mappings and spaces, Russian Math. Surveys, 21(1966), 115-162
- [2] Arhangel'skii, A. and Franklin, S., Ordinal invariants for topological spaces, Michigan Math. J., 15(1968), 313-320
- [3] Blacar, B., and Simon, P., Disjoint refinement, In: Handbook of Boolean Algebras, V2. Amsterdam: North-Holland, 332-386
- [4] Boone, J.R. and Siwec, F., Sequentially quotient mappings, Czech. Math. J., 26(1976), 174-182
- [5] Chaber, J., Conditions which imply compactness in countably compact spaces, Bull. Pol. Acad. Math., 24(1976), 993-998
- [6] Davis, S.W., More on Cauchy condition, Topology Proc., 9(1984), 31-36
- [7] Engelking, R., General Topology(revised and completed edition), Berlin: Heldermann, 1989
- [8] Franklin, S.P., Spaces in which sequence suffice, Fund. Math., 57(1965), 107-115
- [9] Gale, D., Compact sets of functions and function rings, Proc. Amer. Math. Soc., 1(1950), 303-308
- [10] Gao, Z., \aleph -space is invariant under perfect mappings, Q and A in General Topology, 5(1987), 271-279
- [11] Ge, Y., On sn -metrizable spaces, Acta Math. Sinica, 45(2002), 355-360
- [12] Ge, Y., Characterizations of sn -metrizable spaces, Publ. Inst. Math., Nouv. Ser., 74(88)(2003), 121-128

-
- [13] Ge, Y., Spaces with countable sn -networks, *Comment Math. Univ. Carolinae*, 45(2004), 169-176
- [14] Gruenhage, G., Michael, E. and Tanaka, Y., Spaces determined by point-countable covers, *Pacific J. Math.*, 113(1984), 303-332
- [15] Guthrie, J.A., A characterization of \aleph_0 -spaces, *General Topology Appl.*, 1(1971), 105-110
- [16] Lin, S., On normal separable \aleph -space, Q and A in *General Topology*, 5(1987), 249-254
- [17] Lin, S., Spaces with a locally countable k -network, *Northeastern Math. J.*, 6(1990), 39-44
- [18] Lin, S., On g -metrizable spaces, *Chinese Ann. Math., Ser A*, 13(1992), 403-409
- [19] Lin, S., *Generalized Metric Spaces and Mappings*, Chinese Science Press, Beijing, 1995
- [20] Lin, S., A note on the Arens' spaces and sequential fan, *Topology Appl.*, 81(1997), 185-196
- [21] Lin, S., A note on spaces with a locally countable k -network, *J. of Jishou Univ.*, 18(3)(1997), 10-12
- [22] Lin, S., *Point-Countable Covers and Sequence-Covering Mappings*, Beijing: Chinese Science Press, 2002
- [23] Lin, S., and Yan, P., Sequence-covering maps of metric spaces, *Topology. Appl.*, 109(2001), 301-314
- [24] Liu, C., Spaces with a locally countable weak base, *Math. Japonica*, 41(1995), 261-267
- [25] Liu, C., Notes on weak bases, Q and A in *General Topology*, 22(2004), 39-42
- [26] Michael, E.A., \aleph_0 -spaces, *J. Math. Mech.*, 15(1966), 983-1002
- [27] O'Meara, P., On paracompactness in function spaces with the compact-open topology, *Proc. Amer. Math. Soc.*, 29(1971), 183-189
- [28] Roitman, J., Basic S and L , In: Kumen K and Vaughan J E. eds, *Handbook of Set-Theoretic Topology*, Amsterdam: North-Holland, 295-326, 1984
- [29] Siwiec, F., On defining a space by a weak base, *Pacific J. Math.*, 52(1974), 233-245
- [30] Tanaka, Y., Point-countable covers and k -networks, *Topology Proc.*, 12(1987), 327-349
- [31] Tanaka, Y., Theory of k -networks II, Q and A in *General Topology*, 19(2001), 27-46
- [32] Tanaka, Y. and Ge, Y., Around quotient compact images of metric spaces, and symmetric spaces, *Houston J. Math.*(in Press)

Received January, 8, 2007; accepted June, 29, 2007