

Sequential fractional differential equations and unification of anti-periodic and multi-point boundary conditions

Bashir Ahmad, Ahmed Alsaedi, Mohammed H. Aqlan

Nonlinear Analysis and Applied Mathematics (NAAM)-Research Group
Department of Mathematics, Faculty of Science, King Abdulaziz University
P.O. Box 80203, Jeddah 21589, Saudi Arabia

E-mail: bashirahmad_gau@yahoo.com (BA), aalsaedi@hotmail.com (AA), aqlan76@gmail.com (MHA)

Abstract

In this paper, we present a novel idea of unification of anti-periodic and multi-point boundary conditions and develop the existence theory for sequential fractional differential equations supplemented with these new conditions. We apply fixed point theorems due to Banach, Krasnoselskii, Leray-Schauder alternative criterion, and Leray-Schauder degree theory to obtain the desired results. Our results are well illustrated with the aid of examples and correspond to some new special cases for particular choices of parameters involved in the problem.

Keywords: Sequential fractional differential equations; nonlocal; anti-periodic; multi-point; existence; fixed point.

MSC 2000: 34A08, 34B10, 34B15.

1 Introduction

In this paper, we study a nonlinear anti-periodic type multi-point boundary value problems of sequential fractional differential equations given by

$$\left\{ \begin{array}{l} ({}^c D^q + k {}^c D^{q-1})u(t) = f(t, u(t)), \quad 2 < q \leq 3, \quad 0 < t < T, \\ \alpha_1 u(0) + \sum_{i=1}^m a_i u(\eta_i) + \gamma_1 u(T) = \beta_1, \\ \alpha_2 u'(0) + \sum_{i=1}^m b_i u'(\eta_i) + \gamma_2 u'(T) = \beta_2, \\ \alpha_3 u''(0) + \sum_{i=1}^m c_i u''(\eta_i) + \gamma_3 u''(T) = \beta_3 \end{array} \right. \quad (1.1)$$

where ${}^c D^q$ denotes the Caputo fractional derivative of order q , $\alpha_j, \beta_j, \gamma_j \in \mathbb{R}$ ($j = 1, 2, 3$), $a_i, b_i, c_i \in \mathbb{R}$ ($i = 1, 2, \dots, m$), $k \in \mathbb{R}^+$ and f is an appropriately chosen continuous function. The new boundary conditions in (1.1) can be interpreted as the values of the unknown function and its first and second-order derivatives at the end points of the interval under consideration relate to the linear combination of the values of the unknown function, and its first and second-order derivatives at interior points $\eta_i \in (0, T)$.

There has been a great interest in developing theoretical analysis for a variety of boundary value problems of nonlinear fractional order differential equations in the recent years. Anti-periodic and multi-point boundary value problems are two important classes of such problems, which have received considerable attention. Anti-periodic boundary conditions appear in the mathematical modeling of certain physical processes and phenomena, for example, see [1] and the references cited therein. Nonlocal multi-point conditions are regarded as more plausible than the classical initial/boundary conditions

as these condition can describe peculiarities of chemical, physical or other processes happening inside the domain, for instance, see [2]. For some recent works on fractional-order anti-periodic and multi-point boundary value problems, we refer the reader to a series of papers ([3]-[17]) and the references cited therein.

The objective of the present paper is to investigate the existence of solutions for sequential fractional differential equations equipped with a new kind of boundary conditions consisting of a combination of anti-periodic and multi-point boundary conditions. The rest of the paper is organized as follows. Section 2 contains some preliminary concepts of fractional calculus and an auxiliary lemma, which plays a key role in the forthcoming analysis. In Section 3, we prove the uniqueness of solutions for the given problem, while the existence results are presented in Section 4. Illustrative examples are discussed in Section 5. The paper concludes with some interesting remarks described in Section 6.

2 Preliminary work

First of all, we recall some basic definitions [18, 19].

Definition 2.1 *The fractional integral of order r with the lower limit zero for a function f is defined as*

$$I^r f(t) = \frac{1}{\Gamma(r)} \int_0^t \frac{f(s)}{(t-s)^{1-r}} ds, \quad t > 0, \quad r > 0,$$

provided the right hand-side is point-wise defined on $[0, \infty)$, where $\Gamma(\cdot)$ is the gamma function, which is defined by $\Gamma(r) = \int_0^\infty t^{r-1} e^{-t} dt$.

Definition 2.2 *The Riemann-Liouville fractional derivative of order $r > 0$, $n-1 < r < n$, $n \in \mathbb{N}$, is defined as*

$$D_{0+}^r f(t) = \frac{1}{\Gamma(n-r)} \left(\frac{d}{dt} \right)^n \int_0^t (t-s)^{n-r-1} f(s) ds,$$

where the function $f(t)$ has absolutely continuous derivative up to order $(n-1)$.

Definition 2.3 *The Caputo derivative of order r for a function $f : [0, \infty) \rightarrow \mathbb{R}$ can be written as*

$${}^c D^r f(t) = D^r \left(f(t) - \sum_{k=0}^{n-1} \frac{t^k}{k!} f^{(k)}(0) \right), \quad t > 0, \quad n-1 < r < n.$$

Remark 2.4 *If $f(t) \in C^n[0, \infty)$, then*

$${}^c D^r f(t) = \frac{1}{\Gamma(n-r)} \int_0^t \frac{f^{(n)}(s)}{(t-s)^{r+1-n}} ds = I^{n-r} f^{(n)}(t), \quad t > 0, \quad n-1 < r < n.$$

To define a solution for the given problem, we need the following lemma.

Lemma 2.5 *Let $h \in C[0, T]$, $u \in C^3[0, T]$. Then the following linear problem*

$$\left\{ \begin{array}{l} ({}^c D^q + k {}^c D^{q-1})u(t) = h(t), \quad 2 < q \leq 3, \quad 0 < t < T, \\ \alpha_1 u(0) + \sum_{i=1}^m a_i u(\eta_i) + \gamma_1 u(T) = \beta_1, \\ \alpha_2 u'(0) + \sum_{i=1}^m b_i u'(\eta_i) + \gamma_2 u'(T) = \beta_2, \\ \alpha_3 u''(0) + \sum_{i=1}^m c_i u''(\eta_i) + \gamma_3 u''(T) = \beta_3, \end{array} \right. \quad (2.1)$$

is equivalent to the fractional integral equation

$$\begin{aligned}
u(t) &= \nu_1(t) + \int_0^t e^{-k(t-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} h(x) dx \right) ds \\
&+ \sum_{i=1}^m \omega_i(t) \int_0^{\eta_i} e^{-k(\eta_i-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} h(x) dx \right) ds \\
&+ \nu_2(t) \int_0^T e^{-k(T-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} h(x) dx \right) ds \\
&+ \sum_{i=1}^m \psi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-2}}{\Gamma(q-1)} h(s) ds + \nu_3(t) \int_0^T \frac{(T-s)^{q-2}}{\Gamma(q-1)} h(s) ds \\
&+ \sum_{i=1}^m \varphi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-3}}{\Gamma(q-2)} h(s) ds + \nu_4(t) \int_0^T \frac{(T-s)^{q-3}}{\Gamma(q-2)} h(s) ds,
\end{aligned} \tag{2.2}$$

where

$$\begin{aligned}
\nu_1(t) &= \frac{\beta_1}{\lambda_1} - \left(\frac{\sum_{i=1}^m a_i \eta_i + \gamma_1 T}{\lambda_1 \lambda_2} \right) \beta_2 - \frac{\beta_3}{k^2 \lambda_1 \lambda_2 \delta_3} \left(\lambda_2 \delta_1 + k \delta_2 \left(\sum_{i=1}^m a_i \eta_i + \gamma_1 T \right) \right) + \frac{\beta_3 e^{-kt}}{k^2 \delta_3} \\
&+ \frac{t}{\lambda_2} \left(\beta_2 + \frac{\beta_3 \delta_2}{k \delta_3} \right), \\
\omega_i(t) &= \frac{c_i \delta_1}{\delta_3 \lambda_1} - \frac{a_i}{\lambda_1} - \left(\frac{k \left(\sum_{i=1}^m a_i \eta_i + \gamma_1 T \right)}{\lambda_1 \lambda_2 \delta_3} \right) \left(b_i \delta_3 - \delta_2 c_i \right) - \frac{c_i e^{-kt}}{\delta_3} + \frac{kt}{\lambda_2 \delta_3} \left(b_i \delta_3 - \delta_2 c_i \right), \\
\nu_2(t) &= \frac{\gamma_3 \delta_1}{\delta_3 \lambda_1} - \frac{\gamma_1}{\lambda_1} - \left(\frac{k \left(\sum_{i=1}^m a_i \eta_i + \gamma_1 T \right)}{\lambda_1 \lambda_2 \delta_3} \right) \left(\gamma_2 \delta_3 - \delta_2 \gamma_3 \right) - \frac{\gamma_3 e^{-kt}}{\delta_3} + \frac{kt}{\lambda_2 \delta_3} \left(\gamma_2 \delta_3 - \delta_2 \gamma_3 \right), \\
\psi_i(t) &= \frac{-\delta_1 c_i}{k \lambda_1 \delta_3} - \left(\frac{\sum_{i=1}^m a_i \eta_i + \gamma_1 T}{\lambda_1 \lambda_2 \delta_3} \right) \left(\delta_2 c_i - b_i \delta_3 \right) + \frac{c_i e^{-kt}}{k \delta_3} + \frac{t}{\lambda_2 \delta_3} \left(\delta_2 c_i - b_i \delta_3 \right), \\
\nu_3(t) &= \frac{-\delta_1 \gamma_3}{k \lambda_1 \delta_3} - \left(\frac{\sum_{i=1}^m a_i \eta_i + \gamma_1 T}{\lambda_1 \lambda_2 \delta_3} \right) \left(\delta_2 \gamma_3 - \gamma_2 \delta_3 \right) + \frac{\gamma_3 e^{-kt}}{k \delta_3} + \frac{t}{\lambda_2 \delta_3} \left(\delta_2 \gamma_3 - \gamma_2 \delta_3 \right), \\
\varphi_i(t) &= \frac{c_i}{k^2 \delta_3 \lambda_1 \lambda_2} \left(\lambda_2 \delta_1 + \delta_2 k \left(\sum_{i=1}^m a_i \eta_i + \gamma_1 T \right) \right) - \frac{c_i e^{-kt}}{k^2 \delta_3} - \frac{c_i \delta_2 t}{k \delta_3 \lambda_2}, \\
\nu_4(t) &= \frac{\gamma_3}{k^2 \delta_3 \lambda_1 \lambda_2} \left(\lambda_2 \delta_1 + \delta_2 k \left(\sum_{i=1}^m a_i \eta_i + \gamma_1 T \right) \right) - \frac{\gamma_3 e^{-kt}}{k^2 \delta_3} - \frac{\gamma_3 \delta_2 t}{k \delta_3 \lambda_2}, \\
\delta_1 &= \alpha_1 + \sum_{i=1}^m a_i e^{-k \eta_i} + \gamma_1 e^{-kT}, \quad \delta_2 = \alpha_2 + \sum_{i=1}^m b_i e^{-k \eta_i} + \gamma_2 e^{-kT}, \\
\delta_3 &= \alpha_3 + \sum_{i=1}^m c_i e^{-k \eta_i} + \gamma_3 e^{-kT} \neq 0, \quad \lambda_1 = \alpha_1 + \sum_{i=1}^m a_i + \gamma_1 \neq 0, \quad \lambda_2 = \alpha_2 + \sum_{i=1}^m b_i + \gamma_2 \neq 0.
\end{aligned} \tag{2.3}$$

Proof. We know that the general solution of (2.1) can be expressed in term of an integral equation as

$$u(t) = A_0 e^{-kt} + A_1 + A_2 t + \int_0^t e^{-k(t-s)} I^{q-1} h(s) ds, \tag{2.4}$$

where A_0, A_1 and A_2 are arbitrary constants and

$$I^{q-1} h(t) = \int_0^t \frac{(t-x)^{q-2}}{\Gamma(q-1)} h(x) dx.$$

Differentiating (2.4) with respect to t , we obtain

$$u'(t) = -kA_0e^{-kt} + A_2 - k \int_0^t e^{-k(t-s)} I^{q-1} h(s) ds + I^{q-1} h(t), \quad (2.5)$$

$$u''(t) = k^2 A_0 e^{-kt} + k^2 \int_0^t e^{-k(t-s)} I^{q-1} h(s) ds - k I^{q-1} h(t) + I^{q-2} h(t). \quad (2.6)$$

Using the boundary conditions given by (2.1) in (2.4)-(2.6), together with the notations (2.3), we get

$$\begin{aligned} & \delta_1 A_0 + \lambda_1 A_1 + A_2 \left(\sum_{i=1}^m a_i \eta_i + \gamma_1 T \right) + \sum_{i=1}^m a_i \int_0^{\eta_i} e^{-k(\eta-s)} I^{q-1} h(s) ds \\ & + \gamma_1 \int_0^T e^{-k(T-s)} I^{q-1} h(s) ds = \beta_1, \end{aligned} \quad (2.7)$$

$$\begin{aligned} & -k\delta_2 A_0 + \lambda_2 A_2 + \sum_{i=1}^m b_i \left(-k \int_0^{\eta_i} e^{-k(\eta_i-s)} I^{q-1} h(s) ds + I^{q-1} h(\eta_i) \right) \\ & + \gamma_2 \left(-k \int_0^T e^{-k(T-s)} I^{q-1} h(s) ds + I^{q-1} h(T) \right) = \beta_2, \end{aligned} \quad (2.8)$$

$$\begin{aligned} & A_0 k^2 \delta_3 + \sum_{i=1}^m c_i \left(k^2 \int_0^{\eta_i} e^{-k(\eta_i-s)} I^{q-1} h(s) ds - k I^{q-1} h(\eta_i) + I^{q-2} h(\eta_i) \right) \\ & + \gamma_3 \left(k^2 \int_0^T e^{-k(T-s)} I^{q-1} h(s) ds - k I^{q-1} h(T) + I^{q-2} h(T) \right) = \beta_3. \end{aligned} \quad (2.9)$$

Solving the system (2.7)-(2.9) for A_0, A_1 and A_2 , we obtain

$$\begin{aligned} A_0 &= \frac{1}{k^2 \delta_3} \left\{ \beta_3 - \sum_{i=1}^m c_i \left(k^2 \int_0^{\eta_i} e^{-k(\eta_i-s)} I^{q-1} h(s) ds - k I^{q-1} h(\eta_i) + I^{q-2} h(\eta_i) \right) \right. \\ & \left. - \gamma_3 \left(k^2 \int_0^T e^{-k(T-s)} I^{q-1} h(s) ds - k I^{q-1} h(T) + I^{q-2} h(T) \right) \right\}, \\ A_1 &= \frac{\beta_1}{\lambda_1} - \left(\frac{\sum_{i=1}^m a_i \eta_i + \gamma_1 T}{\lambda_1 \lambda_2} \right) \beta_2 - \left(\frac{\delta_1}{k^2 \delta_3 \lambda_1} + \frac{\delta_2 (\sum_{i=1}^m a_i \eta_i + \gamma_1 T)}{k \delta_3 \lambda_1 \lambda_2} \right) \beta_3 \\ & + \sum_{i=1}^m \left(\frac{\delta_1 c_i}{\delta_3 \lambda_1} - \left(\frac{\sum_{i=1}^m a_i \eta_i + \gamma_1 T}{\lambda_1} \right) \left(\frac{b_i k}{\lambda_2} - \frac{k \delta_2 c_i}{\delta_3 \lambda_2} \right) - \frac{a_i}{\lambda_1} \right) \int_0^{\eta_i} e^{-k(\eta_i-s)} I^{q-1} h(s) ds \\ & + \left(\frac{\delta_1 \gamma_3}{\delta_3 \lambda_1} - \left(\frac{\beta_1 \eta + \gamma_1 T}{\lambda_1} \right) \left(\frac{\gamma_2 k}{\lambda_2} - \frac{k \delta_2 \gamma_3}{\delta_3 \lambda_2} \right) - \frac{\gamma_1}{\lambda_1} \right) \int_0^T e^{-k(T-s)} I^{q-1} h(s) ds \\ & - \sum_{i=1}^m \left(\frac{\delta_1 c_i}{k \delta_3 \lambda_1} + \left(\frac{\sum_{i=1}^m a_i \eta_i + \gamma_1 T}{\lambda_1} \right) \left(\frac{\delta_2 c_i}{\delta_3 \lambda_2} - \frac{b_i}{\lambda_2} \right) \right) I^{q-1} h(\eta_i) \\ & - \left(\frac{\delta_1 \gamma_3}{k \delta_3 \lambda_1} + \left(\frac{\beta_1 \eta + \gamma_1 T}{\lambda_1} \right) \left(\frac{\delta_2 \gamma_3}{\delta_3 \lambda_2} + \frac{\gamma_2}{\lambda_2} \right) \right) I^{q-1} h(T) \\ & + \sum_{i=1}^m \left(\frac{\delta_1 c_i}{k^2 \delta_3 \lambda_1} + \frac{\delta_2 c_i (\sum_{i=1}^m a_i \eta_i + \gamma_1 T)}{k \delta_3 \lambda_1 \lambda_2} \right) I^{q-2} h(\eta_i) \end{aligned}$$

$$\begin{aligned}
& + \left(\frac{\delta_1 \gamma_3}{k^2 \delta_3 \lambda_1} + \frac{\delta_2 \gamma_3 (\beta_1 \eta + \gamma_1 T)}{k \delta_3 \lambda_1 \lambda_2} \right) I^{q-2} h(T), \\
A_2 &= \frac{\beta_2}{\lambda_2} + \frac{\delta_2 \beta_3}{k \delta_3 \lambda_2} + \sum_{i=1}^m \left(\frac{b_i k}{\lambda_2} - \frac{k \delta_2 c_i}{\delta_3 \lambda_2} \right) \int_0^{\eta_i} e^{-k(\eta_i-s)} I^{q-1} h(s) ds \\
& + \left(\frac{\gamma_2 k}{\lambda_2} - \frac{k \delta_2 \gamma_3}{\delta_3 \lambda_2} \right) \int_0^T e^{-k(T-s)} I^{q-1} h(s) ds + \sum_{i=1}^m \left(\frac{\delta_2 c_i}{\delta_3 \lambda_2} - \frac{b_i}{\lambda_2} \right) I^{q-1} h(\eta_i) \\
& + \left(\frac{\delta_2 \gamma_3}{\delta_3 \lambda_2} - \frac{\gamma_2}{\lambda_2} \right) I^{q-1} h(T) - \sum_{i=1}^m \frac{\delta_2 c_i}{k \delta_3 \lambda_2} I^{q-2} h(\eta_i) - \frac{\delta_2 \gamma_3}{k \delta_3 \lambda_2} I^{q-2} h(T).
\end{aligned}$$

Substituting the values of A_0, A_1 and A_2 in (2.4), we get the desired solution (2.2). The converse of the lemma follows by direct computation. This completes the proof. $\square$

3 Uniqueness result

Let $\mathcal{P} = C([0, T], \mathbb{R})$ denote the Banach space of all continuous functions from $[0, T] \rightarrow \mathbb{R}$ endowed with the norm defined by $\|u\| = \sup\{|u(t)|, t \in [0, T]\}$.

In view of Lemma 2.5, we transform problem (1.1) into an equivalent fixed point problem as

$$u = \mathcal{H}u, \quad (3.1)$$

where $\mathcal{H} : \mathcal{P} \rightarrow \mathcal{P}$ is defined by

$$\begin{aligned}
(\mathcal{H}u)(t) &= \nu_1(t) + \int_0^t e^{-k(t-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} f(x, u(x)) dx \right) ds \\
& + \sum_{i=1}^m \omega_i(t) \int_0^{\eta_i} e^{-k(\eta_i-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} f(x, u(x)) dx \right) ds \\
& + \nu_2(t) \int_0^T e^{-k(T-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} f(x, u(x)) dx \right) ds \\
& + \sum_{i=1}^m \psi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-2}}{\Gamma(q-1)} f(s, u(s)) ds + \nu_3(t) \int_0^T \frac{(T-s)^{q-2}}{\Gamma(q-1)} f(s, u(s)) ds \\
& + \sum_{i=1}^m \varphi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-3}}{\Gamma(q-2)} f(s, u(s)) ds + \nu_4(t) \int_0^T \frac{(T-s)^{q-3}}{\Gamma(q-2)} f(s, u(s)) ds.
\end{aligned} \quad (3.2)$$

Observe that problem (1.1) has solutions if the operator equation (3.1) has fixed points.

For computational convenience, we set

$$\begin{aligned}
Q &= \sup_{t \in [0, T]} \left\{ \frac{t^{q-1}(1-e^{-kt})}{k\Gamma(q)} + \frac{|\sum_{i=1}^m \omega_i(t) \eta_i^{q-1}(1-e^{-k\eta_i})|}{k\Gamma(q)} + \frac{|\nu_2(t)| T^{q-1}(1-e^{-kT})}{k\Gamma(q)} \right. \\
& \left. + \frac{|\sum_{i=1}^m \psi_i(t) \eta_i^{q-1}|}{\Gamma(q)} + \frac{|\nu_3(t)| T^{q-1}}{\Gamma(q)} + \frac{|\sum_{i=1}^m \varphi_i(t) \eta_i^{q-2}|}{\Gamma(q-1)} + \frac{|\nu_4(t)| T^{q-2}}{\Gamma(q-1)} \right\}.
\end{aligned} \quad (3.3)$$

Now we are in a position to discuss the existence of a unique solution for the problem (1.1) via Banach's contraction mapping principle.

Theorem 3.1 Assume that $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ is a continuous functions satisfying the Lipschitz condition:

(A₁) there exists a positive number ℓ such that $|f(t, u) - f(t, v)| \leq \ell|u - v|$, $\forall t \in [0, T]$, $u, v \in \mathbb{R}$.

Then the boundary value problem (1.1) has a unique solution on $[0, T]$ if $\ell < 1/Q$, where Q is given by (3.3).

Proof. Consider a set $B_r = \{u \in \mathcal{P} : \|u\| \leq r\}$, where $r \geq \frac{QM + \|\nu_1\|}{1 - \ell Q}$, $\sup_{t \in [0, T]} |f(t, 0)| = M$, and Q, ν_1 are respectively given by (3.3) and (2.3). In the first step, we show that $\mathcal{H}B_r \subset B_r$, where the operator $\mathcal{H}$ is defined by (3.2). For any $u \in B_r, t \in [0, T]$, observe that

$$\begin{aligned} |f(t, u(t))| &= |f(t, u(t)) - f(t, 0) + f(t, 0)| \leq |f(t, u(t)) - f(t, 0)| + |f(t, 0)| \\ &\leq \ell\|u\| + M \leq \ell r + M. \end{aligned}$$

Then, for $u \in B_r$, we obtain

$$\begin{aligned} &\|(\mathcal{H}u)\| \\ &\leq \sup_{t \in [0, T]} \left\{ |\nu_1(t)| + \int_0^t e^{-k(t-s)} \left(\int_0^s \frac{(s-x)^{\alpha-2}}{\Gamma(\alpha-1)} |f(x, u(x))| dx \right) ds \right. \\ &\quad + \left| \sum_{i=1}^m \omega_i(t) \int_0^{\eta_i} e^{-k(\eta_i-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} |f(x, u(x))| dx \right) ds \right. \\ &\quad + |\nu_2(t)| \int_0^T e^{-k(T-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} |f(x, u(x))| dx \right) ds \\ &\quad + \left| \sum_{i=1}^m \psi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-2}}{\Gamma(q-1)} |f(s, u(s))| ds + |\nu_3(t)| \int_0^T \frac{(T-s)^{\alpha-2}}{\Gamma(\alpha-1)} |f(s, u(s))| ds \right. \\ &\quad \left. \left. + \left| \sum_{i=1}^m \varphi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-3}}{\Gamma(q-2)} |f(s, u(s))| ds + |\nu_4(t)| \int_0^T \frac{(T-s)^{q-3}}{\Gamma(q-2)} |f(s, u(s))| ds \right\} \right. \\ &\leq (\ell r + M) \sup_{t \in [0, T]} \left\{ \int_0^t e^{-k(t-s)} \left(\int_0^s \frac{(s-x)^{\alpha-2}}{\Gamma(\alpha-1)} dx \right) ds \right. \\ &\quad + \left| \sum_{i=1}^m \omega_i(t) \int_0^{\eta_i} e^{-k(\eta_i-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} dx \right) ds + |\nu_2(t)| \int_0^T e^{-k(T-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} dx \right) ds \right. \\ &\quad + \left| \sum_{i=1}^m \psi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-2}}{\Gamma(q-1)} ds + |\nu_3(t)| \int_0^T \frac{(T-s)^{\alpha-2}}{\Gamma(\alpha-1)} ds + \left| \sum_{i=1}^m \varphi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-3}}{\Gamma(q-2)} ds \right. \right. \\ &\quad \left. \left. + |\nu_4(t)| \int_0^T \frac{(T-s)^{q-3}}{\Gamma(q-2)} ds \right\} + \|\nu_1\| \\ &\leq (\ell r + M)Q + \|\nu_1\| \leq r. \end{aligned}$$

This shows that $\mathcal{H}B_r \subset B_r$. Next we show that the operator $\mathcal{H}$ is a contraction. Let $u, v \in \mathcal{P}$. Then

$$\|\mathcal{H}u - \mathcal{H}v\|$$

$$\begin{aligned}
&\leq \sup_{t \in [0, T]} \left\{ \int_0^t e^{-k(t-s)} \left(\int_0^s \frac{(s-x)^{\alpha-2}}{\Gamma(\alpha-1)} |f(x, u(x)) - f(x, v(x))| dx \right) ds \right. \\
&+ \left| \sum_{i=1}^m \omega_i(t) \int_0^{\eta_i} e^{-k(\eta_i-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} |f(x, u(x)) - f(x, v(x))| dx \right) ds \right. \\
&+ |\nu_2(t)| \int_0^T e^{-k(T-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} |f(x, u(x)) - f(x, v(x))| dx \right) ds \\
&+ \left| \sum_{i=1}^m \psi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-2}}{\Gamma(q-1)} |f(s, u(s)) - f(s, v(s))| ds \right. \\
&+ |\nu_3(t)| \int_0^T \frac{(T-s)^{q-2}}{\Gamma(q-1)} |f(s, u(s)) - f(s, v(s))| ds \\
&+ \left| \sum_{i=1}^m \varphi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-3}}{\Gamma(q-2)} |f(s, u(s)) - f(s, v(s))| ds \right. \\
&+ \left. |\nu_4(t)| \int_0^T \frac{(T-s)^{q-3}}{\Gamma(q-2)} |f(s, u(s)) - f(s, v(s))| ds \right\} \\
&\leq \ell \|u - v\| \sup_{t \in [0, T]} \left\{ \int_0^t e^{-k(t-s)} \left(\int_0^s \frac{(s-x)^{\alpha-2}}{\Gamma(\alpha-1)} dx \right) ds \right. \\
&+ \left| \sum_{i=1}^m \omega_i(t) \int_0^{\eta_i} e^{-k(\eta_i-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} dx \right) ds + |\nu_2(t)| \int_0^T e^{-k(T-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} dx \right) ds \right. \\
&+ \left| \sum_{i=1}^m \psi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-2}}{\Gamma(q-1)} ds + |\nu_3(t)| \int_0^T \frac{(T-s)^{\alpha-2}}{\Gamma(\alpha-1)} ds + \left| \sum_{i=1}^m \varphi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-3}}{\Gamma(q-2)} ds \right. \right. \\
&+ \left. |\nu_4(t)| \int_0^T \frac{(T-s)^{\alpha-3}}{\Gamma(\alpha-2)} ds \right\} \\
&\leq \ell Q \|u - v\|,
\end{aligned}$$

where we have used (3.3). By the given assumption: $\ell < 1/Q$, it follows that the operator $\mathcal{H}$ is a contraction. Thus, by Banach's contraction mapping principle, we deduce that the operator $\mathcal{H}$ has a fixed point, which corresponds to a unique solution of the problem (1.1) on $[0, T]$. $\square$

4 Existence results

In this section, we obtain different criteria for the existence of solutions for the problem (1.1) under different constraints on the nonlinearity involved in the problem. The first results relies on Krasnoselskii's fixed point theorem.

Lemma 4.1 (Krasnoselskii's fixed point theorem) [20]. Let Y be a closed bounded, convex and nonempty subset of a Banach space X . Let B_1, B_2 be the operators such that (i) $B_1 y_1 + B_2 y_2 \in Y$ whenever $y_1, y_2 \in Y$; (ii) B_1 is compact and continuous and (iii) B_2 is a contraction mapping. Then there exists $z \in Y$ such that $z = B_1 z + B_2 z$.

Theorem 4.2 Let $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be continuous functions satisfying the condition (A_1) and that $|f(t, x)| \leq g(t)$, $\forall (t, x) \in [0, T] \times \mathbb{R}$ with $g \in C([0, T], \mathbb{R}^+)$, and $\sup_{t \in [0, T]} |g(t)| = \|g\|$. In addition, it is assumed that $\ell Q_1 < 1$, where

$$Q_1 = \sup_{t \in [0, T]} \left\{ \frac{|\sum_{i=1}^m \omega_i(t) \eta_i^{q-1} (1 - e^{-k\eta_i})|}{k\Gamma(q)} + \frac{|\nu_2(t)| T^{q-1} (1 - e^{-kT})}{k\Gamma(q)} \right. \\ \left. + \frac{|\sum_{i=1}^m \psi_i(t) \eta_i^{q-1}|}{\Gamma(q)} + \frac{|\nu_3(t)| T^{q-1}}{\Gamma(q)} + \frac{|\sum_{i=1}^m \varphi_i(t) \eta_i^{q-2}|}{\Gamma(q-1)} + \frac{|\nu_4(t)| T^{q-2}}{\Gamma(q-1)} \right\}. \quad (4.1)$$

Then the problem (1.1) has at least one solution on $[0, T]$.

Proof. Consider $B_{\bar{r}} = \{u \in \mathcal{P} : \|u\| \leq \bar{r}\}$, where $\bar{r} \geq Q\|g\| + \|\nu_1\|$ and Q is given by (3.3). Introduce the operators $\mathcal{H}_1$ and $\mathcal{H}_2$ on $B_{\bar{r}}$ as

$$(\mathcal{H}_1 u)(t) = \int_0^t e^{-k(t-s)} \left(\int_0^s \frac{(s-x)^{\alpha-2}}{\Gamma(\alpha-1)} f(x, u(x)) dx \right) ds, \\ (\mathcal{H}_2 u)(t) = \nu_1(t) + \sum_{i=1}^m \omega_i(t) \int_0^{\eta_i} e^{-k(\eta_i-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} f(x, u(x)) dx \right) ds \\ + \nu_2(t) \int_0^T e^{-k(T-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} f(x, u(x)) dx \right) ds \\ + \sum_{i=1}^m \psi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-2}}{\Gamma(q-1)} f(s, u(s)) ds + \nu_3(t) \int_0^T \frac{(T-s)^{q-2}}{\Gamma(q-1)} f(s, u(s)) ds \\ + \sum_{i=1}^m \varphi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-3}}{\Gamma(q-2)} f(s, u(s)) ds + \nu_4(t) \int_0^T \frac{(T-s)^{q-3}}{\Gamma(q-2)} f(s, u(s)) ds.$$

For $u, v \in B_{\bar{r}}$, it is easy to verify that $\|\mathcal{H}_1 u + \mathcal{H}_2 v\| \leq Q\|g\| + \|\nu_1\|$. Thus, $\mathcal{H}_1 u + \mathcal{H}_2 v \in B_{\bar{r}}$. Using the assumption (A_1) and (4.1), we can get $\|\mathcal{H}_2 u - \mathcal{H}_2 v\| \leq \ell Q_1 \|u - v\|$, which implies that $\mathcal{H}_2$ is a contraction in view of the given condition: $\ell Q_1 < 1$.

Notice that continuity of f implies that the operator $\mathcal{H}_1$ is continuous. Also, $\mathcal{H}_1$ is uniformly bounded on $B_{\bar{r}}$ as

$$\|\mathcal{H}_1 u\| \leq \frac{(1 - e^{-kT}) T^{q-1} \|g\|}{k\Gamma(q)}.$$

In the last step, it will be shown that the operator $\mathcal{H}_1$ is compact. Fixing $\sup_{(t, u) \in [0, T] \times B_{\bar{r}}} |f(t, u)| = f_{\bar{r}}$ and for $t_1, t_2 \in [0, T]$, we obtain

$$|(\mathcal{H}_1 u)(t_2) - (\mathcal{H}_1 u)(t_1)| \\ = f_{\bar{r}} \left| (e^{-kt_2} - e^{-kt_1}) \int_0^{t_1} e^{ks} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} dx \right) ds + \int_{t_1}^{t_2} e^{-k(t_2-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} dx \right) ds \right| \\ \leq \frac{|1 - e^{-k(t_2-t_1)}|}{k\Gamma(q)} \left(|t_1^{q-1} (1 - e^{-kt_1})| + t_2^{q-1} \right) f_{\bar{r}} \rightarrow 0 \text{ as } t_2 - t_1 \rightarrow 0, \text{ independently of } u \in B_{\bar{r}}.$$

This implies that $\mathcal{H}_1$ is relatively compact on $B_{\bar{r}}$. Hence, by the Arzelà-Ascoli Theorem, the operator $\mathcal{H}_1$ is compact on $B_{\bar{r}}$. Thus all the assumptions of Lemma 4.1 are satisfied. In consequence, the conclusion of Lemma 4.1 applies and hence the problem (1.1) has at least one solution on $[0, T]$. $\square$

Our second existence result is based on the following (Schaefer like) fixed point theorem.

Lemma 4.3 [20] Let X be a Banach space. Assume that $\Omega : X \rightarrow X$ is a completely continuous operator and the set $Y = \{u \in X \mid u = \mu\Omega u, 0 < \mu < 1\}$ is bounded. Then Ω has a fixed point in X .

Theorem 4.4 Let $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function. Assume that there exists a positive constant L_1 such that $|f(t, u(t))| \leq L_1$ for $t \in [0, T]$, $u \in \mathbb{R}$. Then the boundary value problem (1.1) has at least one solution on $[0, T]$.

Proof. In the first step, we show that the operator $\mathcal{H}$ defined by (3.2) is completely continuous. Observe that continuity of $\mathcal{H}$ follows from the continuity of f . For a positive constant r , let $B_r = \{u \in \mathcal{P} : \|u\| \leq r\}$ be a bounded ball in $\mathcal{P}$. Then for $t \in [0, T]$, we have

$$\begin{aligned} \|(\mathcal{H}u)\| &\leq \sup_{t \in [0, T]} \left\{ |\nu_1(t)| + L_1 \left[\int_0^t e^{-k(t-s)} \left(\int_0^s \frac{(s-x)^{\alpha-2}}{\Gamma(\alpha-1)} dx \right) ds \right. \right. \\ &\quad + \left| \sum_{i=1}^m \omega_i(t) \int_0^{\eta_i} e^{-k(\eta_i-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} dx \right) ds \right. \\ &\quad + |\nu_2(t)| \int_0^T e^{-k(T-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} dx \right) ds \\ &\quad + \left| \sum_{i=1}^m \psi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-2}}{\Gamma(q-1)} |f(s, u(s))| ds + |\nu_3(t)| \int_0^T \frac{(T-s)^{\alpha-2}}{\Gamma(\alpha-1)} ds \right. \\ &\quad \left. \left. + \left| \sum_{i=1}^m \varphi_i(t) \int_0^{\eta_i} \frac{(\eta_i-s)^{q-3}}{\Gamma(q-2)} ds + |\nu_4(t)| \int_0^T \frac{(T-s)^{q-3}}{\Gamma(q-2)} ds \right] \right\} \leq \|\nu_1\| + L_1 Q, \end{aligned}$$

where Q is defined by (3.3).

Next we show that the operator $\mathcal{H}$ maps bounded sets into equicontinuous sets of $\mathcal{P}$. Let $\tau_1, \tau_2 \in [0, T]$ with $\tau_1 < \tau_2$ and $u \in B_r$. Then we have

$$\begin{aligned} &|(\mathcal{H}u)(\tau_2) - (\mathcal{H}u)(\tau_1)| \\ &\leq \left| \frac{\beta_3}{k^2 \delta_3} (e^{-k\tau_2} - e^{-k\tau_1}) \right| + \left| \frac{(\tau_2 - \tau_1)}{\lambda_2} \left(\beta_2 + \frac{\beta_3 \delta_2}{k \delta_3} \right) \right| \\ &+ \left| \frac{1 - e^{-k(\tau_2 - \tau_1)}}{k \Gamma(q)} \left(|t_1^{q-1} (1 - e^{-k t_1})| + t_2^{q-1} \right) L_1 \right| \\ &+ \sum_{i=1}^m \left[\left| \frac{c_i (e^{-k\tau_2} - e^{-k\tau_1})}{\delta_3} \right| + \left| \frac{k(\tau_2 - \tau_1)}{\lambda_2 \delta_3} (b_i \delta_3 - \delta_2 c_i) \right| \right] \frac{L_1 \eta_i^{q-1}}{k \Gamma(q)} (1 - e^{-k \eta_i}) \\ &+ \left[\left| \frac{\gamma_3 (e^{-k\tau_2} - e^{-k\tau_1})}{\delta_3} \right| + \left| \frac{k(\tau_2 - \tau_1)}{\lambda_2 \delta_3} (\gamma_2 \delta_3 - \delta_2 \gamma_3) \right| \right] \frac{L_1 T^{q-1}}{k \Gamma(q)} (1 - e^{-k T}) \\ &+ \sum_{i=1}^m \left[\left| \frac{c_i (e^{-k\tau_2} - e^{-k\tau_1})}{k \delta_3} \right| + \left| \frac{(\tau_2 - \tau_1)}{\lambda_2 \delta_3} (b_i \delta_3 - \delta_2 c_i) \right| \right] \frac{L_1 \eta_i^{q-1}}{\Gamma(q)} \\ &+ \left[\left| \frac{\gamma_3 (e^{-k\tau_2} - e^{-k\tau_1})}{k \delta_3} \right| + \left| \frac{(\tau_2 - \tau_1)}{\lambda_2 \delta_3} (\gamma_2 \delta_3 - \delta_2 \gamma_3) \right| \right] \frac{L_1 T^{q-1}}{\Gamma(q)} \\ &+ \sum_{i=1}^m \left[\left| \frac{c_i (e^{-k\tau_2} - e^{-k\tau_1})}{k^2 \delta_3} \right| + \left| \frac{c_i \delta_2 (\tau_2 - \tau_1)}{k \lambda_2 \delta_3} \right| \right] \frac{L_1 \eta_i^{q-2}}{\Gamma(q-1)} \\ &+ \left[\left| \frac{\gamma_3 (e^{-k\tau_2} - e^{-k\tau_1})}{k^2 \delta_3} \right| + \left| \frac{\gamma_3 \delta_2 (\tau_2 - \tau_1)}{k \lambda_2 \delta_3} \right| \right] \frac{L_1 T^{q-2}}{\Gamma(q-1)}. \end{aligned}$$

As $\tau_2 - \tau_1 \rightarrow 0$, the right-hand side of the above inequality tends to zero independently of $u \in B_r$. Therefore, by the Arzelà-Ascoli theorem, the operator $\mathcal{H} : \mathcal{P} \rightarrow \mathcal{P}$ is completely continuous.

Finally, we consider the set $V = \{u \in \mathcal{P} : u = \mu \mathcal{H}u, 0 < \mu < 1\}$ and show that V is bounded. For $u \in V$ and $t \in [0, T]$, we get

$$\|u\| \leq L_1 Q + \|\nu_1\|.$$

Therefore, V is bounded. Hence, by Lemma 4.3, the problem (1.1) has at least one solution on $[0, T]$. $\square$

Now we show the existence of solutions for the problem (1.1) via Leray-Schauder nonlinear alternative for single valued maps.

Lemma 4.5 (Leray-Schauder alternative [21]). *Let E be a Banach space, C a closed, convex subset of E , U an open subset of C and $0 \in U$. Suppose that $\mathcal{H} : \overline{U} \rightarrow C$ is a continuous, compact (that is, $\mathcal{H}(\overline{U})$ is a relatively compact subset of C) map. Then either*

- (i) $\mathcal{H}$ has a fixed point in $\overline{U}$, or
- (ii) there is a $u \in \partial U$ (the boundary of U in C) and $\lambda \in (0, 1)$ with $u = \lambda \mathcal{H}(u)$.

Theorem 4.6 *Let $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function. Assume that*

- (A₃) *there exist functions $p \in C([0, T], \mathbb{R}^+)$, and nondecreasing function $\Psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $|f(t, u)| \leq p(t)\Psi(\|u\|)$, $\forall (t, u) \in [0, T] \times \mathbb{R}$;*
- (A₄) *there exists a constant $\overline{M} > 0$ such that*

$$\frac{\overline{M}}{\|\nu_1\| + \|p\|\Psi(\|\overline{M}\|)Q} > 1. \quad (4.2)$$

Then the boundary value problem (1.1) has at least one solution on $[0, T]$.

Proof. We establish the proof in several steps. Firstly we show that the operator $\mathcal{H} : \mathcal{P} \rightarrow \mathcal{P}$ defined by (3.2) maps bounded sets into bounded sets in $C([0, T], \mathbb{R})$. For the positive number r , let $B_r = \{u \in C([0, T], \mathbb{R}) : \|u\| \leq r\}$, be a bounded set in $C([0, T], \mathbb{R})$. Then

$$\begin{aligned} |\mathcal{H}u(t)| &\leq |\nu_1(t)| + \int_0^t e^{-k(t-s)} \left(\int_0^s \frac{(s-x)^{\alpha-2}}{\Gamma(\alpha-1)} p(x)\Psi(\|u\|)dx \right) ds \\ &\quad + \left| \sum_{i=1}^m \omega_i(t) \right| \int_0^{\eta_i} e^{-k(\eta_i-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} p(x)\Psi(\|u\|)dx \right) ds \\ &\quad + |\nu_2(t)| \int_0^T e^{-k(T-s)} \left(\int_0^s \frac{(s-x)^{q-2}}{\Gamma(q-1)} p(x)\Psi(\|u\|)dx \right) ds \\ &\quad + \left| \sum_{i=1}^m \psi_i(t) \right| \int_0^{\eta_i} \frac{(\eta_i-s)^{q-2}}{\Gamma(q-1)} p(s)\Psi(\|u\|)ds + |\nu_3(t)| \int_0^T \frac{(T-s)^{\alpha-2}}{\Gamma(\alpha-1)} p(s)\Psi(\|u\|)ds \\ &\quad + \left| \sum_{i=1}^m \varphi_i(t) \right| \int_0^{\eta_i} \frac{(\eta_i-s)^{q-3}}{\Gamma(q-2)} p(s)\Psi(\|u\|)ds + |\nu_4(t)| \int_0^T \frac{(T-s)^{q-3}}{\Gamma(q-2)} p(s)\Psi(\|u\|)ds \\ &\leq \|\nu_1\| + \|p\|\Psi(\|u\|)Q, \end{aligned}$$

which, on taking the norm for $t \in [0, T]$, yields $\|\mathcal{H}u\| \leq \|\nu_1\| + \|p\|\Psi(r)Q$.

Next we show that $\mathcal{H}$ maps bounded sets into equicontinuous sets of $C([0, T], \mathbb{R})$. Let $t_1, t_2 \in [0, T]$ with $t_1 < t_2$ and $u \in B_r$, where B_r is a bounded set of $C([0, T], \mathbb{R})$. Then we obtain

$$\begin{aligned} |\mathcal{H}u(t_2) - \mathcal{H}u(t_1)| &\leq |\nu_1(t_2) - \nu_1(t_1)| + \|p\|\Psi(r) \left[\int_0^{t_1} \left| e^{-k(t_2-s)} - e^{-k(t_1-s)} \right| \frac{s^{q-1}}{\Gamma(q)} ds + \int_{t_1}^{t_2} e^{-k(t_2-s)} \frac{s^{q-1}}{\Gamma(q)} ds \right] \end{aligned}$$

$$\begin{aligned}
& + \sum_{i=1}^m |\omega_i(t_2) - \omega_i(t_1)| \int_0^{\eta_i} e^{-k(\eta_i-s)} \frac{s^{q-1}}{\Gamma(q)} ds + |\nu_2(t_2) - \nu_2(t_1)| \int_0^T e^{-k(T-s)} \frac{s^{q-1}}{\Gamma(q)} ds \\
& + \sum_{i=1}^m |\psi_i(t_2) - \psi_i(t_1)| \frac{\eta_i^{q-1}}{\Gamma(q)} + |\nu_3(t_2) - \nu_3(t_1)| \frac{T^{q-1}}{\Gamma(q)} \\
& + \sum_{i=1}^m |\varphi_i(t_2) - \varphi_i(t_1)| \frac{\eta_i^{q-2}}{\Gamma(q-1)} + |\nu_4(t_2) - \nu_4(t_1)| \frac{T^{q-2}}{\Gamma(q-1)} \Big].
\end{aligned}$$

Obviously the right hand side of the above inequality tends to zero independently of $u \in B_r$ as $t_2 - t_1 \rightarrow 0$. As $\mathcal{H}$ satisfies the above assumptions, therefore it follows by the Arzelà-Ascoli theorem that $\mathcal{H} : C([0, T], \mathbb{R}) \rightarrow C([0, T], \mathbb{R})$ is completely continuous.

The result will follow from the Leray-Schauder nonlinear alternative (Lemma 4.5) once we have proved the boundness of the set of all solutions to the equation $u = \lambda \mathcal{H}u$ for $\lambda \in [0, 1]$.

Let u be a solution. Then, for $t \in [0, T]$, and using the computations in proving that $\mathcal{H}$ is bounded, we have

$$|u(t)| = |\lambda(\mathcal{H}u)(t)| \leq \|\nu_1\| + \|p\|\Psi(\|r\|)Q,$$

which implies that

$$\frac{\|u\|}{\|\nu_1\| + \|p\|\Psi(r)Q} \leq 1.$$

In view of (A₄), there exists $\overline{M}$ such that $\|u\| \neq \overline{M}$. Let us set

$$U = \{u \in C([0, T], \mathbb{R}) : \|u\| < \overline{M}\}.$$

Note that the operator $\mathcal{H} : \overline{U} \rightarrow C([0, T], \mathbb{R})$ is continuous and completely continuous. From the choice of U , there is no $u \in \partial U$ such that $u = \lambda \mathcal{H}(u)$ for some $\lambda \in (0, 1)$. Consequently, by the nonlinear alternative of Leray-Schauder type (Lemma 4.5), we deduce that $\mathcal{H}$ has a fixed point $u \in \overline{U}$ which is a solution of the problem (1.1). This completes the proof. $\square$

In our final existence result for the problem (1.1), we apply Leray-Schauder degree theory [21].

Theorem 4.7 *Let $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$. Assume that there exist constants $0 \leq \overline{k} < \frac{1}{Q}$, where Q is given by (3.3) and $\overline{M} > 0$ such that $|f(t, u(t))| \leq \overline{k}\|u\| + \overline{M}$ for all $t \in [0, T], u \in \mathbb{R}$. Then the boundary value problem (1.1) has at least one solution.*

Proof. In view of the fixed point problem (3.1), we just need to show that there exists at least one solution $u \in \mathbb{R}$ for (3.1). Define a suitable ball $B_R \subset C([0, T])$ with radius $R > 0$ as

$$B_R = \{u \in C([0, T]) : \|u\| \leq R\},$$

where R will be fixed later. Then, it is sufficient to show that $\mathcal{H} : \overline{B}_R \rightarrow C$ satisfies

$$u \neq \lambda \mathcal{H}u, \forall u \in \partial B_R \text{ and } \forall \lambda \in [0, 1]. \quad (4.3)$$

Let us set

$$\Phi(\lambda, u) = \lambda \mathcal{H}u, u \in C, \lambda \in [0, 1].$$

Then, by the Arzelà-Ascoli Theorem, $\omega_\lambda(u) = u - \Phi(\lambda, u) = u - \mathcal{H}u$ is completely continuous. If (4.3) is true, then the following Leray-Schauder degrees are well defined and by the homotopy invariance of topological degree, it follows that

$$\begin{aligned}
deg(\omega_\lambda, B_R, 0) &= deg(I - \lambda \mathcal{H}, B_R, 0) = deg(\omega_1, B_R, 0) \\
&= deg(\omega_0, B_R, 0) = deg(I, B_R, 0) = 1 \neq 0, 0 \in B_R,
\end{aligned} \quad (4.4)$$

where I denotes the unit operator. By the nonzero property of Leray-Schauder degree, $\omega_1(t) = u - \lambda \mathcal{H}u = 0$ for at least one $u \in B_R$. To prove (4.3), we assume that $u = \lambda \mathcal{H}u = 0$ for some $\lambda \in [0, 1]$ and for all $t \in [0, 1]$. Then, as before, one can obtain

$$|u(t)| = |\lambda(\mathcal{H}u)(t)| \leq \|\nu_1\| + (\bar{k}\|u\| + \bar{M})Q,$$

which implies that

$$\|u\| \leq \frac{\|\nu_1\| + \bar{M}Q}{1 - \bar{k}Q}.$$

Letting $R = \frac{\|\nu_1\| + \bar{M}Q}{1 - \bar{k}Q} + 1$, (4.3) holds. This completes the proof. $\square$

5 Examples

Example 5.1 Consider the following anti-periodic multi-point fractional boundary value problem:

$$\begin{cases} ({}^c D^{5/2} + 2{}^c D^{3/2})u(t) = f(t, u(t)), & t \in [0, 2], \\ u(0) + u(1/4) + 1/2u(3/4) - u(5/4) + u(7/4) - u(2) = 1, \\ u'(0) - u'(1/4) + u'(3/4) - 1/2u'(5/4) + 1/4u'(7/4) - 1/2u'(2) = 2, \\ 1/2u''(0) + 1/4u''(1/4) - u'(3/4) + u''(5/4) + u''(7/4) + u''(2) = 1. \end{cases} \quad (5.1)$$

$T = 2$, $0 < \eta_1 = 1/4$, $\eta_2 = 3/4$, $\eta_3 = 5/4$, $\eta_4 = 7/4 < T$, $k = 2$, $\alpha_1 = 1$, $\gamma_1 = -1$, $\alpha_2 = 1$, $\gamma_2 = -1/2$, $\alpha_3 = 1/2$, $\gamma_3 = 1$, $a_1 = 1$, $a_2 = 1/2$, $a_3 = -1$, $a_4 = 1$, $b_1 = -1$, $b_2 = 1$, $b_3 = -1/2$, $b_4 = 1/4$, $c_1 = 1/4$, $c_2 = -1$, $c_3 = 1$, $c_4 = 1$, $\beta_1 = 1$, $\beta_2 = 2$, $\beta_3 = 1$, $Q = 193.8001283$, $Q_1 = 192.7557672$, where Q and Q_1 are respectively given by (3.3) and (4.1).

- For the applicability of Theorem 3.1 to the problem (5.1), let $f(t, u(t)) = \frac{1}{200} \tan^{-1} u(t) + \cos t$ in (5.1). It is easy to find that $\ell = 1/200$ as $|f(t, u) - f(t, v)| \leq \frac{1}{200} |u - v|$ and that $\ell Q \approx 0.9690006415 < 1$. Clearly all the conditions of Theorem 3.1 are satisfied. Hence we deduce by the conclusion of Theorem 3.1 that there exists a unique solution for problem (5.1) on $[0, 2]$.
- For the illustration of Theorem 4.2, we take $f(t, u(t)) = \frac{1}{200} \tan^{-1} u(t) + \cos t$ in (5.1). Obviously $|f(t, u)| \leq \pi/400 + \cos t = g(t)$, $\|g\| = \frac{\pi+400}{400}$ and $\ell Q_1 \approx 0.9637788360 < 1$. Thus all the conditions of Theorem 4.2 are satisfied. Hence the conclusion of Theorem 4.2 applies to the problem (5.1).
- Let us take $f(t, u(t)) = \frac{1}{2} \sin u(t) + e^{-t} + 2$ in (5.1) such that $|f(t, u)| \leq 7/2$. Thus, by the conclusion of Theorem 4.4, we deduce that that that problem (5.1) has at least one solution on $[0, 2]$.
- For the elaboration of Theorem 4.6, let $f(t, u(t)) = \frac{1}{\sqrt{t^2+4}} \frac{|u|}{1+|u|}$ in (5.1). Clearly $|f(t, u)| \leq \frac{1}{\sqrt{t^2+4}}$. Fixing $\Psi(\|u\|) = 1$, $p(t) = \frac{1}{\sqrt{t^2+4}}$, we find by (4.2) that $\bar{M} > 136.8973405$. In consequence, it follows by the conclusion of Theorem 4.6 that the problem (5.1) has at least one solution on $[0, 2]$.
- In order to demonstrate the application of Theorem 4.7, we take $f(t, u(t)) = \frac{1}{400} \sin(2u) + 1$ in (5.1) and note that $|f(t, u)| \leq \|u\|/200 + 1$. Clearly $\bar{k} = \frac{1}{200}$, $\bar{M} = 1$ and $0 < \bar{k} = \frac{1}{200} < \frac{1}{193.8001283} = \frac{1}{Q}$. As all the conditions of Theorem 4.7 hold true, so the conclusion of Theorem 4.7 applies and hence there exists at least one solution for the problem (5.1) on $[0, 2]$.

6 Concluding remarks

In this paper, we have discussed the existence and uniqueness of solutions for a new class of boundary value problems consisting of sequential fractional differential equations and a combination of anti-periodic and multi-point boundary conditions by means of appropriate tools of the fixed point theory. Our results are not only new in the given configuration but also correspond to the ones for certain new problems for particular values of the parameters involved in the problem at hand. For example, by taking $a_i, b_i, c_i = 0, i = 1, 2, \dots, m$, $\beta_j = 0, \alpha_j = 1 = \gamma_j, j = 1, 2, 3$ in the results of this paper, we obtain the new ones for the problem of sequential fractional differential equations with anti-periodic boundary conditions:

$$\begin{aligned}({}^c D^q + k {}^c D^{q-1})u(t) &= f(t, u(t)), \quad 2 < q \leq 3, \quad 0 < t < T, \\ u(0) + u(T) &= 0, \quad u'(0) + u'(T) = 0, \quad u''(0) + u''(T) = 0.\end{aligned}$$

Similarly the results for nonlocal multi-point boundary value problems associated with either left-end point ($x = 0$) or right-end point ($x = T$) of the interval $[0, T]$ can be obtained by taking $\gamma_j = 0$ and $\alpha_j = 0$ ($j = 1, 2, 3$) respectively in our results. So the present work is a useful contribution to the existing literature on the topic.

References

- [1] A. Alsaedi, S. Sivasundaram, B. Ahmad, On the generalization of second order nonlinear anti-periodic boundary value problems, *Nonlinear Stud.* **16** (2009), 415-420.
- [2] A. Bitsadze, A. Samarskii, On some simple generalizations of linear elliptic boundary problems, *Russian Acad. Sci. Dokl. Math.* **10** (1969), 398-400.
- [3] M. Klimek, Sequential fractional differential equations with Hadamard derivative, *Commun. Nonlinear Sci. Numer. Simul.* **16** (2011), 4689-4697.
- [4] B. Ahmad, J.J. Nieto, Anti-periodic fractional boundary value problems with nonlinear term depending on lower order derivative, *Fract. Calc. Appl. Anal.* **15** (2012), 451-462.
- [5] J. Cao, Q. Yang, Z. Huang, Existence of anti-periodic mild solutions for a class of semilinear fractional differential equations, *Commun. Nonlinear Sci. Numer. Simul.* **17** (2012), no. 1, 277-283.
- [6] B. Ahmad, J.J. Nieto, Sequential fractional differential equations with three-point boundary conditions, *Comput. Math. Appl.* **64** (2012), no. 10, 3046-3052.
- [7] M. Benchohra, N. Hamidi, J. Henderson, Fractional differential equations with anti-periodic boundary conditions, *Numer. Funct. Anal. Optim.* **34** (2013), 404-414.
- [8] B. Ahmad, J.J. Nieto, A class of differential equations of fractional order with multi-point boundary conditions, *Georgian Math. J.* **21** (2014), 243-248.
- [9] J. Tariboon, T. Sitthiwiratham, S.K. Ntouyas, Existence results for fractional differential inclusions with multi-point and fractional integral boundary conditions, *J. Comput. Anal. Appl.* **17** (2014), 343-360.
- [10] J.R. Wang, Y. Zhou, M. Feckan, On the nonlocal Cauchy problem for semilinear fractional order evolution equations, *Cent. Eur. J. Math.* **12** (2014), 911-922.
- [11] B. Ahmad, S.K. Ntouyas, On a coupled system of fractional differential equations with coupled nonlocal and integral boundary conditions, *Chaos Solitons Fractals* **83** (2016), 234-241.
- [12] S. Aljoudi, B. Ahmad, J.J. Nieto, A. Alsaedi, A coupled system of Hadamard type sequential fractional differential equations with coupled strip conditions, *Chaos Solitons Fractals* **91** (2016), 39-46.
- [13] S. Xie, Y. Xie, Positive solutions of a system for nonlinear singular higher-order fractional differential equations with fractional multi-point boundary conditions, *Bound. Value Probl.* (2016), 2016:134.
- [14] L. Peng, Y. Zhou, Bifurcation from interval and positive solutions of the three-point boundary value problem for fractional differential equations, *Appl. Math. Comput.* **257** (2015), 458-466.
- [15] J. Jiang, Solvability of anti-periodic boundary value problem for coupled system of fractional p -Laplacian equation, *Adv. Difference Equ.* **2015:305** (2015), 11 pp.

- [16] Y. Li, A. Qi, Positive solutions for multi-point boundary value problems of fractional differential equations with p-Laplacian, *Math. Methods Appl. Sci.* **39** (2016), 1425–1434.
- [17] M.H. Aqlan, A. Alsaedia, B. Ahmad, J.J. Nieto, Existence theory for sequential fractional differential equations with anti-periodic type boundary conditions, *Open Mathematics*, accepted.
- [18] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, 204. Elsevier Science B.V., Amsterdam, 2006.
- [19] Y. Zhou, Yong. *Basic theory of fractional differential equations*, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2014.
- [20] Smart D.R., *Fixed point theorems*, Cambridge University Press, 1980
- [21] Granas A., Dugundji J., *Fixed point theory*, Springer-Verlag, New York, 2003