

УДК 517.53

DOI 10.46698/p5608-0614-8805-b

UNICITY ON ENTIRE FUNCTIONS CONCERNING  
THEIR DIFFERENCE OPERATORS AND DERIVATIVES

Rajeshwari, S.<sup>1</sup> and Sheebakousar, B.<sup>2</sup>

<sup>1</sup>Department of Mathematics, Bangalore Institute of Technology,  
Vishweshwarapura, Basavanagudi, Bangalore-560004, India;

<sup>2</sup>Presidency University, School of Engineering,  
Itagalpura, Rajanakunte, Yelahanka, Bangalore-560 064, India

E-mail: rajeshwari.s@presidencyuniversity.in,  
rajeshwaripreetham@gmail.com; sheeba.buzurg@gmail.com

**Abstract.** In this paper we study the uniqueness of entire functions concerning their difference operator and derivatives. The idea of entire and meromorphic functions relies heavily on this direction. Rubel and Yang considered the uniqueness of entire function and its derivative and proved that if  $f(z)$  and  $f'(z)$  share two values  $a, b$  counting multiplicities then  $f(z) \equiv f'(z)$ . Later, Li Ping and Yang improved the result given by Rubel and Yang and proved that if  $f(z)$  is a non-constant entire function and  $a, b$  are two finite distinct complex values and if  $f(z)$  and  $f^{(k)}(z)$  share  $a$  counting multiplicities and  $b$  ignoring multiplicities then  $f(z) \equiv f^{(k)}(z)$ . In recent years, the value distribution of meromorphic functions of finite order with respect to difference analogue has become a subject of interest. By replacing finite distinct complex values by polynomials, we prove the following result: Let  $\Delta f(z)$  be transcendental entire functions of finite order,  $k \geq 0$  be integer and  $P_1$  and  $P_2$  be two polynomials. If  $\Delta f(z)$  and  $f^{(k)}$  share  $P_1$  CM and share  $P_2$  IM, then  $\Delta f \equiv f^{(k)}$ . A non-trivial proof of this result uses Nevanlinna's value distribution theory.

**Key words:** difference operator, shared values, finite order, uniqueness, entire function, polynomials.

**AMS Subject Classification:** 30D35, 39A32.

**For citation:** Rajeshwari, S. and Sheebakousar, B. Unicity on Entire Functions Concerning Their Difference Operators and Derivatives, *Vladikavkaz Math. J.*, 2023, vol. 25, no. 1, pp. 81–92. DOI: 10.46698/p5608-0614-8805-b.

## 1. Introduction and Main Results

The reader is presumed to be familiar with the fundamental notations and conclusions of Nevanlinna's value distribution theory of meromorphic functions [1, 2].  $S(r, f)$  means that  $S(r, f) = o(T(r, f))$  as  $r \rightarrow \infty$  outside of a possible exceptional set of finite logarithmic measure, and

$$\lambda(f) = \limsup_{r \rightarrow \infty} \frac{\log^+ N(r, \frac{1}{f})}{\log r}, \quad \delta(P_1, f) = \liminf_{r \rightarrow \infty} \frac{m(r, \frac{1}{f-a})}{T(r, f)} = 1 - \limsup_{r \rightarrow \infty} \frac{N(r, \frac{1}{f})}{T(r, f)}$$

stand for the exponents of convergence of zero sequence of  $f$  and the deficiency of  $f$  at the point  $a$ , respectively. For a nonconstant meromorphic function  $h$ , we denote by  $T(r, h)$  the Nevanlinna characteristic of  $h$  and by  $S(r, h)$  any quantity satisfying  $S(r, h) = o(T(r, h))$ , as  $r$  runs to infinity outside of a set  $E \subset (0, +\infty)$  of finite linear measure. We say that  $h$  is a small

function of  $f$ , if  $T(r, h) = S(r, f)$ . In the sequel, we denote by  $I$  a set of infinite linear measure not necessarily the same in all its occurrences.

We say that  $f$  and  $g$  share the value  $a$  IM (ignoring multiplicities), if  $f$  and  $g$  have the same  $a$  point. If  $f$  and  $g$  have the same  $a$  point with the same multiplicities, then we say  $f, g$  share the value  $a$  CM (counting multiplicities).

**DEFINITION 1** [3]. Let  $k$  be a nonnegative integer or infinity. For  $a \in \mathbb{C} \cup \{\infty\}$ , we denote by  $E_k(a; f)$  the set of all  $a$  points of  $f(z)$  where an  $a$  point of multiplicity  $m$  is counted  $m$  times if  $m \leq k$  and  $k + 1$  times if  $m > k$ . If  $E_k(a; f) = E_k(a; g)$ , then we say that  $f, g$  share the value  $a$  with weight  $k$ .

We write  $f$  and  $g$  share  $(a, k)$  to mean that  $f, g$  share the value  $a$  with weight  $k$ .

Rubel and Yang Chung-Chun [4] considered the uniqueness of an entire function and its derivative. They proved the following.

**Theorem 1.** *Let  $f(z)$  be a non-constant entire function, let  $a, b$  be two finite distinct values. If  $f(z)$  and  $f'(z)$  share  $a, b$  CM, then  $f(z) \equiv f'(z)$ .*

Li Ping and Yang Chung-Chun [5] improved Theorem 1 and proved.

**Theorem 2.** *Let  $f(z)$  be a non-constant entire function, and let  $a, b$  be two finite distinct complex values. If  $f(z)$  and  $f^{(k)}(z)$  share  $a$  CM, and share  $b$  IM. Then  $f(z) \equiv f^{(k)}(z)$ .*

The value distribution of meromorphic functions of finite order with respect to difference analogue has become a subject of some interests, see [6–16].

**Theorem 3** [17]. *Suppose that  $f(z)$  and  $g(z)$  are nonconstant meromorphic functions. If  $f, g$  share  $0, 1, \infty$  CM and  $\overline{N}(r, \frac{1}{f}) + \overline{N}(r, f) < (d + o(1))T(r, f)$  for  $r \in I$  and  $r \in \infty$ , where  $d$  is a positive number satisfying  $0 < d < \frac{1}{2}$ , which  $I \subset (0, +\infty)$  is a subset of infinite linear measure, then  $f = g$  or  $f.g = 1$ .*

**Theorem 4** [4]. *Let  $f$  be a nonconstant entire function. If  $f$  shares two distinct finite values CM with  $f'$ , then  $f \equiv f'$ .*

More results on uniqueness of  $f'$  with its  $n$ -th derivative  $f^{(n)}$  were obtained by several authors (see [18–20]). In view of the progress on the difference analogues of classical Nevanlinna theory of meromorphic functions [21, 22], it is quite natural to investigate the uniqueness problems of meromorphic functions and their difference operators (see [23–26]).

**EXAMPLE 1.** Let  $f(z) = e^{Az}$ , where  $A \neq 0$  is a constant. Then  $f^{(k)} = A^k e^{Az}$  and  $\Delta f = f(z+1) - f(z) = (e^A - 1)e^{Az}$ . Clearly,  $\Delta(f)$  and  $f^{(k)}$  share 0 CM and  $\infty$  IM, and that  $\mu = 1$ . We can choose  $A$  such that  $e^A - 1 \neq A^k$ , and so  $f' \not\equiv \Delta(f)$ .

**Theorem 5** [27]. *Let  $f(z)$  be a transcendental entire function of finite order, let  $\eta \neq 0$  be a finite complex number,  $n \geq 1, k \geq 0$  be two integers and let  $a, b$  be two distinct finite complex values. If  $f(z)$  and  $(\Delta_\eta^n f(z))^{(k)}$  share  $a$  CM and share  $b$  IM, then  $f(z) \equiv (\Delta_\eta^n f(z))^{(k)}$ .*

**Lemma 1** [9]. *Let  $\Delta f$  be a nonconstant meromorphic function of finite order, let  $\eta \neq 0$  be a finite complex number. Then*

$$m\left(r, \frac{\Delta f(z + \eta)}{\Delta f}\right) = S(r, \Delta f),$$

for all  $r$  outside of a possible exceptional set  $E$  with finite logarithmic measure.

**Lemma 2** [28, Lemma 4.3]. *Let  $\Delta f$  be a nonconstant meromorphic function. Suppose that the polynomials  $P_j, j = 0, 1, \dots, q, q > p$ , and let  $P(\Delta f) = a_0(\Delta f)^p + a_1(\Delta f)^{p-1} + \dots + a_p$  ( $a_0 \neq 0$ ) is a polynomial of degree  $p$  with constant coefficient  $a_j, j = 0, 1, \dots, p$ . Then*

$$m\left(r, \frac{P(\Delta f)(\Delta f)'}{(\Delta f - P_1)(\Delta f - P_2) \dots (\Delta f - P_q)}\right) = S(r, \Delta f).$$

**Lemma 3.** Let  $\Delta f$  and  $\Delta g$  be two non constant entire functions, and let  $P_1, P_2$  be two polynomials. If

$$H = \frac{\Delta f'}{(\Delta f - P_1)(\Delta f - P_2)} - \frac{\Delta g'}{(\Delta g - P_1)(\Delta g - P_2)} \equiv 0,$$

and  $\Delta f$  and  $\Delta g$  share  $P_1$  CM, and share  $P_2$  IM, then either

$$2T(r, \Delta f) \leq \bar{N}\left(r, \frac{1}{\Delta f - P_1}\right) + \bar{N}\left(r, \frac{1}{\Delta f - P_2}\right) + S(r, \Delta f)$$

or

$$\Delta f \equiv \Delta g.$$

◁ Integrating  $H$  which leads to

$$\frac{\Delta g - P_2}{\Delta g - P_1} = C \frac{\Delta f - P_2}{\Delta f - P_1},$$

where  $C$  is a nonzero constant.

If  $C = 1$ , then  $\Delta f \equiv \Delta g$ . If  $C \neq 1$ , then from above, we have

$$\frac{P_1 - P_2}{\Delta g - P_1} \equiv \frac{(C - 1)\Delta f - CP_2 + P_1}{\Delta f - P_1}$$

and

$$T(r, \Delta f) = T(r, \Delta g) + S(r, \Delta f) + S(r, \Delta g).$$

Obviously,  $\frac{CP_1 - P_2}{C - 1} \neq a$  and  $\frac{CP_1 - P_2}{C - 1} \neq b$ . It follows that  $N\left(r, \frac{1}{\Delta f - \frac{CP_1 - P_2}{C - 1}}\right) = 0$ . Then by the Second Fundamental Theorem,

$$\begin{aligned} 2T(r, \Delta f) &= \bar{N}(r, \Delta f) + \bar{N}\left(r, \frac{1}{\Delta f - P_1}\right) + \bar{N}\left(r, \frac{1}{\Delta f - P_2}\right) + \bar{N}\left(r, \frac{1}{\Delta f - \frac{CP_1 - P_2}{C - 1}}\right) \\ &+ S(r, \Delta f) \leq \bar{N}\left(r, \frac{1}{\Delta f - P_1}\right) + \bar{N}\left(r, \frac{1}{\Delta f - P_2}\right) + S(r, \Delta f), \end{aligned}$$

that is

$$2T(r, \Delta f) \leq \bar{N}\left(r, \frac{1}{\Delta f - P_1}\right) + \bar{N}\left(r, \frac{1}{\Delta f - P_2}\right) + S(r, \Delta f). \triangleright$$

**Lemma 4.** Let  $\Delta f$  be a transcendental entire function of finite order,  $k$  be positive integer, let  $P_1$  be a nonzero complex value or constant. If  $\Delta f$  and  $f^{(k)}$  share  $P_1$  CM, and  $N\left(r, \frac{1}{f^{(k)}}\right) = S(r, \Delta f)$ , then one of the following cases must occur:

- $f^{(k)} = He^p$ , where  $p$  is a polynomial, and  $H \not\equiv 0$  is a small function of  $e^p$ .
- $T(r, e^p) = S(r, \Delta f)$ .

◁ Since  $\Delta f$  is a transcendental entire function of finite order,  $\Delta f$  and  $f^{(k)}$  share  $P_1$  CM, then there is a polynomial  $p$  such that

$$\Delta f - P_1 = e^p(f^{(k)}) - P_1e^p. \tag{1}$$

Set  $g = f^{(k)}$ . It follows by (1) that

$$g = (ge^p)^{(k)} - (P_1e^p)^{(k)}. \tag{2}$$

Then we rewrite (2) as

$$1 + \frac{(P_1 e^p)^{(k)}}{g} = D e^p, \quad (3)$$

where

$$D = \frac{(g e^p)^{(k)}}{g e^p}. \quad (4)$$

Note that  $N\left(r, \frac{1}{(f)^{(k)}}\right) = N\left(r, \frac{1}{g}\right) = S(r, f)$ , then by Lemma 1 we have

$$T(r, D) = T\left(r, \frac{(g e^p)^{(k)}}{\Delta g e^p}\right).$$

Next we discuss two cases.

Case 1:  $e^{-p} - D \neq 0$ . Rewrite (3) as

$$g e^p (e^{-p} - D) = (P_1 e^p)^{(k)}. \quad (5)$$

When  $D \equiv 0$ , (5) implies

$$g = H e^p. \quad (6)$$

Here  $H \neq 0$  is a small function of  $e^p$ .

When  $D \neq 0$ , it follows from (5) that  $N\left(r, \frac{1}{e^{-p}-D}\right) = S(r, f)$ . Then use the Second Fundamental Theorem to  $e^p$  we can obtain

$$T(r, e^p) = T(r, e^{-p}) + O(1) \leq \bar{N}(r, e^p) + \bar{N}\left(r, \frac{1}{e^{-p}}\right) + \bar{N}\left(r, \frac{1}{e^{-p}-D}\right) + O(1) = S(r, \Delta f).$$

Case 2:  $e^{-p} - D \equiv 0$ . It implies that  $T(r, e^p) = T(r, e^{-p}) + O(1) = S(r, \Delta f)$ , a contradiction. From above discussions, we get  $T(r, e^p) = S(r, \Delta f)$ .  $\triangleright$

**Theorem 6.** Let  $\Delta f(z)$  be a transcendental entire functions of finite order,  $k$  be integer such that  $k \geq 0$  and let  $P_1$  and  $P_2$  be two polynomials. If  $\Delta f(z)$  and  $f^{(k)}$  share  $P_1$  CM and share  $P_2$  IM, then  $\Delta f \equiv f^{(k)}$ .

$\triangleleft$  If  $\Delta f \equiv f^{(k)}$ , there is nothing to prove. Solve  $\Delta f \not\equiv f^{(k)}$ . Since  $\Delta f$  is a transcendental entire function of finite order,  $\Delta f$  and  $f^{(k)}$  share  $P_1$  CM, then we get

$$\frac{f^{(k)} - P_1}{\Delta f - P_1} = e^Q, \quad (7)$$

where  $Q$  is a polynomial.

Since  $\Delta f$  and  $f^{(k)}$  share  $P_1$  CM and share  $P_2$  IM, then by second fundamental theorem and Lemma 1 we have

$$\begin{aligned} T(r, \Delta f) &\leq \bar{N}\left(r, \frac{1}{\Delta f - P_1}\right) + \bar{N}\left(r, \frac{1}{\Delta f - P_2}\right) + S(r, \Delta f) \\ &\leq \bar{N}\left(r, \frac{1}{\Delta f - P_2}\right) + \bar{N}\left(r, \frac{1}{f^{(k)} - P_2}\right) \leq N\left(r, \frac{1}{\Delta f - f^{(k)}}\right) + S(r, \Delta f) \\ &\leq T(r, \Delta f - f^{(k)}) + S(r, \Delta f) \leq m(r, \Delta f - f^{(k)}) + S(r, \Delta f) \\ &\leq m(r, \Delta f) + m\left(r, 1 - \frac{f^{(k)}}{\Delta f}\right) + S(r, \Delta f) \leq T(r, f) + S(r, f), \end{aligned}$$

$$T(r, \Delta f) = \overline{N} \left( r, \frac{1}{\Delta f - P_1} \right) + \overline{N}(\Delta f - P_2) + S(r, \Delta f). \quad (8)$$

According to (7) and (8) we have

$$T(r, \Delta f) = T(r, \Delta f - f^{(k)}) + S(r, \Delta f) \leq N \left( r, \frac{1}{\Delta f - f^{(k)}} \right) + S(r, \Delta f) \quad (9)$$

and

$$T(r, e^Q) = m(r, e^Q) = m \left( r, \frac{f^{(k)} - P_1}{\Delta f - P_1} \right) \leq m \left( r, \frac{1}{\Delta f - P_1} \right) + S(r, \Delta f). \quad (10)$$

Then it follows from (7) and (9) that

$$\begin{aligned} m \left( r, \frac{1}{\Delta f - P_1} \right) &= m \left( r, \frac{e^Q - 1}{\Delta f - f^{(k)}} \right) \leq m \left( r, \frac{1}{\Delta f - f^{(k)}} \right) + m(r, e^Q - 1) \\ &\leq T(r, e^Q) + S(r, \Delta f). \end{aligned} \quad (11)$$

Then by (10) and (11)

$$T(r, e^Q) = m \left( r, \frac{1}{\Delta f - P_1} \right) + S(r, \Delta f). \quad (12)$$

On the other hand, (1) can be rewritten as

$$\frac{f^{(k)} - \Delta f}{\Delta f - P_1} = e^Q - 1, \quad (13)$$

which implies

$$\overline{N} \left( r, \frac{1}{\Delta f - P_2} \right) \leq \overline{N} \left( r, \frac{1}{e^Q - 1} \right) = T(r, e^Q) + S(r, \Delta f). \quad (14)$$

Then by (8), (12) and (14)

$$\begin{aligned} m \left( r, \frac{1}{\Delta f - P_1} \right) + N \left( r, \frac{1}{\Delta f - P_1} \right) &= \overline{N} \left( r, \frac{1}{\Delta f - P_1} \right) + \overline{N} \left( r, \frac{1}{\Delta f - P_2} \right) + S(r, \Delta f) \\ &\leq \overline{N} \left( r, \frac{1}{\Delta f - P_1} \right) + \overline{N} \left( r, \frac{1}{e^Q - 1} \right) + S(r, \Delta f) \\ &\leq \overline{N} \left( r, \frac{1}{\Delta f - P_1} \right) + m \left( r, \frac{1}{\Delta f - P_1} \right) + S(r, \Delta f), \\ N \left( r, \frac{1}{\Delta f - P_1} \right) &= \overline{N} \left( r, \frac{1}{\Delta f - P_1} \right) + S(r, \Delta f), \end{aligned} \quad (15)$$

and then

$$\overline{N} \left( r, \frac{1}{\Delta f - P_2} \right) = T(r, e^Q) + S(r, \Delta f). \quad (16)$$

Set

$$\phi = \frac{(\Delta f)'(\Delta f - f^{(k)})}{(\Delta f - P_1)(\Delta f - P_2)} \quad (17)$$

and

$$\psi = \frac{f^{(k+1)}(\Delta f - f^{(k)})}{(f^{(k)} - P_1)(f^{(k)} - P_2)}. \quad (18)$$

Easy to know that  $\phi$  is an entire function by Lemma 1 and Lemma 2 we have

$$\begin{aligned} T(r, \phi) &= m(r, \phi) = m\left(r, \frac{(\Delta f)'(\Delta f - f^{(k)})}{(\Delta f - P_1)(\Delta f - P_2)}\right) + S(r, \Delta f) \\ &\leq m\left(r, \frac{(\Delta f)'(\Delta f)}{(\Delta f - P_1)(\Delta f - P_2)}\right) m\left(r, 1 - \frac{f^{(k)}}{\Delta f}\right) + S(r, \Delta f) = S(r, \Delta f), \end{aligned}$$

that is

$$T(r, \phi) = S(r, \Delta f). \quad (19)$$

Obviously Let  $d = P_1 - k(P_1 - P_2)$ ,  $k \neq 0$ , by Lemmas 1 and 2, we obtain

$$\begin{aligned} &m\left(r, \frac{1}{\Delta f}\right) + m\left(r, \frac{1}{(P_2 - P_1)\phi} \left(\frac{(\Delta f)'}{\Delta f - P_1} - \frac{(\Delta f)'}{\Delta f - P_2}\right) \left(\frac{1 - (\Delta f)'}{\Delta f}\right)\right) \\ &\leq m\left(r, \frac{1}{\phi}\right) + m\left(r, \left(\frac{(\Delta f)'}{\Delta f - P_1} - \frac{(\Delta f)'}{\Delta f - P_2}\right)\right) + m\left(r, \frac{(\Delta f)'}{\Delta f}\right) + S(r, \Delta f) = S(r, \Delta) \end{aligned} \quad (20)$$

and

$$\begin{aligned} &m\left(r, \frac{1}{\Delta f - d}\right) = m\left(r, \frac{(\Delta f)'(\Delta f - f^{(k)})}{(\Delta f - P_1)(\Delta f - P_2)(\Delta f - d)}\right) \\ &\leq m\left(r, \frac{1 - f^{(k)}}{\Delta f}\right) + m\left(r, \frac{(\Delta f)'(\Delta f - f^{(k)})}{(\Delta f - P_1)(\Delta f - P_2)(\Delta f - d)}\right) + S(r, \Delta f) = S(r, \Delta f). \end{aligned} \quad (21)$$

Set

$$\phi = \frac{(f)^{(k+1)}}{(f^{(k)} - P_1)(f^{(k)} - P_2)} - \frac{(\Delta f)'}{(\Delta f - P_1)(\Delta f - P_2)}. \quad (22)$$

We discuss two cases

Case 1:  $\phi \equiv 0$ . Integrating both side of (22) which leads to

$$\frac{\Delta f - P_2}{\Delta f - P_1} = c \frac{f^{(k)} - P_2}{f^{(k)} - P_1}, \quad (23)$$

where  $c$  is a non zero constant. Then by Lemma 3 we see that

$$2T(r, \Delta f) \leq \bar{N}\left(r, \frac{1}{\Delta f - P_1}\right) + \bar{N}\left(r, \frac{1}{\Delta f - P_2}\right) + S(r, \Delta f), \quad (24)$$

which contradicts with (8).

Case 2:  $\phi \not\equiv 0$ . By (9), (19) and (22) we can obtain

$$\begin{aligned} m(r, \Delta f) &= m(r, \Delta f - f^{(k)}) + S(r, \Delta f) = m\left(r, \frac{\phi(\Delta f - f^{(k)})}{\phi}\right) + S(r, \Delta f) \\ &= m\left(r, \frac{\psi - \phi}{\phi}\right) + S(r, \Delta f) \leq T\left(r, \frac{\phi}{\psi - \phi}\right) + S(r, \Delta f) \\ &\leq T(r, \psi - \phi) + T(r, \phi) + S(r, \Delta f) \leq T(r, \psi) + \bar{N}\left(r, \frac{1}{\Delta f - P_2}\right) + S(r, \Delta f). \end{aligned} \quad (25)$$

On the other hand

$$\begin{aligned}
 T(r, \psi) &= T\left(r, \frac{f^{(k+1)}(\Delta f - f^{(k)})}{(f^{(k)} - P_1)(f^{(k)} - P_2)}\right) \\
 &= m\left(r, \frac{f^{(k+1)}(\Delta f - f^{(k)})}{(f^{(k)} - P_1)(f^{(k)} - P_2)}\right) + S(r, \Delta f) \\
 &\leq m\left(r, \frac{f^{(k+1)}}{f^{(k)} - P_2}\right) + m\left(r, \frac{\Delta f - f^{(k)}}{f^{(k)} - P_1}\right) \\
 &\leq m\left(r, \frac{1}{\Delta f - P_1}\right) + S(r, \Delta f) = \overline{N}\left(r, \frac{1}{\Delta f - P_2}\right) + S(r, \Delta f).
 \end{aligned} \tag{26}$$

Hence combining (25) and (26) we obtain

$$T(r, \Delta f) \leq 2\overline{N}\left(r, \frac{1}{\Delta f - P_2}\right) + S(r, \Delta f). \tag{27}$$

Next, case 2 is divided into two subcases.

Subcase 2.1.  $P_1 = 0$ . Then by (7) and Lemma 1 we get

$$m(r, e^Q) = m\left(r, \frac{f^{(k)}}{\Delta f}\right) = S(r, \Delta f). \tag{28}$$

Then by (16), (27) and (28) we can have  $T(r, \Delta f) = S(r, \Delta f)$  a contradiction.

Subcase 2.2.  $P_2 = 0$ . Then by (16), (27) and (28) and Lemma 1 we get

$$\begin{aligned}
 T(r, \Delta f) &\leq m\left(r, \frac{1}{\Delta f - P_1}\right) + \overline{N}\left(r, \frac{1}{f^{(k)}}\right) + S(r, \Delta f) \\
 &\leq m\left(r, \frac{1}{f^{(k)}}\right) + \overline{N}\left(r, \frac{1}{f^{(k)}}\right) + S(r, \Delta f) \leq \left(T(r, f^{(k)})\right) + S(r, \Delta f).
 \end{aligned} \tag{29}$$

From the fact that

$$T(r, f^{(k)}) \leq T(r, \Delta f) + S(r, \Delta f), \tag{30}$$

which follows from (29) that

$$T(r, \Delta f) = T(r, f^{(k)}) + S(r, \Delta f), \tag{31}$$

By second Nevanlinna Fundamental theorem, Lemma 1, (8) and (31) we have

$$\begin{aligned}
 2T(r, \Delta f) &\leq 2T(r, f^{(k)}) + S(r, \Delta f) \\
 &\leq \overline{N}\left(r, \frac{1}{f^{(k)} - P_1}\right) + \overline{N}\left(r, \frac{1}{f^{(k)}}\right) + \overline{N}\left(r, \frac{1}{f^{(k)} - d}\right) + S(r, \Delta f) \\
 &\leq \overline{N}\left(r, \frac{1}{\Delta f - P_1}\right) + \overline{N}\left(r, \frac{1}{\Delta f}\right) + T\left(r, \frac{1}{f^{(k)} - d}\right) - m\left(r, \frac{1}{f^{(k)} - d}\right) + S(r, \Delta f) \\
 &\leq 2T(r, \Delta f) - m\left(r, \frac{1}{f^{(k)} - d}\right) + S(r, \Delta f).
 \end{aligned}$$

Thus

$$m\left(r, \frac{1}{f^{(k)} - d}\right) = S(r, \Delta f). \quad (32)$$

From the First Fundamental Theorem, Lemma 1, (20) to (21), (31), (32) and  $\Delta f$  is a transcendental entire function of finite order, we obtain

$$\begin{aligned} m\left(r, \frac{\Delta f - d}{f^{(k)} - d}\right) &\leq m\left(r, \frac{\Delta f}{f^{(k)} - d}\right) + m\left(r, \frac{d}{f^{(k)} - d}\right) + S(r, \Delta f) \\ &\leq T\left(r, \frac{\Delta f - d}{f^{(k)} - d}\right) - N\left(r, \frac{\Delta f - d}{f^{(k)} - d}\right) + S(r, \Delta f) \\ &= m\left(r, \frac{f^{(k)} - d}{f}\right) + N\left(r, \frac{f^{(k)} - d}{f}\right) - N\left(r, \frac{\Delta f}{f^{(k)} - d}\right) + S(r, \Delta f) \\ &\leq N\left(r, \frac{1}{\Delta f}\right) - N\left(r, \frac{1}{f^{(k)} - 1}\right) + S(r, \Delta f) = T\left(r, \frac{1}{\Delta f}\right) - T\left(r, \frac{1}{f^{(k)} - d}\right) + S(r, \Delta f) \\ &= T(r, \Delta) - T(r, f^{(k)}) + S(r, \Delta f) = S(r, \Delta f). \end{aligned}$$

Thus, we get

$$m\left(r, \frac{\Delta f - d}{f^{(k)} - d}\right) = S(r, \Delta f). \quad (33)$$

It's easy to see that  $N(r, \psi) = S(r, \Delta f)$  and ((12)) can be rewritten as

$$\psi = \left[ \frac{P_1 - d}{P_1} \frac{(f)^{(k+1)}}{f^{(k)} - P_1} + \frac{df^{(k+1)}}{f^{(k)}} \right] \left[ \frac{f - d}{f^{(k)} - d} - 1 \right]. \quad (34)$$

Then by (33) and (34) we can get

$$T(r, \psi) = m(r, \psi) + N(r, \psi) = S(r, \Delta f). \quad (35)$$

By (7), (25) and (35) we get

$$\overline{N}\left(r, \frac{1}{\Delta f - P_1}\right) = S(r, \Delta f). \quad (36)$$

Moreover, by (7), (31) and (36), we have

$$m\left(r, \frac{1}{f^{(k)}}\right) = S(r, \Delta f), \quad (37)$$

which implies

$$\overline{N}\left(r, \frac{1}{\Delta f}\right) = m\left(r, \frac{1}{\Delta f - P_1}\right) \leq m\left(r, \frac{1}{f^{(k)}}\right) = S(r, \Delta f). \quad (38)$$

Then by (7) we obtain  $T(r, \Delta f) = S(r, \Delta f)$ , a contradiction. So, by (12), (16), (27) and the Second Fundamental Theorem of Nevanlinna, we can get

$$\begin{aligned} T(r, \Delta f) &\leq 2m\left(r, \frac{1}{\Delta f - P_1}\right) + S(r, \Delta f) \leq 2m\left(r, \frac{1}{f^{(k)}}\right) + S(r, \Delta f) \\ &\leq 2T(r, f^{(k)}) - 2N\left(r, \frac{1}{f^{(k)}}\right) + S(r, \Delta f) \leq \overline{N}\left(r, \frac{1}{\Delta f - P_1}\right) + \overline{N}\left(r, \frac{1}{\Delta f - P_1}\right) \\ &\quad + \overline{N}\left(r, \frac{1}{f^{(k)}}\right) - 2N\left(r, \frac{1}{f^{(k)}}\right) + S(r, \Delta f)T(r, \Delta f) - N\left(r, \frac{1}{f^{(k)}}\right) + S(r, f), \end{aligned}$$

which deduces that

$$N\left(r, \frac{1}{f^{(k)}}\right) = S(r, \Delta f). \tag{39}$$

It follows from the second theorem of Nevanlinna that

$$\begin{aligned} T(r, f^{(k)}) &\leq \overline{N}\left(r, \frac{1}{f^{(k)}}\right) + \overline{N}\left(r, \frac{1}{f^{(k)} - P_1}\right) + S(r, \Delta f) \\ &\leq \overline{N}\left(r, \frac{1}{f^{(k)} - P_1}\right) + S(r, \Delta f) \leq T(r, f^{(k)}) + S(r, \Delta f), \end{aligned}$$

which implies that

$$T(r, f^{(k)}) = \overline{N}\left(r, \frac{1}{f^{(k)} - P_1}\right) + S(r, \Delta f). \tag{40}$$

Similarly

$$T(r, f^{(k)}) = \overline{N}\left(r, \frac{1}{f^{(k)} - P_2}\right) + S(r, \Delta f). \tag{41}$$

Then, by (27), we get

$$T(r, \Delta f) = 2T(r, f^{(k)}) + S(r, \Delta f). \tag{42}$$

By (25) and (26) we have

$$T(r, \phi) = T(r, f^{(k)}) + S(r, \Delta f). \tag{43}$$

When case 1 occurs, we apply Lemma 4 and obtain

$$f^{(k)} = He^t. \tag{44}$$

Here  $H \neq 0$  is a small function of  $e^t$ . Rewrite (16) as

$$\phi = \frac{f^{(k+1)}(\Delta f - P_1)(\Delta f - P_2) - \Delta f(f^{(k)} - P_1)(f^{(k)} - P_2)}{(\Delta f - P_1)(\Delta f - P_2)(f^{(k)} - P_1)(f^{(k)} - P_2)}. \tag{45}$$

Combining (27) with (44) we get

$$X = f^{(k+1)}(\Delta f - P_1)(\Delta f - P_2) - \Delta f'(f^{(k)} - P_1)(f^{(k)} - P_2) = \sum_{i=0}^5 \delta_i e^{it}, \tag{46}$$

and

$$Y = (\Delta f - P_1)(\Delta f - P_2) - \Delta f'(f^{(k)} - P_1)(f^{(k)} - P_2) = \sum_{j=0}^5 \gamma_j e^{jt}, \tag{47}$$

where  $\delta_i$  and  $\gamma_j$  are small functions of  $e^t$ ,  $\delta_5 \neq 0$  and  $\gamma_6 \neq 0$ .

If  $X$  and  $Y$  are two mutually prime polynomials in  $e^t$ , then we can get  $T(r, \phi) = 6T(r, e^t) + S(r, \Delta f)$ . It follows from (16), (41)–(43) that  $T(r, \Delta f)$ , a contradiction.

If  $X$  and  $Y$  are not two mutually prime polynomials in  $e^t$ , it's easy to see that the degree of  $Y$  is large than  $X$ . Then submitting (38) into (12) implies

$$H = P_2 \tag{48}$$

and

$$t = P_1 z + P_2, \tag{49}$$

where  $P_1 \neq 0$  and  $P_2$  are polynomials.

According to (45), (48), (49) and by simple calculation, we must have

$$\phi = \frac{C}{f^{(k)} - P_2}, \quad (50)$$

where  $C$  is a non-zero constant. Put (44) into (16) we have

$$\frac{c((f)^{(k)} - f^{(k+1)} - P_1)}{(f^{(k)} - P_1)(f^{(k)} - P_2)} = \frac{-\Delta f'}{(\Delta f - P_1)(\Delta f - P_2)}. \quad (51)$$

We claim that  $f^{(k)} \equiv f^{(k+1)}$ .

Otherwise, combining (22), (44) and (51) we can get  $T(r, e^t) = S(r, \Delta f)$ . It follows from (16) and (27) that  $T(r, \Delta f) = S(r, \Delta f)$ , a contradiction. Hence, it is a easy work to verify that

$$P_1 = 1 \quad (52)$$

and

$$f^{(k)} = P_2 e^{z-P_2} = A e^z, \quad (53)$$

where  $A$  is a nonzero constant and furthermore

$$\Delta f = A e^{2z} - P_1 A e^z + P_1. \quad (54)$$

Then rewrite (27) as

$$\frac{\Delta f - f^{(k)}}{f^{(k)} - P_1} = e^t - 1. \quad (55)$$

Put (49), (52)–(54) into (55) and a direct calculation deduces

$$A = P_2 = e^{P_1} = 1. \quad (56)$$

It follows from (1), (28), (52) and (56) that

$$H = -P_1(e^{\varpi} - 1)^n = 1. \quad (57)$$

Since  $\Delta f$  and  $f^{(k)}$  share  $P_2$  IM and (41), (42) and (56) we get

$$e^{2z} - P_1 e^z + (P_1 - 1) = (e^z - 1)^2, \quad (58)$$

i. e.,

$$P_1 = 2. \quad (59)$$

It follows from (57) that

$$e^{\varpi} = (-2)^{-1/n} + 1. \quad (60)$$

But we cannot get (2) from (60), a contradiction. When case 2 occurs we know that  $m(r, e^t) = m(r, e^Q) + O(1) = S(r, \Delta f)$ . Then by (16) and (27) we deduce  $T(r, \Delta f) = S(r, \Delta f)$  a contradiction.  $\triangleright$

## References

1. Haymann, W. K. *Meromorphic Function*, Oxford, Clarendon Press, 1964.
2. Zhang, J. J. and Liao, L. W. Entire Functions Sharing Some Values with their Difference Operators, *Science China Mathematics*, 2014, vol. 57, pp. 2143–2152. DOI: 10.1007/s11425-014-4848-5.

3. Lahiri, I. Weighted Sharing and Uniqueness of Meromorphic Functions, *Nagoya Mathematical Journal*, 2001, vol. 161, no. 3, pp. 193–206. DOI: 10.1017/S0027763000027215.
4. Rubel, L. A. and Yang, C. C. Values Shared by an Entire Function and its Derivative, *Complex Analysis*, Lecture Notes in Mathematics, vol. 599, Berlin, Springer, 1977, pp. 101–103. DOI: 10.1007/BFb0096830.
5. Li, P. and Yang, C. C. Value Sharing of an Entire Function and its Derivatives, *Journal of the Mathematical Society of Japan*, 1999, vol. 51, no. 4, pp. 781–799. DOI: 10.2969/JMSJ/05140781.
6. Chiang, Y. M. and Feng, S. J. On the Nevanlinna Characteristic of  $f(z + \eta)$  and Difference Equations in the Complex Plane, *The Ramanujan Journal*, 2008, vol. 16, no. 1, pp. 105–129. DOI: 10.1007/s11139-007-9101-1.
7. Chiang, Y. M. and Feng, S. J. On the Growth of Logarithmic Differences, Difference Quotients and Logarithmic Derivatives of Meromorphic Functions, *Transactions of the American Mathematical Society*, 2009, vol. 361, pp. 3767–3791. DOI: 10.1090/S0002-9947-09-04663-7.
8. Chen, Z. X. and Yi, H. X. On Sharing Values of Meromorphic Functions and Their Differences, *Results in Mathematics*, 2013, vol. 63, pp. 557–565. DOI: 10.1007/s00025-011-0217-7.
9. Halburd, R. G. and Korhonen, R. J. Nevanlinna Theory for the Difference Operator, *Annales Academiæ Scientiarum Fennicæ. Mathematica*, 2006, vol. 31, pp. 463–478.
10. Heittokangas, J., Korhonen, R., Laine, I. and Rieppo, J. Uniqueness of Meromorphic Functions Sharing Values with their Shifts, *Complex Variables and Elliptic Equations*, 2011, vol. 56, no. 1–4, pp. 81–92. DOI: 10.1080/17476930903394770.
11. Huang, H. X. and Fang, M. L. Unicity of Entire Functions Concerning their Shifts and Derivatives, *Computational Methods and Function Theory*, 2021, vol. 21, pp. 523–532. DOI: 10.1007/s40315-020-00358-1.
12. Li, S., Duan, M. and Chen, B. Q. Uniqueness of Entire Functions Sharing Two Values with their Difference Operators, *Advances in Difference Equations*, 2017, Article no. 390. DOI: 10.1186/s13662-017-1444-3.
13. Liu, D., Yang, D. G. and Fang, M. L. Unicity of Entire Functions Concerning Shifts and Difference Operators, *Abstract and Applied Analysis*, 2014, Article ID 380910. DOI: 10.1155/2014/380910.
14. Liu, K. and Dong, X. J. Some Results Related to Complex Differential-Difference Equations of Certain Types, *Bulletin of the Korean Mathematical Society*, 2014, vol. 51, pp. 1453–1467. DOI: 10.4134/BKMS.2014.51.5.1453.
15. Qi, X. G. Value Distribution and Uniqueness of Difference Polynomials and Entire Solutions of Difference Equations, *Annales Polonici Mathematici*, 2011, vol. 102, pp. 129–142. DOI: 10.4064/ap102-2-3.
16. Qi, X., Li, N. and Yang, L. Uniqueness of Meromorphic Functions Concerning Their Differences and Solutions of Difference Painlevé Equations, *Computational Methods and Function Theory*, 2018, vol. 18, pp. 567–582. DOI: 10.1007/s40315-018-0241-7.
17. Yi, H. X. Meromorphic Functions that Share Two or Three Values, *Kodai Mathematical Journal*, 1990, vol. 13, pp. 363–372. DOI: 10.2996/kmj/1138039280.
18. Al-Khaladi, A. H. H. Meromorphic Functions that Share one Finite Value CM or IM with their  $k$ -th Derivatives, *Results in Mathematics*, 2013, vol. 63, pp. 95–105. DOI: 10.1007/s00025-011-0163-4.
19. Frank, G. and Weissenborn, G. Meromorphic Funktionen, die Mit Einer Ihrer Ableitungen Werte Teilen, *Complex Variables and Elliptic Equations*, 1986, vol. 67, pp. 33–43. DOI: 10.1080/17476938608814184.
20. Wang, J. Uniqueness of Entire Function Sharing a Small Function with its Derivative, *Journal of Mathematical Analysis and Applications*, 2010, vol. 362, no. 2, pp. 387–392. DOI: 10.1016/j.jmaa.2009.09.052.
21. Chiang, Y. M. and Feng, S. J. On the Nevanlinna Characteristic of  $f(z + \eta)$  and Difference Equations in the Complex Plane, *The Ramanujan Journal*, 2008, vol. 16, pp. 105–129. DOI: 10.1007/s11139-007-9101-1.
22. Halburd, R. G. and Korhonen, R. J. Difference Analogue of the Lemma on the Logarithmic Derivative with Applications to Difference Equations, *Journal of Mathematical Analysis and Applications*, 2006, vol. 314, pp. 477–487. DOI: 10.1016/j.jmaa.2005.04.010.
23. Chen, B., Li, S. and Chen, Z. X. Uniqueness of Difference Operators of Meromorphic Functions. *Journal of Inequalities and Applications*, 2012, Article no. 48. DOI: 10.1186/1029-242X-2012-48.
24. Deng, B. M., Lei, C. L. and Fang, M. L. Meromorphic Function Sharing Sets with its Difference Operator or Shifts, *Chinese Annals of Mathematics, Series B*, 2019, vol. 40, no. 3, pp. 331–338.
25. Heittokangas, J., Korhonen, R., Laine, I., Rieppo, J. and Zhang, J. Value Sharing Results for Shifts of Meromorphic Function and Sufficient Conditions for Periodicity, *Journal of Mathematical Analysis and Applications*, 2009, vol. 355, no. 1, pp. 352–363. DOI: 10.1016/j.jmaa.2009.01.053.

26. Zhang, J. J. and Liao, L. W. Entire Functions Sharing Some Values with their Difference Operators, *Science China Mathematics*, 2014, vol. 57, pp. 2143–2152. DOI: 10.1007/s11425-014-4848-5.
27. Huangm, X. H. Unicity on Entire Function Concerning its Differential-Difference Operators, *Results in Mathematics*, 2021. DOI: 10.1007/s00025-021-01461-y.
28. Yang, C.-C. and Yi, H. X. *Uniqueness Theory of Meromorphic Functions*, Dordrecht, Kluwer Academic Publishers, 2003.

Received November 13, 2021

RAJESHWARI SRINIVAS

Department of Mathematics, Bangalore Institute of Technology,  
Vishweshwarapura, Basavanagudi, Bangalore-560004, India,  
Assistant Professor

E-mail: rajeshwari.s@presidencyuniversity.in, rajeshwaripreetham@gmail.com

<https://orcid.org/0000-0002-7681-0830>

SHEEBAKOUSAR BUZURG

Presidency University, School of Engineering,  
Itagalpura, Rajanakunte, Yelahanka, Bangalore-560 064, India,  
Research Scholar

E-mail: sheeba.buzurg@gmail.com

<https://orcid.org/0000-0001-9210-6707>

Владикавказский математический журнал  
2023, Том 25, Выпуск 1, С. 81–92

## ЕДИНСТВЕННОСТЬ ЦЕЛЫХ ФУНКЦИЙ ОТНОСИТЕЛЬНО ИХ РАЗНОСТНЫХ ОПЕРАТОРОВ И ПРОИЗВОДНЫХ

Раджешвари С.<sup>1</sup>, Шибакусар Б.<sup>2</sup>

<sup>1</sup> Бангалорский технологический институт,

Индия, Бангалор-560004, Вишвешварapura, Басаванагуди;

<sup>2</sup> Президентский университет, Инженерная школа,

Индия, Бангалор-560 064, Итагалпура, Раджанакунте, Елаханка

E-mail: rajeshwari.s@presidencyuniversity.in, rajeshwaripreetham@gmail.com;

sheeba.buzurg@gmail.com

**Аннотация.** В этой статье мы изучаем единственность целых функций относительно их разностного оператора и производных. Представление о целых и мероморфных функциях сильно зависит от этого направления. Рубель и Янг рассмотрели единственность целой функции и ее производных; они доказали, что если  $f(z)$  и  $f'(z)$  разделяют два значения  $a, b$  с учетом кратностей, то  $f(z) \equiv f'(z)$ . Позже Ли Пинг и Янг улучшили результат Рубеля и Янга: если  $f(z)$  — непостоянная целая функция, а  $a$  и  $b$  — два конечных различных комплексных значения, и если  $f(z)$  и  $f^{(k)}(z)$  разделяют  $a$  с учетом кратностей и  $b$  — без учета кратностей, то  $f(z) \equiv f^{(k)}(z)$ . В последние годы проявляется значительный интерес к распределению значений мероморфных функций конечного порядка относительно разностного аналога. Заменяя различные конечные комплексные значения многочленами, устанавливается следующий результат: пусть  $\Delta f(z)$  — трансцендентная целая функция конечного порядка,  $k \geq 0$  — целое число, а  $P_1$  и  $P_2$  — два многочлена; если  $\Delta f(z)$  и  $f^{(k)}$  разделяют  $P_1$  с учетом кратностей и  $P_2$  игнорируя кратности, то  $\Delta f \equiv f^{(k)}$ . Нетривиальное доказательство этого результата использует теорию распределения значений Неванлинны.

**Ключевые слова:** разностный оператор, разделяемые значения, конечный порядок, единственность, целая функция, многочлены.

**AMS Subject Classification:** 30D35, 39A32.

**Образец цитирования:** Rajeshwari S. and Sheebakousar B. Unicity on Entire Functions Concerning Their Difference Operators and Derivatives // Владикавк. мат. журн.—2023.—Т. 25, № 1.—С. 81–92 (in English). DOI: 10.46698/p5608-0614-8805-b.