



## PROPER AND ADMISSIBLE TOPOLOGIES IN THE SETTING OF CLOSURE SPACES

MILA MRŠEVIĆ

ABSTRACT. A Čech closure space  $(X, u)$  is a set  $X$  with a (Čech) closure operator  $u$  which need not be idempotent. Many properties which hold in topological spaces hold in Čech closure spaces as well.

The notions of proper (splitting) and admissible (jointly continuous) topologies are introduced on the sets of continuous functions between Čech closure spaces. It is shown that some well-known results of Arens and Dugundji [1] and Iliadis and Papadopoulos [7] are true in this setting.

We emphasize that Theorems 1–10 encompass the results of A. di Concilio [3] and Georgiou and Papadopoulos [5, 6] for the spaces of continuous-like functions as  $\theta$ -continuous, strongly and weakly  $\theta$ -continuous, weakly and super-continuous.

### 1. ČECH CLOSURE SPACES

An operator  $u : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$  defined on the power set  $\mathcal{P}(X)$  of a set  $X$  satisfying the axioms:

- (C1)  $u(\emptyset) = \emptyset$ ,
- (C2)  $A \subset u(A)$  for every  $A \subset X$ ,
- (C3)  $u(A \cup B) = u(A) \cup u(B)$  for all  $A, B \subset X$ ,

is called a *Čech closure operator* and the pair  $(X, u)$  is a *Čech closure space*. For short, the space will be noted by  $X$  as well, and called a *closure space*.

A subset  $A$  is *closed* in the closure space  $(X, u)$  if  $u(A) = A$  holds. It is *open* if its complement is closed. The empty set and the whole space are both open and closed.

The *interior operator*  $\text{int}_u : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$  is defined by means of the closure operator in the usual way:  $\text{int}_u = c \circ u \circ c$ , where  $c : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$  is the complement operator. A subset  $U$  is a *neighbourhood* of a point  $x$  (subset  $A$ ) in  $X$  if  $x \in \text{int}_u U$  ( $A \subset \text{int}_u U$ ) holds. We denote by  $\mathcal{N}(x)$  the collection of all neighbourhoods (the *neighbourhood system*) at the point  $x$ .

---

2000 *Mathematics Subject Classification.* 54A05, 54A10, 54C05, 54C10.

*Key words and phrases.* Čech closure space, function space, proper (splitting) topology, admissible (jointly continuous) topology,  $\theta$ -closure,  $\theta$ -continuous function, strongly  $\theta$ -continuous, weakly  $\theta$ -continuous, weakly continuous, super-continuous function.

By (C3), the intersection of two (and thus finitely many) neighbourhoods at  $x$  is a neighbourhood at  $x$  again. The condition (C1) is equivalent to  $\text{int}_u X = X$ , that is to  $X \in \mathcal{N}(x)$  for every  $x \in X$ , and  $\text{int}_u A \subset A$  for every  $A \subset X$  is equivalent to (C2).

In a closure space  $(X, u)$  a family  $\mathcal{U}(x) \subset \mathcal{N}(x)$  is a neighbourhood (local) base at a point  $x$  if the following axioms are satisfied:

- (Nb1)  $\mathcal{U}(x) \neq \emptyset$  for every  $x \in X$ ,
- (Nb2)  $x \in U$  for every  $U \in \mathcal{U}(x)$ ,
- (Nb3)  $U_1, U_2 \in \mathcal{U}(x) \Rightarrow (\exists U \in \mathcal{U}(x)) U \subset U_1 \cap U_2$ .

A family  $\mathcal{U}(x) \subset \mathcal{N}(x)$  is a neighbourhood (local) subbase at a point  $x$  if the conditions (Nb1) and (Nb2) are fulfilled.

If a collection  $\{\mathcal{U}(x) \mid x \in X\}$  of filters on  $X$  satisfies the conditions (Nb1)–(Nb3), then there is exactly one closure operator  $u$  for  $X$  such that  $\mathcal{U}(x)$  is a neighbourhood base at  $x$  for each  $x \in X$ . The operator  $u$  is defined by:

$$u(A) = \{x \in X \mid U \in \mathcal{U}(x) \Rightarrow U \cap A \neq \emptyset\}.$$

Let  $(X, u_1)$  and  $(X, u_2)$  be closure spaces. The closure  $u_1$  is *coarser* than the closure  $u_2$ , or  $u_2$  is *finer* than  $u_1$ , denoted by  $u_1 \leq u_2$ , if  $u_1(A) \supset u_2(A)$  for every  $A \subset X$ . So defined relation  $\leq$  is a partial order on the set of all closure spaces.

Let  $\{u_\alpha\}$  be a collection of closure operators on a set  $X$ . The infimum (meet) and supremum (join) operators for  $\{u_\alpha\}$  are the operators  $u_0 = \wedge u_\alpha$  and  $u = \vee u_\alpha$  respectively, defined by:  $\mathcal{U}_0(x) = \bigcap_\alpha \mathcal{N}_\alpha(x)$  is a neighbourhood base (system) and  $\mathcal{U}(x) = \bigcup_\alpha \mathcal{N}_\alpha(x)$  is a neighbourhood subbase at  $x \in X$ , for  $u_0$  and  $u$  respectively.

Many topological notions can be defined in the class of closure spaces by means of neighbourhoods.

Let  $M$  be a directed set and  $(x_\mu)_{\mu \in M}$  a net in  $(X, u)$ . The net  $(x_\mu)$  converges to a point  $x \in X$  if for every neighbourhood  $U$  of  $x$  there is a  $\mu \in M$  such that for every  $\mu' \in M$ ,  $\mu' \geq \mu \Rightarrow x_{\mu'} \in U$ . Similarly,  $x$  is an *accumulation point* of the net  $(x_\mu)$  if for every neighbourhood  $U$  of  $x$  and every  $\mu \in M$  there is a  $\mu' \in M$  such that  $\mu' \geq \mu$  and  $x_{\mu'} \in U$ . For every point  $x$  the neighbourhood system  $\mathcal{N}(x)$  is a filter on  $X$  such that  $x \in \bigcap \mathcal{N}(x)$ . Moreover it is a set directed by the inverse inclusion  $\supset$  and every net  $(x_U)_{U \in \mathcal{N}(x)}$  with  $x_U \in U$ , converges to  $x$ .

Let  $(X, u)$  and  $(Y, v)$  be two closure spaces. A function  $f : (X, u) \rightarrow (Y, v)$  is *continuous at*  $x \in X$  if “close points are mapped into close ones”, that is if the following holds

$$A \subset X \wedge x \in u(A) \Rightarrow f(x) \in v(f(A)).$$

This condition is equivalent to:

- (i) the inverse image of every neighbourhood of  $f(x)$  is a neighbourhood of  $x$ ;

- (ii) for every net  $(x_\mu)$  that converges to  $x$ , the net  $(f(x_\mu))$  converges to  $f(x)$ ;
- (iii) if  $x$  is an accumulation point of a net  $(x_\mu)$ ,  $f(x)$  is an accumulation point of the net  $(f(x_\mu))$ .

A function  $f : (X, u) \rightarrow (Y, v)$  is *continuous* if it is continuous at every point of  $X$ . This condition is equivalent to:

- (i)  $f(u(A)) \subset v(f(A))$  for every  $A \subset X$ ;
- (ii)  $u(f^{-1}(B)) \subset f^{-1}(v(B))$  for every  $B \subset Y$ .

The *product of a family*  $\{(X_\alpha, u_\alpha) \mid \alpha \in A\}$  of closure spaces, denoted by  $\Pi(X_\alpha, u_\alpha)$ , is the set  $X = \Pi_{\alpha \in A} X_\alpha$  endowed with the closure operator  $u$  defined by means of neighbourhoods: for every  $x \in X$  the family

$$\mathcal{U}(x) = \{\pi_\alpha^{-1}(V) \mid \alpha \in A, V \in \mathcal{N}_\alpha(x_\alpha)\}$$

is a neighbourhood subbase at  $x$  in  $(X, u)$ . Here  $\pi_\alpha$  are the projections, while  $\mathcal{N}_\alpha(x_\alpha)$  is the neighbourhood system at  $x_\alpha = \pi_\alpha(x)$  in  $X_\alpha$ . There exists exactly one closure operator  $u$  such that  $\mathcal{U}(x)$  is a local subbase at  $x$  in  $(X, u)$  for every  $x \in X$ . Canonical neighbourhoods of  $x$  are of the form  $\bigcap_{i=1}^k \pi_{\alpha_i}^{-1}(V_i)$ .

The projections are continuous as well as the restrictions of continuous functions. The composition of two continuous mappings is continuous and a mapping  $f : (X, u) \rightarrow \Pi(Y_\alpha, v_\alpha)$  is continuous at  $x \in X$  if and only if each composition  $\pi_\alpha \circ f$  is continuous at  $x$ . Also the product  $f : \Pi(X_\alpha, u_\alpha) \rightarrow \Pi(Y_\alpha, v_\alpha)$  of a family  $\{f_\alpha\}$  of continuous mappings is continuous.

A well-known example of a Čech closure operator which is not a Kuratowski closure operator in general, is the so called  $\theta$ -closure. It was defined by Veličko [10] in the following way: Let  $(X, \mathcal{T})$  be a topological space and let  $A \subset X$ . A point  $x \in X$  is in the  $\theta$ -closure of  $A$ , denoted by  $\text{cl}_\theta A$  (or  $\mathcal{T}\text{cl}_\theta A$ ), if each closed neighbourhood of  $x$  intersects  $A$ . Neighbourhood bases in  $(X, \text{cl}_\theta)$  consist of closed neighbourhoods (or closures of open neighbourhoods) in  $(X, \mathcal{T})$  at every point  $x$ .

Let  $(X, \mathcal{T})$  be the product space of a family  $\{(X_\alpha, \mathcal{T}_\alpha)\}$  of topological spaces. The  $\theta$ -closure space of  $(X, \mathcal{T})$  is the product of the  $\theta$ -closure spaces of  $(X_\alpha, \mathcal{T}_\alpha)$ , i.e.  $(X, \mathcal{T}\text{cl}_\theta) = \Pi(X_\alpha, \mathcal{T}_\alpha\text{cl}_\theta)$ .

A function  $f : (X, \mathcal{T}) \rightarrow (Y, \mathcal{V})$  is  $\theta$ -continuous at  $x \in X$  if for every neighbourhood  $V$  of  $f(x)$  there is a neighbourhood  $U$  of  $x$  such that  $f(\overline{U}) \subset \overline{V}$ . A function  $f : (X, \mathcal{T}) \rightarrow (Y, \mathcal{V})$  is  $\theta$ -continuous if it is  $\theta$ -continuous at each of its points. Every continuous function is  $\theta$ -continuous, but the converse does not hold in general.

$\theta$ -continuity is not a continuity concept in the class of topological spaces, but it is in the class of Čech closure spaces. Namely,

**Proposition.** *A function  $f : (X, \mathcal{T}) \rightarrow (Y, \mathcal{V})$  is  $\theta$ -continuous if and only if the function  $f : (X, \mathcal{T}\text{cl}_\theta) \rightarrow (Y, \mathcal{V}\text{cl}_\theta)$  is a continuous mapping of (Čech) closure spaces.*

Hence the following characterizations of  $\theta$ -continuity:

**Proposition.** A function  $f : (X, \mathcal{T}) \rightarrow (Y, \mathcal{V})$  is  $\theta$ -continuous if and only if:

- (1)  $f(\mathcal{T}\text{cl}_\theta A) \subset \mathcal{V}\text{cl}_\theta f(A)$  for every  $A \subset X$ ;
- (2)  $\mathcal{T}\text{cl}_\theta f^{-1}(B) \subset f^{-1}(\mathcal{V}\text{cl}_\theta B)$  for every  $B \subset Y$ .

The next statement follows from the definitions and the properties of  $\theta$ -closure.

**Proposition.** Let  $X, Y, Z$  be topological spaces. A function  $g : Z \times X \rightarrow Y$  is  $\theta$ -continuous if and only if the function  $g : (Z, \text{cl}_\theta) \times (X, \text{cl}_\theta) \rightarrow (Y, \text{cl}_\theta)$  is continuous.

A closure space  $(X, u)$  is:

- (i) *regular* if for each point  $x$  and each subset  $A$  such that  $x \notin u(A)$ , there exist neighbourhoods  $U$  of  $x$  and  $V$  of  $A$  such that  $U \cap V = \emptyset$ ;
- (ii) *compact* if each net in  $(X, u)$  has an accumulation point.

A closure space  $(X, u)$  is regular if and only if for each point  $x$  and each neighbourhood  $U$  of  $x$ , there is a neighbourhood  $U_1$  of  $x$  such that  $u(U_1) \subset U$ .

Compactness can be characterized by means of covers. [2, 41 A.9. Theorem] An *interior cover* of  $(X, u)$  is a cover  $\{G_\alpha\}$  such that the collection  $\{\text{int}_u G_\alpha\}$  covers  $X$ . The space is compact if and only if every interior cover has a finite subcover.

We give the following

**Definition.** A collection  $\{G_\alpha\}$  is an *interior cover* of a set  $A$  in  $(X, u)$  if the collection  $\{\text{int}_u G_\alpha\}$  covers  $A$ . A subset  $A$  is *compact* if every interior cover of  $A$  has a finite subcover.

All notions not explained here can be found in [2].

## 2. PROPER AND ADMISSIBLE TOPOLOGIES IN THE SETTING OF CLOSURE SPACES

Let  $X, Y$  and  $Z$  be three nonempty sets. For every function  $g : Z \times X \rightarrow Y$  there is a function  $E(g)$  or  $g^*$  from  $Z$  to  $Y^X$ , the set of all functions from  $X$  to  $Y$ , defined by  $(g^*(z))(x) = g(z, x)$ . The mapping  $E : Y^{Z \times X} \rightarrow (Y^X)^Z$  is called the *exponential function*. By  $\varepsilon$  we denote the *evaluation mapping* from  $Y^X \times X$  to  $Y$  defined by  $\varepsilon(f, x) = f(x)$ .

If  $X, Y$  and  $Z$  are topological or closure spaces, in particular sets of continuous functions can be considered. Now on  $Y^X$  will mean the set of *all continuous functions* from  $X$  to  $Y$ . The set  $Y^X$  can be endowed with different topologies. The question is: Find the topologies on the set of functions such that

- (1)  $E(g) = g^* \in (Y^X)^Z$  for every  $g \in Y^{Z \times X}$ , that is, for every continuous  $g : Z \times X \rightarrow Y$  the function  $g^*$  is continuous; and conversely,
- (2)  $g \in Y^{Z \times X}$  for every  $g^* \in (Y^X)^Z$ , that is, for every continuous  $g^* : Z \rightarrow Y^X$  the function  $g$  is continuous.

Following the definitions and notations used by Arens and Dugundji [1], Kuratowski [9] and Iliadis and Papadopoulos [7] for the sets of continuous functions defined in the setting of topological spaces, we give the following definitions.

**Definition 1.** Let  $(X, u)$  be a closure space and  $(A_\lambda)_{\lambda \in \Lambda}$  be a net in  $\mathcal{P}(X)$ . The *upper limit* of the net  $(A_\lambda)$ , denoted by  $\overline{\lim}_\Lambda A_\lambda$ , is the set of all points  $x \in X$  such that for every  $\lambda_0 \in \Lambda$  and every neighbourhood  $U$  of  $x$  in  $X$ , there is a  $\lambda \in \Lambda$  such that  $\lambda \geq \lambda_0$  and  $A_\lambda \cap U \neq \emptyset$ . (See, for example, [1] and [7].)

**Definition 2.** Let  $(X, u)$  and  $(Y, v)$  be closure spaces and  $Y^X$  be the collection of all continuous functions  $f : (X, u) \rightarrow (Y, v)$ . A closure operator  $\sigma$  on  $Y^X$  is called *proper (splitting)* if for any closure space  $(Z, w)$

- (1)  $g : (Z, w) \times (X, u) \rightarrow (Y, v)$  is continuous  $\Rightarrow E(g) = g^* : (Z, w) \rightarrow (Y^X, \sigma)$  is continuous;

$\sigma$  is called *admissible (jointly continuous)* if for every space  $(Z, w)$

- (2)  $g^* : (Z, w) \rightarrow (Y^X, \sigma)$  is continuous  $\Rightarrow g : (Z, w) \times (X, u) \rightarrow (Y, v)$  is continuous.

Let  $f, f_\lambda \in Y^X$ ,  $\lambda \in \Lambda$ , where  $\Lambda$  is a directed set. The net  $(f_\lambda)$  *converges continuously* to  $f$ , denoted by  $f_\lambda \xrightarrow{cc} f$ , if

- (3) the net  $f_\lambda(x_\mu)$ ,  $(\lambda, \mu) \in \Lambda \times M$ , converges to  $f(x)$  in  $(Y, v)$  whenever the net  $(x_\mu)$  converges to  $x$  in  $(X, u)$ .

The convergence of a net  $(f_\lambda)$  to  $f$  in the space  $(Y^X, \sigma)$  will be denoted by  $f_\lambda \xrightarrow{\sigma} f$ .

The next results follow from definitions and the proofs are analogous to the corresponding for the topological case. (See [1] and [7]).

**Theorem 1.** A closure operator  $\sigma$  on  $Y^X$  is admissible if and only if the evaluation mapping  $\varepsilon : (Y^X, \sigma) \times (X, u) \rightarrow (Y, v)$  is continuous.

**Theorem 2.** Let  $(f_\lambda)_{\lambda \in \Lambda}$  be a net in  $Y^X$ . The net  $(f_\lambda)_{\lambda \in \Lambda}$  converges continuously to  $f \in Y^X$  if and only if for every  $x \in X$  and every neighbourhood  $V$  of  $f(x)$  there is a neighbourhood  $U$  of  $x$  and a  $\lambda_0 \in \Lambda$  such that  $f_\lambda(U) \subset V$  for all  $\lambda \geq \lambda_0$ .

**Theorem 3.** Let  $(f_\lambda)_{\lambda \in \Lambda}$  be a net in  $Y^X$ . The net  $(f_\lambda)_{\lambda \in \Lambda}$  converges continuously to  $f \in Y^X$  if and only if the following holds:

- (4)  $\overline{\lim}_\Lambda f_\lambda^{-1}(B) \subset f^{-1}(v(B))$  for every subset  $B$  in  $Y$ .

*Proof.* Let  $f_\lambda \xrightarrow{cc} f$ ,  $B \subset Y$  and  $x \in \overline{\lim}_\Lambda f_\lambda^{-1}(B)$ . For every neighbourhood  $V$  of  $f(x)$ , by Theorem 2, there is a neighbourhood  $U$  of  $x$  and a  $\lambda_0 \in \Lambda$

such that  $f_\lambda(U) \subset V$  for all  $\lambda \geq \lambda_0$ . By the assumption, there is a  $\lambda' \geq \lambda_0$  such that  $f_{\lambda'}^{-1}(B) \cap U \neq \emptyset$ . It follows that  $f_{\lambda'}(U) \cap B \neq \emptyset$ . Thus

$$(\forall V \in \mathcal{N}(f(x))) V \cap B \neq \emptyset$$

implies  $f(x) \in v(B)$ , hence  $x \in f^{-1}(v(B))$ .

For the converse, let  $(f_\lambda)_{\lambda \in \Lambda}$  be a net in  $Y^X$  such that the condition (4) holds. For every  $x \in X$  and every neighbourhood  $V$  of  $f(x)$ ,

$$f(x) \in \text{int}_v V = c(v(c(V))) \text{ implies } f(x) \notin v(V^c).$$

Thus  $x \notin f^{-1}(v(V^c))$  implies  $x \notin \varlimsup_{\Lambda} f_\lambda^{-1}(V^c)$ . Hence

$$\begin{aligned} (\exists U \in \mathcal{N}(x))(\exists \lambda_0 \in \Lambda)(\forall \lambda \in \Lambda) \lambda \geq \lambda_0 &\Rightarrow f_\lambda^{-1}(V^c) \cap U = \emptyset \\ &\Rightarrow V^c \cap f_\lambda(U) = \emptyset. \end{aligned}$$

Thus

$$(\exists U \in \mathcal{N}(x))(\exists \lambda_0 \in \Lambda)(\forall \lambda \in \Lambda) \lambda \geq \lambda_0 \Rightarrow f_\lambda(U) \subset V.$$

By Theorem 2,  $f_\lambda \xrightarrow{cc} f$ . □

**Remark.** In Theorem 3 the condition “for every  $B \subset Y$ ” can be replaced by: for every  $B = V^c$ , where  $V$  is a neighbourhood basic element.

**Corollary 1.** *Let  $(f_\lambda)_{\lambda \in \Lambda}$  be a net in  $Y^X$  and  $Y$  be a topological space. The net  $(f_\lambda)_{\lambda \in \Lambda}$  converges continuously to  $f \in Y^X$  if and only if the following holds:*

$$(4^*) \quad \varlimsup_{\Lambda} f_\lambda^{-1}(B) \subset f^{-1}(B) \text{ for every closed subset } B \text{ in } Y.$$

*Proof.* Let a net  $(f_\lambda)$  converges continuously to  $f \in Y^X$  and  $B$  be a closed set in  $Y$ . By (4),

$$\varlimsup_{\Lambda} f_\lambda^{-1}(B) \subset f^{-1}(\overline{B}) = f^{-1}(B).$$

Conversely, suppose that  $(4^*)$  holds and let  $B$  be a subset in  $Y$ . By  $(4^*)$ , for the closed subset  $\overline{B}$ ,  $\varlimsup_{\Lambda} f_\lambda^{-1}(\overline{B}) \subset f^{-1}(\overline{B})$ , and by isotony of the upper limit,  $\varlimsup_{\Lambda} f_\lambda^{-1}(B) \subset \varlimsup_{\Lambda} f_\lambda^{-1}(\overline{B})$ . Hence the statement. □

**Theorem 4.** *Let  $\sigma$  and  $\sigma'$  be two closure operators on  $Y^X$ .*

- (1) *If  $\sigma'$  is finer than  $\sigma$  and  $\sigma'$  is proper, then  $\sigma$  is proper.*
- (2) *If  $\sigma \leq \sigma'$  and  $\sigma$  is admissible, then  $\sigma'$  is admissible.*
- (3) *If  $\sigma$  is proper and  $\sigma'$  is admissible, then  $\sigma \leq \sigma'$ .*
- (4) *If there is a closure operator on  $Y^X$  which is both proper and admissible, it is unique.*

*Proof.* (3). If  $\sigma'$  is admissible, the evaluation mapping

$$\varepsilon : (Y^X, \sigma') \times (X, u) \rightarrow (Y, v)$$

is continuous by Theorem 1. Since  $\sigma$  is proper, for  $Z = (Y^X, \sigma')$ , the identity

$$\varepsilon^* = 1_{Y^X} : (Y^X, \sigma') \rightarrow (Y^X, \sigma)$$

is continuous, hence  $\sigma'$  is finer than  $\sigma$ . □

**Corollary 2.** *Let  $\{\sigma_\alpha\}$  be a collection of closure operators on  $Y^X$ .*

- (1) *If  $\sigma_\alpha$  is proper for every  $\alpha$ , then the infimum and supremum,  $\bigwedge \sigma_\alpha$  and  $\bigvee \sigma_\alpha$ , are proper.*
- (2) *If  $\sigma_\alpha$  is admissible for every  $\alpha$ , then the supremum,  $\bigvee \sigma_\alpha$  is admissible as well.*

*Proof.* For the nontrivial part of (1), let  $\sigma_\alpha$  be proper for every  $\alpha$ . That is, for any space  $(Z, w)$ , continuity of  $g : (Z, w) \times (X, u) \rightarrow (Y, v)$  implies  $g^* : (Z, w) \rightarrow (Y^X, \sigma_\alpha)$  is continuous. In order to prove continuity of  $g^* : (Z, w) \rightarrow (Y^X, \bigvee \sigma_\alpha)$ , for any  $z \in Z$  and every neighbourhood  $G$  of  $g^*(z)$ , there are finitely many  $G_i \in \mathcal{N}_{\sigma_{\alpha_i}}(g^*(z))$  such that  $\bigcap G_i \subset G$ . By continuity of  $g^* : (Z, w) \rightarrow (Y^X, \sigma_{\alpha_i})$ , for every  $i$  there is a  $W_i \in \mathcal{N}(z)$  such that  $g^*(W_i) \subset G_i$ . Then

$$W = \bigcap W_i \in \mathcal{N}(z)$$

and

$$g^*(W) \subset \bigcap g^*(W_i) \subset \bigcap G_i \subset G$$

holds. Thus  $g^* : (Z, w) \rightarrow (Y^X, \bigvee \sigma_\alpha)$  is continuous at  $z$ .  $\square$

**Theorem 5.** *A closure operator  $\sigma$  on  $Y^X$  is proper (splitting) if and only if continuous convergence of a net implies its convergence in  $(Y^X, \sigma)$ , and it is admissible (jointly continuous) if and only if the reverse holds, that is, convergence of a net in  $(Y^X, \sigma)$  implies its continuous convergence.*

**Theorem 6.** *A closure operator  $\sigma$  on  $Y^X$  is:*

- (1) *proper if and only if continuity of a mapping  $g : Z \times (X, u) \rightarrow (Y, v)$  implies continuity of the mapping  $g^* : Z \rightarrow (Y^X, \sigma)$  for every topological space  $Z$  being either a  $T_1$ -space having at most one non-isolated point or the Sierpinski space;*
- (2) *admissible if and only if the reverse holds, that is, continuity of a mapping  $g^* : Z \rightarrow (Y^X, \sigma)$  implies continuity of the mapping  $g : Z \times (X, u) \rightarrow (Y, v)$  for every topological space  $Z$  being either a  $T_1$ -space having at most one non-isolated point or the Sierpinski space.*

**Corollary 3.** *In Theorem 6 the conditions on the space  $Z$  can be replaced by:  $Z$  is a topological space having at most one non-isolated point.*

In the sequel the finest proper topology on  $Y^X$ , which exists by Corollary 2, is characterized by means of convergence classes and upper limits, analogously to the topological situation. (Cf. [7].)

Let  $\mathcal{C}(\sigma)$  be the *convergence class* of the closure space  $(Y^X, \sigma)$ , that is

$$\mathcal{C}(\sigma) = \{((f_\lambda)_{\lambda \in \Lambda}, f) \mid f_\lambda, f \in Y^X \text{ and } f_\lambda \xrightarrow{\sigma} f\}.$$

It can be easily seen that  $\mathcal{C}(\sigma)$  satisfies the following axioms: (cf. [2, 35 A.2. Theorem] and [8, 2.9 Theorem])

- (i) **(CONSTANTS):** If  $(f_\lambda)_{\lambda \in \Lambda}$  is a net such that  $f_\lambda = f$  for every  $\lambda \in \Lambda$ , then  $(f_\lambda)$  converges to  $f$ , that is,  $((f_\lambda)_{\lambda \in \Lambda}, f) \in \mathcal{C}(\sigma)$ ;

- (ii) **(SUBNETS)**: If a net  $(f_\lambda)$  converges to  $f$ , so does each subnet of  $(f_\lambda)$ , i.e. if  $((f_\lambda)_{\lambda \in \Lambda}, f) \in \mathcal{C}(\sigma)$ , then  $((g_\mu)_{\mu \in M}, f) \in \mathcal{C}(\sigma)$  for every subnet  $(g_\mu)$  of  $(f_\lambda)$ ;
- (iii) **(DIVERGENCE)**: If a net  $(f_\lambda)$  does not converge to  $f$ , then there is a subnet  $(g_\mu)$  of  $(f_\lambda)$  no subnet of which converges to  $f$ , i.e.  $((f_\lambda)_{\lambda \in \Lambda}, f) \notin \mathcal{C}(\sigma)$ , then there is a subnet  $(g_\mu)$  of  $(f_\lambda)$  such that  $((h_\nu)_{\nu \in N}, f) \notin \mathcal{C}(\sigma)$  for every  $(h_\nu)$  subnet of  $(g_\mu)$ .

*Proof.* (iii). Let  $(f_\lambda)$  be a net in  $Y^X$ ,  $f \in Y^X$  such that  $((f_\lambda)_{\lambda \in \Lambda}, f) \notin \mathcal{C}(\sigma)$ . It means that

$$(\exists G_0 \in \mathcal{N}(f))(\forall \lambda \in \Lambda)(\exists \lambda' \in \Lambda)\lambda' \geq \lambda \wedge f_{\lambda'} \notin G_0.$$

Thus there is a cofinal subset  $M \subset \Lambda$  such that  $f_\mu \notin G_0$  for every  $\mu \in M$ .  $(f_\mu)_{\mu \in M}$  is a subnet of  $(f_\lambda)$ , no subnet of which converges to  $f$ .  $\square$

The space  $(Y^X, \sigma)$  is topological if and only if its convergence class satisfies the axiom of (ITERATED LIMITS). (Cf. [8, 2.9 Theorem] and [2, 15 B.13. and 35 A.3. Theorems].)

Denote by  $\mathcal{C}^*$  the class of all pairs  $((f_\lambda)_{\lambda \in \Lambda}, f)$  such that  $(f_\lambda)$  is a net in  $Y^X$  which converges continuously to  $f \in Y^X$ , i.e.

$$\mathcal{C}^* = \{((f_\lambda)_{\lambda \in \Lambda}, f) \mid f_\lambda, f \in Y^X \text{ and } f_\lambda \xrightarrow{cc} f\}.$$

By Theorem 5,  $\sigma$  is proper if and only if  $\mathcal{C}^* \subset \mathcal{C}(\sigma)$  and it is admissible if and only if the reverse inclusion holds:  $\mathcal{C}(\sigma) \subset \mathcal{C}^*$ .

**Theorem 7.** *The class  $\mathcal{C}^*$  satisfies the axioms (CONSTANTS), (SUBNETS) and (DIVERGENCE).*

*Proof.* For (CONSTANTS) and (SUBNETS) is clear.

(DIVERGENCE). Let  $(f_\lambda)$  be a net in  $Y^X$ ,  $f \in Y^X$  and let

$$((f_\lambda)_{\lambda \in \Lambda}, f) \notin \mathcal{C}^*.$$

By Theorem 3 there is a subset  $B \subset Y$  such that  $\overline{\lim}_\Lambda f_\lambda^{-1}(B)$  is not contained in  $f^{-1}(v(B))$ . Let

$$x \in \overline{\lim}_\Lambda f_\lambda^{-1}(B) \setminus f^{-1}(v(B)).$$

Let  $\mathcal{N}(x)$  be the set of all neighbourhoods of  $x$  directed by inverse inclusion and let  $M = \Lambda \times \mathcal{N}(x)$ . If  $\mu = (\lambda, U) \in \Lambda \times \mathcal{N}(x)$ , let  $\varphi : M \rightarrow \Lambda$  be defined by  $\varphi(\mu) \in \Lambda$  such that  $\varphi(\mu) = \varphi(\lambda, U) \geq \lambda$  and  $f_{\varphi(\mu)}^{-1}(B) \cap U \neq \emptyset$ . The net  $(g_\mu)_{\mu \in M}$ , where  $g_\mu = f_{\varphi(\mu)}$ , is a subnet of  $(f_\lambda)$ .

Let  $(h_\nu)$  be a subnet of  $(g_\mu)$  and  $\psi : N \rightarrow M$  be the corresponding map. In order to prove that  $((h_\nu)_{\nu \in N}, f) \notin \mathcal{C}^*$ , let  $\nu_0 \in N$  and  $U \in \mathcal{N}(x)$ . If  $\psi(\nu_0) = (\lambda_0, U_0) \in M$ , set  $\hat{U} = U_0 \cap U \in \mathcal{N}(x)$  and  $\mu_0 = (\lambda_0, \hat{U})$ . There is  $\nu_1 \in N$  such that  $\nu_1 \geq \nu_0$  and  $\nu \geq \nu_1 \Rightarrow \psi(\nu) \geq \mu_0$ . For any  $\nu \geq \nu_1$  and  $\psi(\nu) = (\lambda, \tilde{U})$  we have

$$h_\nu^{-1}(B) \cap U = f_{\varphi(\psi(\nu))}^{-1}(B) \cap U \supset f_{\varphi(\psi(\nu))}^{-1}(B) \cap \hat{U} \supset f_{\varphi(\psi(\nu))}^{-1}(B) \cap \tilde{U} \neq \emptyset.$$



It means that  $x \in \overline{\lim}_N h_\nu^{-1}(B)$  and hence  $\overline{\lim}_N h_\nu^{-1}(B)$  is not contained in  $f^{-1}(v(B))$ . Thus the axiom (DIVERGENCE) is satisfied.  $\square$

**Corollary 4.**  $\mathcal{C}^*$  is the convergence class of the finest proper topology on  $Y^X$  if and only if  $\mathcal{C}^*$  satisfies the axiom (ITERATED LIMITS).

**Theorem 8.** A subset  $G$  in  $Y^X$  is open in the finest proper topology if and only if for every  $f \in G$  and for every net  $(f_\lambda)_{\lambda \in \Lambda}$  in  $Y^X$  such that (4) holds, there exists a  $\lambda_0 \in \Lambda$  such that  $f_\lambda \in G$  for every  $\lambda \geq \lambda_0$ .

*Proof.* ( $\Leftarrow$ .) Let  $\tau$  be the collection of subsets in  $Y^X$  with the given property.

$\tau$  is a topology on  $Y^X$ : for  $G_1, G_2 \in \tau$ ,  $f \in G_1 \cap G_2$  and a net  $(f_\lambda)_{\lambda \in \Lambda}$  satisfying (4), there are  $\lambda_1, \lambda_2 \in \Lambda$  such that  $f_\lambda \in G_i$  for all  $\lambda \geq \lambda_i$ ,  $i = 1, 2$ . Then  $f_\lambda \in G_1 \cap G_2$  for all  $\lambda \geq \lambda_0 = \max\{\lambda_1, \lambda_2\}$ . It follows that  $G_1 \cap G_2 \in \tau$ . Similarly, if  $\{G_\alpha\} \subset \tau$  and  $G = \bigcup_\alpha \{G_\alpha\}$ , let  $f \in G$  and  $(f_\lambda)_{\lambda \in \Lambda}$  be a net which satisfies (4). There is an  $\alpha_0$  such that  $f \in G_{\alpha_0}$ , and since (4) holds, there exists a  $\lambda_0 \in \Lambda$  such that  $f_\lambda \in G_{\alpha_0} \subset G$  for every  $\lambda \geq \lambda_0$ . Thus  $G \in \tau$ .

$\tau$  is proper: let  $(f_\lambda)_{\lambda \in \Lambda}$  be a net such that  $f_\lambda \xrightarrow{cc} f$ . By Theorem 3,  $\overline{\lim}_\Lambda f_\lambda^{-1}(B) \subset f^{-1}(v(B))$  for every subset  $B$  in  $Y$ . Let  $f \in G \in \tau$ . By the assumption, there exists a  $\lambda_0 \in \Lambda$  such that  $f_\lambda \in G$  for every  $\lambda \geq \lambda_0$ , that is  $f_\lambda \xrightarrow{\tau} f$ . By Theorem 5,  $\tau$  is proper.

$\tau$  is the finest proper topology: let  $\sigma$  be a proper topology on  $Y^X$  and  $H \in \sigma$ . Let  $f \in H$  and a net  $(f_\lambda)_{\lambda \in \Lambda}$  satisfy (4). By Theorem 3,  $f_\lambda \xrightarrow{cc} f$ . Since  $\sigma$  is proper,  $f_\lambda \xrightarrow{\sigma} f$ . By definition of convergence, there exists a  $\lambda_0 \in \Lambda$  such that  $f_\lambda \in H$  for every  $\lambda \geq \lambda_0$ . By definition of  $\tau$ ,  $H \in \tau$ . Thus  $\sigma \subset \tau$ .

( $\Rightarrow$ .) Let a subset  $G$  in  $Y^X$  be open in the finest proper topology  $\tau$ , let  $f \in G$  and  $(f_\lambda)$  be a net satisfying (4). By Theorem 3,  $f_\lambda \xrightarrow{cc} f$ . Since  $\tau$  is proper,  $f_\lambda \xrightarrow{\tau} f$ . By definition of convergence, there exists a  $\lambda_0 \in \Lambda$  such that  $f_\lambda \in G$  for every  $\lambda \geq \lambda_0$ .  $\square$

In order to give nontrivial examples of admissible and proper topologies and to get results analogous to Theorems 4.1 and 4.21 in [1], we consider the following sets. Let

$$\mathcal{V} = \{V \subset Y \mid \text{int}_v V \neq \emptyset\}.$$

For  $A \subset X$  and  $V \in \mathcal{V}$ , set

$$(A, V) = \{f \in Y^X \mid f(A) \subset V\}.$$

Let  $\mathcal{A}$  be a family of subsets of  $X$ . The collection

$$\{(A, V) \mid A \in \mathcal{A}, V \in \mathcal{V}, V = \text{int}_v V\},$$

is a subbase for a topology on  $Y^X$ , which will be called the  $\mathcal{A}$ -topology.

Let  $\mathcal{C}$  be an interior cover of  $X$ . The collection

$$\{(u(K), V) \mid V \in \mathcal{V} \text{ and } K \subset X \text{ is such that } u(K) \subset C \text{ for some } C \in \mathcal{C}\},$$

is a subbase for a topology on  $Y^X$ , which will be called the  $\mathcal{C}$ -topology.

**Theorem 9.** *Let  $(X, u)$  be a regular closure space and  $(Y, v)$  be arbitrary. For every interior cover  $\mathcal{C}$  of  $X$ , the  $\mathcal{C}$ -topology is admissible.*

*Proof.* By Theorem 1, it is enough to prove that the evaluation mapping is continuous. Let  $f \in Y^X$ ,  $x \in X$  and  $V \in \mathcal{N}(f(x))$ ,  $f(x) = \varepsilon(f, x)$ . By continuity of  $f$ , the set  $U = f^{-1}(V) \in \mathcal{N}(x)$ . Choose a  $C \in \mathcal{C}$  so that  $x \in \text{int}_u C$ . Then  $U \cap C \in \mathcal{N}(x)$  and by regularity of  $X$ , there is a  $U_1 \in \mathcal{N}(x)$  such that

$$x \in \text{int}_u U_1 \subset U_1 \subset u(U_1) \subset U \cap C.$$

For the subbasic element  $(u(U_1), V)$  in the  $\mathcal{C}$ -topology,  $\varepsilon((u(U_1), V), U_1) \subset V$  since for every  $f_1 \in (u(U_1), V)$  and each  $x \in U_1$ ,  $\varepsilon(f_1, x_1) = f_1(x_1) \in V$ .  $\square$

**Theorem 10.** *Let  $(X, u)$  and  $(Y, v)$  be closure spaces and  $\mathcal{A}$  be a collection of compact subsets in  $(X, u)$ . The  $\mathcal{A}$ -topology is always proper.*

*Proof.* Let  $g : (Z, w) \times (X, u) \rightarrow (Y, v)$  be a continuous function. In order to prove continuity of the mapping  $g^* : (Z, w) \rightarrow (Y^X, \sigma)$ , where  $Y^X$  is endowed with the  $\mathcal{A}$ -topology, let  $z \in Z$  and  $f = g^*(z)$ . For a subbasic element  $(K, V)$  containing  $f$ , where  $K$  is a compact set in  $X$ ,

$$f(K) = g(\{z\} \times K) \subset \text{int}_v V = V.$$

By continuity of  $g$ ,

$$(\forall x \in K) g(z, x) \in \text{int}_v V \Leftrightarrow V \in \mathcal{N}(g(z, x))$$

implies

$$(\forall x \in K)(\exists W_x \in \mathcal{N}(z))(\exists U_x \in \mathcal{N}(x)) g(W_x \times U_x) \subset V.$$

$(\forall x \in K) U_x \in \mathcal{N}_x$  implies  $\{U_x \mid x \in K\}$  is an interior cover of the compact set  $K$ , so there is a finite subcover  $\{U_{x_i} \mid i = 1, \dots, k\}$ . Set  $W = \bigcap_{i=1}^k W_{x_i}$ . Then  $W \in \mathcal{N}(z)$ . It follows that

$$g(W \times K) \subset g\left(\bigcup_{i=1}^k (W_{x_i} \times U_{x_i})\right) \subset V.$$

Thus

$$(\forall z' \in W) g(z', K) \subset V \Rightarrow g^*(W) \subset (K, V).$$

$\square$

### 3. SOME SPECIAL CASES INCLUDING $\theta$ -CLOSURE

3.1. Let  $(X, u)$  be a closure space and  $(Y, \mathcal{V})$  be a topological space. A function  $f : (X, u) \rightarrow (Y, \mathcal{V})$  is continuous if and only if the function  $f : (X, \hat{u}) \rightarrow (Y, \mathcal{V})$  between topological spaces is continuous, where  $\hat{u}$  is the topological modification of the closure operator  $u$ . In that case the problem is reduced to the topological case since  $Y^X = C((X, \hat{u}), Y)$  and the topological modification of the product of closure spaces is the product of topological modifications.

3.2. It was already remarked that  $\theta$ -continuous functions are continuous functions of the corresponding Čech closure spaces. Compact sets in  $(X, \text{cl}_\theta)$  are (quasi-)H-closed (g-H-closed) in  $(X, \mathcal{U})$ . Thus Theorems 3.1–3.6 and 4.2 in [3] are special cases of Theorems 2, 1, 4, 5 and 10 respectively.

If  $(Y, \mathcal{V})$  is regular, the  $\theta$ -topology  $\mathcal{V}_\theta = \mathcal{V}$ . Then for a topological space  $(X, \mathcal{U})$ , a function  $f : (X, \mathcal{U}) \rightarrow (Y, \mathcal{V})$  is  $\theta$ -continuous if and only if  $f : (X, \text{cl}_\theta) \rightarrow (Y, \mathcal{V})$  is continuous, which is equivalent to  $f : (X, \mathcal{U}_\theta) \rightarrow (Y, \mathcal{V})$  be continuous [2, Thm 16.B.4]. Note that the topological modification of  $\mathcal{U}\text{cl}_\theta$  is  $\mathcal{U}_\theta$ , the topology of  $\theta$ -open sets in  $(X, \mathcal{U})$ .

3.3. A large number of continuous-like mappings between topological spaces is known in the literature. Recently, Georgiou and Papadopoulos [5, 6] considered some of them and investigated splitting and jointly continuous topologies on the sets of these functions. Let us remark that all these examples and the main results are special cases of our subjects of investigations. For, let  $(X, \mathcal{U})$  and  $(Y, \mathcal{V})$  be topological spaces. It follows from the definitions that a function  $f : (X, \mathcal{U}) \rightarrow (Y, \mathcal{V})$  is:

- (1) *strongly  $\theta$ -continuous* (cf. [5]) at a point  $x$  (on the set  $X$ ) if and only if  $f : (X, \mathcal{U}\text{cl}_\theta) \rightarrow (Y, \mathcal{V})$  is continuous at  $x$  (on the set  $X$ );
- (2) *super-continuous* (cf. [6]) at  $x$  (on the set  $X$ ) if and only if  $f : (X, \mathcal{U}_s) \rightarrow (Y, \mathcal{V})$  is continuous at  $x$  (on the set  $X$ ), where  $\mathcal{U}_s$  is the semi-regularization topology of  $\mathcal{U}$  (see [4] for example);
- (3) *weakly continuous* (cf. [6]) at  $x$  (on the set  $X$ ) if and only if  $f : (X, \mathcal{U}) \rightarrow (Y, \mathcal{V}\text{cl}_\theta)$  is continuous at  $x$  (on the set  $X$ );
- (4) *weakly  $\theta$ -continuous* (cf. [6]) at a point  $x$  (on  $X$ ) if and only if  $f : (X, \mathcal{U}_s) \rightarrow (Y, \mathcal{V}\text{cl}_\theta)$  is continuous at  $x$  (on  $X$ ).

Also  $\theta$ -convergence of a net  $(x_\mu)$  in  $(X, \mathcal{U})$  means convergence of  $(x_\mu)$  in the corresponding closure space  $(X, \mathcal{U}\text{cl}_\theta)$ , while *weak  $\theta$ -convergence* of a net  $(x_\mu)$  in  $(X, \mathcal{U})$  is convergence of  $(x_\mu)$  in  $(X, \mathcal{U}_s)$ .

Similarly,  $\theta$ -continuous convergence (respectively: *strongly  $\theta$ -continuous convergence*, *weakly  $\theta$ -continuous convergence*, *weakly continuous convergence* and *super continuous convergence*) of a net  $(f_\lambda)$  in  $Y^X$  is continuous convergence of  $(f_\lambda)$  for the corresponding closure spaces. Thus we are concerned with a change of topology, better to say: change of closure operator, technique. So the main results in [5] and [6] are special cases of the above Theorems 1–10.

## REFERENCES

1. Richard Arens and James Dugundji, *Topologies for function spaces*, Pacific J. Math. **1** (1951), 5–31. MR 13,264d
2. Eduard Čech, *Topological spaces*, Publishing House of the Czechoslovak Academy of Sciences, Prague, 1966. MR 35 #2254
3. Anna Di Concilio, *On  $\theta$ -continuous convergence in function spaces*, Rend. Mat. (7) **4** (1984), no. 1, 85–94 (1985). MR 87d:54024

4. D. B. Gauld, M. Mršević, I. L. Reilly, and M. K. Vamanamurthy, *Continuity properties of functions*, Topology, theory and applications (Eger, 1983), North-Holland, Amsterdam, 1985, pp. 311–322. MR 863 913
5. D. N. Georgiou and B. K. Papadopoulos, *Strongly  $\vartheta$ -continuous functions and topologies on function spaces*, Papers in honour of Bernhard Banaschewski (Cape Town, 1996), Kluwer Acad. Publ., Dordrecht, 2000, pp. 433–444. MR 2002b:54012
6. ———, *Weakly continuous, weakly  $\vartheta$ -continuous, super-continuous and topologies on function spaces*, Sci. Math. Jpn. **53** (2001), no. 2, 233–246. MR 1 828 262
7. S. D. Iliadis and Basil K. Papadopoulos, *The continuous convergence on function spaces*, Panamer. Math. J. **4** (1994), no. 3, 33–42. MR 95k:54025
8. John L. Kelley, *General topology*, D. Van Nostrand Company, Inc., Toronto-New York-London, 1955. MR 16,1136c
9. Casimir Kuratowski, *Sur la notion de limite topologique d'ensembles*, Ann. Soc. Polon. Math. **21** (1948), 219–225 (1949). MR 11,120d
10. N. V. Veličko, *H-closed topological spaces*, Mat. Sb. (N.S.) **70 (112)** (1966), 98–112, English translation in *American Mathematical Society Translations. Series 2, Vol. 78*, Amer. Math. Soc., Providence, R.I., 1968; MR 39 #1253; ISBN 0-8218-1778-7. MR 33 #6576

FACULTY OF MATHEMATICS, UNIVERSITY OF BELGRADE, BEOGRAD, YUGOSLAVIA  
E-mail address: mrsevic@matf.bg.ac.yu