

ON THE BIANCHI IDENTITIES IN A GENERALIZED WEYL SPACE*

GÜLÇİN ÇIVI

*Faculty of Science and Letters, Istanbul Technical University
 80626 Maslak-Istanbul, Turkey*

Abstract. In this paper, we show that the first Bianchi identity is valid for a generalized Weyl space having a semi-symmetric E -connection and that the second Bianchi identity is satisfied for a recurrent generalized Weyl space provided that the recurrence vector ψ_l and the Vranceanu vector Ω_l are related by $\psi_l = \frac{2}{n-1} \Omega_l$.

1. Introduction

An n -dimensional differentiable manifold W_n^* having an asymmetric connection ∇^* and asymmetric conformal metric tensor g^* preserved by ∇^* is called a **generalized Weyl space** [1]. For a such a space, in local coordinates, we have the compatibility condition

$$\nabla_k^* g_{ij}^* - 2T_k^* g_{ij}^* = 0, \quad (1.1)$$

where T_k^* are the components of a covariant vector field called the complementary vector field of the generalized Weyl space.

The coefficients L_{jk}^i of the connection ∇^* are obtained from the compatibility condition as [2]

$$L_{jk}^i = \Gamma_{jk}^i + \frac{1}{2} \left[\Omega_{kl}^h g_{(jh)}^* + \Omega_{jl}^h g_{(hk)}^* + \Omega_{jk}^h g_{(hl)}^* \right] g^{*(li)} \quad (1.2)$$

or, putting

$$Q_{jk}^i = \frac{1}{2} \left[\Omega_{kl}^h g_{(jh)}^* + \Omega_{jl}^h g_{(hk)}^* + \Omega_{jk}^h g_{(hl)}^* \right] g^{*(li)} \quad (1.3)$$

* This work is supported by TUBITAK, Research Center of Turkey.